



ISSN NO. 2320-5407

Journal homepage: <http://www.journalijar.com>

INTERNATIONAL JOURNAL
OF ADVANCED RESEARCH

RESEARCH ARTICLE

Effects of licorice extract, probiotic, potassium chloride and sodium bicarbonate on productive performance of broiler chickens exposed to heat stress

Hazim J. Al-Daraji

Department of Animal Production, College of Agriculture, University of Baghdad, Baghdad, Iraq

Manuscript Info

Manuscript History:

Received: 17 May 2013
Final Accepted: 27 May 2013
Published Online: June 2013

Key words:

Liquorice, probiotic, potassium chloride, sodium bicarbonate, broiler, productive performance, heat stress

Abstract

This study was undertaken to compare the effect of licorice extract, probiotic, potassium chloride and sodium bicarbonate on productive performance of broilers exposed to heat stress. A total of 900 Fawbro broiler chicks were allocated at random into 6 treatments groups of 3 replicates each. Experimental treatments were as follows: Treatment 1: control group was subjected to heat stress; Treatment 2: control group was not subjected to heat stress; Treatment 3: Probiotic group were early exposed to 10^6 Colony Forming Unit (CFU) of *Lactobacilli* at their first day of age, and then subjected to heat stress; Treatment 4: heat distressed birds and supplemented with 0.5 % potassium chloride in their drinking water; Treatment 5: heat distressed birds and supplemented with 0.5% sodium bicarbonate in their drinking water; and Treatment 6: heat distressed birds and supplemented with 450 mg / liter licorice extract in their drinking water. At the third week of age, birds in treatments 2, 3, 4, 5 and 6 were exposed to heat stress (38 – 43 °C) for 6 hours(1200 – 1800 h) each day until the end of experiment (8th week of age).

Results revealed that treated the birds with probiotic or licorice extract resulted in a significant improvement in mean body weight, feed conversion ratio, water consumption, cumulative body weight, cumulative weight gain, cumulative feed conversion ratio, livability, productive index, economic figure and dressing percentage with or without viscera as compared with control group which was subjected to heat stress, potassium chloride treatment and sodium bicarbonate treatment. Furthermore, licorice extract treatment surpasses probiotic treatment in respect to mean body weight, water consumption, final body weight, cumulative feed consumption, livability, and dressing percentage with or without gible.

In conclusion, licorice extract and probiotic can be used as an efficient tools in alleviate the deleterious effects of heat stress on productive performance of broiler chickens.

Copy Right, IJAR, 2013., All rights reserved.

Introduction

Like all animals, poultry are most comfortable in a specific temperature range called the thermoneutral zone. The ideal environment for high growth rate after three weeks of age is 15 to 20°C with a relative humidity of 50 % or less; however, their actual thermoneutral zone may extend up 27°C, depending upon the relative humidity and the age of the bird. Birds kept in ambient temperatures outside their

thermoneutral zone have poorer productive efficiency (11).

Heat stress causes a decrease in plasma sodium and increase in plasma chloride. An alteration in sodium: chloride ratio resulted in blood alkalosis which also has a detrimental effect on productive performance (29). Al-Daraji et al. (1) concluded that supplementation of sodium bicarbonate and potassium chloride to drinking water had a positive role of relieve the effect of heat stress on broilers as

indicated by positive changes in physiological traits. In another study (30), the effect of heat stress on growth and feed efficiency was partially alleviated by potassium chloride supplementation to drinking water. Naji et al. (20) showed that early microbial exposure of broilers by *Lactobacilli* resulted in a significant improvement in body weight and feed conversion and significant decrease in mortality rate compared with the addition of salts or vinegar to drinking water. Al-Daraji et al. (2) reported that early microbial exposure of heat stressed broiler to *Lactobacilli* or adding sodium bicarbonate and potassium chloride salts into the drinking water resulted in a significant improvement in all blood and plasma parameters compared with control group. However, *Lactobacilli* treatment surpasses the treatments of salts with regard to all hematological traits included in their study.

Licorice is a group of plant that has a marked effect upon the endocrine system. The glucosides present have a structure that is similar to the natural steroids of the body. They explain the beneficial action that liquorice has in the treatment of adrenal gland problems such as Addison's disease (18). Bernardi et al. (4) stated that licorice root contains glycyrrhizin, 50 times sweeter than sucrose, which encourages the production of hormones such as hydrocortisone. This help to explain its anti – stress and anti – inflammatory action and also its role in stimulating the adrenal cortex after steroid therapy. Kageyama et al. (15) found that glycyrrhizic acid caused a similar mineralcorticoid effect on reversing stress induced hypotension. It stimulates the adrenal gland and helps the body cope with stress (17). Other studies have shown that glycyrrhizin has a similar chemical structure to corticosteroids released by the adrenals and stimulates the excretion of hormones by the adrenal cortex; therefore it is a possible drug to prolong the action of cortisone (27). Baker (3) reported that glycyrrhetic acid, the steroid like constituent of glycyrrhizic acid, inhibits an enzyme responsible for inactivating cortisol in the kidney. The treatment with licorice essentially extends the lifetime of cortisol and hence reinforces the resistance of body to stress. Both liquoriceextract and glycyrrhetic acid are shown to have desoxycorticosterone and ACTH – like effects, though with less toxicity than cortisone, encouraging its uses as an anti – stress agent (10).

The aim of this study was to evaluate the probable role of licorice extract in counteracting the detriments of heat stress and make a comparison for this herb with certain materials (probiotic, potassium chloride and sodium bicarbonate) which have previously proved as good tools to alleviate the

harmful effect of heat stress on productive performance of broiler chickens.

Material and Methods

A total of 900 Fawbro broiler chicks, one day old were used in this study. Birds were fed a starter diet (22.7 % crude protein and 3067.4 Kcal ME / kg of diet) during the first 3 weeks of age and finisher diet (20.6 % crude protein and 2922 Kcal ME / kg of diet) until the termination of the experiment (8 weeks of age). Chicks were allocated at random into 6 treatments groups of 3 replicates per group; each replicate constitutes 50 chicks (150 chicks per each treatment). Experimental treatments were as follows: Treatment 1: Control group which were subjected to heat stress (38 -43 °C for 6 hours); Treatment 2: Control group which were not subjected to heat stress; Treatment 3: Probiotic group which were early exposed to 10⁶ Colony Forming Unit (CFU) of *Lactobacilli* at their first day of age, and then subjected to acute heat stress. The isolates of *Lactobacilli* bacteria were taken from crops of broilers and then activated by infusion it in the MRS (De Man Rogosa and Sharp) broth at 37°C. After 24 h these media were mixed with tap water which contains 20 % skim milk. This probiotic product was sprayed on the chicks at the first day of age by allocate 0.2 ml per each chick. This amount of probiotic was determined to contain 10⁶ colony units of bacteria per chick (2); Treatment 4: Heat – distressed birds and supplemented with 0.5 % potassium chloride in their drinking water; Treatment 5: Heat – distressed birds and supplemented with 0.5 % sodium bicarbonate in their drinking water; Treatment 6: Heat – distressed birds and supplemented with 450 mg / liter licorice extract in their drinking water.

At the third week of age, birds in treatments 2, 3, 4, 5 and 6 were exposed to heat stress (38 – 43 °C) for 6 hours daily (1200 h – 1800h) until the end of experiment (8th week of age). Gaseous brooders were used as a source of heat exposure in the experimental house. On the other hand, the supplementation of potassium chloride, sodium bicarbonate and licorice extract to the drinking water of birds were started at the third week of age and continued until 56 d. of age.

Productive characteristics included in this study were: body weight (BW), weight gain (WG), feed consumption (FC), feed conversion ratio (FCR), water consumption (WC) and mortality (M) rate. However, Productive Index (PI) and Economic Figure (EF) were also determined. At the end of experiment, 18 birds from each treatment (6 birds per each replicate) were sacrificed to determine dressing percentage with giblet (DPG) or without giblet

(DPWV), and weights of visceral organs, viz. liver (L), heart (H), gizzard (G), and spleen (S) in addition to abdominal fat (AFT). The data obtained were statistically analyzed used Duncan's multiple range test (26).

Results and Discussion

Results of this study evinced a highly significant ($p < 0.01$) increase in mean BW during the whole experimental period (4 – 8 weeks of age) in birds receiving licorice extract in their drinking water as compared to all other treatments (Table 1). Results also revealed that probiotic treatment surpasses other treatments (control which was subjected to heat stress, potassium chloride treatment and sodium bicarbonate treatment) as concerns this trait, while there were no significant differences between probiotic treatment and control group which was not subjected to heat stress).

Compared with control group, the addition of probiotic, potassium chloride or licorice extract resulted in a significant ($p < 0.05$) improvement in WG during all weeks of the experiment, while sodium bicarbonate treatment was superior to control group which was subjected to heat stress just during

4, 5, and 6 weeks of experiment (Table 2). However, there were no significant differences between licorice extract treatment and control group which was not subjected to heat stress during all weeks of experiment, whereas there were no significant differences between probiotic treatment and control group which was not subjected to heat stress only during the 6th, 7th and 8th weeks of age regarding WG (Table 2). Besides, there were no significant differences between all treatments groups involved in the present study with relation to FC (Table 3).

Results denoted that the addition of licorice extract results in a significant ($p < 0.05$) improvement in FCR during 4th, 5th, 6th and 7th weeks of age in comparison with control which was subjected to heat stress, potassium chloride treatment and sodium bicarbonate treatment (Table 4). Furthermore, there were no significant differences between licorice treatment, probiotic and control group which was not subjected to heat stress during 5th, 6th, 7th and 8th weeks of age as regards this characteristic. Moreover, there were no significant differences between all treatment groups during the 8th week of age in relation to FCR (Table 4).

Table 1. The effect of licorice extract, probiotic, potassium chloride and sodium bicarbonate and licorice extract on mean body weight (g) of broilers exposed to heat stress.

Treatment Age (weeks)	Control	Control + heat stress	Licorice extract	Probiotic	Potassium chloride	Sodium bicarbonate
4	CD 538.42 ± 35.43	B 567.40 ± 34.09	A 581.78 ± 30.82	B 563.65 ± 36.40	C 553.67 ± 32.88	C 546.69 ± 33.16
5	E 732.53 ± 38.39	B 778.47 ± 36.40	A 794.10 ± 38.91	B 771.96 ± 40.11	C 761.58 ± 34.18	D 747.82 ± 37.52
6	E 933.46 ± 51.19	B 996.91 ± 50.12	A 1014.17 ± 47.47	B 993.34 ± 49.42	C 978.82 ± 44.03	D 963.17 ± 45.85
7	E 1154.92 ± 59.87	B 1235.19 ± 55.14	A 1251.26 ± 56.40	B 1230.86 ± 60.32	C 1206.88 ± 54.07	D 1186.26 ± 57.47
8	E 1415.04 ± 72.81	B 1506.81 ± 67.53	A 1522.53 ± 64.33	B 1501.39 ± 70.80	C 1478.30 ± 65.20	D 1454.56 ± 69.64

* Values in a row with different superscripts differ significantly ($P < 0.05$).

Table 2. The effect of licorice extract, probiotic, potassium chloride and sodium bicarbonate on weight gain (g) of broilers exposed to heat stress.

Treatment Age (weeks)	Control	Control + heat stress	Licorice extract	Probiotic	Potassium chloride	Sodium bicarbonate
4	C 170.28 ± 17.75	A 185.69 ± 15.33	A 187.47 ± 16.06	B 180.65 ± 18.04	B 179.13 ± 16.91	B 177.29 ± 17.75
5	C 194.11 ± 18.73	A 211.07 ± 16.68	A 212.32 ± 17.20	B 208.31 ± 19.07	B 207.91 ± 17.0	B 201.13 ± 18.47
6	B 200.93 ± 22.07	A 218.44 ± 19.74	A 220.07 ± 20.25	A 221.38 ± 19.15	A 217.24 ± 19.01	A 216.0 ± 18.44
7	C 221.46 ± 26.09	A 238.28 ± 24.36	A 237.09 ± 23.18	A 237.52 ± 23.25	B 228.06 ± 25.17	BC 223.09 ± 24.0
8	B 260.12 ± 28.85	A 271.62 ± 27.76	A 272.27 ± 27.14	A 270.53 ± 28.03	A 271.42 ± 26.19	B 268.30 ± 26.49

* Values in a raw with different superscripts differ significantly ($P < 0.05$).

Table 3. The effect of licorice extract, probiotic, potassium chloride and sodium bicarbonate on daily feed consumption (g) of broilers exposed to heat stress.

Treatment Age (weeks)	Control	Control + heat stress	Licorice extract	Probiotic	Potassium chloride	Sodium bicarbonate
4	A 72.97 ± 4.23	A 75.27 ± 3.82	A 74.12 ± 4.35	A 74.83 ± 4.38	A 73.81 ± 4.76	A 73.56 ± 4.32
5	A 81.31 ± 5.27	A 82.86 ± 4.93	A 83.08 ± 5.12	A 82.73 ± 5.67	A 82.96 ± 4.92	A 82.34 ± 5.42
6	A 90.34 ± 5.90	A 91.25 ± 5.24	A 92.44 ± 6.33	A 91.16 ± 5.34	A 92.14 ± 5.70	A 92.01 ± 5.26
7	A 102.98 ± 7.45	A 105.21 ± 6.93	A 104.93 ± 7.03	A 104.65 ± 6.71	A 103.87 ± 7.54	A 103.41 ± 7.52
8	A 110.42 ± 8.46	A 112.47 ± 8.29	A 112.22 ± 8.16	A 112.17 ± 6.62	A 112.44 ± 7.82	A 112.31 ± 8.51
Total feed consumption (g)	A 458.02 ± 13.8	A 467.06 ± 9.63	A 465.54 ± 12.0	A 465.22 ± 12.3	A 463.63 ± 7.30	A 466.79 ± 8.80

Table 4. The effect of licorice extract, probiotic, potassium chloride and sodium bicarbonate on feed conversion efficiency (FC / WG) of broilers exposed to heat stress.

Treatment Age (weeks)	Control	Control + heat stress	Licorice extract	Probiotic	Potassium chloride	Sodium bicarbonate
4	A 3.00 ± 0.041	C 2.84 ± 0.044	D 2.76 ± 0.033	B 2.89 ± 0.056	BC 2.88 ± 0.037	B 2.91 ± 0.054
5	A 2.93 ± 0.056	C 2.75 ± 0.052	C 2.74 ± 0.040	C 2.78 ± 0.061	AB 2.88 ± 0.041	B 2.87 ± 0.047
6	A 3.14 ± 0.051	C 2.92 ± 0.055	C 2.94 ± 0.052	C 2.89 ± 0.054	B 2.97 ± 0.047	B 2.98 ± 0.050
7	A 3.26 ± 0.077	C 3.10 ± 0.064	C 3.10 ± 0.060	C 3.08 ± 0.076	B 3.19 ± 0.069	AB 3.24 ± 0.071
8	A 2.97 ± 0.053	B 2.90 ± 0.061	B 2.89 ± 0.053	B 2.90 ± 0.061	B 2.90 ± 0.051	AB 2.93 ± 0.054

* Values in a row with different superscripts differ significantly ($P < 0.05$).

Table 5. The effect of licorice extract, probiotic, potassium chloride and sodium bicarbonate on daily water consumption (ml) of broilers exposed to heat stress.

Treatment Age (weeks)	Control	Control + heat stress	Licorice extract	Probiotic	Potassium chloride	Sodium bicarbonate
4	D 170.55 ± 20.78	E 157.54 ± 16.44	A 192.61 ± 18.52	B 183.42 ± 17.32	C 178.56 ± 19.47	C 174.82 ± 20.18
5	D 195.93 ± 22.18	E 174.72 ± 18.50	A 229.46 ± 20.47	B 213.62 ± 19.22	C 206.47 ± 20.09	C 200.33 ± 21.27
6	E 284.56 ± 26.18	F 225.56 ± 23.18	A 297.76 ± 25.19	B 286.75 ± 24.47	C 272.47 ± 29.02	D 261.11 ± 28.43
7	D 299.52 ± 30.22	E 265.20 ± 27.41	A 329.61 ± 30.36	B 317.33 ± 29.90	BC 314.54 ± 32.25	C 308.65 ± 30.11
8	D 320.94 ± 30.76	E 302.73 ± 33.68	A 351.71 ± 36.12	B 340.39 ± 34.77	C 333.60 ± 32.49	C 329.84 ± 35.62

* Values in a row with different superscripts differ significantly ($P < 0.05$).

Results of this study also manifested that licorice extract treatment recorded the highest values of WC followed by the results of probiotic group, whereas the lowest values were noticed with control group which was not subjected to heat stress (Table 5).

Data in Table 6 revealed that probiotic and licorice treatments transcend other treatments groups (control which was subjected to heat stress, potassium chloride treatment and sodium bicarbonate treatment) concerning cumulative BW, WG, FCR,

livability, PI and EF. In addition, licorice treatment overtops probiotic treatment as regards cumulative BW, FC and livability.

Results of dressing percentage and relative weights of visceral organs are seen in Table (7). It was noticed that licorice treatment excel other treatments (control which was subjected to heat stress, probiotic treatment, potassium chloride treatment and sodium bicarbonate treatment) in relation to DPG and DPWG characters. However, licorice and probiotic treatments recorded the lowest means of AFT, L and S in comparison with control

which was subjected to heat stress, potassium chloride treatment and sodium bicarbonate treatment. While, there were no significant differences among all treatments groups included in this study as concerns H and G traits.

The positive result which achieved by using the probiotic in this study as indicated by counteracting the noxious effect of heat stress on productive performance of broilers is in accordance with the previous study of Al-Daraji et al. (2) who found that early treatment with probiotic significantly contributed in easement the effect of heat stress on broiler chickens, as indicated by the preeminence of this treatment on other salt treatments regarding all physiological parameters included in their study. Furthermore, these authors concluded that these advantageous changes may be in part due to the production of acids by these bacteria during the fermentation of sugars which reflected in upshot by the suppression of the effects of heat stress on birds. The acids produced by these bacteria exert their role through the neutralizing of respiratory alkalosis that occurs when birds are housed at a high temperature. The hyperventilation and respiratory alkalosis are reflected in a loss of carbon dioxide from the blood and associated losses of bicarbonate from the blood and body fluids (29). Deyhim and Teeter (8) pointed out that heat – distressed birds exhibit increased respiration rate and respiratory alkalosis. The alkalosis induced by heat distress has been related to negative mineral balance for K^+ as well as Na^+ and elevated blood corticosterone. Other explanations for positive changes that occurred in productive traits and associated with probiotic treatment may be due to the competitive inhibition of *Lactobacilli* bacteria against several pathological bacteria (32) and increase in the activity of digestive enzymes (34) which led to the amelioration in digestive coefficient and increase in the availability of many nutrients (16). These changes were reflected in an improvement of productive performance and reinforcement in the general health of broiler. Oyetayo et al. (21) stated that *Lactobacilli* are important for the maintenance of the intestinal microbial ecosystem, growth enhancement of farm animals, protection from pathogens including many harmful bacteria, viruses and fungi, alleviation of lactose intolerance, relief of constipation, anti cholesterolaemic effect and immunostimulation. *Lactobacilli* play their protective therapeutic effect through production of antimicrobial compounds, reduction of gut pH by stimulating the lactic acid producing microflora, competition for binding of receptor sites that pathogens occupy, liver function improvement, stimulation of immunomodulatory cells and competition with pathogens for available nutrients (24). However, the use of probiotics in

order to competitively exclude the colonization of intestinal pathogens has been proposed for poultry, especially after the European Commission banned certain antibiotics frequently included in feeding stuffs as growth promoters (22).

The improvement of productive performance of broiler exposed to heat stress and supplemented with licorice extract in their drinking water may be attributed to that licorice has properties similar to corticosterone and ACTH hormones and hence its role in enhancement the resistance of body to stress (28). Edens and Siegel (9) found that corticosterone concentration reached a peak after 70 min of heat exposure of broilers to 45 °C and afterward rapidly reduced. This reduction in corticosterone level was accompanied by the reduction in survivability of birds which indicated by the depression of adrenal efficiency that ultimately reflected in the occurrence of heat prostration which led at last to death of all the birds with the continuance of heat exposure. Freeman (13) reported that the third stage of stress which called exhaustion phase is characterized by the depletion of birds' body stores and insufficiency of adrenal gland to produce glucocorticoid hormones in suitable quantity. Thus the birds were surrender to the stressor factor and the death will happened.

Some compounds found in licorice are thought to help the adrenal gland function more smoothly in conditions of stress and exhaustion. The adrenal glands are responsible for hormones that keep the body systems balanced. Naturopaths have used licorice in treating hypoglycemia, diabetes and Addison's disease, which is a malfunction of the adrenal glands (25). By enhancing cortisol activity, glycyrrhizin helps to increase energy, ease stress, and reduce the symptoms of ailments sensitive to cortisol levels, such as chronic fatigue syndrome and fibromyalgia. In the 1800s, licorice extract was a common remedy for a type of persistent fatigue known as neurasthenia, the condition known as chronic fatigue syndrome (31). Ferrari et al. (12) reported that activity of 11 – β – hydroxysteroid dehydrogenase type 2 (11 β – HSD – 2) potently is blocked *in vivo* and *in vitro* by glycyrrhetic acid (GA), the active compound of licorice, by 2 mechanisms, direct competitive inhibition and pretranslational inhibition. When 11 β – HSD activity is decreased, urinary excretion of active cortisol (tetrahydrocortisol) to inactive cortisone (tetrahydrocortisone) metabolites increased. It was reported in early literature that GA enhances endogenous glucocorticoid action by suppressing the metabolism of glucocorticoids (19), and a recent report showed that GA administration to normal mice inhibits 11 β – HSD activities in immune tissue such as thymus, spleen, and peripheral lymph nodes in a

dose – dependent manner (14). Quaschnig et al. (23) pointed out that in GA – induced hypertension, both aldosterone receptor antagonism and endothelin receptor antagonism normalize blood pressure and improve renovascular function and, thus, may represent a new therapeutic approach in cardiovascular disease associated with impaired 11β – HSD – 2 activity. In addition to its established role as a competitive inhibitor of 11β – HSD, licorice result in pretranslational inhibition of 11β – HSD both *in vitro* and *in vivo*. 11β – HSD is clearly an important mechanism in regulating tissue levels of active glucocorticoid and, hence, ligand supply to the glucocorticoid receptor (33).

On the other hand, the beneficial role of licorice extract that found in the present study in which it reinforced the resistance of birds to heat stress as indicated by the amelioration in the productive performance of broiler chickens exposed to heat stress could be also explained by that licorice treatment resulted in a significant increase in water consumption during the entire period of the

experiment in comparison with other treatments included in this study. Cooney and Fitzsimons (6) found that treatment the rat with licorice extract, glycyrrhizic acid or glycyrrhetic acid resulted in a significant increase in sodium appetite, thirst and water consumption. Simmons et al. (29) found that mortality during exposure to high temperature was reduced when birds consuming more water which was partly attributed to the wetting of the litter from increased urination and the possible cooling of the bird upon burrowing into the litter. Branton et al. (5) concluded that the increase in fluid consumption was determined to contribute more to increase survival during exposure to high temperature than did the reduction in blood pH. Deaton (7) stated that increased fluid consumption decreased mortality from heat prostration. Subsequently, decreased fluid consumption increased mortality. It was noticeable that with increased fluid consumption more water was excreted into the litter making a difference in litter moisture.

Table 6. The effect of licorice extract, probiotic, potassium chloride and sodium bicarbonate on certain productive characteristics of broilers exposed to heat stress.

Treatment Traits	Control	Control + heat stress	Licorice extract	Probiotic	Potassium chloride	Sodium bicarbonate
Final body weight (gm)	E 1415.04 ± 72.81	B 1506.81 ± 67.53	A 1522.53 ± 64.33	B 1501.39 ± 70.80	C 1478.30 ± 65.20	D 1454.56 ± 69.64
Mean weight gain (4 – 8; g)	C 209.38 ± 23.45	A 225.02 ± 23.81	A 225.84 ± 22.34	A 223.68 ± 22.35	B 220.75 ± 22.18	B 217.16 ± 23.04
Mean feed consumption (4 – 8; g)	D 3206.14 ± 235.28	A 3269.42 ± 230.07	A 3267.53 ± 245.60	B 3258.78 ± 257.53	B 3256.54 ± 261.76	C 3245.41 ± 259.79
Mean feed conversion efficiency (4 – 8 weeks)	A 3.06 ± 0.044	C 2.90 ± 0.058	C 2.88 ± 0.049	C 2.91 ± 0.055	B 2.96 ± 0.050	AB 2.99 ± 0.047
Total mortality (%)	A 11.11 ± 0.036	D 1.94 ± 0.019	D 1.97 ± 0.022	C 3.01 ± 0.035	B 5.16 ± 0.028	B 5.21 ± 0.030
Productive Index	C 119.11 ± 14.32	A 145.32 ± 14.73	A 147.56 ± 15.77	A 142.97 ± 15.63	B 125.37 ± 13.47	BC 123.39 ± 14.12
Economic Figure	C 118.93 ± 13.49	A 145.11 ± 15.18	A 147.26 ± 16.34	A 142.85 ± 14.76	B 125.19 ± 14.56	BC 123.07 ± 14.17

* Values in a raw with different superscripts differ significantly ($P < 0.05$).

Table 7. The effect of licorice extract, probiotic, potassium chloride and sodium bicarbonate on dressing percentage and organ weights of broilers exposed to heat stress.

Treatment Traits	Control	Control + heat stress	Licorice extract	Probiotic	Potassium chloride	Sodium bicarbonate
Dressing percentage (with giblet)	C 72.06 ± 7.83	A 76.76 ± 6.93	A 76.70 ± 7.63	B 74.63 ± 7.14	C 73.24 ± 6.90	C 73.20 ± 7.11
Dressing percentage (without giblet)	D 62.49 ± 6.67	A 67.95 ± 5.28	A 67.91 ± 6.46	B 65.82 ± 6.17	C 63.97 ± 5.19	C 63.89 ± 5.42
Abdominal fat (%)	A 1.83 ± 0.037	C 1.49 ± 0.029	C 1.51 ± 0.033	C 1.50 ± 0.036	B 1.67 ± 0.027	B 1.68 ± 0.031
Liver weight (%)	A 3.95 ± 0.077	C 3.62 ± 0.066	C 3.61 ± 0.071	C 3.61 ± 0.067	B 3.84 ± 0.073	AB 3.89 ± 0.070
Gizzard weight (%)	A 2.70 ± 0.066	A 2.69 ± 0.069	A 2.68 ± 0.057	A 2.69 ± 0.054	A 2.70 ± 0.055	A 2.69 ± 0.062
Spleen weight (%)	A 0.39 ± 0.001	A 0.33 ± 0.005	B 0.31 ± 0.006	B 0.32 ± 0.005	A 0.37 ± 0.006	A 0.37 ± 0.004
Heart weight (%)	A 0.70 ± 0.015	A 0.68 ± 0.011	A 0.68 ± 0.009	A 0.69 ± 0.010	A 0.69 ± 0.008	A 0.68 ± 0.009

* Values in a row with different superscripts differ significantly ($P < 0.05$).

References

- Al-Daraji, H. J., I. A. Al-ANI, J. K. Minati and H. E. Al-Heeti. 2003. Sodium bicarbonate, potassium chloride and vinegar drinking water supplementation effects on physiological traits of broilers reared in heat – distressed environment. *Iraqi J. Agric. Sci.* 34 (10): 151 – 160.
- Al-Daraji, H. J., I. A. Al-ANI, J. K. Minati and H. E. Al-Heeti. 2005. The use of *Lactobacilli* and certain salts in alleviation the effect of heat stress in broiler. Accepted for publication in *Iraqi J. Agric. Sci.* 36 (1).
- Baker, M. E. 1994. Licorice and enzymes other than 11 – beta – hydroxysteroid dehydrogenase: an evolutionary perspective. *Steroids* 59 (2): 136 – 141.
- Bernardi, M., P. E. D'Intino and F. Trevisani. 1994. Effects of prolonged ingestion of graded doses of licorice by healthy volunteers. *Life Sci.* 55 (11): 863 – 872.
- Branton, S. L., F.N. Reece and J. W. Deaton. 1986. The use of ammonium chloride and sodium bicarbonate in acute heat stress of broilers. *Poultry Sci.* 65: 1659 – 1663.
- Cooney, A. S. and J. T. Fitzsimons. 1996. Increased sodium appetite and thirst in rat induced by the ingredients of licorice, glycyrrhizic acid and glycyrrhetic acid. *Regul. Pept.* 66 (1 – 2): 127 – 133.
- Deaton, J. W. 1994. Acclimation to heat stress in birds. U. S. D. A., Agric. Res. Service, South Central Poultry Research Laboratory, Mississippi State, MS 39762. (Personal Communication).
- Deyhim, F. and R. G. Teeter. 1991. Sodium and potassium chloride drinking water supplementation effects on acid – base balance and plasma corticosterone in broilers reared in thermoneutral and heat – distressed environments. *Poultry Sci.* 70: 2551 – 2553.
- Edens, F. W. and H. S. Siegel. 1976. Modification of corticosterone and glucose response by sympatholytic agents in young chickens during acute heat exposure. *Poultry Sci.* 55: 1704 – 1712.
- Farese, R. V., E. G. Biglieri and C. H. L. Shakelton. 1990. Licorice – induced hypermineralocorticoidism. *N. Engl. J. Med.* 325 (17): 1223 – 1227.
- Ferket, P. R. 1992. Hot weather management and feeding of poultry. Takeda, U. S. A. INC., Corporate Drive, Orangeburg, New York.

12. Ferrari, P., A. Sansonnens, B. Dick and F. J. Frey. 2001. *In vivo* 11 β – HSD – 2 activity, variability, salt – sensitivity, and effect of licorice. *Hypertension* 38: 1330 – 1336.
13. Freeman, B. M. 1988. The domestic fowl in biochemical research: Physiological effects of the environment. *Wld's Poultry Sci. J.* 44: 41 – 60.
14. Hennebold, J. D., S. Y. Ryu, H. H. Mu, A. Galbraith and R. A. Daynes. 1996. 11 β – hydroxysteroid dehydrogenase modulation of glucocorticoid activities in lymphoid organs. *Am. J. Physiol.* 270: R1296 – R1306.
15. Kageyama, Y., H. Suzuki and T. Saruta. 1994. Role of glucocorticoid in the development of glycyrrhizin – induced hypertension. *Clin. Exp. Hypertens.* 16: 761 – 778.
16. Kalenhammer, T. R. and M. J. Kullen. 1999. Selection and design of probiotic: A review. *J. Am. Diet. Assoc.* 101(2): 229 – 241.
17. Kato, H., M. Kaneka and S. Yano. 1995. 3 – Monoglucuronyl – glycyrrhetic acid a major metabolite that causes licorice – induced pseudoaldosteronism. *J. Clin. Endocrin. Metab.* 80 (6): 1929 – 1933.
18. Kerstens, M. N. and R. Dullaart. 1999. 11 Beta – hydroxysteroid dehydrogenase: characteristics and the clinical significance of a key enzyme in cortisol metabolism. *Ned. Tijdschr. Geneesk.* 143(10): 509 – 514.
19. Kumagi, A., S. Yano and M. Otomo. 1957. Study on the corticoid – like action of glycyrrhizin and the mechanism of its action. *Endocrinol. Japon.* 4: 17 – 27.
20. Naji, S. A., I. A. Al-Ani, J. K. Minati and S. A. Mukhlis. 2004. The effect of *Lactobacilli* spraying and the addition of salts and vinegar to drinking water on the performance of heat stressed broiler. Submitted to Iraqi Agric. J. for publication.
21. Oyetayo, V. O., F. C. Adetuyi and F. A. Akinyosoye. 2003. Safety and protective effect of *Lactobacillus acidophilus* and *Lactobacillus casei* used as probiotic agent *in vivo*. *Afric. J. Biotech.* 2 (11): 448 – 452.
22. Pascual, M., M. Hygas, J. I. Badiola, J. M. Monfort and M. Garriaga. 1999. *Lactobacillus salivarius* CTC 2197 prevents *Salmonella enteritidis* colonization in chickens. *Appl. Environ. Microbiol.* 65: 4981 – 4986.
23. Quaschnig, T., F. Ruschitzka, S. Shaw and T. F. Lüscher. 2001. Aldosterone receptor antagonism normalizes vascular function in liquorice – hypertension. *Hypertension* 37: 801 – 813.
24. Rolfe, R. D. 2000. The role of probiotic cultures in the control of gastrointestinal health. *J. Nutr.* 130 (2S): 396S – 402S.
25. Ruszymah, B. H., B. M. Nabishah, S. Aminuddin and B. A. Khalid. 1995. Mineralocorticoid and glycyrrhizic acid blocks stress induced hypotension in rats. *Clin. Exp. Pharmacol. Physiol.* 22 (1): 35 – 39.
26. SAS. 1996. SAS User's Guide: Statistics Version, 6th edn., SAS Institute Inc., Cary, NC.
27. Schambelan, M. 1994. Licorice ingestion and blood pressure regulating hormones. [Review]. *Steroids* 59 (2): 127 – 130.
28. Shibata, S. A. 2000. A drug over the millennia: pharmacognosy, chemistry, and pharmacology of licorice. [Review]. *Yakugaku Zasshi.* 120 (10): 849 – 862.
29. Simmons, J. D., S. L. Branton and J. W. Deaton. 1989. Survivability resulting from increased fluid consumption with broiler exposed to high temperature. *Translocations of the ASAE* 32 (1): 238 – 240.
30. Smith, M. O. and R. G. Teeter. 1987. Effect of ammonium chloride and potassium chloride on survival of broiler chicks during acute heat stress. *Nutr. Res.* 7: 677 – 681.
31. Snow, J. M. 1996. *Glycyrrhiza glabra* L. (Leguminaceae). *Protocol J. Botan. Med.* 1: 9 – 14.
32. Stern, N. J., N. H. Cox and M. T. Musgrover. 2001. Comparison of mucosal competitive exclusion to reduce *Salmonella* and *camylobacter* spp. Colonized in broiler. *Poultry Sci.* 80: 156 – 160.
33. Whorewood, C. B., M.C. Shppard and P. M. Stewart. 1993. Licorice inhibits 11 β – hydroxysteroid dehydrogenase messenger ribonucleic acid levels and potentiates glucocorticoid hormone action. *Endocrinology* 132: 2287 – 2292.
34. Yeo, J. and K. I. Kim. 1997. Effect of feeding diets containing an antibiotic, a probiotic, or yucca extracts on growth and intestinal urease activity in broiler chicks. *Poultry Sci.* 76: 381 – 385.
