



Journal Homepage: -www.journalijar.com
**INTERNATIONAL JOURNAL OF
 ADVANCED RESEARCH (IJAR)**

Article DOI:10.21474/IJAR01/7353
 DOI URL: <http://dx.doi.org/10.21474/IJAR01/7353>



RESEARCH ARTICLE

INTRODUCTION TO PLANNER HARMONIC MAPPINGS.

Dr. Noohi Khan (AP II).

1. Amity School of Applied Sciences.
2. Amity University Lucknow, UP, INDIA.

Manuscript Info

Manuscript History

Received: 05 May 2018
 Final Accepted: 07 June 2018
 Published: July 2018

Abstract


In this paper we will study about the planner harmonic mapping starting from basic definition and theorem of harmonic starlike and convex function, harmonic univalent and multivalent function which is very useful for new researchers.

Copy Right, IJAR, 2018,. All rights reserved.

Introduction:-

Complex-valued function $f=u+iv$ in a simply connected domain D is called a **planar harmonic mapping** in complex domain D if the two conditions holds :

1. u, v are real harmonic in D
2. f is univalent in D .



In any simply connected domain D we can write $f = h + \bar{g}$, where h and g are analytic in D and h is called analytic part of f , g is called co-analytic part of f .

Clunie and Sheil-Small [15] observed that a necessary and sufficient condition for f to be locally univalent and sense-preserving in D is that $|h'(z)| > |g'(z)|, z \in D$. A family of all harmonic complex-valued, sense-preserving univalent functions normalized with the condition $h(0) = 0 = h'(0) - 1$ is denoted by SH.

SH- CLASS:-

Analogous to subclasses of S, various subclasses of SH have been defined and studied.

Note that if the co-analytic part of $f = h + \bar{g}$ is identically zero i.e. $g \equiv 0$ then harmonic functions reduces to analytic functions.

The SH denotes the family of all harmonic, complex valued, orientation-preserving normalized univalent functions defined on Δ . Thus the function f in SH admits the representation $f = h + \bar{g}$, where,

$$h(z) = z + \sum_{n=2}^{\infty} a_n z^n, \text{ and } g(z) = \sum_{n=1}^{\infty} b_n z^n; |b_1| < 1 \text{ are analytic functions in } \Delta.$$

It follows from the orientation-preserving property that $|b_1| < 1$. Therefore, $(f - \overline{b_1 f}) / (1 - |b_1|^2) \in SH$ whenever $f \in SH$. Thus a subclass SH^0 of SH is defined by $SH^0 = \{f \in SH : g'(0) = b_1 = 0\}$.

Note that $S \subset SH^0 \subset SH$. Both families SH and SH^0 are normal families. That is every sequence of functions in SH (or SH^0) has a subsequence that converges locally uniformly in Δ .

It is noted that $SH \equiv S$ if $g=0$.

TH -CLASS

Let TH denote the sub class of SH with negative coefficients whose members $f = h + \bar{g}$ where h and g are of the form

$$h(z) = z - \sum_{n=2}^{\infty} |a_n| z^n \text{ and } g(z) = \sum_{n=1}^{\infty} |b_n| z^n, |b_1| < 1, z \in \Delta.$$

Results:-

sense-preserving harmonic mapping $f \in SH$ (or $f \in SH^0$) is said to be harmonic starlike mapping in Δ if the range $f(\Delta)$ is starlike with respect to the origin. Likewise a function $f \in SH$ (or $f \in SH^0$) is said to be harmonic convex in Δ if $f(\Delta)$ is a convex domain.

Necessary and sufficient condition for a function $f(z) \in SH$ to be harmonic starlike univalent in Δ is that

$$(3) \quad \frac{\partial}{\partial \theta} \{\arg f(re^{i\theta})\} > 0, z \in \Delta, (z = re^{i\theta}, 0 \leq \theta \leq 2\pi, 0 \leq r \leq 1).$$

S*H-CLASS

S^*H denotes the class of harmonic starlike univalent functions. Note that if $b_1 = 0$ then $S^*H \equiv S^*H^0$.

T^*H denotes the class of harmonic starlike univalent functions $f \in TH$.

4. A necessary and sufficient condition for a function $f(z) \in SH$ to be harmonic convex univalent in Δ is that

$$(4) \quad \frac{\partial}{\partial \theta} \left\{ \arg \left(\frac{\partial}{\partial \theta} \arg f(re^{i\theta}) \right) \right\} > 0, z \in \Delta, (z = re^{i\theta}, 0 \leq \theta \leq 2\pi, 0 \leq r \leq 1).$$

KH CLASS:-

The class of harmonic convex univalent functions is denoted by KH.

Note that if $b_1 = 0$ then $KH \equiv KH^0$.

T^*KH denotes the class of harmonic convex univalent functions if f is of the form (2).

Results:-

1. If $f = h + \bar{g}$ with h and g are of the form (1) satisfies

$$(5) \quad \sum_{n=2}^{\infty} n |a_n| + \sum_{n=1}^{\infty} n |b_n| \leq 1$$

then, $f \in S^*H$.

2. If $f = h + \bar{g}$ with h and g are of the form (1.2.4) satisfies

$$(6) \quad \sum_{n=2}^{\infty} n^2 |a_n| + \sum_{n=1}^{\infty} n^2 |b_n| \leq 1$$

then, $f \in KH$.

3. Let $f = h + \bar{g}$ with h and g are of the form (2), then $f \in T^*H$ if and only if

$$(7) \quad \sum_{n=2}^{\infty} n |a_n| + \sum_{n=1}^{\infty} n |b_n| \leq 1.$$

4. Let $f = h + \bar{g}$ with h and g are of the form (2), then $f \in KH$ if and only if

$$(8) \quad \sum_{n=2}^{\infty} n^2 |a_n| + \sum_{n=1}^{\infty} n^2 |b_n| \leq 1.$$

RESULTS Of Uniformly starlike and convex

1. A function $f \in SH$ is said to be in the class $k\text{-USH}(\alpha)$ if it satisfies the following condition

$$(9) \quad \operatorname{Re} \left\{ (1 + ke^{i\phi}) \frac{zf'(z)}{f(z)} - ke^{i\phi} \right\} \geq \alpha, \quad 0 \leq \alpha < 1, \quad \phi \in \mathbb{R}.$$

Function $f \in SH$ given by (1.2.4) is said to be in the class $k\text{-HCV}(\alpha)$ if it satisfies the following condition

$$(10) \quad \operatorname{Re} \left\{ 1 + (1 + ke^{i\phi}) \frac{z^2 h''(z) + \overline{2zg'(z) + z^2 g''(z)}}{zh'(z) - \overline{zg'(z)}} \right\} \geq \alpha, \quad 0 \leq \alpha < 1, \quad \phi \in \mathbb{R}$$

SH (m) -CLASS

Let f be a harmonic function in a Jordan domain D with boundary C . Suppose f is continuous in \bar{D} and $f(z) \neq 0$ on C . Suppose f has no singular zeros in D , and let m to be sum of the orders of the zeros of f in D . Then $\Delta_c \arg(f(z)) = 2\pi m$, where $\Delta_c \arg(f(z))$ denotes the change in argument of $f(z)$ as z traverses C .

It is also shown that if f is sense-preserving harmonic function near a point z_0 , where $f(z_0) = \omega_0$ and if $f(z) - \omega_0$ has a zero of order m ($m \geq 1$) at z_0 , then to each sufficiently small $\epsilon > 0$ there corresponds a $\delta > 0$ with the property: "for each $\alpha \in N_\delta(\omega_0) = \{\omega : |\omega - \omega_0| < \delta\}$, the function $f(z) - \alpha$ has exactly m zeros, counted according to multiplicity, in $N_\epsilon(z_0)$ ". In particular, f has the open mapping property that is, it carries open sets to open sets.

Let Δ be the open unit disc $\Delta = \{z : |z| < 1\}$ also let $a_k = b_k = 0$ for $0 \leq k < m$ and $a_m = 1$. Ahuja and Jahangiri [5], [9] introduce and studied certain subclasses of the family $SH(m)$, $m \geq 1$ of all multivalent harmonic and orientation preserving functions in Δ . A function f in $SH(m)$ can be expressed as $f = h + \bar{g}$, where h and g are of the form

$$(11) \quad h(z) = z^m + \sum_{n=2}^{\infty} a_{n+m-1} z^{n+m-1}$$

$$g(z) = \sum_{n=1}^{\infty} b_{n+m-1} z^{n+m-1}, \quad |b_m| < 1.$$

According to Theorem and above argument, functions in $SH(m)$ are harmonic and sense-preserving in Δ if $J_f > 0$ in Δ . The class $SH(1)$ of harmonic univalent functions was studied in details by Clunie and Sheil Small. It was observed that m -valent mapping need not be orientation-preserving.

TH(m) CLASS

Let $TH(m)$ denotes the subclass of $SH(m)$ whose members are of the form

$$(12) \quad h(z) = z^m - \sum_{n=2}^{\infty} |a_{n+m-1}| z^{n+m-1}$$

$$g(z) = \sum_{n=1}^{\infty} |b_{n+m-1}| z^{n+m-1}, \quad |b_m| < 1.$$

Results:-

1. [5], [9] A function $f(z) \in SH(m)$ is said to be multivalent harmonic starlike if and only if

$$(13) \quad \frac{\partial}{\partial \theta} \left\{ \arg(f(re^{i\theta})) \right\} > 0, \quad z \in \Delta, \quad m \geq 1.$$

S^*H_m denotes the class of multivalent harmonic starlike functions

It is clear that $S^*H_m \subset SH(m)$ and $S^*H_1 \equiv S^*H$.

Also, T^*H_m denotes the class of harmonic starlike m -valent functions $f \in TH(m)$.

2.[8] A function $f(z) \in SH(m)$ is said to be multivalent harmonic convex if and only if

$$(14) \quad \frac{\partial}{\partial \theta} \left\{ \arg \left(\frac{\partial}{\partial \theta} f(re^{i\theta}) \right) \right\} > 0, \quad z \in \Delta, \quad m \geq 1.$$

KH_m denotes the class of multivalent harmonic convex functions. It is clear that $KH_m \subset SH(m)$ and $KH_1 \equiv KH$.

References:-

1. Ahuja, O.P., Jahangiri, J.M., and Silverman, H., Convolutions for special classes of Harmonic Univalent Functions, Appl. Math. Lett, 16(6) (2003), 905-909.
2. Ahuja, O.P. and Silverman, H., Extreme Points of families of Univalent Functions with fixed second coefficient. Colloq. Math. 54 (1987), 127-137.
3. Ahuja, O.P. and Jahangiri, J.M., Harmonic Univalent Functions with fixed second coefficient, Hakkoido Mathematical Journal, Vol. 31(2002) p-431-439.
4. Ahuja, O.P. and Silverman, H., Inequalities associating hypergeometric Functions with planar harmonic mappings, Vol 5, Issue 4, Article 99, 2004.

5. Ahuja,O.P. and Jahangiri,J.M., Multivalent Harmonic starlike Functions, Ann. Univ Mariae Curie – Skłodowska, Section A, 55 (1) (2001), 1-13.
6. Ahuja,O.P. and Jahangiri,J.M., A subclass of Harmonic Univalent Functions, J.Natural geometry, 20, (2001), 45-56.
7. Ahuja,O.P. and Jahangiri,J.M., Noshiro Type Harmonic Univalent Functions, Math. Japonica, 56, No. 1(2002), 1-7.
8. Ahuja,O.P. and Jahangiri,J.M., Multivalent Harmonic convex Functions, submitted for publication.
9. Ahuja,O.P. and Jahangiri,J.M., Errata to “Multivalent Harmonic starlike Functions, Ann. Univ. Mariae Curie-Skłodowska, Vol LV., 1 Sectio A 55 (2001), 1-3, Ann. Univ. Mariae Curie – Skłodowska, Sectio A, 56(1) (2002), 105.
10. Khan,Noohi “Multivalent Harmonic functions with a fixed coefficient”, International Journal of Mathematics, Science & Engineering Applications (IJMSEA) Vol.3 No. IV (2009), PP-305-316.
11. Khan,Noohi “Harmonic Multivalent functions involving a Linear Operator”, International Journal of Mathematical Analysis, Vol.3, 2009, No.6, 295 – 308 Hikari Publication.(Scopus indexed journal)
12. Khan,Noohi “Inequalities of a Generalized class of k-uniformly harmonic univalent functions”Published in IOSR Journal of Mathematics(IOSR-JM)e-ISSN:2278-5728.Volume 11,issue 1,Ver.VI(Jan-feb 2015),www.iosrjournals.org.
13. Khan,Noohi “A Class of Multivalent Harmonic Functions involving a Salagean operator published in GJSFR-Global journal of Mathematics,Volume 15-F Issue 2 Version 1.0,in feb 2015.
14. Khan,Noohi “A Generalized Class of k-Uniformly harmonic functions based on Salagean operators, published online in SAJM south asian journal of mathematics in march 2015 Vol.5 (2)89-96.ISSN 2251-1512.
15. Khan,Noohi “Inequalities on Multivalent harmonic starlike functions involving hypergeometric functions” published in IOSR-JM in may 2016,vol 12,issue 2,ISSN:2319-765X.