




Journal Homepage: -www.journalijar.com
**INTERNATIONAL JOURNAL OF
 ADVANCED RESEARCH (IJAR)**

Article DOI: 10.21474/IJAR01/6975
 DOI URL: <http://dx.doi.org/10.21474/IJAR01/6975>



RESEARCH ARTICLE

PHASE TRANSFER CATALYTIC SYNTHESIS, SPECTRAL CHARACTERIZATION AND ANTIMICROBIAL EVALUATION OF 2-(PHENYL SELENO) ETHANOIC ACID AND 3-(PHENYL SELENO) PROPANOIC ACID.

Richa Saxena¹ and Ambika Chopra².

1. Department of Chemistry, ITM University, Gwalior.
2. SOS in Chemistry, Jiwaji University, Gwalior.

Manuscript Info

Manuscript History

Received: 17 February 2018
 Final Accepted: 19 March 2018
 Published: April 2018

Keywords:-

α -Chloroacetic acid, Phase transfer catalyst, ¹H-¹H COSY NMR.

Abstract

2-(phenyl seleno) ethanoic acid (**1**) and 3-(phenyl seleno) propanoic acid (**2**) has been synthesized in more economical and efficient method using phase transfer catalyst, tetraethylammonium bromide. Compound **1** & **2**, has been synthesized by reacting nucleophile, PhSe⁻ (generated *insitu* by borohydride reduction of Ph₂Se₂) with α -chloro acetic acid & 3-chloro propanoic acid via phase transfer catalysis. The presence of catalyst improves the yield of the products and lowers the reaction time, however in absence of the catalyst yield was very poor. Elemental analysis, IR, ¹H, ¹³C, ¹H-¹H COSY NMR, ESI mass and molar conductance data has been used to authenticate the new species. The antimicrobial evaluation of both the compounds indicates their significant activity at different concentrations against selected bacterial and fungal strains. These Compound bear an acidic group, showed higher activities due to increase in the lipophilicity and easier penetration of the compounds into the outer cell wall of the microorganisms, which causes death due to cell membrane rupturing thereby boosting the bioactivity of these compounds.

Copy Right, IJAR, 2018,. All rights reserved.

Introduction:-

The past two decades have witnessed a remarkable growth in the field of organoselenium/tellurium chemistry. They comprises a wide variety of structures that display a remarkable range of properties. During the last few years, a tremendous efforts has been directed towards the synthesis of stable organoselenium compounds. Such compounds have been found to function as antioxidants, chemoprotectors, apoptosis inducers and chemopreventors in several organs such as brain, liver, skin, colon, lung and prostate^{1,2}.

Besides this organoselenium/tellurium compounds have also showed a great potential for application in biological systems³, organic synthesis⁴, pharmaceutical⁵⁻⁷, semiconducting material⁸⁻¹⁰, ligand chemistry¹¹, biochemistry¹², spectroscopic studies¹³ and have wide applications as homogenous catalyst¹⁴. Recently a novel organotellurium compound (RT-01), organotellurane, has been identified as a new antileishmanial agent¹⁵ besides this certain selenium derivatives like benzeneselenazoles show antibacterial activities against gram-negative E-coli K-12 ROW and gram-positive S.aureus 209P bacteria strains. Even 4H-5,6-dihydro-1,3-selenazine derivatives also shows antibacterial activity against E.coli and S.aureus¹⁶.

Corresponding Author:- Richa Saxena.

Address:- Department of Chemistry, ITM University, Gwalior.

The phase transfer catalyst like cetyltriethyl ammonium bromide or tetraethyl ammonium bromide is believed to be used in reactions where the two phases are mutually immiscible, the catalyst continuously transfers reacting anions in to the organic phase in form of lipophilic ion pairs and helps in proceeding the reaction, that too at faster rate. Transformations of starting materials into desired final products usually require a number of chemical operations in which additional reagents, catalysts, solvents, etc. are used. Thus in course besides desired products, many waste materials are produced. These wastes should be regenerated, destroyed and disposed consuming much energy and creating heavy burden in the environment. Thus the most general and efficient methodologies that fulfill the requirement is phase transfer catalysis¹⁷. The methodology is efficiently applicable to a plethora of other base induced reactions of organic anions particularly carbanions.

Prompted by the above reports, herein we report an efficient method for design and synthesis of selenium bearing compound having [Se, O] donor functionalities through phase transfer catalyst. It is worthwhile to mention that use of tetraethyl ammonium bromide as catalyst is especially useful in green chemistry. These compounds have been characterized by various spectroscopic techniques. The antimicrobial & antioxidant activity of the same are also reported in the following pages.

Experimental:-

Diphenyl diselenide¹⁸ was prepared according to the published procedures. α -chloro acetic acid, 3-chloro propanoic acid and tetraethyl ammonium bromide were purchased from CDH (India). Common organic solvents (procured from Qualigens, CDH, Spectrochem and Merck) viz. chloroform, n-hexane, petroleum ether (40-60°C and 60-80°C), ethanol, methanol, dimethyl sulphoxide (DMSO), dimethylformamide (DMF) etc were used after further purification and drying by standard methods¹⁹. Peptone, beef extract, dextrose, NaCl and agar were purchased from CDH India Ltd. were used as such without any further purification.

The antimicrobial activity has been assessed by using disc diffusion technique for bacterial and fungal strains. Chloramphenicol and fluconazole were used as standard drugs respectively²⁰⁻²².

Synthesis of 2 -(phenylseleno) ethanoic acid (1):-

To the solution of diphenyl diselenide (3.12g, 10mmol) in dry ethanol (15mL) under dry nitrogen atmosphere, sodium borohydride (0.74g, 20mmol) in 2 ml of 1M NaOH was added dropwise until the solution become colorless. To this reaction mixture tetraethyl ammonium bromide (2.10g, 10mmol), a phase transfer catalyst was added. The ethanolic solution of chloroacetic acid (1.89g, 20mmol) was then added to the mixture dropwise and was allowed to stir overnight. It was washed with 100 mL of dil. HCl solution and extracted into 100 mL (25 x 4) of chloroform. The extract was dried over anhydrous magnesium sulphate and concentrated to 5mL on a rotary evaporator under reduced pressure. It was purified by TLC through a mixture of hexane and chloroform (2:1) to yield α -(phenylseleno) acetic acid in the form of yellow viscous oil.

Synthesis of α -(phenylseleno) propanoic acid (2):-

This was synthesized using the above mentioned method except that here 3-chloro propanoic acid was used to produce the compound in form of orange viscous oil.




Fig. 1 :- Zone of inhibition surrounding the disc of compound 1 at 20 $\mu\text{g/mL}$ and 40 $\mu\text{g/mL}$, against *E.coli.*, *B.anthraxis*, *S.aureus* resp.





Fig. 2:- Zone of inhibition surrounding the disc of compound 3 at 20 $\mu\text{g/mL}$ and 40 $\mu\text{g/mL}$ against *T.rubrum*.

Results and Discussion:-

Compound **1** and **2**, were synthesized by the interaction of nucleophile, sodium phenylselenolate (NaPhSe^-) with α -chloroacetic acid & 3-chloropropanoic acid using tetraethylammonium bromide as a phase transfer catalyst²³ in dry ethanol results in the formation of **1** & **2**, as yellow & orange viscous oil in high yield. However in the absence of tetraethyl ammonium bromide the yield was only about 2%.



Scheme 1

Elemental analysis of compounds **1** and **2** provides supporting evidence of their stoichiometry. The compounds are stable and can be stored under ambient conditions upto six months. They have good solubility in CHCl_3 , CH_2Cl_2 , CH_3OH , $\text{C}_2\text{H}_5\text{OH}$. The molar conductance data of **1** and **2** have been determined in Dimethyl formamide at ~1Mm concentration level and the values are given in the **Table 1**. The conductance value of **1** and **2** have been found to be much lower than the value expected for any electrolyte and hence they may be considered to behave as non electrolyte or non-ionic²⁴ (**Table 2**).

Various spectroscopic studies reveals the synthesis of said compounds (**Table 3-5**). The assignments have been made on the basis of literature reports of related compounds and additivity principles²⁵. The results obtained are reported in the Table which agrees well with the reported values²⁶.

Antimicrobial activity:-

The antimicrobial activity of compounds **1** and **2** were tested at two different concentrations by disc diffusion method on bacterial and fungal strains namely *Staphylococcus aureus*, *Bacillus anthracis*, *Escherchia coli* and *Trichophyton rubrum*, *Candida tropicalis*, *Aspergillus niger* respectively. The media used for studies are mentioned in **Table 6**. The compounds show significant activity against all the organisms of bacteria and fungi. However, their activity was found to be minimum against *Aspergillus niger*. It was also concluded that the compounds are slightly more effective against the organism at 40 $\mu\text{g/L}$ then at 20 $\mu\text{g/L}$ (**Table 7 & 8**). It is evident from the data, obtained from antifungal studies that few fungal strains are slightly harder to treat, in comparison to bacteria, because of their cell wall, which is made up of chitin. Thus, the values of the zones of inhibition (**Table 8**) were found to be marginally lower than the values, obtained against bacterial strains, which are easier to penetrate. Both the compound showed minimum activity against A.Niger, which may be due to the carboxylic groups, which are also components of the cell wall of this species. Therefore this makes it difficult for compounds, to easily penetrate into the cell wall of the strain²⁷.


Conclusion:-

2-(phenyl seleno) ethanoic acid (**1**) & 3-(phenyl seleno) propanoic acid (**2**) has been synthesized in high yield (90%) by the reaction of phenylselenolate anions generated *insitu* by borohydride reduction of diphenyldiselenide, with chloroacetic acid and chloro propanoic acid using tetraethyl ammonium bromide as phase transfer catalyst. However in the absence of catalyst, the yield of **2** was about 2%. The catalyst is ecofriendly and improves the yield

of the products. The compounds were found to be significantly effective against certain selected bacterial and fungal strains.

Structure:-

From the preceding discussions mainly based on molar conductance, elemental analysis, ESI mass, IR, ^1H , $^{13}\text{C}\{^1\text{H}\}$ and support from the previous work, the most possible structure of **1** and **2** have been proposed.



Acknowledgements:-

Author acknowledges Prof. S. K. Srivastava, School of studies in Chemistry, Jiwaji University, Gwalior, for his valuable suggestions and encouragement as well as for providing all the facilities which were required to carry out the given piece of work. The author is also thankful to the Director, Central Drug Research Institute, Lucknow, India and Chairman CISC, Banaras Hindu University, Varanasi, India for C, H, N analysis, ESI measurements and ^1H , ^{13}C NMR spectra, respectively.

References:-

1. J.M. Lochowski, K.Kloc, R.Lisiak, P.Potaezek and H.W. Wojtowicz, *Arkivoc VI* (2007) 14.
2. D.M. Driscoll and P.R. Copeland, *Annu. Rev. Nutr.*, 23 (2003) 17.
3. C. Reilly, *Selenium in Food and Health*, Blackie Academic and Professional, London (1996).
4. A.Krief, W.W. Abel, F.G.A. Stone and G.Wilkinson (Eds.), *Comprehensive Organometallic Chemistry*, Pergamon, Oxford, U.K, (1995) 515.
5. M. Carland, T. Fenner, M. Gielen and E.R.T. Tiekling (Eds.), *Medicine in Metallotherapeutic Drugs and metal Based Diagnostic Agents*, Wiley, Chichister, (2005).
6. S. Gracia, *Curr. Med. Chem.*, 11(2004) 1657.
7. A.E. Okoronkwo, B. Godoi, R.F. Schumacher, J.S Santos neto, C. Luchese, M. Prigol, C.W. Nogueira and G. Zeni, *Tet.Lett.*, 50 (2009) 909.
8. H.B. Singh and N. Sudha, *Polyhedron*, (1996) 745.
9. M. Bochman, *Chem. Vapor Depos.*, 2 (1996) 85.
10. S. Mishra, E. Jeanneau and S. Daniele, *Polyhedron*, 29 (2010) 500.
11. L.R. Gray, D.J. Gulliver, W. Levason, M. Webster, *J. Chem. Soc., Dalton Trans.*, (1983) 133.
12. G. Muges, W.-W. duMont and H. Sies, *Chem. Rev.*, 101(2001) 2005.
13. R. Wu, L.A. Silks, J.D. Odom, R.B. Dunlap, *Spectroscopy*, 11 (1996) 37.
14. A. Kumar, M. Agarwal, A.K. Singh and R.J. Butcher, *Inorg. Chim. Acta.*, 362 (2009) 3208.
15. R.Cunha and S.Giorgio, *Korean J. Parasitol.*, 47(3) (2009) 213.
16. M. Koketsu, H. Ishihara and M. Hatsu, *Res. Commun. Mol. Path. Pharmacol*, 179 (1998) 101.
17. M. Makosza and M. Fedorynski. *Adv. Catal.* 35, (1987); E.V. and Dehmlow, S.S. *Demlow. Phase-Transfer Catalysis*, 3rd ed.;Verlag Chemie, Weinheim 1993.
18. K.B. Sharpless and M.W. Young, *J. Org. Chem.*, 40 (1975) 947.
19. W.L.F. Armarego and D.D. Perrin, "Purification of Laboratory Chemicals", 4th Ed., (1998).
20. Bauer AW, Perry D M, Kirby W M M Single disc antibiotic sensitivity testing of Staphylococci. *A.M.A.*

- Arch., Intern. Med 104 (1959) 208-216.
21. Bauer A W, Kirby W M M, Sherris J C, Turck M, Am. J. Clin. Pathol 36 (1996) 493.
 22. Jorgenson J H, Turnidge J D Manual of Clinical Microbiology 9th Ed. ASM Press, Washington. D.C 1152-1172.
 23. M. Makoszka, Pure Appl. Chem., 72 (2000) 1399.
 24. Geary W J Coord. Chem. Rev. Edit. lever 7 (1971) 110.
 25. Silverstein R M, Bassler G C and Morrill T C 1998 Spectrometric Identification of Organic Compounds (6th Edn) (John Wiley) N.Y.
 26. A. Khanna, A. Bala and B.L. Khandelwal, J. Organomet. Chem.. 494 (1995) 199.
 27. Akhtar M., Sastry K., Mohan P., Mechanism of metal ion biosorption by fungal Biomass, Biometals 9 (1996) 21-28.