



ISSN NO. 2320-5407

Journal homepage: <http://www.journalijar.com>  
 Journal DOI: [10.21474/IJAR01](https://doi.org/10.21474/IJAR01)

INTERNATIONAL JOURNAL  
 OF ADVANCED RESEARCH

## RESEARCH ARTICLE

## A Study of Knowledge &amp; Awareness of Radiation Exposure Risk in Makkah, Saudi Arabia.

Nizar Almaghrabi

MS, Umm Al-Qura University – Makkah – Saudi Arabia

**Manuscript Info****Manuscript History:**

Received: 15 May 2016  
 Final Accepted: 22 June 2016  
 Published Online: July 2016

**Key words:**

knowledge and awareness; radiation dose; patient; cross-sectional study; online questionnaire; radiological imaging; Makkah; Saudi Arabia.

**\*Corresponding Author**

Almaghrabi. N.

**Abstract**

**Background:** Radiological procedures are being used much more these days. Patients must be given sufficient information, in a way that they can understand, to be able to make the right decisions about their care.

**Objectives:** To assess the awareness and associated risks caused by medical radiation among the general population of Makkah, Saudi Arabia

**Methods:** A pilot study among the general population of Makkah, Saudi Arabia, 2016.

**Results:** A total of 100 subjects answered the questionnaires. 56% females and 44% male respondents; mean age were 23 years, ranged from 16 to 45. Of these, 60% had attended college, 34% had completed high school, and 5% completed intermediate school, and 1% completed primary school. 69% have experience with medical imaging. Most subjects underwent X-rays (80%), MRI (30%), US (26%), CT (22%) and others (1%).

Seventy-one% and 30% were aware of the radiation exposure involved in plain X-ray and CT scans, respectively. Furthermore, 32% and 17% were not aware of the free nature of MRI and US from radiation, respectively. Further, 30% incorrectly thought that barium studies, do not involve radiation.

Moreover, 67% and 63%, think that they are not vulnerable to radiation on a plane and at home, respectively. 85% were told about the indication, and 9% told about the risks, and 13% were told the associated radiation dosage. 15.6% incorrectly thought that cancer risk does not increase with repeated radiation exposure. Education affect radiation knowledge significantly.

**Conclusion:** Radiation awareness is poor among the general population. They need to be provided with the necessary information to improve their radiation awareness.

Copy Right, IJAR, 2013. All rights reserved.

**Introduction: -**

Radiological procedures are being used much more these days with the advance in technology. It helps in the diagnosis and management of many medical conditions. Requesting imaging modalities compose risk to the patients (ionizing radiation cancer-causing biological effects).

Every year there is increasing in the number of patients who stand for diagnostic radiology<sup>1,2</sup>, Especially CT scanning. Radiation doses have increased up to 40% per scan during the last few years<sup>3</sup>. Radiation exposure repeatedly increases the risk of cancer.

The lowest dosage of radiation for which there is a real proof of cancer-causing is around 10–50 mSv. The regular exposure dosage for one chest radiograph taken is around 0.02 mSv, and for an abdominal CT is around 9 mSv. The radiation from CXR is probably less than background radiation received in a whole year (0.01 mSv daily). Around 0.015 mSv is received during a three-hour airline flight<sup>4,5</sup>.

It is essential that doctors who request imaging to be well trained in determining whether diagnostic imaging is required, but also they need to be aware of the associated risk. It has appeared in different studies that medical professional's knowledge on radiation dangers and dosage is inadequate<sup>6,7</sup>.

In 2006 a study published in the Pediatric Radiology Journal, 87% of pediatricians underestimated the radiation dosage from a chest radiograph and 94% underestimated the radiation dosage from a CT<sup>8</sup>.

Not only doctors are inadequate in knowledge on radiation dosage and risks. A study in 2010 showed, half of the intern doctors and senior medical students underestimated the radiation doses from usually requested radiological procedures. Some of them incorrectly think that ultrasound and MRI produce ionizing radiation<sup>9</sup>.

A study conducted in 2009 shows a significant proportion of physicians anticipate that informed consent should be obtained from patients undergoing radiological exams and the information about cancer-related risks involved should be provided by the radiological department<sup>10</sup>.

Most participants did not talk about the associated risks with patients. Patients must be given sufficient information, in a way that they can understand, to be able to make the right decisions about their care.

Few Projects are focusing on the patient's knowledge about radiation risk in the literature. Two recent studies demonstrated that the majority of patients (74%) would believe that having their condition diagnosed with CT is more important than worrying about radiation and patients had insufficient knowledge about radiation protection<sup>7,11</sup>.

Surprisingly, a number of articles have appeared in the literature that predict hundreds of cancers and cancers mortality per year in the U.S. and U.K. caused by ionizing radiation from medical imaging procedures. It was estimated 100-250 deaths occur per year from cancers directly related to exposure to medical radiation in U.K<sup>12</sup>. In U.S, the estimated number of fatalities attributable to CT was 700-1800 during a year<sup>13</sup>.

Our aim of this study is to assess the awareness of radiation danger and associated risks among the general population. It's hoped that the results of our study would prove that patient education is necessary these days.

### **Material and methods: -**

A pilot questionnaire study conducted in Makkah, Saudi Arabia, 2016, of 100 subjects. The subjects were given an online questionnaire randomly. They were stratified by age, sex, and education. The online questionnaire was in Arabic, multiple choice format and consisted of 19 questions. Divided into three sections inspecting demographic information, radiation knowledge/awareness, and expectations.

In producing the questionnaire, we drew much inspiration from Robert H. Corbett's pilot study<sup>14</sup>.

Firstly, the subjects asked about his demographic and if they had any medical imaging before. If they answer yes, they asked about which imaging modality they were received and which part of their body examine and whether the doctor had explained to them the reason for performing the test, and the associated radiation dose and risk.


In the second section, which examined knowledge, there were ten questions. All subjects answer this part, whether they had medical imaging before or not. All of these were multiple choice questions. This assures all answers were either right or wrong, and there would be no ambivalence in assessing subjects knowledge. The answers to questions were based on the cited references<sup>4,15</sup>.

Finally, in the last section, patient's expectations were examined. The subjects were asked whether or not they expect that point to be explained.

### Results: -

A total of 100 subjects answered an online questionnaire. The age ranged from 16 to 45. The mean was 23. A good balance of the sexes was achieved, with 56% of the subjects being female and 44% males. The study population represents a highly educated group of people, with 60% having attended university or college, 34% having completed high school and 5% completed intermediate school and 1% completed primary school.

Sixty-nine% have experience with medical imaging. Most subjects underwent X-rays (80%), MRI (30%), US (26%), CT (22%) and others (1%), (Figure1).



(Figure1)


A purpose of this study is to assess the target population's knowledge on radiation associated with medical imaging. Thus, 10 questions were devised to accomplish this task (correct answers are marked with ✓). A score of 1 was given to a correct answer and 0 otherwise. For each subject, a maximum score of 10 was calculated. The mean was 5.4 and median was 4.9 scores for all subjects.

Seventy-one% and 30% were aware of the radiation exposure involved in plain X-ray and CT scans, respectively. Furthermore, 32% and 17% were not aware of the free nature of MRI and US from radiation, respectively. Further, 30% incorrectly thought that barium studies, do not involve radiation.

The analysis was done by using SPSS 24 (SPSS Inc., Chicago, IL, USA). There's was no significant differ between male and female  $P > .05$ . Education achieved statistical significance  $p < .05$ .

General knowledge on radiation was unsatisfactory, 67% and 63%, think that they are not vulnerable to radiation on a plane and at home, respectively. 15.6% incorrectly thought that cancer risk does not increase with repeated radiation exposure. 85% of subjects told of the reason for their being subjected to radiological imaging, and 9% told about the risks, and 13% were told the Associated radiation dosage.

Sixteen% of the subject think that it's not necessary to be told about the radiation dosage. 48% of the subjects agreeing that their doctor should tell them about the radiation dosage while 43% thought the radiographer should tell them the radiation dosage and 3% thought the nurse should tell them that.



(Figure2)

**Discussion: -**

Our sample population represents a highly educated group, and may not necessarily be representative of the target population. This study has limitations regarding generalization. Regardless of the above limitations, this study shows that general population awareness is unsatisfactory on radiation.

Radiation is the primary risk involved in most diagnostic imaging. Patients need to be provided with the necessary information to increase their radiation awareness.

Besides, there is a clear mismatch between actual practice and patient expectation. Patient education about radiation should be part of the responsibility of healthcare providers. All health providers should equip themselves with the appropriate information about ionizing and non-ionizing radiation.

**Conclusion: -**

Radiation is the primary risk involved in most diagnostic imaging. Radiation awareness is poor among the general population. The patients need to be provided with the necessary information to increase their radiation awareness.

**References: -**

1. Brenner DJ, Hall EJ. Computed tomography—an increasing source of radiation exposure. *New England Journal of Medicine*. 2007;357(22):2277-2284.
2. Broder J, Fordham LA, Warshauer DM. Increasing utilization of computed tomography in the pediatric emergency department, 2000–2006. *Emergency radiology*. 2007;14(4):227-232.
3. Golding S, Shrimpton P. Radiation dose in CT: are we meeting the challenge? *The British journal of radiology*. 2002;75(889):1-4.
4. Ritenour ER, Geise RA. Radiation sources: medicine. *Health Effects of Exposure to Low-Level Ionizing Radiation, Philadelphia, Institutes of Physics Publishing*. 1996;441.
5. Radiation UNSCotEoA. *Sources and effects of ionizing radiation: sources*. Vol 1: United Nations Publications; 2000.
6. Fartum AR, Gjertsen J-E, Larsen JL. [Patients' knowledge of the effects of X-rays]. *Tidsskrift for den Norske laegeforening: tidsskrift for praktisk medicin, ny raekke*. 2000;120(28):3427-3428.
7. Düzeyleri RKKB, Çalışması KBA. Knowledge About Ionizing Radiation and Radiation Protection Among Patients Awaiting Radiological Examinations: A cross-sectional survey.
8. Thomas KE, Parnell-Parmley JE, Haidar S, et al. Assessment of radiation dose awareness among pediatricians. *Pediatric radiology*. 2006;36(8):823-832.
9. Zhou G, Wong D, Nguyen L, Mendelson R. Student and intern awareness of ionising radiation exposure from common diagnostic imaging procedures. *Journal of medical imaging and radiation oncology*. 2010;54(1):17-23.
10. İKarsli T, Kalra MK, Self JL, Rosenfeld JA, Butler S, Simoneaux S. What physicians think about the need for informed consent for communicating the risk of cancer from low-dose radiation. *Pediatric radiology*. 2009;39(9):917-925.
11. Takakuwa KM, Estepa AT, Shofer FS. Knowledge and attitudes of emergency department patients regarding radiation risk of CT: effects of age, sex, race, education, insurance, body mass index, pain, and seriousness of illness. *American Journal of Roentgenology*. 2010;195(5):1151-1158.
12. Quinn A, Taylor C, Sabharwal T, Sikdar T. Radiation protection awareness in non-radiologists. *The British journal of radiology*. 1997;70(829):102-106.
13. Brenner DJ, Elliston CD, Hall EJ, Berdon WE. Estimated risks of radiation-induced fatal cancer from pediatric CT. *American journal of roentgenology*. 2001;176(2):289-296.
14. Corbett RH. What do patients really know or want to know about X-rays. Paper presented at: Proceedings of 10th International Congress of the International Radiation Protection Association, Hiroshima, Japan 2000.
15. Friedberg W, Copeland K, Duke FE, O'Brien III K, Darden Jr EB. Radiation exposure during air travel: guidance provided by the Federal Aviation Administration for air carrier crews. *Health physics*. 2000;79(5):591-595.