



Journal Homepage: -www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/11496
DOI URL: <http://dx.doi.org/10.21474/IJAR01/11496>



RESEARCH ARTICLE

An overview on *Spilanthes acmella*

Mohd. Naeem Najam Ali Khan and Mohd. Abid*

School of Pharmaceutical Sciences, IFTM University, Lodhipur Rajput, Delhi Road (NH-24), Moradabad (UP), India.

Manuscript Info

Manuscript History

Received: 10 June 2020
Final Accepted: 14 July 2020
Published: August 2020

Key words:-

Sphilanthesacmella, Anti-oxidant,
Immunomodulatory, Anti-Pain
Relieving, Vasorelaxant Movement And
Neuroprotective Etc

Abstract

Spilanthes acmella, (Family-Asteraceae) is normally known as akarkara. *Spilanthesacmella* (syn) *Acmellaoleracea* is a typical elaborate plant developed in Brazil. It is otherwise called eyeball plant because of its trademark appearance of the bloom head and furthermore known as against toothache plant. It is utilized therapeutically in Indochina, Philippine islands, Lareunion and Madagascar; *Spilanthesacmella*, presented from Brazil and regularly developed in garden in numerous piece of India. A concentrate of the leaves and blossoms is customarily utilized for the cure of toothache due to sedative properties, stomatitis, influenza, hack, rabies infections, tuberculosis, throat grievances, ailment and fever. Different kinds of bioactive mixes have been disconnected from the plant occasionally. It has been accounted for exceptionally viable in different illnesses including, Anti-pyretic, Anti-pain relieving, Vasorelaxant movement, Local sedative action, Anti-contagious, Anti-ulcer, Anti-viral, Anti-diuretic, Anti-incendiary, Anti-oxidant, Immunomodulatory, Hepatoprotective, Anti-disease, AntiAIDS and hostile to toothache.

Copy Right, IJAR, 2020,. All rights reserved.

Introduction:-

The leaves and underlying foundations of plant *Sphilanthesacmella*, Family-Asteraceae was researched for its physicochemical and phytochemical screening. The entire plant of *Sphilanthesacmella* is accounted for to show great therapeutic qualities in conventional arrangement of meds particularly for the prophylaxis of different ailments. The current examination manages phytochemical examinations of assurance of concentrates (Petroleum ether remove, ethanol separate, Methanol extricate, and fluid concentrate). Complete sugar and tannin substance was considered. Ethanol solvent extractive worth was discovered higher in the leaves while fluid dissolvable extractive worth was found to higher in the roots. All out starch rate was discovered higher in the leaves. Be that as it may, tannin substance was found at more significant levels in the root bit of the plant. The primer phytochemical screening of *Sphilanthesacmella* leaves and roots indicated the nearness of Alkaloids, Glycosides, Tannins, Flavonoids, Terpenoids and Phlobatannis in various concentrates of *Sphilanthesacmella*^[1].

The most widely recognized alkamide present in *Sphilanthesacmella* is called *Sphilanthol*. In excess of 45 different mixes have been separated, the overwhelming ones being hexanol, tridecanone, germacrene, hexanol, caryophyllene and sequesterpenes^[2].

Corresponding Author:- Mohd.Abid

Address:- School of Pharmaceutical Sciences, IFTM University, Lodhipur Rajput, Delhi Road (NH-24), Moradabad (UP)-240 192 India.

This plant is famous among the old ancestral network; exceptional food thing is set up from this plant in strict celebration. The needy individuals offered this plant alongside the "Ajeng Dues" in DoburUie^[3].

Specifically, this plant is celebrated as an old stories solution for toothache and for throat and gum contaminations^[4].

The blossoms are squashed and applied at the site of toothache, especially in "Irula clan of Hasanur slopes in Erode region of Tamilnadu" where it is known by the nearby name "Mandal Poo Chedi"^[5].

Aside from Tamil Nadu, root glue of the plant is utilized in throat issues in Chindwara and Betul area of Madhya Pradesh^[6].

The plant is additionally referred to be utilized as panacea (Sumatra), as energizer, for toothache (Sudan), for stomatitis (Java), and for wound mending (India)^[7].

In Cameroon, the plant is utilized as a snakebite cure and in the treatment of articular stiffness^[8]. It should be helpful in instances of tuberculosis^[9].

In India, *S. acmella* blossom heads are utilized to treat stammering in youngsters. Leaves and blossoms of the plant are likewise used to treat leucorrhoea in females among individuals of clans in Bangladesh^[7].

Scientific Classification/Taxonomical (According to Bentham and Hooker Classification):

Kingdom-Plantae
 Subkingdom-Phanerogamia
 Class-Dicotyledonae, Magnoliopsida
 Subclass-Gamopetalae
 Arrangement-Inferae
 Order – Asterales
 Family-Asteraceae
 Subfamily-Mimosoideae
 Variety-Sphilanthes
 Species-acmella

Vernacular names:

S. acmella is generally known as Toothache Plant, Paracress, Eyeball Plant, and Spot Plant. Distinctive vernacular names of the plant are:

Languages Vernacular names:

Marathi Akkalkara, pipulka
 Hindi Akarkara, Pipulka
 Kannada Vanmugali, hemmugulu
 Telugu Maratimogga, maratiteega
 Assamese Pirazha
 Malayalam Kuppumaanjel
 Punjabi Akarkarha, pokarmul
 Mundari Raipuru
 Adi Marsan
 Tangkhul Ansahan

Distribution:

This plant is broadly disseminated in the tropical and sub-tropical districts including America, North Australia, Africa, Malaya, Borneo, India and Sri Lanka. In India, it is bound to South and focal piece of India, and Jharkhand^[10].

Plant description:

Spilanthesacmella is equivalent with spilantholeracea. The family Spilanthes contains 35 tropical species, of which three of them are accounted for from India. Spilanthesacmella is extremely delightful, erect or climbing strong spices, 20-50 cm high and can be developed as a yearly in many atmospheres. It is ice delicate however perpetual in hotter atmospheres. A little, erect plant, it develops rapidly and sends up gold and red bloom inflorescences. It very well may be developed in the ground or as a pruned. A rich soil with fertilizer is reasonable and keeps up a temperature of around 70 °F. Stems are glandular bristly with sharp taste. The entire plant is bitter in taste. It has striking cone-like blossoms. Spilanthesacmella has no bloom petals, yet rather, shows brilliant "buds" with a rustredfocus^[11].

Bio active compounds:

Phytochemicals are dynamic parts present in plants that have defensive or sickness preventive properties. The most widely recognized alkamide present in Spillanthescmella is called Sphilanthol. In excess of 45 different mixes have been detached, the overwhelming ones being hexanol, tridecanone, germacrene, hexanol, caryophyllene and sequesterpenes^[2].

Medicinal uses:

A concentrate of the leaves and blossoms is customarily utilized for the cure of toothache in view of sedative properties, stomatitis, influenza, hack, rabies illnesses and tuberculosis and throat grumblings^[12].

It has additionally utilized in cure of stiffness and fever^[13,14].

It has solid diuretic action and the capacity to break up urinary calculi^[10].

It additionally displays antimalarial, disinfectant, hostile to bacterial properties^[12].

The leaves are utilized as immunomodulatory, adaptogenic, toothpaste, lithotriptic, antiscorbutic, ailagogine and stomach related^[11].

Sphilanthol, the most dynamic germicide alkaloid removed from this plant, is discovered successful at amazingly low focuses against blood parasites, and without a doubt is a toxin to most spineless creatures while staying innocuous to warm-blooded animals^[10].

The blossom heads of *S. acmella* can be bitten to diminish toothache and furthermore as a haemostatic and pain relieving^[15].

S. acmella additionally had incredible enemy of microbial exercises against red halophillicocci from salt relieved fish^[12].

Its concentrate is a functioning part utilized in magnificence care makeup as a quick acting muscle relaxant to quicken fix of useful wrinkles. The plant extricate was additionally utilized for invigorating, revamping and reinforcing the collagen arrange in hostile to age applications, for example in antiwrinkle cream plans^[16].

Sphilantholutilized as bug spray it shows strong ovicidal, larvicidal and pupicidal action^[17].

An Indian clan utilized *S. acmella* to treat contagious skin conditions, for example, competitor's foot, ringworm and nail contaminations^[18].

Table 1:- Traditional utilizations and uses of various pieces of Spilanthesacmella plant:

S. No.	Health Care	Treatment	Plant part Used
1	Medical	Rheumatism, Fever, Diuretics, Flu, hack Rabies, Tuberculosis, Antimalarial, Antibacterial, Throat objections, Headache, Leucorrhoea in female and so on.	Leaves, Flower
		Weight control(lipase inhibitors), scurvy	Flower
		GIT disturbances	Roots
		Antifungal, skin illnesses, Immunomodulatory, Antiscorbutic, Local sedatives, Stomachic etc.	Leaves

		Snake bite	Whole plant
2	Dental	Toothache	Leaves, Flower
		Toothpaste	Leaves
		Periodontal disease	Flower head, Roots
		Intermittent poignancy stomatitis	Leaves
3	Beauty consideration cosmetics	Fast acting muscle relaxant Anti-wrinkle	Whole plant

Phytochemistry:

It is important to investigate the phytochemical constituents of any restorative plant to build up a connection among pharmacology and science of the plant. Numerous investigations have been done for substance examination and auxiliary assurance of sharp alkaloids from *Spilanthes acmella*. The major sharp constituent detailed in this plant *S. acmella* is "spilanthol," which is an isobutylamide and is notable for its insecticidal properties^[19].

The bloom head and root some portion of the plant have been accounted for to be the rich wellspring of dynamic standards. Triterpenoids have likewise been found in the plant. Spilanthol has a solid impactful taste; it might create neighborhood astringency and sedative impacts. *Spilanthes acmella* contains optional metabolites. Spilanthol can be moved in the ethanol remove, which has once been found to contain 9.04% of all out Nalkylamides yet 88.84% spilanthol^[20].

Spilanthol is artificially Nisobutylamide which is unpleasant in taste and could invigorate salivation. The sub-atomic equation of spilanthol was resolved as (2E,6Z,8E)- N-isobutylamide-2,6,8-decatrienamide^[21].

Pharmacological Activity:

Anti-inflammatory and Analgesic:

Fluid concentrate of ariel part of *S. acmella*, in exploratory creature models demonstrated portion subordinate restraint of paw edema and expanded torment limit showing huge calming and pain relieving properties^[22].

Spilanthol shows significant calming action on lipopolysaccharide-enacted murine macrophage model RAW 264.7, mostly from inactivation of NF-KAPPA B which contrarily controls creation of favorable to fiery go between^[23].

Various dosages of fluid concentrate of new blossoms was orally regulated to male rodents and their pain relieving potential was determined at various post treatment periods by utilizing hot plate and tail flick tests. The pain relieving movement is interceded supra-spinally went with sedation^[24].

Antifungal activity:

Various groupings of *S. acmella* blossom head extricate (dried bloom heads removed with oil ether) was assessed for antifungal action (0.1 to 2.0 mg). The measurement of restraint zones went from 0.1 to 2.3 cm with the expansion in grouping of test arrangement. In all the living beings, the most extreme zone of restraint was seen at 2000 mg focus^[25].

Antibacterial movement:

The rough ethyl acetic acid derivation concentrate of *S. acmella* indicated promising antibacterial exercises against 27 strains of microorganisms considered. Among the all portions, the division E3 totally hindered the development of *Corynebacterium diphtheria* with MIC estimation of 128µg/mL^[26].

Anticancer Activity:

S. spirulina was tried for anticancer action (10 µg/mL–5mg/mL) by tumoricidal impacts in an unfading neuroblastoma of unconstrained harmful root. The discoveries showed no example of tumoricidal impacts with anticancer screen classification 5 and consequently are viewed as powerless^[27].

Bioinsecticide and convulsant movement:

Around 42 realized species had been accounted for the class *Spilantes*. *Spilantesalba*, *Spilantesmauritiana*, *Spilantesmauritianaoleracea*, *Spilantesmauritianaocymifolia*, and *S. acmella* were found to comprise of a few insecticidal mixes^[3, 28].

Epilepsy in rodents was prompted by the hexanic concentrate of *S. acmella* when the trademark electrographic seizures in the electroencephalogram were went with^[29].

Diuretic:

The cool water concentrate of bloom of *S. acmella*, demonstrated a checked increment in pee yield, likewise stamped increment in urinary Na⁺ and K⁺ levels and decrease of pee osmolarity proposing that it is predominantly going about as a circle diuretic movement. It might likewise hinder ADH discharge and additionally activity^[30].

Ethanol concentrate of leaves of *S. acmella* likewise essentially expanded the urinary yield (by 223%) and electrolytic discharge of Na⁺ (by 136%) and K⁺ (by 172%)^[31].

Hepatoprotective Activity:

Liver plays an astounding cluster of imperative capacities in the body. No particular allopathic medications utilized as hepatoprotective. Home grown medications are all the more broadly utilized for hepatoprotectives. Hepatoprotective movement of *S. acmella* extricates against CCl₄-instigated liver harmfulness in rodents was resolved^[32, 33].

Directed the examination on antihepatotoxic movement of *Spilantes ciliate*. Pretreatment of the rodents with oral organization of the plant ethanolic separates *Ixoracoccinea*, *Rhinacanthusnasuta*, *Spilantes ciliate*. *Spilantes ciliate* preceding Aflatoxin B1 was found to give noteworthy security against poison instigated liver harm.

Immunomodulatory movement:

The ethanol concentrate of leaves demonstrated critical actuation of macrophages and improved their capacity when contrasted with control, proposing the spice as an expected common medication for immunostimulant impact^[34].

Insecticidal movement:

The rough seed concentrate of methanol, hexane, and deltamethrin of *S. acmella* indicated huge insecticidal movement against *Plutellaxylostella*^[35].

Local sedative movement:

Diverse creature models had been utilized for the estimation of the nearby sedative demonstration of *S. acmella*. Nupercaine was utilized as a norm to create intracutaneous wheals in guinea pigs (suitable to decide the proportion of sedation) and Cocaine was utilized as a norm to deliver plexus sedation in frog (utilized to decide initiation of sedation). The neighborhood sedative activity which was powerful, may maybe ascribe to the nearness of alkylamides in *S. acmella*^[36].

Larvicidal action:

Spilanthol, a significant constituent of ethanolic concentrate of blossom heads of *Spilantesacmella* is having powerful ovidal, larvicidal and pupicidal action. Maximum 7.5 ppm focus caused 100% motility of eggs, hatchlings and pupae of *Anopheles*, *Culex* and *Aedes* mosquito. Spilanthol is more successful even at low portions against eggs and pupae. In pupae, it appears to chip away at sensory system which was obvious by strange development like bastards, turning and clumsy strong movement proposing along these lines that it upsets nerve conduction^[17].^[37] found the ethanolic concentrate of *Spilantesacmella* as one tenth dynamic when contrasted with DDT against *Anopheles* hatchling. The bloom tops and airborne parts have been seen as harmful to mosquito hatchlings and *Periplanata*. The compound Spilanthol has been distinguished as having larvicidal action^[38].

Antioxidant action:

The cancer prevention agent movement of methanol concentrate of stem and leaves of *S. acmellawas* estimated utilizing DPPH and superoxide radical rummaging examines ^[39].

The outcome indicated the methanol concentrate of stem of *S. acmella* to have the most noteworthy superoxide radical rummaging movement while leaves demonstrated greatest DPPH searching action ^[34].

In superoxide radical rummaging examine, most elevated radical searching action was seen in stem and callus, while least superoxide radical rummaging movement was found in roots . In DPPH radical rummaging action was discovered most extreme in leaf and least in root. Callus demonstrated noteworthy DPPH radical searching action ^[39].

Antiobesity Activity:

Fluid ethanol (70%) concentrates of *S. acmella* blossom buds indicated pancreatic lipase inhibitory exercises in a fixation (0.75–2.0mg/mL) subordinate way under in vitro conditions. The extricate likewise restrained lipase, and this plant has potential as a possibility for weight decrease and corpulence control ^[40].

Antipyretic impacts:

S. acmella was assessed by yeast initiated strategies, an overall technique utilized for the enlistment of pyrexia ^[41].

Hindering properties of lipo-oxygenase and cyclo-oxygenase was because of the nearness of flavonoids had been recommended by reports.

Various examinations was extended varying portions of yeast which brought about shifted amounts of flavonoids extricated ^[42].

Aphrodisiac Action (Interaction with Testosterone and Sexuality):

Love potion impact of the plant remove has been concentrated in male rodents by ^[3]. They expressed that mount inactivity, intromission inertness, discharge recurrence, and postejaculatory span were expanded in a portion subordinate way after oral organization of concentrate. Albeit definite measurement of these enhancements was not given, estimation got from diagrams proposed that following 28 days of 150mg/kg portion, the upgrades were diminished in mount idleness, intromission inertness, and post ejaculatory dormancy. These advantages were more noteworthy 28 days after supplementation comparative with 14 days, recommending a development impact. The plant end up being better than Viagra in all angles concentrated aside from proerectile properties.

Pancreatic Lipase Inhibition:

Ethanollic concentrates of the blossoms of *Spilanthesacmella* are shown to hinder pancreatic lipase movement (40% at 2mg/mL fixation in vitro) ^[40].

The action was contrasted and *Aframomummeleguetta* (90% hindrance) and end up being second rate, while 0.75mg/mL remove inhibited more pancreatic lipase than *Spilanthes*.

Antiulcer Activity:

Spilanthes plants are utilized to treat different sorts of ulcers, and studies with *S. filicaulis* watery concentrate demonstrated total mucosal cytoprotection at portions of 500, 1000, and 1500mg/kg, separately, in HCl/EtOH incited gastric sores in male Wistar rodents ^[43].

Vasorelaxant movement:

The chloroform and ethylacetate concentrate of *S. acmella* on phenylephrine applies maximal vasorelaxation in a portion subordinate way, albeit not as much as acetylcholine-incited nitric oxide (NO) vasorelaxtion. Chloroform separate indicated the most elevated vasorelaxation and cancer prevention agent movement ^[44].

Antiviral Activity:

There are just restricted reports with respect to the antiviral action of this class. Leaves of *S. mauritiana* were investigated for antiviral movement with moderate action appeared against HSV, herpes simplex infection, Cox,

Coxsackie B2 infection and noteworthy action against measles, measles edmonston A, polio, poliomyelitis infection type 1 strain 1A/S3,SF, Semliki Forest infection A7 and VSV, and vesicular stomatitis infection T2^[45].

Neuroprotective effect

Neuroprotective effect of *S. acmella* Murr. extracts was resulted from their antioxidant properties. Bioactive compounds presented in these plant extracts were triterpenoids (hexane extract), triterpenoidsglucosides (chloroform extract), phenolic acid and coumarin derivatives (ethyl acetate and methanol extracts). Triterpenoids such as β -sitosterol was reported to exert neuroprotective effect Phenolic compound, tran-ferulic acid and coumarin namely scopoletin displayed neuroprotective activity. The highest neuroprotective effect was noted for the hexane extract of *S. acmella* Murr. It may be due to the nonpolar hexane extract which has better penetration to the cells compared with other polar *S. acmella* Murr. extracts (chloroform, ethyl acetate and methanol). Thus, the neuroprotection was resulted from the decreased oxidative stress and cell death. It remains unclear how the *S. acmella* Murr. extracts ameliorate the neurotoxicity.

Cytotoxic activity:

Cytotoxic activity of the plant extracts was determined by a slightly modified method described previously. Briefly, the confluent cell monolayers were trypsinized and diluted with appropriate culture medium to a final concentration of 3×10^5 cells/mL. Portions (100 μ L) containing approximately 3×10^4 cells were distributed into 96-well flat-bottomed tissue culture plates and incubated overnight at 37°C in a humidified 5% CO₂ incubator. Solutions (100 μ L) containing different concentrations of tested extracts (0.001–10 μ g/mL) or taxol (0.012–1.2 μ g/mL) were added to each well and the plates were incubated as above for an additional 48 h. After the incubation, each well was washed (x 3) with phosphate-buffered saline (pH 7.2) and then stained with Crystal Violet. After the excess dye was removed, the stained cells were lysed with 100 mM HCl (100 μ L) in absolute methanol and the optical density was determined by a microtitre plate reader (Titertek, Multiskan MCC/340) set to read at a wavelength of 540 nm. All tests were carried out in quadruplicate and the mean value was calculated. The activity was expressed as ED₅₀ (the effective dose that inhibits 50% of cell growth).

As marker of osteoblast differentiation cells:

The hexane, ethyl acetate, n-butanol and water fractions from *Graptophyllum pictum* and *S. acmella* were evaluated the stimulative activity on alkaline phosphatase (ALP) of MC3T3-E1 osteoblast cells. ALP activity is a marker of osteoblast differentiation. Among the tested, the n-butanol and water fractions of *G. pictum* showed the activity to 112% and 122% respectively, otherwise the n-butanol and water fractions of *S. acmella* showed the activity to 126% and 127% respectively

Conclusion:-

Spilanthes acmella could be a notable plant in Indian standard arrangement of medication with numerous medical specialty activity and minor reactions during this audit; we tend to finished up ethnobotany, phytochemistry, medical specialty, associate degreed pharmacology in an elucidating manner. It's terribly fascinating that *S. acmella*, ranging from the easy ant toothache plant to extremely valuable annual herb, possesses multifunctional roles as autochthonic medication for medical specialty in health care. The foremost well-endowed isolates of the plant species were lipid alkalamides, especially, the spilanthal together with different bioactive metabolites e.g. phenolic, flavonoid, coumarin and triterpenoid compounds.

S. acmella extracts and its constituting compounds like spilanthal and flavonoids are shown to possess restrictive activity toward PG synthesis. It may be presumptively projected that these compounds share a typical purposeful cluster with electrophilic center, interacting with COX enzymes through nucleophilic addition of aminoalkanoic acid residues. As a result, the syntheses of PG were reserved later on contributive to the discovered antiinflammatory, antipyretic and analgesic activities. Additionally, spilanthal has been shown to cut back NO unharms and thereby inhibit inflammatory mediators and attenuating the expression of Cox-2 and iNOS. This might be attributed to the immunostimulant activity of *S. acmella* in its ancient usages.

S. acmella exerted vasorelaxant and inhibitor activities that is helpful for its lifting impact as quick acting relaxant in opposed wrinkle and opposed aging applications. The participation of NO in vasorelaxation makes *S. acmella* robust aphrodisiac in ancient medication for rising sexual performance in men.

In brief, it may be incontestable that *S. acmella* could be a healthful plant enriched with compounds having high therapeutic price that may be additional developed for applications in medicines, health care, cosmetics, supplements and food.

Acknowledgement:-

The authors are grateful to Vice chancellor M.P. Pandey and Dr. Sushilkumar Director of School of Pharmaceutical sciences IFTM University for providing constant encouragement, valuable guidance and facilities at all stages of this work.

Conflicts of interest:

There are no conflicts of interest.

References:-

1. Sangeeta S, Veena D, Shagufta K and Soni UN, (2016). Phyto-Chemical Investigations from Leaves and Flowers of *Spilanthes acmella*. *Ijsrm.Human*, 4 (1), 70-76.
2. Jirovetz, Leopold, Gerhard B, Andrea W, Mohamed P S, George T (2005). Essential oil analysis of *Spilanthes acmella* Murr. Fresh plants from. *Journal of Essential Oil Research*, 17, 429-431.
3. Sharma U and Pegu S, s(2011). "Ethnobotany of religious and supernatural beliefs of the Mising tribes of Assam with special reference to the 'Dobur Uie. *Journal of Ethnobiology and Ethnomedicine*, 7 (16), 1-13.
4. Chung K, Kono Y, Wang C and Peng C, (2008). Notes on *acmella* (Asteraceae): *helianthaceae* in Taiwan. *Botanical Studies*, 49 (1), 73-82.
5. Revathi P and Parimelazhghan T (2010). Traditional knowledge on medicinal plants used by Irula tribe of Hasanur hills, Erode. *Ethnobotanical Leaflets*, 14, 136-160.
6. Vijendra N and Kumar K, (2010). Traditional knowledge on ethnomedicinal uses prevailing in tribal pockets of Chhindwara and Betul Districts. *African Journal of Pharmacy and Pharmacology*, 4 (9), 662-670.
7. Hossan M, Hanif A and Agarwala B, (2010). Traditional use of medicinal plants in Bangladesh to treat urinary tract infections and sexually transmitted diseases. *Ethnobotany Research and Applications*, 8, 61-74.
8. Santesson C, (1926). Several drugs of the Cameroon District and their native uses. *Archiv Furdie Botanik A*, 20 (3), 1-34.
9. Storey C and Salem J, (1997). Lay use of Amazonian plants for the treatment of tuberculosis. *Acta Amazonica*, 27 (3), 175-182.
10. Yadav K, Singh N. Micropropagation of *Spilanthesacmella*Murr. – An Important Medicinal Plant. *Nature and Science* 2010; 8(9): 5-11.
11. Savadi RV, yadav R, yadav N. Study on immunomodulatory activity of ethanolic extract *Spilanthesacmella*Murr. Leaves. *Indian journal of natural Products and Resources* 2010; 1(2): 204-207.
12. Haw AB, Keng CL. Micropropagation of *Spilanthesacmella*L., a bio-insecticide plant, through proliferation of multiple shoots. *J. Appl. Hort.* 2003; 5(2): 65-68.
13. Bunyaphatsara N, Chocheareunporn O. Tradition medicinal plants. Prachachon: Bangkok. 1999.
14. Farnsworth NR, Bunyaphatsara N. Thai medicinal plants recommended for primary healthcare system; Prachachon: Bangkok. 1992.
15. Leng TC, Ping NS, Lim BP, Keng CL. Detection of bioactive compounds from *Spilanthesacmella* (L.) plants and its various in vitro culture products. *Journal of Medicinal Plants Research* 2011; 5(3): 371-378.
16. Schubnel L. A different approach to lifting efficacy based on a natural active ingredient. *SOFW J.* 2007; 133: 34-39.
17. Saraf DK, Dixit VK. *Spilanthesacmella*Murr. Study on its Extract *Spilanthol* as Larvicidal Compound, *Asian J. Exp. Sci* 2002; 16 (1-2): 9-19.
18. http://www.avresources.my.com/contact_us.htm.
19. Kadir H. A., Zakaria M. B., Kechil A.A. and Azirum M. S. (1989): Toxicity and electrophysiological effects of *Spilanthesacmella*(Murr.). *Pestic. Sci.* 25,329-335.
20. D. K. Mukharya and A. H. Ansari, "Olean-12-en-3-O-beta- D-galactopyranosyl (1-4)-O-alpha-L-rhamnopyranoside: a new triterpenoidalsaponin from the roots of *Spilanthesacmella*(Murr.)," *Indian Journal of Chemical Biology*, pp. 26-86, 1987.
21. Yasuda, K. Takeya, and H. Itokawa, "The geometric structure of *spilanthol*," *Chemical & Pharmaceutical Bulletin*, vol. 28, no. 7, pp. 2251-2253, 1980.

22. Chakraborty A, Devi BRK, Sanjebam R, Khumbong S, Thokchom IS. Priliminary studies on local anesthetic and antipyretic activities of *SpilanthescmellaMurr.* in experimental animals models. *Indian Journal of Pharmacology* 2010; 42(5): 277-279.
23. Wu LC, Fan NC, Lin MH, Chu IR, Huang SJ, Hu CY, Han SY. Antiinflammatory effect of spilanthol from *Spilanthescmella* on murine macrophage by down-regulating LPS-induced inflammatory mediators. *J Agric Food Chem.* 2008; 9:56(7):2341-2349.
24. Peiris KPP, Silva GKJ, Ratnasooriya WD. Analgesic activity of water extract of *Spilanthescmella* flowers on rats. *Journal of Tropical Medicinal Plants* 2002; 2(2): 120-126.
25. Rani SA, Murty SU. Antifungal potential of flower head extract of *Spilanthescmella*. *African Journal of Biomedical Research* 2006; 9: 67-69.
26. Sahu J, Jain K, Jain B, Sahu RK. A review on phytopharmacology and micro propagation of *Spilanthescmella*. *Pharmacologyonlinenewslett* 2011;2:1105-10.
27. Paulraj Jayaraj, Govindarajan Raghavan and Palpu Pushpangadan The Genus *Spilanthes* Ethnopharmacology, Phytochemistry, and Pharmacological Properties: A Reviess [Journal] / ed. Munoz Eduardo. - India : Hindawi Publishing Corporation, September 23, 2013. - Vol. 2013. - p. 22
28. Shelton AM, Wyman JA, Cushing NL, Apfelbeck K, Dennehy TJ, Mahr SE, et al. Insecticide resistance of diamondback moth, *Plutellaxylostella* (Lepidoptera: Plutellidae), in North America. *J Econ Entomol* 1993;86:11-9.
29. Fabry W, Okemo P, Ansorg R. Fungistatic and fungicidal activity of east African medicinal plants. *Mycoses* 1996;39:67-70.
30. Ratnasooriya WD, Pieris KP, Samaratunga U, Jayakody JR. Diuretic activity of *Spilanthescmella* flowers in rats. *J Ethnopharmacol.* 2004; 91(2-3):317-320.
31. Yadav R, Kharya MD, Yadav N, Savadi R. Diuretic activity of *SpilanthescmellaMurr.* Leaves extract in rats. *IJRPC* 2011; 1(1): 57-61.
32. Ganga Rao, Y.VenkateswaraRao and T. MallikarjunaRao. Hepatoprotective activity of *Spillanthescmella* Extracts against CCl4-inducedliver toxicity in rats. *B. MedicalDevices.* 2012; 5-7
33. Suja S R, Rajasekharan S, Pushpangadan P. Tropical Botanic Garden and Research Institute. 2003; 41:536-541.
34. www.stuartxchange.org/Biri.html.
35. Anuradha Sharma, Vishal Kumar, Rameshwar Singh Rattan, Neeraj Kumar, Bikram Singh , Insecticidal Toxicity of Spilanthol from *SpilanthescmellaMurr.* against *Plutellaxylostella* L. *American Journal of Plant Sciences*, 2012, 3, 1568-1572
36. Martin R, Becker H. Spilanthol-related amides from *Acmella* ciliate. *Phytochemistry* 1984;23:1781-83.
37. Pendse G. S., Gokhale V. G., Phalnikar N. L. and Bhide B. V. (1946): Investigation on new plant larvicides with special references to *Spilanthescmella*. *J. Univ. Bombay Sect. A.*, 15, Pt. 3, Sci. 20, 26-30.
38. Kadir H. A., Zakaria M. B., Kechil A.A. and Azirum M. S. (1989): Toxicity and electrophysiological effects of *Spilanthescmella(Murr.)*. *Pestic. Sci.* 25,329-335.
39. Tanwer BS, Choudhary RK, Vijayvergia R. In vitro and in vivo Comparative Study of Primary Metabolites and Antioxidant Activity in *SpilanthescmellaMurr.* *International Journal of Biotechnology and Biochemistry* 2010; 6 (5): 819–825
40. Tanwer BS, Choudhary RK, Vijayvergia R. In vitro and in vivo Comparative Study of Primary Metabolites and Antioxidant Activity in *SpilanthescmellaMurr.* *International Journal of Biotechnology and Biochemistry* 2010; 6 (5): 819–825
41. P. Ekanem, M. Wang, J. E. Simon, and D. A. Moreno, “Antiobesity properties of two African plants (*Afromommumleguetta* and *Spilanthescmella*) by pancreatic lipase inhibition,” *Phytotherapy Research*, vol. 21, no. 12, pp. 1253–1255, 2007.
42. Martin R, Becker H. Amides and Other constituents from *Acmella* ciliate. *Phytochemistry* 1985;24:2295-300.
43. Broussalis AM, Ferraro GE, Martino VS, Pin-Zon R, Coussio JD, Alvarez JC. Argentine plants as potential source of insecticidal compounds. *J Ethnopharmacol* 1999;67:219-23.
44. P. V. Tan, C. K. Njimi, and J. F. Ayafor, “Screening of some African medicinal plants for antiulcerogenic activity,” *PhytotherapyResearch*, vol. 11, no. 1, pp. 45–47, 1997.
45. Wongsawatkul O, Prachayasittikul S, Ayudhya CI, Satayavivad J, Ruchirawat S, Prachayasittikul V. Vasorelaxant and Antioxidant Activities of *SpilanthescmellaMurr.* *Int J Mol Sci.* 2008; 9(12): 2724–2744.
46. P. Cos, N. Hermans, T. De Bruyne et al., “Further evaluation of Rwandan medicinal plant extracts for their antimicrobial and antiviral activities,” *Journal of Ethnopharmacology*, vol. 79, no. 2, pp. 155–163, 2002.
47. Wilasinee S, Bongkot K, Sujitra S, Supaluk P, VirapongP. Neuroprotective effect of *SpilanthescmellaMurr.* on pesticide-induced neuronal cells death. *Asian Pacific Journal of Tropical Medicine* 2017 ; 10 (1): 35-41

48. Supaluk P, SaowapaS, Apilak W, Ratana L, Somsak R and Virapong P. Bioactive Metabolites from *Spilanthescmella*Murr.first_pagesettings. Bioactive Metabolites from *Spilanthescmella*Murr. *Molecules* 2009, 14(2), 850-867.
49. Retnowidyowati. Alkaline phosphatase activity of *graptophyllumpictum* and *sphilanthescmella* fractions against mc3t3e1 cells as marker of osteoblast differentiation cells. *Int J Pharm PharmSci*, 2011; Vol 3, Suppl 1, 3437.