



ISSN NO. 2320-5407

Journal homepage: <http://www.journalijar.com>

INTERNATIONAL JOURNAL
OF ADVANCED RESEARCH

RESEARCH ARTICLE

STUDY OF EFFECT OF pH ON ENZYME ACTIVITY OF PROTEASE PRODUCED BY
BACTERIAL ISOLATES COLLECTED FROM SOIL SAMPLES OF LACHIWALA, DOON VALLEY

CHARU LATA, Prof. D.S.NEGI, Prof.VINIT MEHROTRA

Manuscript Info**Manuscript History:**

Received: 26 August 2014
Final Accepted: 25 September 2014
Published Online: October 2014

Key words:

Protease, bacterial pure cultures,
spectrophotometrically, azocasein

***Corresponding Author**

CHARU LATA

Abstract

Proteases are typical enzymes that are being investigated for their potential usage in many areas of application, such as detergent, brewing meat, photography, leather and dairy. A large number of proteases have been purified and characterized from bacteria and few from fungi. Bacteria were isolated from the soil sample of Lachiwala, Doon Valley. Out of 64 bacterial pure cultures, 10 best protease producing microorganisms were isolated through activity zone methods. Protease activity was checked spectrophotometrically using azocasein dye as a substrate. 1 bacteria was selected from them. The maximum enzyme activity was obtained at near neutral pH (6.5).

Copy Right, IJAR, 2014,. All rights reserved

Introduction

Proteases (E.C.3.4) are the single class of enzymes which occupy a pivotal position with respect to their applications in both physiological and commercial fields. Proteases are hydrolytic enzymes that hydrolyze protein by adding water across peptide bonds and break them in smaller peptides in organic solvents (1). Proteases represent one of the three largest groups of industrial enzymes and account for about 60% of the total worldwide sale of enzymes (18) with 2/3 of this amount, being of microbial origin (5). Proteases execute a large variety of functions, extending from the cellular level to the organ and organism level, to produce cascade systems such as homeostasis and inflammation (18). Proteases play an important role in a wide range of industrial process viz., baking, brewing, detergents, leather processing, pharmaceuticals, meat tenderization, cosmetics, peptide synthesis, waste treatment and medical diagnosis (7).

MATERIAL AND METHODS**Study area**

The vegetation at Lachiwala is a mix of moist deciduous and Sal trees. It contains the marshes and grassland along with the Song River. The point coordinates of Lachiwala is E 78° 06' 38.6" and N 30° 13' 37.2" with altitude of 1683 feet above the sea level. The site of Lachiwala with species of Sal, Syzygium, rich in plant litter and content of organic carbon (oc) is low.

The soil surface is compact. The texture of soil is sandy loam. Moisture on the surface is moderate while at sub-surface it is moist. (15)

Sampling

Sediment samples were collected by sterile spatula or by a corer from 6 different locations of Lachiwala area of Doon Valley. For this survey the sampling was done in 3 times a year-in the month of January (S_A), May (S_B) and September (S_C). The collected samples were brought to laboratory under refrigeration and were preserved at -20°C

Sample Preparation and spreading

Samples (S_A, S_B, S_C) were taken out of -20°C freezer and kept at 4°C for 1 h to 2 hrs. Then they were kept at RT for 12hrs to avoid the heat shock. The samples were prepared by serial dilution and spread plate methods. 200uL of each

dilution was spread into the nutrient agar plates (Beef extract 3.0g., Peptone 5.0g., NaCl 5.0g, Glucose 10.0g., Yeast extract 5.0g., Agar agar 20.0g., SDW 1000ml., pH 7.2-7.4) (20).

Primary stock cultures preparation

After 7 days all the plates were taken out and they were then picked and streaked again in the nutrient agar plates with sterile tooth picks. The plates were kept at 30°C in the incubator for 24 h. The bacterial cultures were then inoculated into nutrient broth (beef extract 3.0g, peptone 5.0 g, NaCl 5.0 g, glucose 10 g, yeast extract 5 g, SDW q.s. 1000 ml, pH 7.2 – 7.4) and placed in the orbital shaker at 30°C and 120 r.p.m for 24 h. The cultures were restreaked onto the nutrient agar plates and the process was repeated until pure bacterial cultures were obtained.

Preservation of pure cultures

The selected microorganisms were streaked onto solid nutrient agar medium (1 organism in 1 plates) in order to make final and pure working cultures and at the same time glycerol stock were prepared for the final selected bacteria for long term preservation and preserved in -70°C freezer. The entire procedure was repeated until a certain number of pure bacterial cultures (64) were found to carry on the further experiments.

Screening of microorganisms for proteolytic activity

The selection of most potent isolates were performed by two steps:

Activity zone technique according to Sonia Sethi et al., 2012(16) and assay of enzymes according to Kole et al., 1988(11).

Activity zone technique

Determination of the most potent isolates was performed by clear cut zone technique. 64 bacterial colonies (from each sample s_a, s_b, s_c) were screened for proteolytic activity in Petri dishes with production medium PM1 which contains: Beef extract 3.0 g, Peptone 5.0 g, NaCl 5.0 g, Glucose 10 g, Yeast extract 5.0 g, Gelatin 4.0g, Casein 10.0 g, Agar agar 20.0 g, SDW 1000 ml, pH 7.2 – 7.4

The plates were incubated at room temperature for 24 hour. At the end of incubation period, the remainder supernatant was removed and plates were flooded with 1% tannic acid (12).

The isolates were evaluated for their proteolytic activity by measuring a clear zone of gelatin hydrolysis formed on milky agar. A total of 10 isolates were selected.

The activity was then detected by the appearance of clear-cut clearing zones around the bacterial growth indicated the ability of the isolate to hydrolyze gelatin due to the biosynthesis of the protease enzyme. The resulting diameters of the clearing zones were recorded.

Enzyme assay

Protease activity in microbial culture is estimated by different ways. Estimation of Protease activity by Spectrophotometer is the cheap and best method.

Spectrophotometric assay of protease could be done in many ways. But azoprotein substrates would be the best choice for specific as accurate enzymatic reaction (6).

Azocasein is a chemically modified protein containing sulphanilamide group covalently linked to peptide bond of casein. During incubation for 30 min. proteases hydrolyse peptide bond, liberating shorter peptide and amino acid from the chain, which is orange in colour. The intensity of colour, is measured spectrophotometrically to determine protease activity.

One unit of enzyme activity was defined as the amount of enzyme required to solubilize 1 µg. of substrate/min under assay condition.

Culture preparation: Microorganisms were collected from respective working culture plates and inoculated into the production medium. The fermentation was carried out at 30°C with orbital shaking at 120 rpm for 48 h in 100 ml. Erlenmeyer flasks containing 20 ml Lauria- Bertani (LB) medium. It contains: Yeast extract 5.0 g, Peptone 10.0 g, NaCl 10.0 g, SDW q.s. 1000 ml, pH 7.2 – 7.4.

Reagents and Instruments

1. 0.1 M Tris- HCl (pH 8.5)

Preparation: 0.1 M of Tris- HCl solution was prepared in water and the desired pH was adjusted as required with 0.1 M Hydrochloric acid (HCl).

2. Azocasein (2500 µg.ml⁻¹ solution): Substrate

Preparation: In accurately weighed azocasein half if the required amount of water was mixed in a container and it was dissolved by very slow and gentle shaking. After a part of azocasein was dissolved it was kept in refrigerator at 4°C for 30 min. Next the solution was taken out and rest half of water was mixed and remaining undissolved azocasein was dissolved with slow shaking of the container and kept at 4°C for few min before use. It should be freshly prepared before use and should not be shaken vigorously during preparation, which might lead to degradation of azocasein.

3. 10% tri- chloro acetic acid (TCA) solution

Preparation: 10.0 g of TCA was dissolved in distilled water and kept at 4°C before use.

4. Spectrophotometer

5. Centrifuge

Procedure: Reaction mixture was prepared with 100 μL culture supernatant (1.5ml culture centrifuged at 10×10^3 r.c.f. for 5 min at 4°C to separate supernatant and cells), 125 μL azocasein solution, 50 μL 0.1 M Tris-HCl and 225 μL distilled water. The entire mixture was incubated for 30 min at 37°C and then the assay was terminated by 500 μL ice cold 10% TCA solution (18). TCA was precipitated by centrifugation at 10×10^3 r.c.f. for 5 min at 4°C . Finally the O.D. was measured with required dilution at λ_{440} .

Control: Media filtrate without the substrate was taken as blank. Volume was made up by adding 0.1 M Tris-HCl buffer only.

Precaution: Azocasein being a very delicate and light substrate should be shaken gently.

Study of enzyme activity -Effect of pH

Any enzyme has an optimum pH requirement for its activity. Like time course study here also two types of fermentation media were set. A study has been carried out in different pH from 6.0 to 10.0 in order to find the activity profile of the enzyme at various pH. The crude sample (microbial culture supernatant) was assayed for the enzyme activity spectrophotometrically in the same procedure as mentioned earlier. Only the pH of the reaction mixture was altered. 2 the same of buffer were taken for this study, viz, phosphate buffer (for pH 6.0,6.5,7.0,7.5,and 8.0)and Tris-HCL buffer(for pH 8.5,9.0,9.5,10.0). The activity found in various pH were tabulated graphically.

RESULTS

Results of effect of pH on enzyme activity by LW-1, LW-2 and LW-3




Fig. No. 01: Effect of pH on enzyme activity by microorganism LW-1




Fig. No. 02: Effect of pH on enzyme activity by microorganism LW-2




Fig. No. 03: Effect of pH on enzyme activity by microorganism LW-3

Effect of pH on enzyme production was studied. Fig.01 predicts that when the culture supernatant was collected from the fermentation medium Enzyme produced in the fermentation medium showed that enzyme stability of LW-1 was found to be maximum at pH 6.5 (5.40 U./ml.). Then again with the increase of pH enzyme activity started decreasing up to pH 8 (3.6 U./ml.) on further increasing the pH it also gave a better activity profile as it also gave two optimal pH; 8.5 (4.80 U./ml) and 9.5 (5.29 U./ml). So two sharp peaks were observed, one at pH 6.5 and another at pH 9.5. So the protease is highly active in neutral as well as alkaline pH.

Fig.No.02 depicts that enzyme stability of LW-2 was maximum at pH 6.5 (5.40 U. /ml.). Then again enzyme stability declined and found to be minimum at pH 8.0 (1.75 U. /ml.). Then from pH 8.0 a sharp rise in stability was noticed

and after that LW-2 protease maintained a constant stability. Protease from LW-2 gave 2 sharp peaks, one at pH 6.5 and another at pH 9.5. So the protease was active in neutral as well as alkaline pH.

Fig.No.03 shows that the enzyme stability of LW-3 was found to be maximum at pH 6.5 (5.84 U./ml.) and at 7.0 pH enzyme activity was found to be 2.3 U./ml. Then again with increase of pH enzyme activity started increasing and 2 more peaks were noticed at pH 8.5 (5.09 U./ml.) and 9.5 (4.75 U./ml.). Protease from LW-3 gave 2 sharp peaks, one at pH 6.5 and another at pH 8.5. So the protease was active in neutral as well as alkaline pH.

DISCUSSION

The effect of pH on the growth and protease production by soil bacteria was studied. The maximum production of protease(s) was recorded in a medium at pH 6.5 with maximum yield of 5.40 U./ml., 5.40 U./ml. and 5.84 U./ml. for LW-1, LW-2 and LW-3 respectively. However, the maximum yield of 5.84 U./ml. for LW-3 at pH 6.5 was observed. At pH values above the previously recorded optimum values pH for the tested three strains, the relative production of protease(s) was markedly stable. It is quite obvious that the maximal productivity of protease(s) for the tested three strains could be recorded within the neutral side of pH. It had been reported previously that the highest protease production was determined within this optimum pH range (17). Sumantha et al. (2006) reported that the optimum pH ranges of proteases were 5-7 (21). On the other hand other optimum pH ranges were reported to be suitable for maximum protease production including 6.6 (13), 6.8 (8), 7 (16), 8 (4), 8.5 (19), 10.5 (3), 11 (10) and 12 (14).

CONCLUSION

The enzyme was stable over a broad pH range of 6.5 to 9.5 indicating that it is neutral protease enzyme (8). The results of pH and temperature stability of the investigated neutral protease suggesting that it may have potential application in detergent pharmaceutical formulations, leather, laundry, food (18) and waste processing industries (8). Johnvesly and Naik (2001) reported that the maximum enzyme activity between pH 6.5 and 12.0 indicating its potential for practical use in industrial purpose which require activity over wide pH range (9).

REFERENCES

1. Al-Shehri, M., A.; (2004), Production and some properties of protease produced by *Bacillus licheniformis* isolated from Tihamet Aseer, Saudi Arabia. *Pakistan J. Biol. Sci.* 7:1631-1635.
2. Ammar, M., S., Elwan, S., H. and Meki, M., A.; (1986), Thermostable ureate oxidase from obligate thermophilic bacteria I-Estimation using the uric acid clearing zone (U.C.Z.). *New technique Microbiol. Espaniola*, 39 (1-4), 3.
3. Banik, R., M. and Prakash, M.; (2004), Laundry detergent compatibility of the alkaline protease from *Bacillus cereus*. *Microbiol. Res.*, 159 (2): 135-140.
4. Beg, Q., K. and Gupta, R.; (2003), Purification and characterization of an oxidation-stable, thiol-dependent serine alkaline protease from *Bacillus mojavensis*. *Enz. Microbial Techn.*, 32: 294-304.
5. Cowan, D., A.; (1994), Industrial Enzymes, In *Biotechnology-The science and the business*, edited by V. Moses and R.E. Cape, (Harwood Academic Publishers, Switzerland), 326-328.
6. Ghosh, D.; (2004), Bioprospecting in the deltaic Sundarbans for marine microorganism producing commercially important bioactive compound; PhD thesis; Jadavpur University. Kolkata, India.
7. Gupta, R., Beg, Q., K., Lorenz, P.; (2002), Bacterial alkaline proteases: molecular approaches and industrial applications. *Appl. Microbiol. and Biotechnol.* 59 (1): 15-32.
8. Hazem, A., Farouk, Al-Quadani, Tahani, K., Yousef; (2012), A novel neutral protease from thermophilic *Bacillus* strain HUTBS62, *J. BioSci. Biotech.*, 1(2): 117-123.
9. Johnvesly, B. and Naik, G., R.; (2001), Studies on production of thermostable alkaline protease from thermophilic and alkaliphilic *Bacillus* sp. JB-99 in a chemically defined medium. *Proc. Biochem.*, 37: 139-144.
10. Kobayashi, T., Hakamada, Y., Hitomi, J.; (1996), Purification of alkaline proteases from *Bacillus* strain and their possible interrelationship, *Appl. Microbiol. Biotechnol.*, 45: 63-71.
11. Kole, M., M., Draper, I., Gerson, D., F.; (1988), Production of protease by *Bacillus subtilis* using simultaneous control of glucose and ammonium concentrations, *J. Chem. Tech. Biotechnol.*, 41: 197-206.
12. Kumar, A., Sachdev, A., Balasubramanyam, S., D., Saxena, A., K., Lata; (2002), Optimization of conditions for production of neutral and alkaline protease from species of *Bacillus* and *Pseudomonas*; *Ind. J. of Microbiol.*, 42: 233.
13. Louise tran, xu-chuwu, and sui-lam wong (1991): cloning and expression of a novel protease gene encoding an Extracellular Neutral Protease from *Bacillus Subtilis*, *J. of Bacteriol.*, 6364-6372.
14. Miyaji, T., Ota, Y., Shibata, T., Mitsui, K., Nakagawa, T., Watanabe, T., Nimura, Y. and Tomizuka, N. (2005): Purification and characterization of extracellular alkaline serine protease from *Stenotrophomonas maltophilia* strain. *Letters in Appl. Microbiol.*, 41, 253-257.

15. Mukesh, Manhas R K , Tripathi A K Raina A K , Gupta M K, Kamboj S K :Sand and clay mineralogy of sal forest soils of the Doonsiwalik Himalayas.
16. Palsaniya, P., Mishra, R., Beejawat, N., Sethi, S., Gupta, B., L.; (2012), Optimization of alkaline protease production from Bacteria isolated from soil, *J. Microbiol. Biotech. Res.*, 2 (6): 858-865.
17. Rahman, R., N., Z. , Razak, C., N. , Ampon, K. , Basri, M. , Yunus, W. , M., Z. , W. et al (1994), Purification and characterization of heat stable alkaline protease from *Bacillus stearothermophilus* F1, *Appl. Microbiol. Biotechnol.* , 40 , 822-827.
18. Rao, M., B., Tanksale, A., M., Ghatge, M., S., and Deshpande, V., V.; (1998), Molecular and Biotechnological Aspects of Microbial Proteases, *Microbiol. Mol. Biol. Rev.*, 62:597-635.
19. Sanchez-Porro, C., Mellado, E., Bertoldo, C., Antranikian, G., Ventosa, A.; (2003), Screening and characterization of the protease CP1 produced by the moderately halophilic bacterium *Pseudoalteromonas* sp. strain CP76. *Extremophiles*, 7(3): 221-228.
20. Sevinc, N. and Demirkan, E.; (2011), Production of Protease by *Bacillus* sp. N-40 Isolated from Soil and Its Enzymatic Properties. *J. Biol. Environ. Sci.*, 5(14), 95-103.
21. Sumantha, A., Larroche, C., Pandey, A.; (2006), Microbiology and industrial biotechnology of food-grade proteases: A perspective. *Food Technol. Biotechnol.*, 44(2): 211-220.