



ISSN NO. 2320-5407

Journal homepage: <http://www.journalijar.com>

INTERNATIONAL JOURNAL
OF ADVANCED RESEARCH

RESEARCH ARTICLE

Phytogeography of lesser known *Mimusops andamanensis* King & Gamble (Sapotaceae) with special reference to its occurrence in Little Andaman Island

Sam P. Mathew, C. K. Biju & H. Biju

Jawaharlal Nehru Tropical Botanic Garden and Research Institute, Thiruvananthapuram 695 562, Kerala, India.

Manuscript Info**Manuscript History:**

Received: 14 April 2015
Final Accepted: 29 May 2015
Published Online: June 2015

Key words: *Mimusops andamanensis*, natural history, Andaman-Nicobar Islands

***Corresponding Author**

Sam P. Mathew

Abstract

The Andaman - Nicobar Archipelago in the Bay of Bengal exhibits multi dimensional floristic affinities towards major landmasses of South and Southeast Asia such as Northeast India, Myanmar, Thailand, Malaysia, Indonesia, Philippines, Peninsular India (Western Ghats) and Sri Lanka. From geological point of view, the Andaman - Nicobar Islands are considered as the emergent peaks of a submerged mountain range in continuation with the Arakan Yoma Mountains of the Myanmar towards the Moluccas Island of the Indonesia. *Mimusops andamanensis* King & Gamble, is a critically endangered insular botanical entity with disjunct distribution in Sri Lanka and Andaman-Nicobar Islands. The natural history of *Mimusops andamanensis* King & Gamble is described here with illustrations, based on the rediscovery of this taxon from Andaman Islands

Copy Right, IJAR, 2015, All rights reserved

INTRODUCTION

The terminology “endangerment” currently referred to biological species is obviously becoming an uncritical cliché in many contexts owing to various causes. Endangerment and extinction of biological species in nature has become a matter of curiosity since Darwin’s era and regarded as a natural phenomenon of evolution in regular course of time as denoted by Darwin, “.....*Natural selection almost inevitably causes much extinction of the less improved forms of life and induces what I have called divergence of character.*” (Darwin, 1872). Although, Darwin’s concept on rarity is frequently quoted since then, extinction of species has become a global crisis in due course of time owing to causes other than evolutionary progression. Disjunct geographical distribution or fragmentation of range in distribution of biological entities is recognized as one of the natural motivations for rarity and endangerment. According to Drury (1974) ‘A rare species is one that occurs in widely separated small sub-populations so that, interbreeding between in the sub-populations is seriously affected or reduced or restricted to a single population’. This definition seemingly very prudent on account of several insular species, since they are characterized by small gene pools, geographical isolation and least possibility for out breeding.

Mimusops andamanensis King & Gamble is one of the endangered insular taxa enlisted in the world list of threatened trees (Oldfield *et al.*, 1998) as well as in red list of threatened plants by IUCN and Ministry of Environment and Natural Resources (MENR) (Anonymous, 2007). Interestingly, the phytogeography of the taxon demonstrates disjunct distribution from Sri Lanka towards Andaman-Nicobar Islands, although these regions are widely separated by the Indian Ocean. The Andaman-Nicobar Archipelago located around 700 nautical miles away from Sri Lanka in the Bay of Bengal is the abode of around 2463 Angiosperm species within a limited geographical region of 8249 sq km, while Sri Lanka is apparently larger than the former cover an area of 65610 sq km and is known to host about 4143 Angiosperm taxa (Senarantna, 2001).

Mimusops andamanensis King & Gamble is rather a lesser known species on account of Andaman flora and Interestingly, this taxon is treated as endemic to Andaman Islands among several literature on Andaman-Nicobar floristics (Balakrishnan and Rao, 1983; Pandey & Diwakar, 2008). This species is originally described by King and Gamble in 1906 based on a specimen procured from Port Mouat Hill jungle of South Andaman Island by one of the King's collectors dated as 16/04/92. The type specimens have been deposited at herbaria of Botanical Survey of India, Kolkata (acc. no. CAL 0000017649) and the Royal Botanic Garden, Kew (acc. no. K 000777706).

Unlike Andaman-Nicobar Islands, there are detailed investigation reports on rare and endangered plant species of Sri Lanka. According to an official estimation by IUCN and MENR of Sri Lanka on conservation status of 1099 indigenous Angiosperm species belonging to 68 families, recorded as 675 species including 412 endemics have been assessed to be threatened. There are 72 species including 43 endemics have already been extinct from the island. This document also enumerated 250 species including *Mimusops andamanensis* King & Gamble, under critically endangered category and another 188 insular species as endangered (Anonymous, 2007). The National Red List 2012 of Sri Lanka published by the Biodiversity Secretariat of the Ministry of Environment in collaboration with the National Herbarium also included *Mimusops andamanensis* King & Gamble under endangered category as per the IUCN criteria B1(i,ii,iii) and 2ab (i,ii,iii). Surprisingly, it is found that no detailed investigations on rarity and endangerment of the insular floristic components of the Andaman-Nicobar Islands have so far been done except a preliminary investigation carried out by Balakrishnan and Rao (1983). Balakrishnan and Rao enumerated about 110 endangered endemic Angiosperm taxa from the Andaman-Nicobar Islands; of which 73 species are known only from their type collection and the remaining 37 species reported only from their type locality. According to Balakrishnan and Rao (1983) *Mimusops andamanensis* King & Gamble is a critically endangered 'endemic species' known only from its type collection deposited at CAL and K. A perusal of literature also confirmed that this taxon has not been reported from Andaman-Nicobar islands after type collection and no further indication has been found about its habitat or ecological niches in Andaman rainforests except the reference given in the protologue. The occurrence of this species in Sri Lanka is reported from dry zone of lower altitude. As regards to Andaman-Nicobar Islands the accurate status quo of several threatened species are lesser known or quite little known. Extinction of a species is the condition where the last living specimen of the species dies. Nevertheless, it may be difficult to determine precisely when it did occur in large diverse and complex rainforests found to occur among Andaman-Nicobar Islands. In general, it is better to be presumed that an insular taxon may be extinct, if it failed to locate for an uninterrupted period of 50 years or more and *Mimusops andamanensis* King & Gamble was thought to be extinct from the Andaman-Nicobar Islands until the recent discovery from Little Andaman Island.

During the plant explorations carried out in 2014 among the Andaman-Nicobar Islands, the authors had relocated this taxon from the lowland evergreen forests of the Little Andaman Island. There are three trees found growing apart within an area of one sq km in the reserve forest at southern part of the island. All the trees were under fruiting and bearing ripened fruits during the month of February. Surprisingly, no seedlings of this taxon have been found at the vicinities of these trees, which indicate its natural regeneration has been curtailed owing to certain reasons, so far unknown. The Little Andaman Island constitutes the southernmost island in the Andaman group of islands cover an area of 734.39 sq km, partially denuded the natural forest coverage and substituted with manmade vegetation of oil palm plantation; nevertheless, around 70005 hectares of virgin forest coverage is protected under reserve forest category in this island. Another exploration in Great Nicobar Island during the month of February, 2014 by a group of scientists from the Botanical Survey of India also had located *Mimusops andamanensis* in fruiting stage and collected mature fruits. It indicates relics of this species are extending up to the southernmost island of the Andaman-Nicobar Archipelago.

According to Bustard (1971) "a species is endangered when its numbers are consistently being depleted more rapidly than they can reproduce themselves, absolute number may not be relevant to its threatened status, since large populations can be reduced catastrophically by adverse factors". Range in geographical distribution of many insular taxa in Andaman-Nicobar Archipelago becoming shrunken owing to isolation, catastrophic events, alterations in special ecological micro-niches, barriers in natural regeneration and anthropogenic interventions. Therefore they are presently confined their occurrence to a few isolated fragments of their former range and becoming more vulnerable on the road of extinction. Interestingly, a few individuals of *Mimusops andamanensis* King & Gamble, a taxon thought to be extinct from the insular habitats of the Andaman-Nicobar Islands are still surveying in small population as relics of its glorious past, invites great attention for its long term conservation measures.

Being an interesting rediscovery, a taxonomic description of the species is given below.

Mimusops andamanensis King & Gamble, J. Asiat. Soc. Bengal 74: 201. 1906; Rev. Hbk. Fl. Ceylon 9: 359.1995.

Type: S. Andaman Is., near Port Mouat, *King's Collector s.n.* (CAL, holotype; K, isotype).

Evergreen tree c. 7m tall. Branchlets adpressed puberulous. Leaves 5-10 x 4-7 cm, elliptic to elliptic-obovate or round ovate, cuniate at base, acute to shortly acuminate at apex, pubescent when young, slightly pubescent beneath on mature leaves, coriaceous, glabrous, 10-14 pairs of inconspicuous lateral veins; petiole 1- 2 cm long. Inflorescence axillary with 1- 3 flowered fascicles; flower buds ovoid; pedicel slender 2-3 cm long; calyx lobes 8 in two series of 4 each, c. 7 x 3 mm, lanceolate, acuminate-recurved at apex, corolla lobes 24, white, ca. 8 x 3 mm, oblong lanceolate, obtuse re-curved at apex. Stamens 8, filaments slender, anthers linear-sagittate, staminodes alternate with stamens, pubescent. Ovary pubescent with subulate style. Fruit ellipsoid, 3 x 2 cm or globose up to 2.5 cm across with persistent style, orange turning bright red in colour.

Distribution : Andaman – Nicobar Archipelago (South Andaman, Little Andaman & Great Nicobar) and Sri Lanka.

CONCLUSION

The phytogeography of Andaman-Nicobar Islands characterized with a unique transitional zone of floristic elements with remarkable degree of intraspecific variations between the South Asian and Southeast Asian floristic regions. Interestingly, it also demonstrates multi-dimensional affinities to other nearer and far-off geographical regions such as Myanmar, Northeast India, Thailand, Vietnam, Malesia, Sri Lanka, Peninsular India (Western Ghats) and even towards African Continent to some extent. The climatological features and geographical isolation of these islands from the major landmasses of South and Southeast Asia over millions of years have resulted into remarkable degree of distinct plant diversity and endemism. According to geological evolution, the Andaman-Nicobar Islands are recognized as the continuation of Arakan Yoma Mountains of the Myanmar to the Moluccas Island of the Indonesia. It is believed that these islands have been isolated from the major continental land masses during the course of 'Continental Drift', most probably during the Early Tertiary or Late Cretaceous period (Renvoize, 1979). Therefore the biological entities of the Andaman-Nicobar Islands are obviously 'continental' in origin and have evolved into the present status through evolution over millions of years. Hence, the present insular bio-system may be referred to as a 'sub-continental bio-system'. Melville (1973) suggests that the Indian Ocean Islands were formerly fractions of 'Gondwana Continent' during the Paleozoic and Mesozoic periods. Rao and Chakraborti (1987) had also highlighted the floristic affinities of the littoral species of the Andaman-Nicobar Islands towards Indo-African, Indo-Pacific and Indo-Malaysian regions which obviously highlights its Gondwana connections in the remote past. The Western Ghats of the Peninsular India, Sri Lanka and the Andaman- Nicobar Archipelago have remarkable similarities in climatological features and distribution of floristic elements. The fragmented distribution of several floristic elements such as *Cleidion nitidum* (Müll.Arg.) Thwaites ex Kurz, *Polyalthia rufescens* Hook. f. & Thomson, *Salacia reticulata* Wight, *Mimusops andamanensis* King & Gamble, *Nageia wallichiana* (Presl.) O. Ktze. etc among Andaman-Nicobar Archipelago, Sri Lanka and the Western Ghats demonstrates the previous geological connections of these widely separated regions.

ACKNOWLEDGEMENTS

The authors are thankful to Dr P G Latha, Director, JNTBGRI for encouragement and to Dr C Satheeshkumar, Senior Scientist, JNTBGRI for providing information on concerned species from Sri Lankan herbaria. We are also thankful to forest officials of Andaman-Nicobar Islands for rendering facilities and to DST, New Delhi for the financial assistance for exploration.

REFERENCES

- Anonymous, 2007: The 2007 red list of threatened flora and fauna of Sri Lanka. IUCN and MENR, Sri Lanka.
Anonymous, 2012: The national red list 2012 of Sri Lanka - Conservation status of the fauna and Flora, Biodiversity Secretariat of the Ministry of Environment and National Herbarium, Department of National Botanic Gardens pp 332

Balakrishnan, N. P. and Vasudeva Rao M. K. 1983: (in Jain, S. K. & R. R. Rao, An assessment of threatened plants of India) The dwindling plant species of the Andaman and Nicobar Islands 186 – 201.

Bustard, H. R. 1971: Summary of meeting in crocodiles, vol. 1, paper no. 32, pp 15-30, IUCN publication

Darwin, C. 1872 : On the Origin of Species. A Penn State Electronic Classics Series Publication, Pennsylvania State University *pp* 13.

Drury, W. H. 1974: Rare species. *Biol. Conservation* 6: 162 – 169.

Melville, R. 1973: Continental drift and the distribution of island floras of the Indian Ocean. *J. Mar. Biol. Assoc. India*, 15: 236 – 241.

Oldfield, S., Lusty, C. and MacKinven, A. 1998: *The World List of Threatened Trees*. World Conservation Press, Cambridge, UK.

Pandey, R. P. and Diwakar P. G. 2008: An integrated check-list flora of Andaman and Nicobar Islands, India. *J. Econ. Taxon. Bot.* 32: 403-500.

Rao, T. A. and Chakraborti, S. 1987 : Distributional resume of coastal floristic elements in Andaman and Nicobar Islands. *Curr. Sci.* 56 : 1045 – 1051.

Renvoize, S. A. 1979 : In *Plants and Islands* (ed. Bremwell, D.) The origins of Indian Ocean Island floras, Academic Press Inc., New York, *pp* 107- 127.

Senaratna, L. K., 2001: A checklist of the flowering plant of Sri Lanka. MAB Check List and Hand Book Series Publication No. 22, National Science Foundation 47/5, Maitland Place, Colombo 7, *pp* 333.



A. *Mimusops andamanensis* King & Gamble – A view of live specimen from Little Andaman **B.** Fruits – Variation in fruit shape and size **C.** Ripened fruits **D.** Seeds **E.** Isotype of *Mimusops andamanensis* King & Gamble deposited at Kew

