

# Correlation between determining factors and dietary diversity in children aged 6 to 59 months in market gardening and non-market gardening areas of Niger

*by* Jana Publication & Research

---

**Submission date:** 18-Feb-2025 01:29PM (UTC+0700)

**Submission ID:** 2579267288

**File name:** IJAR-50310.docx (423.67K)

**Word count:** 5593

**Character count:** 31123

## Correlation between determining factors and dietary diversity in children aged 6 to 59 months in market gardening and non-market gardening areas of Niger.

### Summary

An unbalanced diet has been identified as a major cause of morbidity and mortality in children. The general objective is to identify the factors determining the food diversification of children aged 6 to 59 months in two areas of different horticultural specificity. This is a cross-sectional study, conducted in three municipalities, with 214 mothers and or caregivers. A questionnaire evaluating the food diversity was used, followed by statistical analysis of the results. The group of cereals and fruits and vegetables rich in vitamin A is the most consumed. Those of animal origin were low. A significant link between Minimum Food Diversity and certain characteristics such as area, income, mother education, agriculture, gardening and breeding ( $p < 0.01$ ). According to the logistic regression, residence of the child in non-market gardening zone (Dosso commune urbains) (ORa [IC95%] =44.60 [5.90-152.50];  $P < 0.001$ ) and in market gardening zone of Tillabéri (ORa [IC95%] =30 [7.67-259.40],  $p < 0.001$ ) Increases their chances of having a diversified diet by 44 and 30 times respectively compared to a child in the Dosso market zone. Efforts to improve food security, especially in vulnerable households, were needed to ensure a balanced diet for children.

**Keywords:** diversification of food, mothers, children from 6 to 59 months, horticultural specificity.

### INTRODUCTION


Food diversity is the basis of good nutrition, and an individual's nutritional status has important implications for labour productivity, lifetime incomes and economic development across the country (Shively and Evans, 2021). A diversified diet leads to malnutrition in the image of marasmus and kwashiorkor (Houndji et al., 2013; Lourme Ruiz et al., 2016; Sanou et al., 2018). Malnutrition compromises the physical and intellectual growth of young children (Kayodé et al., 2012). In Niger, children suffering from acute malnutrition have a rate above the WHO 10% alert threshold and sometimes even above the emergency threshold (15%) (World Health Organization, 2009). This malnutrition is directly linked to inadequate nutrition. However, for a good balance of food, a diversified diet is necessary. It provides the body with most of the nutrients it needs to function properly (Sanou et al., 2018). Adherence to a minimum acceptable diet is also essential for reducing macronutrient and micronutrient deficiencies (Jones et al., 2014a; Bougma et al., 2022). According to the SMART 2019 survey, the majority of Nigerian children have a low food diversification with 42.1% consuming at least 4 groups of foods per day and only 31.8% having an acceptable minimum diet (INS, 2019). This situation varies greatly from one region to another. The lowest

proportions are in Maradi (3%), Tahoua (8.7%), Tillabéri (9.2%) and <sup>33</sup> refugee camps (4.1%) (INS, 2018). Indeed, several studies have shown that inappropriate **infant and young child feeding practices** are <sup>24</sup> limited to the socio-demographic characteristics of mothers, but little information is available on the dietary diversity of children for better understand the causes of this malnutrition. This study was initiated to identify the key determinants that influence children's dietary diversity.

## <sup>56</sup> MATERIALS AND METHODS

### Study area

The present study concerned three municipalities. This is the commune of Liboré located in the department of Kollo region of Tillabéri, between 13° 24' 17" latitude north and 2° 11' 29" longitude east, with an area of 110 km. The commune of N'Gonga located 30 km south of the chief departmental place Birni N'Gaouré located between latitude 13° 17' 9" and 2.826 13° 10' 48" north longitude and 2° 49' 34" east longitude with 211m of altitude. The urban commune of Dosso occupies the central part between 13° 05' latitudes North, and 1° 30' and 30' 20" longitudes East on national road 1 (axis Niamey-N'Guigmi) 139 km to the East of Niamey (the capital) with an area of 592.6 km<sup>2</sup>. In Figure 1, you can see the map of Niger and the locations of the various communes (HASSANE et al., 2024).



Carte 1: Location map of the study areas (HASSANE et al., 2024).

Description of the study:

The study was conducted from February 20 to August 5, 2022, and involved mothers/caregivers of children aged 6-59 months. Children should be physically healthy and have resided in the study area for at least 6 months. Parental consent is also obtained prior to the inclusion of the mother/guardian and child in the study.

### Sampling

The total number of mother-child couples in the study was determined using ASN SMART 2020 software used in the national nutrition survey to calculate sample size. The formula is  $N = (1 + NR) (D * Z^2 * p (1 - p)) / m^2$ .

The mother/child couples were selected randomly in the three municipalities of Liboré, N'Gonga and Dosso. The study covered a total of 214 mother-child couples.

### tools for collecting data

Data collection, processing and analysis were aligned with WHO guidelines on indicators for assessing infant and young child feeding practices (World Health Organization, 2010) and FAO to measure Food Diversity at household and individual level (Gina et al., 2013). A questionnaire was developed and focused on the collection of demographic and economic information. The Food Diversity questionnaire was developed according to WHO and FAO guidelines in accordance with the study objectives and country.

### Data collection tools

Data collection, processing and analysis were adapted to the WHO guidelines on indicators to assess infant and young child feeding practices (World Health Organization, 2010) and FAO guidelines to measure Dietary Diversity at household and individual levels. A questionnaire was developed and focused on collecting demographic information (age, sex and education level of mothers or caregivers), socio-economic information (income-generating activities), food consumption (different food group consumed) and production systems (crops, livestock and gardening). The Dietary Diversity questionnaire was developed according to WHO and FAO guidelines in accordance with the study objectives and the country. Face-to-face interviews with the mothers of the children were conducted in the households. Based on the open recall, it was checked which food groups were consumed using a predefined list of food groups. A list of 42 food items was used. One point was awarded for each food or food group consumed, and zero was awarded if none were consumed.

### Variables

Variables that include demographic, socioeconomic characteristics, food consumption and production systems. The dependent variable was Minimum Food Diversity with these two

subgroups (low and acceptable). The choice of explanatory variables was based on a prior knowledge of the variables likely to influence the level of dietary diversity. The variables examined as determinants of food diversity were the place of residence, sex of the child; the level of education of the mother/guardian, the practice of breeding; the practice of gardening; farmer; and sources of income. In relation to the children's food diversity score (FDSE), it was equal to the number of food groups consumed by children. Based on WHO recommendations, the FDSE selected 7 groups according to WHO and FAO (Gina et al., 2013; World Health Organization, 2010; Ouédraogo et al., 2019; World Health Organization, 2010). Children who consumed daily foods from at least 4 out of 7 food groups defined during the previous day reach the Minimum Dietary Diversity (MDD). Minimum Dietary Diversity (MDD) and the frequency of eating group are dependent variables and were examined according to the areas and characteristics of households, mothers and children. The DAM is divided into two classes, low food diversity score (LFDS), when  $MDD < 4$  and Minimum Food Diversity acceptable (MFDA) when  $FDS > 4$ . The children's dietary diversity was crossed with the household characteristics to determine their association.

### Statistical analysis

Study data were collected using the KoboCollect software and processed on STATA MP 16 and SPSS 25. Chi-two association analyses were performed to describe the characteristics of mothers, children and households, the minimum food diversity and types of food groups consumed in relation to residential areas, the p-value of 0.05 was considered statistically significant. A multi-variate logistic regression analysis was performed to predict the probability that a child would belong to an adequate or low level of dietary diversity. The dependent variable was the Minimum Food Diversity with these two subgroups (low and acceptable). The choice of explanatory variables was based on a prior knowledge of the variables likely to influence the level of acceptable Dietary Diversity. Those with a p-value of 0.05 are associated with the variable to be explained in other words dependent. The Odds ratios (OR) column indicates how often it influences, positively or negatively, the food diversity and a confidence interval (CI) at 95%. The predictive power of the logistic model, the identification of determinants of children's food diversification was determined to create assumptions and predictions about children's food diversification in the future

### Ethical considerations

This study was granted by the health authorities of the region and approved by the National Ethics Committee for Health Research (CNER) under number 075/2021/CNER of December 09, 2021. The survey content was explained in detail to participants. Participation was voluntary and each mother or caregiver signed a declaration of informed consent. Data collection and Processing was carried out in a confidential manner during the analysis and reporting of the data.

## Results

Distribution of children by age group according to zones.

The distribution of children by age group and area is shown in Table 1.

**Table 1:** Distribution of children 6 to 59 months according to age group according to market gardening and non-market gardening areas.

age groups	Areas			Total	p
	market gardener Dosso	market gardener Tillabéri	non- market gardening Dosso		
	N (%)	N (%)	N (%)	N (%)	<10 <sup>-3</sup>
6 to 23 months	35 (47.3)	34 (48.6)	68 (97.1)	137 (64.0)	
24 to 59 months	39 (52.7)	36 (51.4)	2 (2.9)	77 (36.0)	
Sample size	74 (100.0)	70 (100.0)	70 (100.0)	214 (100.0)	

Table 1 shows that overall, children aged 6 to 23 months are most represented (64%). Specifically, children aged 6 to 23 months or more dominant (97.1%) in the non-market gardening area (Dosso) compared with (52.7%) and (51.4%) children aged 24 to 59 months respectively in the market gardening areas of Dosso and Tillabéri. There is a statistically significant relationship between the zones and age groups ( $p < 0.001$ ).

Table 2 presents an analysis of the sociodemographic and economic characteristics of mothers or caregivers who care for children aged 6 to 59 months (HASSANE S.Z et al., 2024).

**Table 2:** Relationship between the socio-demographic and economic characteristics of mothers or caregivers of children aged 6 to 59 months according to the areas

Variables	Areas			Total	p
	Market gardener	Market gardener	Non-market		

		Dosso N (%)	Dosso N (%)	gardening Dosso N (%)		
Level of education	schooled	8 (10.9)	40 (57.14)	49 (70)	97 (45.33)	<b>&lt; 10<sup>-3</sup></b>
	Out of school	60 (81.08)	17 (24.29)	17 (24.29)	94 (43.93)	
	Koranic School	6 (8.11)	13 (18.57)	4 (5.71)	23 (10.75)	
Total		74 (100)	70 (100)	70 (100)	214 (100)	
Civil status	Bride	70 (94.59)	68 (97.14)	70 (100)	208 (97.2)	<b>0.02</b>
	Divorced	0	2 (2.86)		2 (0.93)	
	Widow	4 (5.41)	0	0	4 (1.87)	
Total		74 (100)	70 (100)	70 (100)	214 (100)	
Source of income	Employees	0 (0.0)	3 (4.29)	10 (14.29)	15 (6.19)	<b>&lt; 10<sup>-3</sup></b>
	Farmers	25 (33.78)	5 (7.14)	2 (2.86)	32 (14.95)	
	traders	29 (39.19)	17 (24.29)	21 (29.9)	65 (31.2)	
	Without income	20 (27.03)	45 (64.29)	37 (52.86)	102 (47.66)	
Total		74 (100)	70 (100)	70 (100)	214 (100.0)	
Gardening practice	Yes	64 (86.49)	17 (24.29)	3 (4.29)	84 (39.25)	<b>&lt; 10<sup>-3</sup></b>
	No	10 (13.51)	53 (75.71)	67 (95.71)	130 (60.75)	
Total		74 (100.0)	70 (100.0)	70 (100.0)	214 (100.0)	


This analysis shows that the majority of mothers were married and there is a strong link between zones and marital status (p 0.05). The rate of no income was 47.66%, the difference was also significant (p < 0.001). The rate of non-income was particularly high (52.86%) in the non-horticultural area of the urban municipality of Dosso and in the horticultural area of Tillabéri (64.29%). In addition, 39.25% are engaged in gardening. The

proportion of mothers engaged in gardening is particularly low in the non-garden sectors (4.29%) and in the Tillabéri market (24.29%), which is also significant ( $p < 0.001$ ). (43.93%) were not in school and this rate was higher at the level of the municipality of Dosso (81.08%), the difference was significant ( $p < 0.001$ ). There was a significant difference between the two study areas for all characteristics studied ( $p > 0.05$ ).

#### **Food profile of children aged 6 to 59 months according to the food consumption score (FCS)**

Figure 1 shows the consumption rate of food groups according to the areas of children aged 6 to 59 months. In all areas, starchy foods (cereals, roots, tubers) and fruits and vegetables rich in vitamin A are the most consumed. Legumes and nuts, dairy products (milk, yogurt, and cheese), meat products (meat, fish, poultry, and liver/offal) and eggs are the least consumed. At the area level, consumption shows disparities depending on the potency and availability of food. Children in the market gardening area of Dosso (N'Gonga) and the non-market gardening area (Dosso urban commune) not only consumed 100% of starchy foods but also fruits and vegetables rich in vitamin A. The difference is being no significant between the areas ( $P > 0.05$ ). Other fruits and vegetables were more consumed (94.4%) in the market gardening area of the commune of Liboré (Tillabéri region), a significant difference was observed according to the areas ( $P < 0.001$ ). In addition, legumes and nuts, dairy products (milk, yogurt, and cheese) and meat products (meat, fish, poultry, and liver/offal) are more consumed in the non-market gardening area of the urban commune of Dosso, so, respectively, 50.0.0% and 50.0%. The proportions were statistically significant according to this area ( $P < 0.05$ ). However, no difference was observed between the areas concerning the consumption of legumes and nuts, ( $P > 0.05$ ).





The percentages of food groups consumed by area: dairy products (milk, yogurt, cheese):  $p < 10^{-3}$ ; meat products (meat, fish, poultry, liver/offal):  $p < 10^{-3}$ ; other fruits and vegetables:  $p < 10^{-3}$

#### Relationships between variables

The chi-two association's results indicate that the dietary diversity of children was statistically associated with certain sociodemographic and economic characteristics. Table 3 provides details of all these results.

**Table 3:** The minimum dietary diversity of children aged 6 to 59 months is influenced by socio-demographic and economic characteristics.

	Less than 4 groups	At least 4 groups	Total	p
areas	N(%)	N(%)	N(%)	
Market gardener Dosso	71(95.9)	3(4.1)	74(100.0)	< 10 <sup>-3</sup>
Market gardener Tillabéri	32(45.7)	38(54.3)	70(100.0)	
Not-market gardening Dosso	24(34.3)	46(65.7)	70(100.0)	

<b>Total</b>	127(59.3)	87(40.7)	214(100.0)	
Educational level				
<b>Higher level</b>	0(0.0)	7(100.0)	7(100.0)	< 10 <sup>-3</sup>
<b>Secondary level</b>	19(46.3)	22(53.7)	41(100.0)	
<b>Primary level</b>	18(36.7)	31(63.3)	49(100.0)	
<b>Koranic School</b>	16(69.6)	7(30.4)	23(100.0)	
<b>Out of school</b>	74(78.7)	20(21.3)	94(100.0)	
<b>Total</b>	127(59.3)	87(40.7)	214(100.0)	
Mains source of income				
<b>Employees</b>	2(15.4)	11(84.6)	13(100.0)	< 10 <sup>-3</sup>
<b>Farmers</b>	28(87.5)	4(12.5)	32(100.0)	
<b>traders</b>	37(56.9)	28(43.1)	65(100.0)	
<b>Without income</b>	61(58.8)	43(41.2)	104(100.0)	
<b>Total</b>	127(59.3)	87(40.7)	214(100.0)	
Agricultural practice				
<b>No</b>	18(34.6)	34(65.4)	52(100.0)	< 10 <sup>-3</sup>
<b>yes</b>	109(67.3)	53(32.7)	162(100.0)	
<b>Total</b>	127(59.3)	87(40.7)	214(100.0)	
Gardening Prattice				
<b>No</b>	60(46.2)	70(53.8)	130(100.0)	< 10 <sup>-3</sup>
<b>Yes</b>	67(79.8)	17(20.2)	84(100.0)	
<b>Total</b>	127(59.3)	87(40.7)	214(100.0)	
Briefing Prattice				
<b>No</b>	40(44.0)	51(56.0)	91(100.0)	< 10 <sup>-3</sup>
<b>Yes</b>	87(70.7)	36(29.3)	123(100.0)	
<b>Total</b>	127(59.3)	87(40.7)	214(100.0)	
Gender of child				
<b>Male</b>	68(59.1)	47(40.9)	115(100.0)	0.940
<b>Female</b>	59(59.6)	40(40.4)	99(100.0)	
<b>Total</b>	127(59.3)	87(40.7)	214(100.0)	


The analysis reveals that the <sup>2</sup> Dietary Diversity of children aged 6 to 59 months is low <sup>5</sup> in all study areas (40.7%). The proportion of children who consumed at least 4 food groups (adequate dietary diversity) increases as we move from the market gardening areas of N'Gonga (Dosso), the commune of Liboré (Tillabéri) to that of the non-market gardening area of the urban commune of Dosso. It is 4.1% respectively; 54.3% and <sup>9</sup> 65.7%. Regarding the level of education of mothers, children from educated mothers were less likely to have a low dietary diversity compared to their counterparts. Children born to mothers whose main source of income is wages were less likely to be food insecure (84.6%). Regarding practices, children from households engaged in agriculture, gardening or livestock farming have a low food

diversity composed of less than four (4) food groups. In terms of sex, it <sup>67</sup>did not influence the Minimum Dietary Diversity of children (P > 0.05).

#### <sup>16</sup>Correlation between food diversity determinants and minimum dietary diversity score of children aged 6-59 months

##### *Relationships between variables*

<sup>61</sup>According to the results of the bivariate analysis in figure 2, some characteristics are associated with an increase in the probability that a child has Minimum Acceptable Food Diversity, it is: the area (Market gardener Tillabéri, Not-marketgardening Dosso), Educational Level (secondary level, primary level) and Mains source of income (Employees), (OR>1).



**Figure 2:** Relationship between sociodemographic and economic characteristics and Minimum Dietary Diversity of children aged 6 to 59 months according to bivariate analysis

<sup>8</sup>To study the determinants of dietary diversity in children, a bivariate analysis of selected explanatory variables (figure 2) was performed. Then, using a logistic model, the simultaneous effects of the variables selected in the first step (household characteristics) on the children's dietary diversity are determined.

Table 4 presents the factors associated with minimum dietary diversity. The model estimates to identify two relevant variables influencing the dietary diversity of children aged 6-59 months. These are the non-market gardening area "Dosso urban commune" and the market gardening area of Tillabéri "commune of Liboré" which have significant links with the minimum acceptable dietary diversity regime of children ( $P < 0.05$ ).

**Table 4:** Logistic regression of variables related to the Minimum Acceptable Dietary Diversity of children aged 6 to 59 months

Variables	ORa [IC95% ]	p
<b>Areas</b>		
Market gardener Dosso	1	-
Market gardener Tillabéri	30.01[5.90-152.50]	$< 10^{-3}$
Non-marketgardening Dosso	44.59752	$< 10^{-3}$
<b>Educational Level</b>		
Higher level	1	-
secondary level	0.93[0.33-2.59]	0.891
primary level	1.64[0.61-4.39]	0.329
Koranic School	0.75[0.21-2.66]	0.659
<b>Mains source of income</b>		
Employees	1	-
Farmers	0.30[0.03-2.90]	0.300
Traders	0.93[0.15-5.82]	0.939
Without income	0.41[0.07-2.26]	0.306
<b>Agricultural practice</b>		
No	1	-
Yes	0.48[0.20-1.17]	0.107
<b>Gardening Praticce</b>		
No	1	-
Yes	2.30[0.62-8.53]	0.212
<b>Briefing Praticce</b>		
No	1	-
Yes	1.38[0.61-3.09]	0.441
<b>Gender of child</b>		
Male	1	-
Female	1.14[0.55-2.38]	0.725

Areas are a very important feature in the analysis of food diversity. Children aged 6 to 59 months in the non-garden area "Urban Dosso " (ORa [IC95%] =44.60 [5.90-152.50];  $p=0.000$ ) and the Tillabéri market gardening area (ORa [IC95%] =30 [7.67-259.40],  $p=0.000$ ) were less likely to have low food diversity compared to their Dosso market gardening area counterpart

#### Predictive power of the logistic model of minimum dietary diversity in children aged 6-59 months

The results of the estimates of the predictive power of the logistic model, the identification of the determinants of the Minimum Dietary Diversity of children as well as their interpretations are presented in Table 6.

**Table 6:** Predictive power of the logistic model of minimum dietary diversity in children aged 6-59 months

Prévisions	Minimal dietary diversity in children			P
	Less than 4 groups n(%)	At least 4 groups n(%)	Total n(%)	
Less than 4 groups	92 (72.44)	35 (27.56)	127 (100)	0.5000
At least 4 groups	18 (20.69)	69 (79.31)	87 (100)	
Total	110 (51.40)	104 (48.60)	214 (100)	

The ranking table (Table 5) shows that the model has a low predictive power as it predicts the dietary diversification of children at 48.60%. However, it appears that the dietary diversity of at least 4 groups is correctly classified (79.31%) than the minimal dietary diversity of less than 4 groups (72.44%).

## DISCUSSION

The results show that the most representative age group is 6-23 months with a proportion of 64.0% ( $p < 0.01$ ). This is in disagreement with the INS (2021). According to the analyses, 43.93% of mothers had no formal education. This very high percentage of uneducated women could be due to the Nigerian education system, which is characterized by a low rate of schooling for girls in rural areas. These results are consistent with the national INS documentation (2017; 2021). Furthermore, 24.29% of mothers in the non-gardening area of Dosso were uneducated. This same percentage was obtained in the market gardening area of Tillabéri. This percentage was significantly lower ( $P < 0.01$ ) than the percentage of mothers interviewed in the market gardening area of Dosso (81.08%). This very high proportion of uneducated women in the market garden area of Dosso could be due to the characteristic of the commune, which is a very rural commune, or in rural areas, from a young age, the girl is perceived as a woman destined to perform specific functions such as reproduction (marriage) and maintenance of family members. Among the respondents, 47.66% were without income, and rates varied considerably between zones ( $p < 0.01$ ). The market gardening area of Dosso had a very low rate of no income, which is 27.03% compared to the non-market gardening area of Dosso and the market gardening area of Tillabéri. These results are different from the national INS documentation (2022). This may be due to the nature of the area, which is a natron production area par excellence where almost all women participate in the production

and are paid. This is comparable to studies conducted by Berde et al. (2019) in 23 countries in sub-Saharan Africa where 43.8% of women had no formal education and 34.5% were without income. Similarly, our results are similar to the studies conducted by Lucha et al. (2022) in Ethiopia where more than one third (48.6%) of mothers had no education. These results are also consistent with the study conducted by Jones et al. (2014b) in Madagascar where they found a significant difference for all socio-demographic characteristics.

Analysis of the dietary profile of children shows that cereals and vitamin A-rich fruits and vegetables were frequently consumed in the areas. The high consumption of these products in this study could be due to their physical availability. These results are consistent with those of Moursi et al. (2008) and several studies on food consumption and the national INS documentation (2015; 2017; 2019; 2020; 2021). In contrast, animal products are consumed sparingly. Meat is eaten either once a week or monthly. Milk and dairy products are also consumed at low levels. The majority of respondents told us that their children almost never eat eggs. The low consumption of these foods and the predominant consumption of cereals in the diets of children living in rural households such as those in our study has already been observed in other rural communities in Sudan, Niger, Burkina Faso, in Madagascar, and Ethiopia (Workicho et al., 2016; Khalid et al., 2017; Rakotonirainy et al., 2018; Sanou et al., 2018; Ouedraogo et al., 2019; INS, 2022). Although consumption of animal products is generally low, it was higher in the non-gardener area of the urban municipality of Dosso. This could be explained by the availability of these products, as the area is not only urban (these products are available and accessible), but also parents are more educated so they know the advantage of their children's consumption of these products.

Diets based on the consumption of cereals or starches as staple foods and low in fruits and animal products are common in low-income countries (Jones et al., 2014b; Sanou et al., 2018). The risk of such a diet is micronutrient deficiency and, as a result, malnutrition and poor health Sanou et al. (2018).

The analysis of variables related to the dietary diversity of children aged 6-59 months shows that the area, the source of income and the level of education of mothers have an influence on the dietary diversity of children. Apart from these characteristics mentioned above, the practice of agriculture, gardening and livestock farming are also associated with food diversity. These results are similar to those of Rakotomanana et al. (2017). They found that the main risk factors for low dietary diversity were low levels of maternal education and socio-economic status. These results are similar to those of Arimond et al. (2004) who

reported, through their bivariate regression analysis of their study carried out on recent data from EDS surveys (Demographic and Health Surveys) from 11 countries, an association between urban/rural conditions and the nutritional status of children (Arimond and Ruel, 2004). They are also similar to the results reported by Jones et al. (2014b); Workicho et al. (2016). Previous studies have reported the positive influence of market gardening on children's food diversity (Sibhatu et al., 2015; Ouedraogo et al., 2019). The results of the logistic regression show that the area was positively associated with high food diversity. It revealed that the food diversity varied according to place of residence.

It appears that children living in the market gardening area of Tillabéri are 30 times more likely to have an adequate minimum food diversity compared to a child living in the market gardening area of Dosso (ORa [IC95%] =30 [7.67-259.40], p=0.000). These results could be due to the availability and accessibility of food to the population, which is explained by the proximity of the municipality to the capital and the river. In addition, the fact that a child is in the non-garden area of Dosso urban commune multiplies by 44 his chance to have a diversified diet compared to a child who is in the market gardening area of Dosso (ORa [IC95%] =44.60 [5.90-152.50]; P=0.000). These results are different from those of H. Rakotomanana et al. (2017). They found that the main risk factors for low food diversity were low levels of education and low socio-economic status. These results are consistent with studies conducted among Chinese preschool children and children from eleven (11) African countries. These previous studies indicated that food diversity is positively influenced by the urban environment (Mary Arimond and Ruel, 2004; Luo et al., 2012). This data is consistent with the results of a study conducted in eight (8) cities in China by Zhao et al. (2017).

## CONCLUSION

In this study, the aim was to establish a correlation between the determining factors of dietary diversification and the Minimum Dietary Diversity of children aged 6 to 59 months in two areas of different horticultural specificity.

The analysis shows that the determining factors are the residential environment (the area) which was correlated with a high dietary diversity for children aged 6 to 59 months. The dietary situation of children aged 6 to 59 months remains a concern. The quality of children's diet during their early years depends mainly on the behavior and decisions of mothers or people who usually take care of the child. Hence the need to develop actions for women's education and awareness-raising integrated into a long-term multi-sectoral development program to increase the diversity of agricultural production and access to nutritious foods.

## COMPETING INTERESTS

The authors declare no competing interest relating to the content of this article.

## AUTHORS' CONTRIBUTIONS.

All authors contributed to the conduct of this work. The authors of this article have read and approved its content.

## ACKNOWLEDGMENTS

We would like to thank all the mothers and/or guardians who contributed to this study. Our thanks also go to all the staff of the traditional chiefdom of Libérée and N'Gonga and the town hall of the urban commune of Dosso and Liboré. We would like to thank all the mothers and/or guardians who contributed to this study. Our thanks also go to all the staff of the traditional chiefdom of Liboré and N'Gonga and the town hall of the urban commune of Dosso and Liboré. Our thanks to the National Nutrition Information Platform (PNIN), particularly Almoustapha Théodore Yatta, for his valuable assistance in data collection. This study was made possible thanks to the financial support of ECOWAS

## REFERENCE

1. Arimond M, Ruel MT. (2004). Dietary Diversity Is Associated with Child Nutritional Status : Evidence from 11 Demographic and Health Surveys. *J. Nutr.* **134**(10): 2579–2585. DOI: <https://doi.org/10.1093/jn/134.10.2579>
2. Berde AS, Bester P, Kruger IM. (2019). Coverage and factors associated with vitamin A supplementation among children aged 6–59 months in twenty-three sub-Saharan African countries. *Public Health Nutr.* **22**(10): 1770–1776. DOI: <https://doi.org/10.1017/S1368980018004056>
3. Bougma S, Hama-ba F, Garanet F, Savadogo A. (2022). Sociodemographic characteristics of mothers and complementary feeding practices among children aged 6 to 23 months in north-central Burkina Faso. *Afr. J. Food Agric. Nutr. Dev.* **22**(10): 22017–22040 DOI: <https://doi.org/10.18697/ajfand.115.22250>
4. Gina K, Terri B, Marie C. (2013). Guidelines for measuring household and individual dietary diversity. Food Agric. Organ. U. N. Rome Italy, p.60.
5. Hassane SZ, Alkassou SI, Mamane AN, Mamane MR, Sabo HS. (2024). Connaissances, perceptions et pratiques sur la carence en vitamine A chez les mères/gardiennes d'enfants âgés de 6 à 59 mois dans des zones maraîchères et non maraîchères au Niger. *Med.Afr. Francoph.* **71**: 501-512. <https://www.santetropicale.com/manelec/fr/index.asp>
6. Houndji BVS, Bodjrenou SF, Londji SB, Ouetchehou RA, Acakpo A, Amouzou K. (2013). Amélioration de l'état nutritionnel des enfants âgés de 6 à 30 mois à Lissèzoun (Centre-Bénin) par la poudre de feuilles de Moringa oleifera (Lam.). *Int. J. Biol. Chem. Sci.* **7**(1): 225–235. DOI : <http://dx.doi.org/10.4314/ijbcs.v7i1i.19>



7. INS. (2022). Enquête retrospective sur la nutrition et la mortalité au Niger, p.103.
8. INS. (2021). Enquête nutritionnelle et de mortalité Rétrospective au Niger (No. rapport finale). Niamey, Niger, p.125.
9. INS. (2020). Enquête Vulnérabilité à l'Insécurité Alimentaire des Ménages (janvier-février 2020) (rapport final). Niamey, Niger; p.139.
10. INS. (2019). Rapport Final de l'Evaluation nationale de la situation nutritionnelle par la méthodologie SMART au Niger (16 août et 25 septembre 2019). Niamey Niger, p.139.
11. INS. (2018). Rapport final de l'évaluation nationale de la situation nutritionnelle par la méthodologie SMART 2018. Niger, p.181.
12. INS. (2017). Résultat préliminaire de l'enquête conjointe sur la vulnérabilité à l'insécurité alimentaire des ménages au Niger (décembre 2017). Niamey, Niger, p.184.
13. INS. (2015). Enquête conjointe sur la vulnérabilité à l'insécurité alimentaire des ménages au Niger (décembre 2014- janvier 2015) (rapport final). Niamey/Niger, p.169.
14. Jones AD, Ickes SB, Smith LE, Mbuya MN, Chasekwa B, Heidkamp RA, Menon P, Zongrone AA, Stoltzfus RJ. 2014a. World Health Organization infant and young child feeding indicators and their associations with child anthropometry : a synthesis of recent findings. *Matern. Child. Nutr.* **10(1)**: 1–17. DOI: [10.1111/mcn.12070](https://doi.org/10.1111/mcn.12070)
15. Jones AD, Shrinivas A, Bezner-Kerr R. 2014b. Farm production diversity is associated with greater household dietary diversity in Malawi: Findings from nationally representative data. *Food Policy* **46** (C): 1–12. DOI: <https://doi.org/10.1016/j.foodpol.2014.02.00>
16. Kayodé AP, Akogou FU, Hounkpatin WA, Hounhouigan DJ. (2012). Effects of processing methods on the nutritional value of sorghum-based supplementary porridge formulations. *Int. J. Biol. Chem. Sci.* **6(5)**: 2192–2201. DOI: <http://dx.doi.org/10.4314/ijbcs.v6i5.25>
17. Khalid FA, Ali AK, Ali SA, Mosmar ZY, Salih SS, Salman TK, Desogi MA, Soghaier MA, Mohammed EE, Mohammed AA. (2017). Households' dietary habits and food consumption patterns in Hamishkoreib locality, Kassala State, Sudan. *J. Ethn. Foods* **4(3)**: 181–186. DOI: <https://doi.org/10.1016/j.jef.2017.08.009>
18. Lourme R A, Dury S, Martin-Prével Y. (2016). Do we consume what we sow? Relation between production diversity, agricultural income and dietary diversity in Burkina Faso. *Cah. Agric.* **25**: 65001. DOI: [10.1051/CAGRI/2016038](https://doi.org/10.1051/CAGRI/2016038)
19. Lucha T, Engida T, Ketsela A. (2022). Assessing the potential determinants of national vitamin A supplementation among children aged 6-35 months in Ethiopia: further analysis of the 2019 Ethiopian Mini Demographic and Health Survey. *BMC Pediatr.* **22(1)**: 439. DOI: <https://doi.org/10.1186/s12887-022-03499-5>
20. Luo R, Shi Y, Zhang L, Liu C, Rozelle S, Sharbono B, Yue A, Zhao Q, Martorell R. (2012). Nutrition and Educational Performance in Rural China's Elementary Schools: Results of a Randomized Control Trial in Shaanxi Province. *Econ. Dev. Cult. Change* **60(4)**: 735–772. DOI: <https://doi.org/10.1086/665606>
21. Moursi MM, Arimond M, Dewey KG, Treche S, Ruel MT, Delpeuch F. (2008). Dietary diversity is a good predictor of the micronutrient density of the diet of 6-to 23-

- month-old children in Madagascar. *J. Nutr.* **138(12)**: 2448–2453. <https://doi.org/10.3945/jn.108.093971>
22. Organization WH. 2010. Indicators for assessing infant and young child feeding practices part 3: country profiles, p.60.
  23. Ouedraogo O, Compaore EW, Amouzou SK, Zeba AN, Dicko M H. (2019). Determination and Characterization of Women, Infants, and Young Children's Dietary Diversity in Agricultural Mitigation Period of Burkina Faso. *J. Nutr. Food Secur.* **4(2)**: 114-125; DOI: <https://doi.org/10.18502/jnfs.v4i2.774>
  24. Ouédraogo O, Wendinpuikondo RCE, Kou'santa SAE, Hama DM. (2019). Toddlers' Dietary Diversity and Its Determinants in Different Agricultural Periods. *Int. J. Nutr. Sci.* **4(3)**:151-161. DOI: <https://doi.org/10.30476/ijns.2019.82283.1018>
  25. Rakotomanana H, Gates GE, Hildebrand D, Stoecker BJ. (2017). Situation and determinants of the infant and young child feeding (IYCF) indicators in Madagascar: analysis of the 2009 Demographic and Health Survey. *BMC Public Health* **17(1)**: 1–9. DOI: [10.1186/s12889-017-4835-1](https://doi.org/10.1186/s12889-017-4835-1)
  26. Rakotonirainy NH, Razafindratovo V, Remonja CR, Rasoloarijaona R, Piola PR, Randremanana RV. (2018). Dietary diversity of 6-to 59-month-old children in rural areas of Moramanga and Morondava districts, Madagascar. *PLoS One* **13(7)**: e0200235. DOI: [10.1371/journal.pone.0200235](https://doi.org/10.1371/journal.pone.0200235)
  27. Sanou S, Ayantunde A, Nianogo AJ. (2018). Household food consumption and determinants of dietary diversity: the case of four communes in the North region, Burkina Faso. *Int. J. Biol. Chem. Sci.* **13(12)**: 1784–1801. DOI: <https://doi.org/10.4314/ijbcs.v12i4.21>
  28. Shively G, Evans A. (2021). Dietary Diversity in Nepal: A Latent Class Approach. *Food Nutr. Bull.* **42(2)**: 259–273. DOI: <https://doi.org/10.1177/0379572121998121>
  29. Sibhatu KT, Krishna VV, Qaim M. (2015). Production diversity and dietary diversity in smallholder farm households. *Proc. Natl. Acad. Sci.* **112(34)**: 10657–10662. DOI: <https://doi.org/10.1073/pnas.1510982112>
  30. Workicho A, Belachew T, Feyissa GT, Wondafrash B, Lachat C, Verstraeten R, Kolsteren P. (2016). Household dietary diversity and Animal Source Food consumption in Ethiopia: evidence from the 2011 Welfare Monitoring Survey. *BMC Public Health.* **16(1)**:1192. DOI: [10.1186/s12889-016-3861-8](https://doi.org/10.1186/s12889-016-3861-8).
  31. World Health Organization. (2010). Indicators for assessing infant and young child feeding practices: part 2: measurement. Indic. Pour Évaluer Prat. Aliment. Nourrisson Jeune Enfant Partie 2 Calc, p.91.
  32. World Health Organization. (2009.) Global prevalence of vitamin A deficiency in populations at risk 1995-2005: WHO global database on vitamin A deficiency. p.68.
  33. Zhao W, Yu K, Tan S, Zheng Y, Zhao A, Wang P, Zhang Y. (2017). Dietary diversity scores: an indicator of micronutrient inadequacy instead of obesity for Chinese children. 2017 May 12 *BMC Public Health*; **17(1)** : 440. DOI: [10.1186/s12889-017-4381-x](https://doi.org/10.1186/s12889-017-4381-x)

# Correlation between determining factors and dietary diversity in children aged 6 to 59 months in market gardening and non-market gardening areas of Niger

## ORIGINALITY REPORT

24%

SIMILARITY INDEX

21%

INTERNET SOURCES

17%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

1

[theses.hal.science](https://theses.hal.science)

Internet Source

2%

2

[hal-riip.archives-ouvertes.fr](https://hal-riip.archives-ouvertes.fr)

Internet Source

1%

3

[journalcra.com](https://journalcra.com)

Internet Source

1%

4

[eujournal.org](https://eujournal.org)

Internet Source

1%

5

[d-nb.info](https://d-nb.info)

Internet Source

1%

6

[works.bepress.com](https://works.bepress.com)

Internet Source

1%

7

[www.frontiersin.org](https://www.frontiersin.org)

Internet Source

1%

8

Mahama Saaka, Shaibu Mohammed Osman, Irmgard Hoeschle-Zeledon. "Relationship between agricultural biodiversity and dietary diversity of children aged 6-36 months in rural areas of Northern Ghana", Food & Nutrition Research, 2017

Publication

1%

9

[www.ssph-journal.org](https://www.ssph-journal.org)

Internet Source

1%

10	Natacha KERE, Ella W. R. COMPAORE, Ousmane OUEDRAOGO, Inoussa KY et al. "Status report on vitamin A deficiency and associated factors in the health area of Sanmatenga province (Burkina Faso)", Springer Science and Business Media LLC, 2024 Publication	1 %
11	easpublisher.com Internet Source	1 %
12	qa1.scielo.br Internet Source	1 %
13	journals.plos.org Internet Source	<1 %
14	www.annexpublishers.com Internet Source	<1 %
15	article.sciencepublishinggroup.com Internet Source	<1 %
16	downloads.hindawi.com Internet Source	<1 %
17	www.mdpi.com Internet Source	<1 %
18	www.ajol.info Internet Source	<1 %
19	edepot.wur.nl Internet Source	<1 %
20	tropmedhealth.biomedcentral.com Internet Source	<1 %
21	Rachel Y. Goldstein, Derek A. Seehausen, Alice Chu, Debra A. Sala, Wallace B. Lehman. "Predicting the Need for Surgical Intervention	<1 %

in Patients With Idiopathic Clubfoot", Journal  
of Pediatric Orthopaedics, 2015

Publication

22

Submitted to Vrije Universiteit Amsterdam

Student Paper

<1 %

23

KilinÃ§, NurgÃ¼l. "Clothing Color Preferences  
of Boys and Girls Aged Between Six and  
Nine", Social Behavior and Personality An  
International Journal, 2011.

Publication

<1 %

24

Pallavi Rajkhowa, Suman Chakrabarti.  
"Temperature and children's dietary diversity:  
Evidence from India", Food Policy, 2024

Publication

<1 %

25

ora.ox.ac.uk

Internet Source

<1 %

26

Alekaw Sema, Yalelet Belay, Yonatan  
Solomon, Assefa Desalew et al. "Minimum  
Dietary Diversity Practice and Associated  
Factors among Children Aged 6 to 23 Months  
in Dire Dawa City, Eastern Ethiopia: A  
Community-Based Cross-Sectional Study",  
Global Pediatric Health, 2021

Publication

<1 %

27

sciencepublishinggroup.com

Internet Source

<1 %

28

Submitted to Kenyatta University

Student Paper

<1 %

29

Submitted to University of South Africa

Student Paper

<1 %

30

www.advancingnutrition.org

Internet Source

<1 %

31 Marius Affonfere, Flora Josiane Chadare, Yann Emeric Madode, Finagnon Toyi Kévin Fassinou, Gudrun B. Keding, Paulin Azokpota. " Consumption and mineral profile of spp. root powder: A traditional food ingredient in Sudanian zone of Benin (West Africa) ", Engineering Reports, 2023  
Publication

---

32 [bmcpediatr.biomedcentral.com](https://bmcpediatr.biomedcentral.com) <1 %  
Internet Source

---

33 [documents.wfp.org](https://documents.wfp.org) <1 %  
Internet Source

---

34 Heidkamp, Rebecca A., Mohamed Ag Ayoya, Ismael Ngnie Teta, Rebecca J. Stoltzfus, and Joseline Pierre Marhone. "Complementary feeding practices and child growth outcomes in Haiti: an analysis of data from Demographic and Health Surveys : Complementary feeding in Haiti", Maternal and Child Nutrition, 2013.  
Publication

---

35 Huguette Abi Khalil, Mariam Hawi, Maha Hoteit. "Feeding Patterns, Mother-Child Dietary Diversity and Prevalence of Malnutrition Among Under-Five Children in Lebanon: A Cross-Sectional Study Based on Retrospective Recall", Frontiers in Nutrition, 2022  
Publication

---

36 Ines Sebai, Stéphane Decelles, Malek Batal. "Determinants of dietary diversity among children 6-23 months: a cross-sectional study in three regions of Haiti", Journal of Human Nutrition and Dietetics, 2022  
Publication

---

37 Jong Hyun Pyun, Sung Gu Kang, Seok Ho Kang, Jun Cheon, Je Jong Kim, Jeong Gu Lee. "Efficacy of holmium laser enucleation of the prostate (HoLEP) in men with bladder outlet obstruction (BOO) and non-neurogenic bladder dysfunction", The Kaohsiung Journal of Medical Sciences, 2017  
Publication

---

38 Mehedi Hasan, Md. Hassan, Md. Shafiqul Khan, Md. Tareq, Mst. Afroj. "Prevalence, knowledge, attitudes and factors associated with exclusive breastfeeding among mothers in Dhaka, Bangladesh: A cross-sectional study", Population Medicine, 2021  
Publication

---

39 Purnima Menon, Apurva Bamezai, Ali Subandoro, Mohamed Ag Ayoya, Victor M. Aguayo. "Age-appropriate infant and young child feeding practices are associated with child nutrition in India: insights from nationally representative data", Maternal & Child Nutrition, 2015  
Publication

---

40 [biblio.ugent.be](http://biblio.ugent.be)  
Internet Source

---

41 [bmcpublichealth.biomedcentral.com](http://bmcpublichealth.biomedcentral.com)  
Internet Source

---

42 "Handbook of public health in natural disasters", Brill, 2015  
Publication

---

43 Stanley Chitekwe, Kedar Raj Parajuli, Naveen Paudyal, Karan Courtney Haag, Andre Renzaho, Abukari Issaka, Kingsley Agho. "Individual, household and national factors

associated with iron, vitamin A and zinc deficiencies among children aged 6–59 months in Nepal", Maternal & Child Nutrition, 2021

Publication

---

44	<a href="http://ageconsearch.umn.edu">ageconsearch.umn.edu</a>	<1 %
	Internet Source	

---

45	<a href="http://bmcnutr.biomedcentral.com">bmcnutr.biomedcentral.com</a>	<1 %
	Internet Source	

---

46	<a href="http://e-journal.unair.ac.id">e-journal.unair.ac.id</a>	<1 %
	Internet Source	

---

47	<a href="http://prh.hec.gov.pk">prh.hec.gov.pk</a>	<1 %
	Internet Source	

---

48	<a href="http://repository.maseno.ac.ke">repository.maseno.ac.ke</a>	<1 %
	Internet Source	

---

49	<a href="http://repository.ruforum.org">repository.ruforum.org</a>	<1 %
	Internet Source	

---

50	<a href="http://ris.cdu.edu.au">ris.cdu.edu.au</a>	<1 %
	Internet Source	

---

51	Neha Kumar, Jody Harris, Rahul Rawat. "If They Grow It, Will They Eat and Grow? Evidence from Zambia on Agricultural Diversity and Child Undernutrition", The Journal of Development Studies, 2015	<1 %
	Publication	

---

52	S. A. Adegoke, O. A. Oyelami, O. S. Olatunya, L. A. Adeyemi. "Effects of Lime Juice on Malaria Parasite Clearance", Phytotherapy Research, 2011	<1 %
	Publication	

---

53	Sossa-Vihotogbe, ANC, BV Anihouvi, HN Akissoe, LG Amadji, and DJ Hounhouigan. "Effect of organic and mineral fertilization on	<1 %
----	---	------



proximate composition of three leafy vegetables harvested at different periods", International Journal of Biological and Chemical Sciences, 2013.

Publication

54	<a href="https://assets.researchsquare.com">assets.researchsquare.com</a> Internet Source	<1 %
55	<a href="https://cgspace.cgiar.org">cgspace.cgiar.org</a> Internet Source	<1 %
56	<a href="https://core-cms.prod.aop.cambridge.org">core-cms.prod.aop.cambridge.org</a> Internet Source	<1 %
57	<a href="https://core.ac.uk">core.ac.uk</a> Internet Source	<1 %
58	<a href="https://curis.ku.dk">curis.ku.dk</a> Internet Source	<1 %
59	<a href="https://docslib.org">docslib.org</a> Internet Source	<1 %
60	<a href="https://doi.org">doi.org</a> Internet Source	<1 %
61	<a href="https://etheses.whiterose.ac.uk">etheses.whiterose.ac.uk</a> Internet Source	<1 %
62	<a href="https://opus.uni-hohenheim.de">opus.uni-hohenheim.de</a> Internet Source	<1 %
63	<a href="https://pmc.ncbi.nlm.nih.gov">pmc.ncbi.nlm.nih.gov</a> Internet Source	<1 %
64	<a href="https://repository.out.ac.tz">repository.out.ac.tz</a> Internet Source	<1 %
65	<a href="https://reserds.vsu.edu.ph">reserds.vsu.edu.ph</a> Internet Source	<1 %
66	<a href="https://www.nutriweb.org.my">www.nutriweb.org.my</a> Internet Source	<1 %

67

Godwin Opoku Agyemang, Samuel Selorm Attu, Reginald Adjetey Annan, Satoru Okonogi, Takeshi Sakura, Odeaf O Asamoah-Boakye. "Factors associated with food consumption and dietary diversity among infants aged 6–18 months in Ashanti Region, Ghana", PLOS ONE, 2023

Publication

&lt;1 %

68

Tewodros Getaneh Alemu, Masresha Asmare Techane, Chalachew Adugna Wubneh, Nega Tezera Assimamaw et al. "Spatial variation and determinates of dietary diversity among children aged 6–23months in Ethiopia: spatial and multilevel analysis using Ethiopian Demography Health Survey (EDHS) 2019", Archives of Public Health, 2022

Publication

&lt;1 %

69

Zhen Yan, Xiangyi Xiao, Jingjing Jiao, Wen Lin. "How does agricultural production diversity nourish household dietary diversity? Evidence from China", Global Food Security, 2024

Publication

&lt;1 %

70

Fikre Moga Lencha, Zenebe Jebero Zaza, Lankamo Ena Digesa, Tegegn Mulatu Ayana. "Minimum dietary diversity and associated factors among children under the age of five attending public health facilities in Wolaita Sodd town, Southern Ethiopia, 2021: a cross-sectional study", BMC Public Health, 2022

Publication

&lt;1 %

71

Mallard, S. R., L. A. Houghton, S. Filteau, A. Mullen, J. Nieuwelink, M. Chisenga, J. Siame, and R. S. Gibson. "Dietary Diversity at 6 Months of Age Is Associated with Subsequent Growth and Mediates the Effect of Maternal

&lt;1 %

Education on Infant Growth in Urban  
Zambia", Journal of Nutrition, 2014.

Publication

72

Ravi Nandi, Swamikannu Nedumaran. "Rural  
Market Food Diversity and Farm Production  
Diversity: Do They Complement or Substitute  
Each Other in Contributing to a Farm  
Household's Dietary Diversity?", Frontiers in  
Sustainable Food Systems, 2022

<1 %

Publication

73

Urbain Zongo. "Effect of Moringa Leaves  
Powder Consumption on Young Children  
Nutritional and Serum Retinol Status in  
Burkina Faso Rural Area", International  
Journal of Nutrition and Food Sciences, 2018

<1 %

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On