Jana Publication & Research

An Adult Presentation of Ebstein's Anomaly of Tricuspid Valve: **Case Report and Review of Literature**

Ê 08

BioTech

Institut Seni Indonesia Surakarta

Document Details

Submission ID 13 Pages trn:oid:::1:3169691771 2,151 Words Submission Date Mar 1, 2025, 12:44 PM GMT+7 Download Date Mar 1, 2025, 3:55 PM GMT+7 File Name IJAR-50445.docx

File Size

2.0 MB

13,121 Characters



32% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- Bibliography
- Quoted Text

Match Groups

- 39 Not Cited or Quoted 21% Matches with neither in-text citation nor quotation marks
- **91 Dissing Quotations 11%** Matches that are still very similar to source material
- 0 Missing Citation 0% Matches that have quotation marks, but no in-text citation

O Cited and Quoted 0% Matches with in-text citation present, but no quotation marks

latches with m-text citation present, but no quotation marks

Top Sources

- 20% (Internet sources
- 32% 🔳 Publications
- 6% **L** Submitted works (Student Papers)

Match Groups **Top Sources 39** Not Cited or Quoted 21% 20% Internet sources Matches with neither in-text citation nor quotation marks 32% Publications **99 21** Missing Quotations 11% Submitted works (Student Papers) 6% Matches that are still very similar to source material = 0 Missing Citation 0% Matches that have quotation marks, but no in-text citation • 0 Cited and Quoted 0% Matches with in-text citation present, but no quotation marks **Top Sources** The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

	Publication		
"Cardio	vascular Medicine	e", Springer Science and Business Media LLC, 2007	3%
2	Internet		
www.nc	bi.nlm.nih.gov		3%
3	Internet		
emedici	ne.staging.meds	cape.com	2%
4	Publication		
Osami H	lonjo, Amine Maz	ine, Lindsay R. Freud. "Chapter 10 Ebstein's Anomaly", S	2%
5	Publication		
Pediatri	c and Congenital	Cardiology Cardiac Surgery and Intensive Care, 2014.	1%
6	Internet		
link.spri	inger.com		1%
7	Student papers		
Aspen U	Jniversity		<1%
8	Internet		
www.st	atpearls.com		<1%
9	Publication		
Ilkun Pa	ark, Tae-Gook Jun	, Ji-Hyuk Yang, I-Seok Kang, June Huh, Jinyoung Song, Ok	<1%
10	Publication		
"Transe	sophageal Echoca	ardiography for Pediatric and Congenital Heart Disease",	<1%

11 Publication	
Celermajer, D.S "Ebstein's anomaly: Presentation and outcome from fetus to ad	<1%
12 Student papers	
Central Piedmont Community College	<1%
13 Publication	
Heidi M. Connolly. "Ebstein's Anomaly", Adult Congenital Heart Disease, 04/17/2009	<1%
14 Publication	
Timothy Gardner, Thomas L. Spray. "Operative Cardiac Surgery, Fifth edition", CR	<1%
15 Internet	
medilib.ir	<1%
16 Internet	
static1.1.sqspcdn.com	<1%
17 Internet	
www.verywellhealth.com	<1%
18 Publication	
"Practical Manual of Tricuspid Valve Diseases", Springer Science and Business Me	<1%
19 Publication	
Aleksandra Cieplucha, Olga Trojnarska, Agnieszka Bartczak-Rutkowska, Anna Koc	<1%
20 Publication	
Critical Care of Children with Heart Disease, 2010.	<1%
21 Student papers	
University of South Australia	<1%
22 Internet	
iris.unisr.it	<1%
23 Internet	
www.jscimedcentral.com	<1%
24 Publication	
Akira Shiina, James B. Seward, William D. Edwards, Donald J. Hagler, Abdul J. Tajik	<1%

Page 5 of 19 - Integr	ty Overview
-----------------------	-------------

25	Publication		
Anatom	ic Basis of Echoca	rdiographic Diagnosis, 2011.	<1%
	Dublication		
Massim	iliano Cantinotti	"Role of magnetic resonance imaging in different ways	<1%
1018351111		Kole of magnetic resonance imaging in unreferit ways	\$170
27	Internet		
journalı	mc.org		<1%
20	Internet		
	rc abaiournals or		<1%
	c.anajournais.org	3	
29	Publication		
"Cardia	c Surgery", Spring	er Science and Business Media LLC, 2017	<1%
	B. L.P. et al.		
Jian Ean	Publication	llans "Intracardias Echocardiographic Imaging in Padio	<106
Jian-rai	ig Kell, David J. Ca		<170
31	Publication		
Joseph /	A. Dearani. "Surgi	cal treatment of Ebstein's malformation: state of the art	<1%
	Dublication		
32 Karl M	Publication	lang Susan Cooko David Hall Mark S Bloiwois "The Sur	<106
	Reyes, Enci-Hun	eng, susan cooke, bavia nan, mark s. bierweis. The sul	<170
33	Publication		
R Kobza	, D J Kurz, E N Oed	chslin, R Prêtre, M Zuber, P Vogt, R Jenni. "Aberrant tendi	<1%
34 Stulek	Publication	non-ment of Chotoin's Anomaly," Sominaws in They are a	<104
Stulak,	I.M Surgical Ma	nagement of Ebstein's Anomaly , Seminars in Thoracic a	<1%
35	Internet		
epdf.tip	S		<1%
36	Publication		
"Clinica	Atlas of Cardiac	and Aortic CT and MRI", Springer Science and Business	<1%
37	Publication		
"Critica	Care of Children	with Heart Disease", Springer Science and Business Med	<1%
38	Publication		
"The Na	tural and Modife	d History of Congenital Heart Disease", Wiley, 2003	<1%





Mathias Possner, Francisco J. Gensini, David C. Mauchley, Eric V. Krieger, Zachary ... <1%



Thomas L. Spray, Michael A. Acker. "Operative Cardiac Surgery", CRC Press, 2019 <1%

An Adult Presentation <mark>of Ebstein's Anomaly</mark> of Tricuspid Valve: Case Report and Review of Literature.

9 Introduction:

21

Ebstein's anomaly is a rare congenital heart condition characterized by a combination of tricuspid
 valve malformation with right ventricular myopathy [1]. It is frequently associated with left-sided
 heart anomalies[2] and excitability disorders. It accounts for less than 1% of all cases of congenital
 heart disease. The main pathophysiological mechanism is a failure of delamination during

- 6 embryogenesis, leading to downward displacement of the functional annulus and division of the right ventricle into a functional and an atrialized ventricle. Its evaluation requires a series of tests for anatomical and clinical classification, assessing disease severity, and ultimately guiding treatment.
- Prognosis and management depend on the size of the functional right ventricle, tricuspid valve anatomy, and left ventricular function.

Case presentation:

We present a clinical case of a 30-year-old female with a recent medical history of hypertension on diuretic medication. The patient was admitted for scheduled hospitalization for the management of severe palpitations, reporting three episodes of dizziness without syncope. At her admission to our service, the patient was stable; slight cyanosis of the lips was noted, with peripheral oximetry around 89%, blood pressure at 117/74 mmHg, and heart rate at 140 bpm. The cardiovascular examination revealed a regular rhythm with no additional sounds or murmurs. The electrocardiogram displayed a regular sinus rhythm with a left deviation of the heart axis and a positive delta wave in the QRS complex.

On a biological level, we note polycythemia with hemoglobin at 17 g/dl without any other biological abnormalities.

The transthoracic echocardiography (TTE) and transesophageal echocardiography (TOE) showed a severe type C Ebstein's anomaly with major predominance of an atrialized right ventricle and a small functional right ventricle. A dilated, non-obstructive right ventricular outflow tract (RVOT) was also noted. A restrictive ventricular septal defect (VSD) and a small atrial septal defect (ASD) of less than 5 mm were found with minimal tricuspid regurgitation (Figure *1A*, *1B*, *1C*)



FIGURE 1: Figure 1: (1A) 2D Transoesophageal Echocardiography (TOE) image showing displacement of the septal leaflet (blue arrow) towards the apical right ventricle, (1B) 2D TOE image showing dilated right ventricular outflow (green arrow), (1C) 3D TOE image showing tunnel- shaped apical portion of the right ventricle (red arrow)

The right heart catheterization demonstrated quite normal pulmonary artery pressures and a low end- diastolic left ventricular pressure. It also noted a high right atrial pressure at 15 mmHg, systemic
 venous desaturation at 85% (Table 1), a low cardiac index at 1.5 l/min/m², and normal pulmonary vascular resistance (PVR) (Table 2).

			ulmonary Artery			LV			Aorta		
	Right Atrium	Systol ic	Diastol ic	Midd le	Capillar y	Systol ic	Diastol ic	Mid dle	Systoli c	Diastol ic	Middl e
Pressures (mmHg)	15	29	14	19	15	140	3	14	143	76	95
Saturations (%)	59%	72%				85%			85%		

Table 1 : Pressures and saturations measured during right heart catheterization

30

31

	Systemic flow (L/mn)	Pulmonary Flow (L/mn)	Qp/Qs	Cardiac Index (I/mn/m ²)
Flow	2,78	3,1	1,16	
Resistance	Systemic Vascular Resistance (SVR) (UW)	Pulmonary Vascular Resistance (PVR) (UW)	PVR/SVR	1,56
	29,13	1,29	0,04	

TABLE 2: Flow and resistances measured during right heart catheterization

During the examination, several episodes of supraventricular and ventricular tachycardias, such as atrial and ventricular tachycardia occurred. The angiography injections revealed a significantly enlarged atrialized right ventricle with a dilated pulmonary artery (Figure 2).



36 FIGURE 2: Injection for angiography in anteroposterior view showing a dilated pulmonary artery (red arrow).

⁵ The final decision of the heart team was to perform an electrophysiological study due to the severe episodes of arrhythmias documented on the 24-hour electrocardiogram. Our patient then underwent tricuspid valve repair with a cone reconstruction using the Da Silva technique. Upon coming off bypass and due to elevated right atrial filling pressures, a partial cavopulmonary diversion was performed due to the favorable anatomy of the pulmonary vasculature.

Discussion:

- Page 10 of 19 Integrity Submission
- 22 The Ebstein anomaly occurs during the phase of heart development between the 7th and 12th
- 39 week of intrauterine life [3]. It results from incomplete tricuspid valve delamination from the
- 12 right endo- myocardium, leading to adhesion of the septal and posterior leaflets to the
- ²⁵ myocardium with fenestration [4] and tethering of the anterior leaflet [5]. A lack of coaptation of
- the tricuspid leaflets is described, causing tricuspid regurgitation that increases with annular
- 6 dilatation [6]. Also, displacement of the tricuspid annulus is reported [7], resulting in splitting of
- the right ventricle into two portions: the "atrialized" right ventricle (aRV), which acts as a passive
 tank during atrial contraction, and the "functional" right ventricle (fRV), which can become
- dysfunctional [8] due to severe tricuspid regurgitation.
- 2 Some pathologies are commonly associated with Ebstein's anomaly; the most frequent are ASD or
- Patent Foramen Ovale (PFO) [9], VSD, or pulmonary stenosis or atresia [10].
- 8 Ebstein's anomaly can present at any age, but a significant number of patients remain asymptomatic and present in adulthood. The most frequent manifestations among adult
- patients are cyanosis, right heart failure, arrhythmias, and sudden cardiac arrest [11]. Cyanosis is
- usually subtle and is due to right-to-left shunting across the atrial defect. Palpitations result from tachyarrhythmias and are the first manifestation among adult patients, whereas sudden cardiac arrest is caused by ventricular arrhythmias [12].
- 8 Cardiac auscultation shows a triple or quadruple rhythm with a loud first heart sound resulting from anterior leaflet closure; a soft holosystolic murmur of tricuspid regurgitation is sometimes audible
 6 [13]. Electrocardiogram findings typically include right bundle branch block (RBBB) associated with right atrial enlargement represented by tall P waves called "Himalayan P waves." Frequently, excitability disorders are found, such as Delta waves due to an accessory pathway or first-degree atrioventricular block [14], which are integral to Ebstein's disease.
- In echocardiography, Ebstein's anomaly is defined as displacement of the tricuspid leaflet greater than 8 mm/m2 from the insertion of the anterior mitral leaflet [15]. Color doppler helps assess the
 degree and site of tricuspid regurgitation. Tethering of the tricuspid valve can also be present [16]. Assessment of both the aRV and the fRV size is essential, especially the RVOT size. It should be noted that the smaller the size of the fRV, the worse the prognosis of the disease. Other important elements to evaluate are left ventricular function and the search for other anomalies often associated with Ebstein's disease. TOE is more efficient in adults in cases of poor echogenicity; intracardiac shunts are better visualized.

Several classification systems exist for Ebstein anomaly, with the Carpentier classification [17] (Figure 3) being the most commonly used, focusing on anatomical categorization. Its initial
assessment relies on echocardiography, where anomalies are graded as mild, moderate, or severe based on the degree of leaflet displacement, tethering, and right ventricular dilation (Table 3)





17

11

FIGURE 3: Carpentier's Classification of Ebstein's Anomaly

Carpentier's classification of Ebstein's anomaly based on the anatomical description of the anomaly [18].

	Type A : Mild	Type B : Moderate	Type C : Severe	Type D :Extremely severe
Anterior leaflet (attachment -	Normal morphology	Abnormal attachments	Partial adhesions to the venticular free wall	Severe adhesions to the ventricular wall
mobility)	Normal mobility	Normal mobility	Restricted motion	-
'Atrialized' ventricle (size - function)	Small ventricle	Moderate- sized ventricle	Large ventricle	Very large ventricle
(,	Preserved function	Reduced function	Reduced function	Minimal function
Functional ventricle (size- function)	Normal- sized ventricle	Reduced ventricle	Small ventricle	Infundibulum
, ,	Preserved function	Preserved function	Reduced function	Depressed function

TABLE 3: Description of the Carpentier Classification of Ebstein's Anomaly

Ebstein's classification into 4 types based on the attachments of the tricuspid valve, the size and function of the atrialized and functional right ventricle[18].

- 26 Cardiac magnetic resonance imaging (CMR) allows for a detailed analysis of tricuspid valve anatomy and accurate assessment of right ventricular function, but it is not used as a first-line approach.
- Right heart catheterization guides surgical treatment. Left ventricular function, anatomy of the pulmonary artery tree, and pulmonary pressures are important parameters for palliative surgery.

If no surgical intervention can be proposed, medical treatment should be optimal, with effective management of heart failure. In cases of significant thromboembolic risk, oral anticoagulation may be considered. Management of arrhythmias can use medical treatment if symptomatic. Assessment of arrhythmia substrates by catheter-based intervention [19] or even catheter ablation can sometimes be considered before surgery.

- The principle of surgical treatment is either biventricular repair , a single ventricle pathway with right ventricular exclusion [20], or heart transplantation. Biventricular repair involves
- 1 reconstruction of the tricuspid valve, sometimes with repair of the ASD. Single ventricle repair is recommended for patients with inadequate right ventricle size and is considered a palliative
- procedure. Cardiac transplantation is reserved for severe forms with significant left ventricular dysfunction.

For surgical treatment in this case, tricuspid valve repair was the first-line approach considered,
 with a Cone repair using the Da Silva technique. These method involve some specific steps (Figure 4)



FIGURE 4: Cone reconstruction of tricuspid valve in Ebstein anomaly : The Da Silva Technique Steps of Da Silva Technique : [21]

A. The right atrium is opened, revealing displacement of the tricuspid valve.

5

TTA: true tricuspid annulus; ASD: atrial septal defect; CS: coronary sinus.

B. A portion of the anterior and posterior leaflet is detached, creating a single unit.

C. The posterior leaflet edge is rotated clockwise and sutured to the septal edge of the anterior leaflet. The true tricuspid annulus is then plicated.

D. The valve is completely reattached to the true tricuspid annulus, and the atrial septal defect is closed with a valve.

In this particular case, due to advanced age and tricuspid annular dilation, augmentation with a prosthetic ring could have been necessary if valve competence was inadequate.

- Upon separation from cardiopulmonary bypass (CPB), the right atrial filling pressures will be monitored. Elevated pressures may necessitate a partial cavopulmonary diversion, achieved by
 suturing the superior vena cava directly to the right pulmonary artery. In our case this technique was performed given the favorable hemodynamics for potential cavopulmonary diversion.
- Prognosis depends on the degree of tricuspid displacement, severity of tricuspid regurgitation, size of the fRV, and degree of right ventricular dysfunction.

Conclusion:

- 27 Ebstein's disease is a rare congenital heart condition typically characterized by
- tachyarrhythmia. Echocardiography serves as the primary diagnostic tool, confirming the condition
 and aiding in severity assessment. Evaluation necessitates meticulous examination of the size of
 the atrialized and functional right ventricles, along with an assessment of left ventricular function,

in order to guide surgical management.

References:

- Dearani JA, Danielson GK: Congenital heart surgery nomenclature and database project: Ebstein's anomaly and Tricuspid valve disease. Ann Thorac Surg. 2000, 69:106-17. 10.1016/s0003-4975(99)01265-5
- Attenhofer Jost CH, Connolly HM, O'Leary PW, et al. Left heart lesions in patients with Ebstein anomaly. Mayo Clin Proc. 2005, 80:361-8. 10.4065/80.3.361
- 3. Lamers WH, Virágh S, Wessels A, et al.: Formation of the Tricuspid valve in the human heart . Circulation. 1995, 91:111-21. 10.1161/01.CIR.91.1.111
- Holst KA, Connolly HM, Dearani JA: Ebstein's Anomaly. Methodist Debakey Cardiovasc J. 2019, 15:138-144. 10.1016/j.athoracsur.2017.09.058
- 5. Anderson KR, Zuberbuhler JR, Anderson RH, Becker AE, Lie JT: Morphologic spectrum of Ebstein's anomaly of the heart: a review. Mayo Clin Proc. 1979, 54:174-80.
- S. A. Usta, A. I. Yıldırım, A. T. Karaağaç, and K. Kirali: 'Ebstein's Anomaly', Structural Insufficiency Anomalies in Cardiac Valves. 6, 03-2018:10.5772/intechopen.78067. 10.5772/intechopen.78067
- Fuchs MM, Connolly HM: Ebstein Anomaly in the Adult Patient . Cardiol Clin. 2020, 38:353-363. 10.1016/j.ccl.2020.04.004
- Qureshi MY, O'Leary PW, Connolly HM: Cardiac imaging in Ebstein anomaly. Trends Cardiovasc Med. 2018, 28:403-409. 10.1016/j.tcm.2018.01.002
- Danielsen GK, Driscoll DJ, Mair DD, Warnes CA, Oliver WC Jr: Operative treatment of Ebstein's anomaly. The. Journal of Thoracic and Cardiovascular Surgery. 1992, 104:1195-1202. 10.1016/S0022-

Page 14 of 19 - Integrity Submission 5223(19)34605-7

- Jost CH A, Connolly HM, Edwards WD, Hayes D, Warnes CA, Danielson GK: Ebstein's anomaly: Review of a multifaceted congenital cardiac condition. Swiss Medical Weekly. 2005, 135:269-281.
- 11. Yuan S-M: Ebstein's anomaly: Genetics, clinical manifestations, and management . Pediatrics and Neonatology. 2017, 58:211-215. 10.1016/j.pedneo.2016.08.004
- 12. Attenhofer Jost CH, Tan NY, Hassan A, et al.: Sudden death in patients with Ebstein anomaly . Eur Heart J. 2018, 01-39. 10.1093/eurheartj/ehx794
- 13. Ciepłucha A, Trojnarska O, Kociemba A, et al.: Clinical aspects of myocardial fibrosis in adults with Ebstein's anomaly. Heart Vessels. 2018, 33:1076-1085. 10.1007/s00380-018-1141-5
- 14. Attenhofer Jost CH, Connolly HM, Dearani JA, Edwards WD, Danielson GK: Ebstein's anomaly. Circulation. 2007, 16:277-85. 10.1161/CIRCULATIONAHA.106.619338
- Shiina A, Seward JB, Edwards WD, Hagler DJ, Tajik AJ: Two-dimensional echocardiographic spectrum of Ebstein's anomaly: detailed anatomic assessment. J Am Coll Cardiol. 1984, 3:356-70. 10.1016/S0735-1097(84)80020-0
- Roberson DA, Silverman NH: Ebstein's anomaly: echocardiographic and clinical features in the fetus and neonate. J Am Coll Cardiol. 1989, 14:1300-1307. 10.1016/0735-1097(89)90432-4
- 17. Alain Carpentier, Sylvain Chauvaud, Loïc Macé, et al.: A new reconstructive operation for Ebstein's anomaly of the tricuspid valve. 1988, 96:92-101. 10.1016/S0022-5223(19)35302-4.
- Alsaied T, Christopher AB, Da Silva J, et al.: Multimodality Imaging in Ebstein Anomaly. 2022, 44:1-9. 10.1007/s00246-022-03011-x
- Khositseth A, Danielson GK, Dearani JA, Munger TM, Porter CJ: Supraventricular tachyarrhythmias in Ebstein anomaly: management and outcome. J Thorac Cardiovasc Surg. 2004, 128:826-33.
 10.1016/j.jtcvs.2004.02.012
- Knott-Craig C.J., Overholt E.D., Ward K.E., et al.: Repair of Ebstein's anomaly in the symptomatic neonate: an evolution of technique with 7-year follow-up. Ann Thorac Surg. 2002, 73:1786-1792.
 10.1016/S0003- 4975(02)03507-5
- 21. da Silva JP, Baumgratz JF, da Fonseca L, et al.: The cone reconstruction of the tricuspid valve in Ebstein's anomaly. The operation: early and midterm results. . 10.1016/j.jtcvs.2006.09.01



Page 15 of 19 - Integrity Submission



Page 16 of 19 - Integrity Submission







Page 18 of 19 - Integrity Submission



