

1                   **CONTRIBUTION TO THE STUDY OF THE EFFECTS OF VARIOUS**  
2                   **ADDITIVES OF ANIMAL ORIGIN ON THE GROWTH OF HYBRID CLARIAS**  
3                   **FRY (MALE *CLARIAS GARIEPINUS* X FEMALE *CLARIAS ANGUILLARIS*)**  
4  
5  
6

7                   **Abstract**

8                   Aquaculture in Senegal faces constraints such as the high cost of fish feed, representing  
9                   a brake on the sector's development. In this study, alternative local meals (cow blood  
10                  meal, chicken viscera meal and black soldier fly larvae meal) were used as feed  
11                  additives for hybrid fry of *Clarias anguillaris* and *Clarias gariepinus*. Four  
12                  experimental diets (R1, R2, R3 and R4) were formulated, in addition to an industrial  
13                  control diet (R0), to assess their effects on growth, feed efficiency and production cost.  
14                  300 fry of initial mean weight of 1.56 g were distributed in 15 tanks of 50 liters. After  
15                  45 days of experimentation, the best zootechnical performance was obtained with diet  
16                  R4, containing the combination of the three additives. This diet showed a relative mean  
17                  weight gain (RMWG) of 521.58% and a feed conversion ratio (FCR) of 1.44, similar to  
18                  the control diet. Economically, the R4 diet had the lowest feed cost of 666.41 FCFA/kg  
19                  and fish cost of 958.94 FCFA/kg, while maintaining a high survival rate (95%). These  
20                  results demonstrate that the use of local ingredients as additives is a sustainable and  
21                  cost-effective alternative for improving the productivity and competitiveness of  
22                  aquaculture in Senegal.

23                  **Keywords:** Cow blood meal, chicken viscera, black soldier fly larvae, growth, feed  
24                  efficiency, cost

25  
26                  **Introduction**

27                  Fishery plays a considerable socio-economic role in Senegal. However, given the  
28                  difficulties associated with effective fisheries management and the growing demand for  
29                  fisheries resources, Senegal has been faced for several decades with the effects of  
30                  overfishing of its ecosystems, climate change and the collapse of fisheries, impacting

local consumption of fisheries production, food security and economic growth (Baldé, 2019). Faced with this situation, the Senegalese government has been committed since 2007 to promote aquaculture as an alternative way of coping with dwindling fisheries resources (Ndiaye, 2019).

Aquaculture plays a crucial role in food security and economic development, particularly in developing countries. However, one of the major obstacles to its expansion remains the high cost of fish feed, representing between 40% and 60% of total production costs (Siddhuraju and Becker, 2003). This dependence on imported feed limits farm profitability and hinders the development of high-potential species such as *Clarias gariepinus* and *Clarias anguillaris*, renowned for their rapid growth and adaptability. In this context, the use of local by-products to formulate less costly diets represents a sustainable and economically viable alternative.

In Senegal, products such as chicken viscera, cow blood and black soldier fly larvae are often under-exploited. Slaughterhouses discharge their effluents, including blood, into drainage channels without pre-treatment, contributing to significant environmental pollution. These by-products, which are rich in essential nutrients, nonetheless offer potential for valorization. Blood, for example, contains high-quality proteins, notably albumin and hemoglobin, and its recovery reduces the biochemical oxygen demand (BOD) of wastewater, thus limiting the degradation of aquatic ecosystems (WHO, 2023). Chicken viscera, also rich in proteins, fats and amino acids, are often regarded as waste, although they can be processed into meal for animal feed. Black soldier fly larvae, on the other hand, are rich in protein and essential fatty acids, and have the advantage of being produced from recycled organic matter, making them an ecological and sustainable source.

The general aim of this study was to evaluate the effects of these ingredients, used as feed additives, on the growth of catfish hybrid fry of male *Clarias gariepinus* and female *Clarias anguillaris*. More specifically, it aims to:

- add value to cow blood, broiler viscera and black soldier fly larvae by incorporating them as additives in *Clarias* hybrid feeds;
- determine the zootechnical performance of *Clarias* hybrids;

- determine the nutritional contribution of foodstuffs on the flesh of *Clarias* hybrids;
- determine variations in rearing water temperature and pH during experimentation;
- determine the production cost of feed and fish.

## **Material and methods**

### **Methods**

#### **Processing of some ingredients used in this study**

Various methods were used in this study. These included processing of raw materials into finished products, feed formulation and manufacture, rearing conditions and determination of zootechnical parameters of *Clarias* hybrid fry.

#### **Blood meal production**

Fresh blood obtained from the SOGAS slaughter house in Dakar was placed in a 20L plastic container and transported to the IUPA aquaculture station. To sterilize the fresh blood, it was heat-treated by boiling at 120 degrees Celsius for a quarter of an hour, then sun-dried for 48 hours and finely ground for use in diets. The blood meal was stored in glass jars until use.

#### **Black soldier fly meal production**

Black soldier fly larvae were harvested when they reached an optimal stage of development, generally after 7 to 14 days of rearing.

After harvesting, the larvae were immobilized with hot water. Once treated, they were spread out in the sun to dry and passed through a press to extract the oil. Finally, the dried and defatted larvae were ground to obtain a fine flour.

### Chicken viscera meal production

Chicken guts obtained fresh from the market were meticulously washed and rinsed. This step was crucial to remove impurities, food residues and contaminants, thus ensuring the cleanliness of the intestines prior to processing. After this stage, they were then subjected to heat treatment by boiling at 100 degrees Celsius for a quarter of an hour. After draining, the viscera were spread out in the sun to dry. Once dried, they were ground into a fine flour and preserved in glass jars.

### Diets formulation

Four experimental diets were formulated using the Pearson square method, and an industrial feed was used as a control. Table I provides information on the centesimal composition of the different ingredients used and the composition of the different ingredients in the industrial feed, as well as their bromatological compositions.

**Table I:** Centesimal composition of the four formulated diets

Ingredients	R1 (viscera)	R2 (viscera + blood)	R3 (viscera +BSF)	R4 (viscera + blood +BSF)	Industrial food
Fish meal	35	35	35	35	Fish meal, Soya cake meal, Wheat meal, Soya oil, Maize meal,
Soybean meal	10	10	10	10	
Peanut meal	10	10	10	10	
Wheat bran meal	10	10	10	10	
Corn flour	10	10	10	10	
Cassava flour	9	9	9	9	Rice bran,

<b>Chicken viscera meal</b>	5	2,5	2,5	1,66	Distillers Dried Grains with Solubles (DDGS) and Maize gluten
<b>Cow's blood meal</b>	0	2,5	0	1,67	
<b>BSF larvae meal</b>	0	0	2,5	1,67	
<b>Fish oil</b>	5	5	5	5	
<b>Premix vitamins<sup>a</sup></b>	1	1	1	1	
<b>Premix minerals<sup>b</sup></b>	1	1	1	1	
<b>Yeast</b>	4	4	4	4	
<b>Total</b>	100	100	100	100	
<b>Crude Protein (%)</b>	37,5	37,5	37,5	37,5	37
<b>Crude lipid (%)</b>	11,8	11,37	11,43	11,44	8
<b>Cost of feed (F CFA)</b>	657,22	654,72	691,43	666,415	1050

a=vit A 250000 UI; vit D3 250000UI; vit E 5000mg; vit B1 100mg; vit B2 400mg; vit B3(pp) 1000mg; vit B5 pantode Ca2000mg; vit B6 300mg; vit K3 1000g; vit C 5000mg; H biotin 15mg; choline 100g; anti-oxydant (BHT), crushed and calcined attapulgitite qs 1000mg;

b=phosphorus 7%; calcium 17%; sodium 1,5%; potassium 4,6%; magnesium 7,5%; manganese 738mg; zinc 3000mg; iron 4000mg; copper 750mg; iodine 5mg; cobalt 208mg; calcined and ground attapulgitite qs 1000g; fluorine 1.5% (approximately).

## Diets manufacturing

The different proportions of ingredients shown in Table III are formulated to produce 1000g of feed. The raw materials for each feed were sieved, weighed and mixed to homogenize them. Water was then added at a rate of 30% of dry matter. This resulted in a malleable dough mixture. This mixture was passed into a Moulinex mincer, from which it emerged in the form of spaghetti filaments. These filaments then underwent fragmentation, followed by sun-drying for two days. Once dried, they were placed in a mortar, then crushed and sieved to obtain flour, which was then preserved in bags.

## Breeding conditions

The hybrids used in this experiment were derived from the artificial reproduction of a male *Clarias gariepinus* and a female *Clarias anguillaris*. Three days after hatching, the larvae were fed artemia SEP for 15 days. During the course of the experiment, 300 fry with an initial average weight of 1.56 g were randomly distributed in 15 aquariums of 50-liter, i.e. 20 fry per tank. These fish were acclimatized for one week.

The experiment lasted 45 days. The fish received a daily ration corresponding to 8% of their live weight. This ration was distributed in the morning at 8 a.m. and in the evening at 5 p.m., after temperature and pH measurements had been taken.

Finally, to maintain water quality, the bottom of the rearing tanks was siphoned off each morning before feeding. The edges were cleaned and rinsed thoroughly to limit the deposit of any fatty matter left by the feed.

## Evaluation of survival growth parameters, feed efficiency and feed costs

Growth checks were carried out early in the morning after every two weeks of the experiment. Using a landing net, the fish were transferred to a plastic pot placed on a scale. After weighing, the fry were returned to the carefully cleaned aquariums. The results obtained were then entered into an Excel spreadsheet to determine zootechnical parameters for assessing fish growth and feed utilization efficiency. The various zootechnical parameters used as indicators in this experiment are:

**Specific Growth rate (SGR, %/d)** =  $(\ln(\text{final average weight}) - \ln(\text{initial average weight})) / \text{experiment duration} \times 100$ .

**Survival Rate (SR, %)** =  $\text{number of final fish} / \text{number of initial fish} \times 100$ .

**Feed Conversion Ratio (FCR)** =  $\text{Feed intake} / \text{weight gain}$ .

**Absolute Mean Weight Gain (AMWG, g/fish)** =  $\text{average final weight} - \text{average initial weight}$

**Relative Mean Weight Gain (RMWG, %)** =  $100 \times ((\text{final average weight} - \text{initial average weight}) / \text{initial average weight})$

**Cost of feed** (FCFA) = Total costs (raw materials, processing, transport) / Quantity of feed produced per kg

**Production cost per kg of fish** (FCFA)= FCR× Feed cost (per kg)

## **Data analysis**

Analysis of statistical differences between groups was carried out using ANOVA and Duncan test, implemented via SAS software, which is a widely used computer program for statistical data analysis and database management. Created in the 1970s by SAS Institute, it is particularly popular in fields as diverse as social sciences, bio statistics. It enabled us to identify homogeneous groups based on their significant differences. Thanks to SAS' advanced features, the averages were automatically grouped into distinct classes, facilitating interpretation of the results. This method proved effective in highlighting significant differences while controlling the risk of Type I error, in line with the study's objectives

## **Results and discussion**

### **Results**

#### **Yield of main ingredients used**

The yields of the main ingredients used as additives are shown in Table II

**TableII :** Yield of main ingredients used as additives

Ingredients	Initial quantity	Final quantity (g)	Yield (%)
Cow's blood	17 L	7000	41,48
Black soldier fly larvae	1.70 kg	465	27,35

Chicken viscera	20 kg	800	4
-----------------	-------	-----	---

Table II shows a significant variation in yields between processed ingredients. Cow's blood shows the highest yield (41.18%), reflecting efficient processing with moderate losses, making it an economically advantageous option. Black soldier fly larvae, with a yield of 27.35%, show losses linked to their high water content, but remain attractive due to their high nutritional value and ecological impact. Chicken intestines, on the other hand, had a very low yield (4%), indicating significant losses during the process, probably due to a high proportion of unusable material. These results suggest that cow's blood and fly larvae should be preferred, while chicken intestine processing needs to be optimized or alternative uses considered to reduce losses and improve recovery

#### **Biochemical composition of main ingredients**

The various ingredients used, in particular cow blood meal, chicken intestine meal and black soldier fly larvae meal, were sent to LANAC for analysis using the AOAC, 1995 method. The results of the biochemical composition are presented in Table III.

**Table III:** Biochemical composition of blood meal from Black Soldier Fly (BSF) larvae and chicken viscera meal

<b>Biochemical composition</b>	<b>Blood meal</b>	<b>BSF meal</b>	<b>Chicken viscera meal</b>
<b>Fat content (%)</b>	1,97	14,32	19,33
<b>Protein (%)</b>	93,81	32,37	61,25

#### **Water quality parameters (Temperature and pH variations)**

**TableIV :** Changes in temperature and pH during experiment

<b>Parameters</b>	<b>R0</b>	<b>R1</b>	<b>R2</b>	<b>R3</b>	<b>R4</b>
<b>Temperature</b>	28,61±0,005 <sup>a</sup>	28,42±00 <sup>c</sup>	28,45±0,01 <sup>c</sup>	28,45±0,03 <sup>c</sup>	28,50±0,04 <sup>b</sup>



<b>pH</b>	7,83±0,005 <sup>a</sup>	8,04±0,02 <sup>b</sup>	7,99±0,01 <sup>c</sup>	7,96±0,02 <sup>d</sup>	7,93±0,01 <sup>e</sup>
-----------	-------------------------	------------------------	------------------------	------------------------	------------------------

Table IV shows the results of statistical analyses of water parameters, namely temperature and pH, for five different feeding diets (R0, R1, R2, R3, R4). Values shown include means and their standard deviations, while letters (a, b, c, d, e) show significant differences between treatments.

For temperature, diet R0 (28.61) had a significantly higher temperature than all other diets. Diets R1, R2, and R3 (28.42; 28.45; 28.45°C) had statistically similar temperatures. Diet R4 (28.50°C) was slightly higher than R1, R2, and R3, but still significantly different from R0 and the others.

In terms of pH, diet R0 (7.83) had the lowest pH and was significantly different from the other diets. Diet R1 (8.04) showed the highest pH and was significantly different from all other treatments. Diets R2 (7.99) and R3 (7.96) had close but statistically distinct values. Diet R4 (7.93) has the lowest pH after R0, but remains significantly different from the others.

### Zootechnical parameters

During the experiment, various zootechnical parameters were studied to determine fish growth in each diet. These results are shown in Table V.

**TableV:** Growth, feed efficiency and survival parameters of catfish hybrids.

Parameters	R0	R1	R2	R3	R4
<b>IMW g</b>	1,56 <sup>a</sup>	1,56 <sup>a</sup>	1,56 <sup>a</sup>	1,56 <sup>a</sup>	1,56 <sup>a</sup>
<b>FMW g</b>	10,62±0,25 <sup>a</sup>	5,65±0,15 <sup>d</sup>	7,85±0,23 <sup>c</sup>	5,82±0,38 <sup>d</sup>	9,69±0,11 <sup>a</sup>

<b>AMWG g</b>	9,07±0,25 <sup>a</sup>	6,09±3,34 <sup>ba</sup>	6,09±0,49 <sup>ba</sup>	4,27±0,39 <sup>b</sup>	8,13±0,11 <sup>ba</sup>
<b>RMWG%</b>	582,76±16,81 <sup>a</sup>	262,46±17,59 <sup>d</sup>	390,73±31,89 <sup>c</sup>	274,56±25,44 <sup>d</sup>	521,58±7,36 <sup>ba</sup>
<b>FCR</b>	1,45±0,05 <sup>b</sup>	2,41±0,20 <sup>a</sup>	1,66±0,18 <sup>b</sup>	2,28±0,36 <sup>a</sup>	1,44±0,06 <sup>b</sup>
<b>SCR%/J</b>	4,27±0,06 <sup>a</sup>	2,86±0,05 <sup>c</sup>	3,53±0,14 <sup>b</sup>	2,93 ±0,15 <sup>c</sup>	4,06±0,03 <sup>a</sup>
<b>SR%</b>	93,33±5,77 <sup>a</sup>	93,33±2,88 <sup>a</sup>	90±0,10 <sup>a</sup>	85±13,23 <sup>a</sup>	95±5 <sup>a</sup>

**NB:** The letters a, b, c and d show the differences between the results of the statistical analyses.

The IMW value was constant at 1.56 g for all diets (R0 to R4). This indicates that all fish groups had a homogeneous initial weight, which is essential to ensure that the differences observed in the other parameters are not due to variations in initial weight.

With regard to growth parameters (FMW, AMWG, RMWG, SGR), no statistically significant differences were observed between fish fed diets R0 and R4. Diets R1 and R3 showed no significant differences between them, but were both significantly lower than R0 and R4. For SGR, no significant differences were observed between fish fed diets R0 and R4, nor between R1 and R3. On the other hand, fish fed the R2 diet showed statistical differences from R0, R4, as well as R1 and R3.

In terms of feed efficiency (FCR), fish fed with R4 and R0 diets recorded the best FCR with values of 1.44 and 1.45 respectively, followed by those fed with R2 diet (1.66). However, these three diets had no significant statistical difference and were better than R1 (2.41) and R3 (2.28), which had no significant difference. On the other hand, fish fed diets R0, R2, R4 and those fed diets R1(2.41), R3(2.28) had FCR that were statistically different.

With regard to survival rates, the results obtained in this study do not show a significant difference for fish fed the R0, R1, R2, R3 and R4 diets. Fish fed the R3 diet had the lowest survival rate, with a value of 85%.

## Economic analysis

**TableVI** : Economic analysis of tested diets

Parameters	R0	R1	R2	R3	R4
Cost of Kg of feed (FCFA)	1050	657,22	654,72	691,43	666,41
Cost of Kg of fish (FCFA)	1523,91±50,26 <sup>a</sup>	1585,86±137,72 <sup>a</sup>	1087,76±119,76 <sup>b</sup>	1579,70±250,33 <sup>a</sup>	958,94±39,58 <sup>b</sup>

**NB:** The letters a and b show the differences between the results of the statistical analyses.

Table VI shows the economic analysis of the diets tested. For feed costs, diet R0 (industrial feed) has the highest cost (1050 FCFA), while diets R1, R2, R3 and R4 have significantly lower costs (around 657 to 691 FCFA).

Production costs per Kg of fish vary between groups, with R4 (958.94 FCFA) and R2 (1087.76 FCFA) showing the lowest costs and no significant statistical difference. Diets R0, R1, and R3 have the highest costs, exceeding 1500 FCFA and showing no significant statistical difference.

However, R4, R2 and R0, R1, R3 are statistically different.

### III.1.6. Fish flesh analysis

**TableVII** : Results of bromatological analysis of fish flesh

Parameters	Initial fish	R0	R1		R3	R4
Crude Protein (%)	16,81±0,10 <sup>a</sup>	17,09±0,46 <sup>a</sup>	16,82±0,10 <sup>a</sup>	16,58±0,20 <sup>a</sup>	16,74±0,84 <sup>a</sup>	17,05±0,76 <sup>a</sup>
Crude Fat (%)	3,94±0,19 <sup>b</sup>	6,40±0,43 <sup>a</sup>	4,31±0,54 <sup>b</sup>	4,22±0,19 <sup>b</sup>	4,57±0,15 <sup>b</sup>	5,70±0,24 <sup>a</sup>

Ash (%)	3.94±0.14 <sup>bac</sup>	3.49±0.26 <sup>dc</sup>	4.25±0.19 <sup>ab</sup>	4.53±0.13 <sup>a</sup>	3.76±0.22 <sup>dc</sup>	3.28±0.31 <sup>d</sup>
---------	--------------------------	-------------------------	-------------------------	------------------------	-------------------------	------------------------

**NB:** The letters a, b, c and d show the differences between the results of the statistical analyses and

Bromatological results show that crude protein remains constant between groups, with values ranging from 16.58% to 17.09%, with no significant difference. On the other hand, Crude fat content varies significantly: group R1 has the highest content (6.40%), while the other groups (R0, R2, R3 and R4) have lower values, ranging from 3.94% to 4.57%, with no significant statistical difference. In terms of ash, groups R2 and R3 stand out with higher contents (4.25% and 4.53%, respectively), indicating better mineralization. Conversely, groups R4 (3.28%) and R1 (3.49%) show the lowest values, statistically different.

### III.2. Discussion

Temperatures recorded during the experiment ranged from 26.6°C to 30.1°C. These results are in line with those of Alofa *et al.* (2016), who argued that *Clarias* can withstand temperatures as low as 6°C and as high as 50°C. However, a temperature range of 28 to 30° C is considered optimal for the growth of *Clarias gariepinus* and *Clarias anguillaris* (Adebayo *et al.*, 2008).

The pH ranged from 7.11 to 8.94. These results fall within the range reported by Geoffroy *et al.* (2019), who summarized the literature on biological and zootechnical parameters of the African catfish *Clarias gariepinus* and argue that *Clarias gariepinus* survives and thrives best in water with a pH between 6 and 9.

In the present study, the highest growth was observed in diet R4 (a mixture of blood meal, chicken viscera meal and black soldier fly larvae meal).

Mbaye (2023) has shown that blood meal and black soldier fly larvae meal represent promising alternative protein sources in animal feed. Protein-rich blood meal and black soldier fly larvae meal were evaluated for their nutritional benefits, but it was observed that the higher the fishmeal substitution, the higher the FCR. In terms of feed efficiency

(FCR), fish fed the diet containing 5% bloodmeal recorded the best FCR. The best growth performance was also noted in fish fed the diet (5% blood meal). These results are similar to those of Olukunle *et al.* (2002), who reported that optimal growth and nutrient utilization in hybrid Clarias (male *Clarias gariepinus* x female *Heterobranchus longifilis*) were achieved with 5% sun-dried bloodmeal.

Similarly, fly larvae meal, recognized for its durability and nutritional value, has shown significant potential as an additive. Study have suggested that the optimal incorporation rate for this insect meal should not exceed 30%, due to its chitin content (Gougbedji, 2022).

In this study, the R1 diet containing 5% chicken viscera meal had the highest FCR, followed by the R3 diet containing 2.5% chicken viscera meal and 2.5% black soldier fly meal. These results concur with those of Amtul *et al.* (2012) who worked on replacing fish meal with poultry by-product meal (chicken intestine) as a protein source in carp fry diets. Diets containing chicken intestine meal, even at the lowest level, significantly limited growth performance and feed efficiency. Values deteriorated as the incorporation rate of chicken intestine meal increased. This was due to the ash and carbohydrate content of the gut meal, producing a faster intestinal transit rate, leading to increased feed intake, associated with poor growth performance and feed efficiency. Similar results were reported by Ronyai *et al.* (2002) and Fagbenro *et al.* (2004), who revealed that deficiency in one or more amino acids limits protein synthesis and affects fish growth

The R2 diet containing 2.5% blood meal and 2.5% chicken viscera meal has a FCR of 1.66, and these results are in line with those of Milliamena (2002) and Guo *et al.* (2007) who obtained better results when using blood meal to balance lysine content by replacing fishmeal with poultry byproducts (including broiler viscera) used alone or in combination with other ingredients of animal origin. According to Hu *et al.* (2008a), methionine and lysine are present in small quantities in chicken viscera (1.43 and 3.51% respectively). This could account for the low growth achieved in fish diets fed chicken viscera. The use of cow blood meal can correct lysine deficiency due to its richness in this amino acid (Hertrampf and Piedad-Pascual, 2000 and Alofa *et al.*, 2016). However,

methionine deficiency remains. Although poultry by-products such as chicken viscera have proven to be nutritionally adequate protein sources for many fish species (Fowler, 1991; Robaina *et al.*, 1997; Abdel-Warith *et al.*, 2001; Wang *et al.*, 2006), they were deficient in essential amino acids such as lysine, methionine that limited their utilization (Glencross *et al.*, 2007). Therefore, it will be beneficial to adjust the level of essential amino acids in chicken viscera through supplementation. Hu *et al.* (2008b) provided 0.49% lysine and 0.14% methionine as supplementation in diets containing chicken viscera with improved growth performance in *Carassius auratus gibelio*. Similar trends in results were obtained by Rawles *et al.* (2006) in the *Morone chrysops* x *Morone saxatilis* hybrid fed a broiler viscera diet supplemented with methionine (0.63%) and lysine (1.55%)

The results of this study show that the diet containing a mixture of chicken viscera meal, cow blood meal and black soldier fly meal, with a total incorporation rate of 5%, offers the best growth performance for catfish hybrid fry of *Clarias gariepinus* and *Clarias anguillaris*. This success can be attributed to the nutritional complementarity of the three ingredients: chicken viscera meal, rich in digestible proteins and deficient in amino acids including lysine and methionine according to Wang *et al.* (2006), cow blood meal which provides iron-rich proteins beneficial to hematopoiesis (Olukunle *et al.*, 2002) and black soldier fly meal, a sustainable source of essential fatty acids and balanced proteins (Gougbedji, 2022; Henry *et al.*, 2015). The synergistic effect observed with this blend, highlighted by Santiago and Lovell (1988) and Youssouf and Sedro (2020), probably results from the combination of diversified nutritional inputs that optimize nutrient digestibility and assimilation. By keeping the incorporation rate low ( $\leq 5\%$ ), negative effects such as reduced palatability or digestibility were avoided, as noted by El Sayed *et al.* (1998). These results also confirm the findings of similar study done by Youssouf and Sédro (2020) who reported that diets combining different animal and insect protein sources improve growth performance and feed conversion in fish.

The results in Table VIII highlight the economic efficiency of treatments R4 and R2, characterized by significantly lower production costs per Kg of fish, respectively 958.94 and 1087.76 F CFA compared to R0, R1 and R3, where these costs exceed 1500 FCFA. These data corroborate the studies by Tacon and Metian (2015), who emphasized the

importance of optimized feed formulation to reduce production costs in aquaculture, by favoring accessible and digestible ingredients. Furthermore, the discrepancy observed between treatments may be explained by variations in feed conversion rate (FCR), a key parameter influenced by feed composition and rearing conditions, as suggested by Boyd and Tucker (2012). Thus, the results reinforce the idea that the use of adapted formulations (R4 and R2) can improve profitability while reducing the economic and environmental impact of fish production.

Results for overall fish body composition showed similar protein content in all treatments.

However, lipid content increased, probably due to the level of chicken viscera meal in the diets. These results are in agreement with Goda *et al.* (2007), Giri *et al.* (2010) and Sugumaran *et al.* (2015), who reported that replacing fish meal with poultry viscera meal in the diets did not affect the body protein content, but increased the whole-body lipid content of the fish.

## **Conclusion**

This study is part of an innovative approach to evaluate the effects of cow blood meal, chicken intestine meal and black soldier fly larvae meal as additives in the feed of hybrid fry of *Clarias gariepinus* and *Clarias anguillaris*. The results obtained show that these ingredients, used together at a total incorporation rate of 5%, offer significantly better growth performance and feed efficiency than the other diets tested.

Diets containing the mixture of these three ingredients were well accepted by the fish, with no pathological symptoms or negative impacts on their health. Environmental conditions, notably temperature and pH, were favorable throughout the experiment. These observations underline the potential of these raw materials as viable and sustainable alternatives to conventional fishmeal.

In order to consolidate the results obtained in the present study, it would be relevant to analyze the impact of the mixture on the fish immune system and disease resistance, and

to study the long-term effects of these additives on the health and overall performance of fish in different production systems.

## References

- Abdel-Warith AA, Russell PM, Davies SJ. Inclusion of a commercial poultry by-product meal as a protein replacement of fish meal in practical diets for African catfish *Clarias gariepinus* (Burchell 1822). *Aquaculture Research*. 2001; 32:296-306.
- Adebayo, O. T., Fagbenro, O. A., & Popoola, O. M. (2008). Effects of temperature on hatching and larval survival of *Clarias gariepinus* (Burchell, 1822). *Journal of Fisheries and Aquatic Science*, **3**(4), 302-307
- Alofa, C.S., & Abou, Y. (2020). A comparison between chicken viscera and housefly maggot cultured from this by products for Nile tilapia Diets: Growth Performance, Feed Utilization and Whole Body Composition. *Asian Journal of Fisheries and Aquatic Research*, 5(3), 1-12. <https://doi.org/10.9734/ajfar/2019/v5i330075>
- Alofa, C.S., Oke V., & Abou, Y. (2016). Effect of replacement of fish meal with broiler chicken viscera on growth, feed utilization and production of African catfish *Clarias gariepinus* (Burchell, 1822). *International Journal of Fisheries and Aquatic Studies*, 4(6), 182-186.
- Amtul, B.T., and Amna, B., 2012. Replacement of Fish Meal with Poultry by-Product Meal (Chicken Intestine) as a Protein Source in Grass Carp Fry Diet. *Pakistan J. Zool.* vol. 44(5), pp. 1373-1381, 2012.
- Baldé, B.S., 2019. Dynamics of small pelagic fish (*Sardinella aurita* and *Ethmalosa fimbriata*) in Senegal in a context of climate change: bioecological diagnosis and synthesis (PhD thesis). Université Cheikh Anta Diop, Dakar, Senegal.
- Barragan-Fonseca, K.B.; Dicke, M.;van Loon J.J.A., 2017. Nutritional value of the black soldier fly (*Hermetia illucens* L.) and its suitability as animal feed - a review. *J. Insects as Food Feed* 3(2), 105-120.
- Bishop, C. D., Angus, R. A. and Watts, S. A. (1995). The use of feather meal as a replacement for fish meal in the diet of *Oreochromis niloticus* fry. *Bioresource Technology*, 54, 291-295.



- 400 - Bloukounon-Goubalan A.Y., Saïdou A., Chrysostome C.A.A.M., Kenis M., Amadji G.L.,  
401 Igué A.M. & Mensah G.A., 2020. Physical and Chemical Properties of the Agroprocessing  
402 By-products Decomposed by Larvae of *Musca domestica* and *Hermetia illucens*. *Waste and*  
403 *Biomass Valorization* 11(6), 2735-2743.
- 404 - Booth, Donald C., and Craig Sheppard. "Oviposition of the black soldier fly, *Hermetia*  
405 *illucens* (Diptera: Stratiomyidae): eggs, masses, timing, and site  
406 characteristics." *Environmental entomology* 13.2 (1984): 421-423.
- 407 - Boyd, C. E., & Tucker, C. S. (2012). Pond aquaculture water quality management. Springer  
408 Science & Business Media, p 699
- 409 - Caruso D., Devic E., Subamia I.W., Talamond P. & Baras E., 2014. *Technical handbook of*  
410 *domestication and production of diptera Black Soldier Fly (BSF)*, IPB Press, Bogor,  
411 Indonesia, 141.
- 412 - Cuvelier C., Cabaraux J., Dufrasne I., Hornick J. & Istasse L., 2004. Fatty acids:  
413 nomenclature and dietary sources. *Ann. Médecine Vétérinaire* 133-140.
- 414 - Debode J., De Tender C., Soltaninejad S., Van Malderghem C., Haegeman A., Van der  
415 Linden I., Cottyn B., Heyndrickx M. & Maes M., 2016. Chitin mixed in potting soil alters  
416 lettuce growth, the survival of zoonotic bacteria on the leaves and associated rhizosphere  
417 microbiology. *Front. Microbiol.* 7(APR), 1-15.
- 418 - Dehaumont P. (1982). Les problèmes posés par la valorisation du sang d'abattoirs d'animaux  
419 de boucherie. *R.T.V.A.*,183: 23-32.
- 420 - Diener S., Zurbrugg C. & Tockner K., 2009. Conversion of organic material by black soldier  
421 fly larvae: establishing optimal feeding rates. *Waste Manag. Res.* 27(6), 603-610.
- 422 - Diener S., Zurbrugg C., Roa Gutiérrez F., Nguyen H.D., Morel A., Koottatep T. & Tockner  
423 K., 2011. Black soldier fly larvae for organic waste treatment - prospects and constraints. *In:*  
424 *WasteSafe 2011 2nd International Conference on Solid Waste Management in Developing*  
425 *Countries 1315 February 2011 Khulna Bangladesh.* 978- 984.
- 426 - El-Sayed, A. F. M. (1998). Total replacement of fish meal with animal protein sources in  
427 Nile tilapia, *Oreochromis niloticus* (L.), feeds. *Aquaculture Research*, 29, 275-280;
- 428 - Ewald N., Vidakovic A., Langeland M., Kiessling A., Sampels S. & Lalander C.,  
429 Fagbenro OA, Balogun AM, Fasakin EA, Bello-Olusoji OA. 1998b. Dietary lysine

requirement of the African catfish *Clarias gariepinus*. Journal of Applied Aquaculture; 8(2):71-77.

- Fagbenro OA. 2004. Soybean meal replacement by arugula (*Eruca sativa* Miller) seed meal as protein feedstuff in diets for African Catfish, *Clarias gariepinus* (Burchell 1822), fingerlings. Aquaculture Resource. 35:917-923.

- FAO, 2020. The State of World Fisheries and Aquaculture. Roma, Italy, ISBN 97892-5132692-3 224p.

Fly Produces a Foodstuff High in Omega-3 Fatty Acids. *J. World Aquac. Soc.* 38(2), 309-313.

- Fowler LG. 1991. Poultry by product meal as a dietary protein source in fall Chinook salmon diets, Aquaculture. 99:309-321.

- Giannetto A., Oliva S., Ceccon Lanes C.F., de Araújo Pedron F., Savastano D., Baviera C., Parrino V., Lo Paro G., Spanò N.C., Cappello T., Maisano M., Mauceri A. & Fasulo S., 2020. *Hermetia illucens* (Diptera: Stratiomyidae) larvae and prepupae: Biomass production, fatty acid profile and expression of key genes involved in lipid metabolism. *J. Biotechnol.* 307: 44-54.

- Gilles S., Dugué R., Slembrouck J., (2001) Manuel pratique de pisciculture du poisson-chat africain (*Clarias gariepinus*) page

- Giri SS, Sahoo SK, Mohanty SN. 2010. Replacement of by catch fishmeal with dried chicken viscera meal in extruded feeds: effect on growth, nutrient utilisation and carcass composition of catfish *Clarias batrachus* (Linn.) fingerlings Aquaculture Int. 18:539-544.

- Glencross BD, Booth M, Allan GL. 2007. A feed is only as good as its ingredients- a review of ingredient evaluation strategies for aquaculture feeds. Aquaculture Nutrition. 3:17-34.

- Goda, A.M., El-Haroun, E.R. & Chowdhury, M.A.K. (2007). Effect of totally or partially replacing fish meal by alternative protein sources on growth of African catfish *Clarias gariepinus* (Burchell, 1822) reared in concrete tanks. Aquaculture Research, 38, 279-287.

- Goddard, S., Al-Shagaa, G. and Ali, A. (2008). Fisheries by-catch and processing waste meals as ingredients in diets for Nile tilapia, *Oreochromis niloticus*. Aquaculture Research, 39, 518-525;

- Guo, J., Wang, Y. & Bureau, D.P. (2007). Inclusion of rendered animal ingredients as fishmeal substitutes in practical diets for cuneate drum, *Nibea miichthioides* (Chu, Lo and Wu). *Aquaculture Nutrition*, 13, 81-87.
- Henry M., Gasco L., Piccolo G. & Fountoulaki E., 2015. Review on the use of insects in the diet of farmed fish: Past and future. *Anim. Feed Sci. Technol.* 203 (1), 1-22.
- Hernandez, C., Olvera-Novoa, M. A., Hardy, R. W., Hermosillo, A., Reyes, C. and Gonzalez, B. (2010). Complete replacement of fish meal by porcine and poultry by-product meals in practical diets for fingerling Nile tilapia *Oreochromis niloticus*: digestibility and growth performance. *Aquaculture Nutrition*, 16, 44-53.
- Hertrampf JW, Piedad-Pascual F. 2000. Handbook on ingredients for Aquaculture Feeds, Kluwer academic publishers, Dordrecht, Netherlands. 482-483.
- Hoc B., Genva M., Fauconnier M.-L., Lognay G., Francis F. & Caparros Megido R., 2020. About lipid metabolism in *Hermetia illucens* (L. 1758): on the origin of fatty acids in prepupae. *Sci. Rep.* 10(1), 11916.
- Holmes L., 2010. Role of Abiotic Factors on the Development and Life History of the Black Soldier Fly, *Hermetia illucens*(L.)(Diptera: Stratiomyidae). *Masters Abstr. Int.*
- Holmes L.A., Vanlaerhoven S.L. & Tomberlin J.K., 2012. Relative Humidity Effects on the Life History of *Hermetia illucens* (Diptera: Stratiomyidae). *Environ. Entomol.* 41(4), 971-978.
- Hu M, Wang Y, Wang Q, Zhao M, Xiong B, Qian X et al. Evaluation of rendered animal protein ingredients for replacement of fish meal in practical diets for gibel carp, *Carassius auratus gibelio* (Bloch), *Aquaculture research*. 2008a; 39:1474-1482.
- Hu M, Wang Y, Wang Q, Zhao M, Xiong B, Qian X et al. Replacement of fish meal by rendered animal protein ingredients with lysine and methionine supplementation to practical diets for gibel carp, *Carassius auratus gibelio*. *Aquaculture*. 2008b; 275:260-265
- Kim W., Bae S., Park H., Park K., Lee S., Choi Y., Han S. & Koh Y., 2010. The Larval Age and Mouth Morphology of the Black Soldier Fly, *Hermetia illucens* (Diptera: Stratiomyidae). *Int. J. Ind. Entomol.* 21(2), 185-187.

- 487 - Lévêque C. & Paugy D., 2006. *Les poissons des eaux continentales africaines: Diversité,*  
488 *écologie, utilisation par l'homme*, 59-74.
- 489 -Liland N.S., Biancarosa I., Araujo P., Biemans D., Bruckner C.G., Waagbø R., Torstensen  
490 B.E. & Lock E.-J., 2017. Modulation of nutrient composition of black soldier fly (*Hermetia*  
491 *illucens*) larvae by feeding seaweed-enriched media. *PLoS One* 12(8), e0183188.
- 492 - Liu X., Chen X., Wang H., Yang Q., ur Rehman K., Li W., Cai M., Li Q., Mazza L.,  
493 Zhang J., Yu Z. & Zheng L., 2017. Dynamic changes of nutrient composition throughout the  
494 entire life cycle of black soldier fly. *PLoS One* 12(8), e0182601.
- 495 - Maier T., Leibundgut M., Boehringer D. & Ban N., 2010. Structure and function of  
496 eukaryotic fatty acid synthases. *Q. Rev. Biophys.* 43(3), 373-422.
- 497 - Milliamena, O.M. (2002). Replacement of fish meal by animal by-product meals in a  
498 practical diet for grow-out culture of grouper *Epinephelus coioides*. *Aquaculture*, 204, 75 84.  
499 [https://doi.org/10.1016/s0044-8486\(01\)00629-9](https://doi.org/10.1016/s0044-8486(01)00629-9)
- 500 - Ndiaye, N. A., Maiguizo-Diagne, H., Diadhiou, H. D., Ndiaye, W. N., Diedhiou, F., Cournac,  
501 L., ... & Brehmer, P. (2020). Methanogenic and fertilizing potential of aquaculture waste:  
502 towards freshwater farms energy self-sufficiency in the framework of blue growth. *Reviews*  
503 *in Aquaculture*, 12(3), 1435-1444.
- 504 - Newton L., Sheppard C., Watson W.D., Burtle G. & Dove R., 2005. Using the black soldier  
505 fly, *Hermetia illucens*, as a value-added tool for the management of swine manure. *J. Korean*  
506 *Entomol. Appl. Sci.* 36(12), 17 pp.
- 507 - Nguyen T.T.X., Tomberlin J.K. & Vanlaerhoven S., 2015. Ability of Black Soldier Fly  
508 (Diptera: Stratiomyidae) Larvae to Recycle Food Waste. *Environ. Entomol.* 44(2), 406410.
- 509 - NRC, 2011. Nutrient requirements of fish and shrimp, Washington, DC, 376.
- 510 - Olukunle , O.A. ,Ogunsanmi,A.O., Taiwo,V.O., and Samuel,A.A. (2002). The nutritional value  
511 of cow blood meal and its effect on growth performance haematology and plasma enzymes of  
512 hybrid catfish. Departments of wildlife and fisheries Management and Veterinary Pathology,  
513 university of Ibadan, Ibadan. Nigeria;
- 514 - Oonincx D.G.A., Laurent S., Veenenbos M.E. & van Loon J.J.A., 2020. Dietary enrichment  
515 of edible insects with omega 3 fatty acids. *Insect Sci.* 27(3), 500-509.

- Oonincx D.G.A.B., van Broekhoven S., van Huis A. & van Loon J.J.A., 2015. Feed Conversion, Survival and Development, and Composition of Four Insect Species on Diets Composed of Food By-Products. *PLoS One* 10(12), e0144601.
- Oonincx D.G.A.B., van Itterbeeck J., Heetkamp M.J.W., van den Brand H., van Loon J.J.A. & van Huis A., 2010. An Exploration on Greenhouse Gas and Ammonia Production by Insect Species Suitable for Animal or Human Consumption. *PLoS One* 5(12), e14445.
- Quilliam R.S., Nuku-Adeku C., Maquart P., Little D., Newton R. & Murray F., 2020. Integrating insect frass biofertilisers into sustainable peri-urban agro-food systems. *J. Insects as Food Feed* 6(3), 315-322.
- Rawles SD, Riche M, Gaylord TG, Webb J, Freeman DW, Davis M. Evaluation of poultry by-product meal in commercial diets for hybrid striped bass (*Morone chrysops* male X *M. saxatilis* female) in recirculated tank production. *Aquaculture*, 2006; 259:377-389.
- Reyes M., Rodríguez M., Montes J., Barroso F.G., Fabrikov D., Morote E. & SánchezMuros M.J., 2020. Nutritional and Growth Effect of Insect Meal Inclusion on Seabass (*Dicentrarchus labrax*) Feeds. *Fishes* (2), 16-0.
- Robaina L, Moyano FJ, Izquierdo MS, Socorro J, Vergera JM, Montero D. Corn gluten meal and meat and bone meals as protein sources in diets for gilthead seabream *Sparus aurata* nutritional and histological implications. *Aquaculture*. 1997; 157:347-359.
- Rónyai A, Csengeri I, Váradi L. Partial substitution of animal protein with full-fat soybean meal and amino acid supplementation in the diet of Siberian sturgeon (*Acipenser baerii*). *J Appl Ichthyol*. 2002; 18:682-684.
- Sanogo S, Compaore I, Senou I, Somda M B, Ouedraogo R B, Ouattara B And Kabre Ta. (2020) Etude comparée de la structuration des macro invertébrés benthiques de cours d'eau urbain et péri-urbain à l'ouest du burkina faso. *International Journal of Development Research* Vol. 11, Issue, 01, pp. 4317;
- Scala A., Cammack J.A., Salvia R., Scieuzo C., Franco A., Bufo S.A., Tomberlin J.K. & Falabella P., 2020. Rearing substrate impacts growth and macronutrient composition of *Hermetia illucens* (L.) (Diptera: Stratiomyidae) larvae produced at an industrial scale. *Sci. Rep.* 10 (1), 19448.
- Sprangers T., Ottoboni M., Klootwijk C., Olyn A., Deboosere S., De Meulenaer B., Michiels J., Eeckhout M., De Clercq P. & De Smet S., 2017. Nutritional composition of

547 black soldier fly (*Hermetia illucens*) prepupae reared on different organic waste substrates. *J.*  
548 *Sci. Food Agric.* (8), 2594-2600.

549 - Stanley-Samuelson D.W., Jurenka R.A., Cripps C., Blomquist G.J. & de Renobales M., 1988.  
550 Fatty acids in insects: Composition, metabolism, and biological significance. *Arch. Insect*  
551 *Biochem. Physiol.* 9(1), 1-33.

552 - St-Hilaire S., Cranfill K., McGuire M.A., Mosley E.E., Tomberlin J.K., Newton L., Sealey  
553 W., Sheppard C. & Irving S., 2007. Fish Offal Recycling by the Black Soldier  
554 - St-Hilaire S., Sheppard C., Tomberlin J.K., Irving S., Newton L., McGuire M.A., Mosley  
555 E.E., Hardy R.W. & Sealey W., 2007. Fly Prepupae as a Feedstuff for Rainbow Trout,  
556 *Oncorhynchus mykiss*. *J. World Aquac. Soc.* 38(1), 59-67.

557 - Sugumaran, E. and Radhakrishnan, M.V. (2015). Feed utilization, growth and carcass  
558 composition of catfish *Clarias batrachus* (L.) fed on fish meal replaced by dried chicken  
559 viscera incorporated diets. *International Journal of Research in Fisheries and Aquaculture* 5,  
560 143-146.

561 - Tacon A.G.J. & Metian M., 2008. Global overview on the use of fish meal and fish oil in  
562 industrially compounded aquafeeds: Trends and future prospects. *Aquaculture* 285(1-4), 146-  
563 158.

564 - Tacon, A. G. J. and Jackson, A. J. (1985). Utilization of conventional and unconventional  
565 protein sources in practical fish feeds. In *Nutrition and Feeding in Fish*, Cowey CB, Mackie  
566 AM, Bell JG. (eds). Academic Press, London, 119 - 145.

567 - Tacon, A. G. J., & Metian, M. (2015). *Feed matters: Satisfying the feed demand of*  
568 *aquaculture*. *Reviews in Fisheries Science & Aquaculture*, 23(1), 1-10.

569 - Teugels G.G. & Thys van den Audenaerde D.F.E., 2003. Cichlidae. In: *The Fresh and*  
570 *Brackish Water Fishes of West Africa*. Paris, 521-600.

571 - Teugels, 1986: A systematic revision of the African species of genus *Clarias* Pices;Clariidae).  
572 *Annales du musée royal d'Afrique centrale* 247, 1-19

573 - Tomberlin J.K., Adler P.H. & Myers H.M., 2009. Development of the Black Soldier Fly  
574 (Diptera: Stratiomyidae ) in Relation to Temperature. *Environ. Entomol.* 38(Hogsette  
575 1992), 930-934.

- 576 - Tomberlin J.K., Sheppard D.C. & Joyce J.A., 2002. Selected Life-History Traits of Black  
577 Soldier Flies (Diptera: Stratiomyidae) Reared on Three Artificial Diets. *Ann. Entomol. Soc.*  
578 *Am.* 95(3), 379-386.
- 579 - Truzzi C., Giorgini E., Annibaldi A., Antonucci M., Illuminati S., Scarponi G., Riolo P.,  
580 Isidoro N., Conti C., Zarantoniello M., Cipriani R. & Olivotto I., 2020. Fatty acids profile of  
581 black soldier fly (*Hermetia illucens*): Influence of feeding substrate based on coffeewaste  
582 silverskin enriched with microalgae. *Anim. Feed Sci. Technol.* 259(October 2019), 114309.
- 583 - Van Huis A. & Tomberlin J.K., 2017. *Insects as food and feed: from production to*  
584 *consumption*. van Huis, A., Tomberlin, J.K. eds, Wageningen Academic Publishers, The  
585 Netherlands, 448.
- 586 - Viveen W.J.A.R., C.J.J. Richter, P.G.W.J. Van Oordt, J.A.L. Janssen, E.A. Huisman (1985).  
587 Practical manual for fish culture of African catfish (*Clarias gariepinus*), Netherlands, 128 .
- 588 - Wang Y, Li K, Han H, Zheng Z, Bureau DP. 2008. Potential of using a blend of rendered  
589 animal protein ingredients to replace fish meal in practical diets for malabar grouper  
590 (*Epinephelus malabaricus*). *Aquaculture* 281:113 117.
- 591
- 592 - Zarantoniello M., Zimbelli A., Randazzo B., Compagni M.D., Truzzi C., Antonucci M.,  
593 Riolo P., Loreto N., Osimani A., Milanović V., Giorgini E., Cardinaletti G., Tulli F.,  
594 Cipriani R., Gioacchini G. & Olivotto I., 2020. Black Soldier Fly (*Hermetia illucens*)  
595 reared on roasted coffee by-product and *Schizochytrium* sp. as a sustainable terrestrial  
596 ingredient for aquafeeds production. *Aquaculture* 518, 734659.
- 597 - Zhou J.S., Liu S.S., Ji H. & Yu H.B., 2018. Effect of replacing dietary fish meal with  
598 black soldier fly larvae meal on growth and fatty acid composition of Jian carp (*Cyprinus*  
599 *carpio* var . Jian). *Aquac. Nutr.* 24(1), 424-433.