

1 **Heavy metal bioaccumulation in *Zea mays* L. i *Medicago sativa* L. in the area of Zenica**

2
3

4 **Abstract**

5 In the period March-September 2018, research was conducted on the content of heavy metals: iron
6 (Fe), manganese (Mn), zinc (Zn), nickel (Ni), lead (Pb), vanadium (V), molybdenum (Mo), cadmium
7 (Cd), chromium (Cr), copper (Cu) and cobalt (Co) in the soil and plants of corn (*Zea mays* L.) and
8 alfalfa (*Medicago sativa* L.) at three locations in the Zenica area (Gradišće, Stranjani and Šerići). Soil
9 analysis shows that the soil's average concentrations of manganese, nickel, vanadium, cadmium, and
10 molybdenum exceed the permitted limits. Based on the measured values, higher concentrations of
11 vanadium (4.93 mg/kg), cadmium (0.251 mg/kg), chromium (5.03 mg/kg), and copper (10.05 mg/kg)
12 are present in the roots of corn. More manganese (170.7 mg/kg) is concentrated in the corn's leaf, and
13 zinc (19.95 mg/kg) in the corn's grain. Most iron (1301.5 mg/kg) is concentrated in alfalfa roots, while
14 nickel (2.43 mg/kg) and lead (4.92 mg/kg) dominate in its above-ground part. The results show that
15 the bioaccumulation of heavy metals is more pronounced in the above-ground part of alfalfa and the
16 roots of corn.

17 Keywords: heavy metals, corn, alfalfa, soil

18 **INTRODUCTION**

19
20 Heavy metals are increasingly present in the environment, especially in industrialized areas. The most
21 common sources of soil contamination with heavy metals in the Zenica area are metallurgical and
22 thermal power plants that have been active for many years. The increased content of heavy metals in
23 the surface part of the soil, plants, and atmosphere in mining and metallurgical areas has been
24 documented in numerous studies such as [1], [2], [3] etc. Combustion of fossil fuels and incineration
25 of waste contributes significantly to the emission of As, Se, Hg, Cd, Cr, Ni, and Zn [4].

26 The content of heavy metals in the soil has increased compared to the natural state and the limit values
27 prescribed by the Rulebook on the determination of permitted amounts of harmful and dangerous
28 substances in the soil and methods of their examination [5], which is a consequence of their
29 anthropogenic redistribution during the integral operation of the Ironworks Zenica. Research
30 conducted by the Federal Institute for Agropedology in Sarajevo [6] and the Metallurgical Institute
31 "Kemal Kapetanović" in Zenica [7] prove that the soil around Ironworks Zenica is heavily
32 contaminated with heavy metals (manganese, nickel, cadmium, and zinc). A study conducted in 2010
33 [8] based on analyses of plant crops (leaves of fruit crops, lettuce, chard, nettles, alfalfa) in the area of
34 Zenica concluded that the plants were contaminated with lead, cadmium, zinc, copper, nickel, cobalt
35 and chromium. Heavy metals from the soil accumulated in plants enter the food chain and their
36 excessive amount can have a toxic effect on humans and animals.

37 This paper presents the results of researching heavy metals concentration and their distribution in
38 different parts of corn and alfalfa collected in polluted sites in the Zenica area, to determine their
39 ability to bioaccumulate heavy metals, and their potential use for soil decontamination.

40 Deficiencies or excesses of certain metals can be reliably determined on a leaf, stem, or root [9],
41 considering their necessary presence in plant tissue. The behavior of heavy metals in the soil and their
42 accumulation in plants depends on many factors, such as the composition and pH value of the soil, the
43 content of organic matter, the adsorption complex, and other factors [10].

44 The absorption and storage of heavy metals from the soil into plants occurs through a series of
45 transport mechanisms that normally serve to absorb micronutrients [9, 11].

46 47 **MATERIALS AND METHODS**

48 The research was carried out in three localities in the area of Zenica, established according to the
49 system of cyclic circles at different distances, from 2.78 to 16.4 km as the crow flies from the

50 dominant industrial sources of dust and heavy metal emissions (metallurgical and thermal energy
51 plants in Zenica). The distances, longitude, latitude, and altitude are shown in Table 1.

52 **Table 1.** Distance longitude, latitude, and altitude of research sites from industrial emission sources

No.	Location	Distance (km)	Latitude	Longitude	Altitude
1.	Gradišće	2,78	44 ⁰ 14' 13" N	17 ⁰ 52' 8" E	528 m
2.	Stranjani	4,88	44 ⁰ 13' 14" N	17 ⁰ 50' 54" E	565 m
3.	Šerići	16,4	44 ⁰ 21' 30" N	17 ⁰ 48' 23" E	809 m

53
54 To obtain a representative sample, from each location, several soil samples were taken from a depth
55 of 25 cm. After drying in the air, a sample was reduced using the quartering method, and then the
56 remaining sample was shredded, sieved, and packed for chemical analysis.

57 To monitor the content of heavy metals in the soil, soil samples were taken before and after planting
58 corn and alfalfa. Initial soil analysis is given in Table 2 and the soil analysis after removing plants is
59 shown in Tables 3, 4, and 5.

60 The plant material was sampled as whole plants with roots, and then the fraction samples were made
61 from root, leaf, and grain in the case of corn and root and the above-ground part of the plant in the
62 case of alfalfa. Fraction samples were air-dried for one month and then crushed. The content of
63 heavy metals in corn and alfalfa samples was done using atomic absorption spectrophotometry
64 according to PERKIN ELMER Analytical Methods for Atomic Absorption Spectrometry.

65 Limit values of heavy metals (HM) depending on soil texture are defined by the Rulebook on the
66 determination of permitted amounts of harmful and dangerous substances in the soil and methods of
67 their examination [5] are shown in Table 2. The textural features of the soil of the investigated
68 locations are powdery-loamy. Limit values for manganese and iron, given in Table 2, are taken from
69 the Federal Institute for Agropedology[6]. These soil limit values refer to soils with an acidic
70 reaction. The specified values can be increased by 25% for alkaline and carbonate soils.

71 Results and discussion

72 The initial soil analysis shows that the average concentrations of manganese, nickel, vanadium, and
73 cadmium are higher than the permitted limit value at all locations in the Zenica area. Elevated
74 molybdenum concentrations in the soil were registered at sites, Gradišće and Stranjani (Table 2).
75

76 **Table 2.** Initial soil analysis at research sites Gradišće, Stranjani, and Šerići

Metal (mg/kg)	Research site			Limit value (mg/kg)
	Gradišće	Stranjani	Šerići	
Fe	42.000 (4.2%)	34.633 (3.5%)	22. 700 (2.3%)	5%
Mn	1417	1667	3063	1.000.
Zn	29	21	19	150
Ni	111	66	46	40
Pb	59	25	23	80
V	104	67	44	40
Mo	20	24	5	15
Cd	1.4	1.36	3	1.0
Cr	78	41	16	80
Cu	29	31	49	65
Co	<1	<1	<1	45
pH in H ₂ O	7.7	7.6	6.2	-
pH in KCl	7.0	6.8	5.6	-

77 Note: Shaded numbers indicate exceeding limit values.

78
79 The content of heavy metals in fraction samples of corn and alfalfa as well as the content of heavy
80 metals in the soil after the removal of plants are presented in the following tables.

81 **Tabela 3.** Content of heavy metals in plants and soil after removal of plants at research site Gradišće (mg/kg)

Metal	Corn				Alfalfa		
	Root	Leaf	Grain	Soil after removal	Root	The above-ground part	Soil after removal

Fe	1146.3	238.8	57.36	45.890 (4.589 %)	462.5	339.1	43.590 (4.359 %)
Mn	27.12	76.32	9.98	1400	18.00	24.49	1440
Zn	4.93	7.39	19.95	10	<0.01	4.92	10
Ni	<0.01	<0.01	<0.01	80	<0.01	<0.01	90
Pb	<0.01	1.97	2.25	90	<0.01	4.92	70
V	4.93	<0.01	<0.01	50	<0.01	<0.01	<1
Mo	<0.01	<0.01	<0.01	<1	<0.01	<0.01	<1
Cd	<0.001	<0.001	<0.001	<0.1	<0.001	<0.001	<0.1
Cr	4.93	2.46	2.49	90	<0.01	<0.01	100
Cu	7.40	4.92	<0.01	50	<0.01	2.45	60
Co	<0.01	<0.01	<0.01	<1	<0.01	<0.01	<1

82

83 **Table 4.** Content of heavy metals in plants and soil after removal of plants at research site Stranjani (mg/kg)

Metal	Corn				Alfalfa		
	Root	Leaf	Grain	Soil after removal	Root	The above-ground part	Soil after removal
Fe	997.5	110.4	30.10	35.500 (3.55%)	943.8	549.5	36.100 (3.61 %)
Mn	105.8	52.9	20.1	1630	32.4	26.7	1630
Zn	2.65	<0.01	<0.01	20	2.49	2.43	20
Ni	<0.01	<0.01	<0.01	60	<0.01	2.43	70
Pb	<0.01	4.60	<0.01	50	<0.01	<0.01	10
V	<0.01	<0.01	<0.01	60	<0.01	2.43	<1
Mo	<0.01	<0.01	<0.01	<1	<0.01	<0.01	<1
Cd	<0.001	<0.001	<0.001	<0.1	<0.001	<0.001	<0.1
Cr	<0.01	2.30	2.51	70	<0.01	2.43	80
Cu	<0.01	4.60	<0.01	40	7.47	7.29	30
Co	<0.01	<0.01	<0.01	<1	<0.01	<0.01	<1

84

85 **Table 5.** Content of heavy metals in plants and soil after removal of plants at research site Šerići (mg/kg)

Metal	Corn				Alfalfa		
	Root	Leaf	Grain	Soil after removal	Root	The above-ground part	Soil after removal
Fe	1100.7	263.6	19.3	24980 (2.498 %)	1301.5	235.1	26980 (2.698 %)
Mn	105.5	170.7	7.24	3000	158.3	67.2	2730
Zn	2.51	5.02	2.41	20	2.36	4.80	20
Ni	<0.01	<0.01	<0.01	40	2.36	<0.01	60
Pb	<0.01	<0.01	<0.01	20	<0.01	<0.01	70
V	2.51	<0.01	<0.01	<1	<0.01	<0.01	20
Mo	<0.01	<0.01	<0.01	<1	<0.01	<0.01	<1
Cd	0.251	<0.001	<0.001	<0.1	<0.001	<0.001	<0.1
Cr	5.03	2.51	2.41	30	2.36	2.40	40
Cu	10.05	2.51	2.41	80	7.09	4.79	80
Co	<0.01	<0.01	<0.01	<1	<0.01	<0.01	<1

86

87 Results from Tables 3, 4, and 5 show that The iron content in the dry samples of plants is mostly
88 between 19.3 (grain of corn) and 1301.5 (the root of alfalfa) mg/kg, which is fairly low concerning
89 that some ferrophilic plants (spinach, oats) can contain up to 3500 mg/kg of iron [12]. According to
90 the author, the distribution of iron in different parts of corn is as follows: root 44.1%, stem 14.6%,
91 leaves 23%, grain 15.7%, cob 2.7%. Lazarevic et al. find that the concentrations of Fe in the roots and
92 leaves of corn increase with the increase in the concentration of Fe in nutrient solutions [13], which is
93 in agreement with the higher concentrations of iron in the roots of corn due to the high content of Fe in
94 the soil at the Gradišće location (Tables 2 and 3).

95 Results in above tables show that the concentration of Iron in corn ranges from 19.3 mg/kg in grain to
96 1146.3 mg/kg in root. The average concentration of iron in corn leaves ranges from 110.4 mg/kg at the
97 Stranjani site to 263.3 mg/kg at Šerići site. The low concentration of Iron in corn leaves on research
98 site Stranjani indicates low mobility of iron in the soil of this research site. Since the critical
99 concentration of iron in leaves is from 50 to 150 mg/kg [14] low mobility of iron could lead to iron
100 deficiency in this research site. The average concentration of Iron in the roots of alfalfa ranges
101 between 460.5 mg/kg and 1143 mg/kg, and from 235.1 to 549.5 mg/kg in the aboveground part of
102 alfalfa (Tables 3, 4, and 5).

103
104 The level of manganese in plants in research sites ranges from 7.24 mg/kg in corn grain to 170.7
105 mg/kg in corn leaves, followed by 158.3 mg/kg in alfalfa roots (Table 3, 4, 5). According to [12],
106 manganese concentration in the above-ground part of alfalfa reaches 86 mg/kg. According to [15], the
107 concentration of manganese in the grain of corn is 5.7 mg/kg, and according to [16], the concentration
108 of manganese in the grain of corn ranges between 2.85 and 4.35 mg/kg. The concentration of
109 manganese in grain of corn on all research sites is higher (Tables 2, 3, and 4) probably due to the
110 significantly high content of manganese in soil (1.43 to 3.06 times higher than the limit value).

111 The highest concentration of manganese, 158.3 mg/kg, is found in the roots of alfalfa on research site
112 Šerići because the soil on this research site has the highest manganese content.

113 Manganese is mobile in the plant and it is found in all parts of the investigated plants. The mobility of
114 manganese increases with soil acidity and its reduction to active manganese Mn^{2+} , which is easily
115 absorbed by plants [17], while more oxidized forms such as Mn^{3+} and Mn^{4+} are inactive forms.

116
117 The natural zinc content in plants varies from 20 to 100 mg/kg [18]. In cabbage leaves it is 35 mg/kg,
118 wheat grains contain 23 mg/kg, and sugar beet roots 17.5 mg/kg of zinc [12]. The mobility of zinc in
119 the plant is moderate, and it is believed that it is in the xylem in the form of citrate, chelate, or as a free
120 ion [19]. The critical limit of zinc deficiency is 15 – 30 mg/kg in dry matter of leaves [17].

121 The maximum measured zinc concentration in corn grain is 19.95 mg/kg. According to research
122 performed by [16], the average zinc value in corn grain varies between 13.6 and 25.6 mg/kg
123 depending on the genotype, weather conditions, and soil properties. Varga [20] states that the zinc
124 concentration in corn grain ranges between 19.09 and 22.60 mg/kg and depends on soil cultivation.

125 The maximum measured zinc concentration in corn leaves is 7.39 mg/kg and in the aboveground part
126 of alfalfa 4.92 mg/kg (Table 3), which indicates a zinc deficiency in the plant that is very important for
127 normal plant development.

128
129 The average concentration of nickel in plants ranges from 1 to 10 mg/kg dry matter [19], while in
130 alfalfa ranges from 1.0 to 4.0 mg/kg [12]. Nickel is considered relatively poorly mobile in the soil
131 because it has a high tendency to bind to clay minerals and iron and manganese oxides. The
132 accessibility of nickel increases when the soil pH is less than 6.5 (Table 2, 4, 5) because under these
133 conditions many Ni-components are relatively soluble [21]. This research showed that higher values of
134 Ni are present in alfalfa (in the root 2.36 mg/kg and the aboveground part 2.43 mg/kg Table 4 and 5)
135 in locations where the pH is less than 6.5, while corn fractions (roots, leaf, grain) contain the same
136 amounts of nickel (less than 0.01 mg/kg). According to a study performed by Federal Institute of
137 Agropedology B&H [8] the nickel content in an alfalfa sample at the Gradišće location was 13.90
138 mg/kg and it can be considered as contaminated. Studies show that the nickel concentration in alfalfa
139 increased proportionally to the increase in the amount of available nickel in the soil [22].

140
141 The natural content of lead in plants varies between 0.1 and 10 mg/kg [23, 24]. Plant roots are able to
142 accumulate lead in plants, but their subsequent translocation to aboveground parts is very limited [25].
143 Scientific research shows the significant accumulation of lead in the leaf [26, 27], and that lead
144 reaches plants primarily through leaves from the air [28, 29]. Wild herbaceous plants such as alfalfa
145 and nettle have significantly higher bioaccumulation of lead in plants than cultivated vegetable and
146 cereal plants, which is a result of the specific genetic conditions of the plants studied and their
147 adaptation to living conditions [8]. The lead content in the alfalfa sample was 12.20 mg/kg at the
148 Gradišće location, which exceeds the limit value for plants.

149 The greatest biostimulation by lead was in the Gradišće area, due to greater exposure to atmospheric
150 lead deposition because it is closest to metallurgical plants. The least accumulation of lead in plants
151 was recorded in the Šerić area, which is the furthest from the pollution source. This research showed
152 that the highest amounts of lead are present in the aboveground part of alfalfa (4.92 mg/kg), and in the
153 corn leaf (4.60 mg/kg) (Tables 3, 4, 5) and it is consistent with research of Federal Institute of
154 Agropedology performed in 2010 [8] at Gradišće site.

155

156 This study shows that vanadium concentration in corn roots ranged between 2.51 and 4.93 mg/kg,
157 while in alfalfa the maximum vanadium concentration of 2.43 mg/kg was recorded in the aboveground
158 part of the plant (Tables 3, 4, and 5).

159

160 Plants contain very little molybdenum, even below 1 mg/kg (0.1-0.5 mg/kg in dry matter), and
161 relatively higher molybdenum content is in plants from the legume and cruciferous families [19]. The
162 molybdenum content in alfalfa hay is 0.98 mg/kg [12]. The mobility of molybdenum in plants is
163 moderate. Plants absorb molybdenum in the form of MoO_4^{2-} and in plants, it exists as an anion, so its
164 accessibility increases with increasing alkalinity, unlike all other microelements, This study shows that
165 the molybdenum content in corn and alfalfa is the same and is less than 0.01 mg/kg (Table 3, 4 and 5).

166 The natural content of cadmium in plants varies between 0.05 and 0.20 mg/kg [23]. It has been
167 determined that 30-60% of cadmium contained in plants comes directly from the atmosphere and 40-
168 60% from the soil. Cadmium ions are mainly retained in the roots, and only small amounts are
169 transported to the aboveground part of the plant [30]. The highest concentrations of cadmium were
170 found in the roots of plants [31], which was also shown in this study, as 0.251 mg/kg was recorded in
171 corn roots at the Šerići location (Table 5) because the pH of the soil at the Šerići location is the lowest
172 and favors cadmium mobility. The cadmium content in other parts of corn and alfalfa is the same and
173 amounts to less than 0.001 mg/kg, which complies with previous research.

174

175 According to [24], the amount of chromium in plants usually varies in the range of 0.02 to 0.2 mg/kg
176 of dry matter [24] while according to [18] the amount of chromium in plants varies between 0.2 and
177 0.4 mg/kg. Plants growing on contaminated soil can accumulate higher amounts of chromium [21].
178 Chromium is preferentially accumulated in corn roots with a low rate of translocation to leaves [32],
179 while [29] believes that plants characterized by iron accumulation also accumulate chromium. Higher
180 values of chromium were recorded in the roots of corn at 4.93 - 5.03 mg/kg and in the aboveground
181 part of alfalfa at 2.45 mg/kg (Table 3,4,5). Given that the critical value of chromium in a plant is 1
182 mg/kg [8], the chromium content in both sown crops is above the permitted critical value.

183

184

185 The natural content of copper in plants is 2-20 mg/kg [18]. Copper translocation is moderate, in the
186 form of Cu complexes, usually with amino acids, so plant roots contain it in significant quantities [19].
187 The amount and availability of copper are significantly affected by the pH reaction of the soil, and its
188 accessibility increases with soil acidity[8,19], which is in agreement with the higher concentration of
189 copper in plants at the Šerići location, where the pH is the lowest of the observed locations. (Tables 2
190 and 5). The concentration of copper in corn roots was 10.05 mg/kg and 7.47 mg/kg in alfalfa roots.
191 The copper content in corn grain ranges from <0.01 to 2.41 mg/kg, while [15] found that the copper
192 content in corn grain ranges from 2.2 to 5.8 mg/kg.

193

194 The amount of cobalt in plants varies from 1 to 40 mg/kg of dry matter [19]. The natural content of
195 cobalt most often varies from 0.02 to 0.4 mg/kg [18]. Alfalfa hay contains 0.38 mg/kg [12]. It is a
196 component of vitamin B12, and is especially important for legumes because it is necessary for N_2
197 fixation, so in the event of a cobalt deficiency, organic production decreases [19]. Due to the poor
198 mobility of cobalt within the plant, only a small amount of cobalt is moved from the stem to the leaves
199 [33]. This research shows that different parts of corn and alfalfa have the same amount of cobalt and it
200 is less than 0.01 mg/kg (Table 3,4,5) probably due to the low cobalt content in the soil (below 1
201 mg/kg).

202

203 The area of Šerić is further away from the emission center, but the land at this location is more
204 sensitive from the point of view of contamination. Heavy metals in more acidic soils are in an active
205 form fixed to soil particles and they represent a potential danger. The largest amounts of iron,
206 manganese, vanadium, cadmium, chromium, and copper were accumulated by the plants at the
207 location of Šerići, while the highest amount of nickel was accumulated at the location of Stranjani,
208 The largest accumulation of zinc and lead was registered at the Gradišće location where the soil is
209 neutral to weakly alkaline. In such soil, heavy metals turn into hydroxides, and with a further increase
210 in the pH value, they turn into insoluble hydroxides, and if the carbonate content in the soil is high,
211 heavy metals are inactivated, i.e. they become difficult for plants to access [8]. The research results
212 show that the concentration of nickel, chromium, copper, manganese, and lead (Table 2, 3, 4, and 5) in
213 the soil is higher after the removal of plants due to the pollution of the Zenica area with an increased
214 amount of sediment containing heavy metals [7].

215
216
217 Alfalfa (*Medicago sativa* L.) shows a greater ability to bioaccumulate iron in the roots, and higher
218 values of nickel and lead are accumulated in the aboveground part. (Table 3, 4 and 5). The same
219 values of molybdenum, cadmium, and cobalt are present in both fractions of alfalfa (root and
220 aboveground part). Previous research has shown that alfalfa accumulates chromium, vanadium, lead,
221 nickel, molybdenum, and iron better than corn [34]. Plant species that do not prevent the entry of
222 metals into the roots and allow the bioaccumulation of higher concentrations of metals in the tissues
223 of their aboveground organs are most suitable for use in phytoremediation [35].

224
225 Maize (*Zea mays* L.) has the ability to accumulate in its roots more vanadium, cadmium, chromium,
226 and copper compared to alfalfa. The translocation of iron, manganese, zinc, chromium, and copper
227 from the roots to the aerial parts of the plant is observed. The leaves have higher values of manganese
228 and lead because heavy metals from the roots and the air accumulate in the leaf. Higher values of zinc
229 are present in corn grains. The same values of nickel, molybdenum, and cobalt are present in the
230 aboveground parts of corn. It can be seen that copper and chromium have high mobility in the soil and
231 are easily taken up by plants.

232 **Conclusion**

233
234 The results of this research conducted in the period March-September 2018 show that the content of
235 iron, chromium, and manganese in corn, as well as iron and chromium in alfalfa, is above the
236 permitted limits for heavy metals in plants growing on polluted land in the area of Zenica. The biggest
237 accumulator of iron is the root of the plant, with values in alfalfa 1301.5 mg/kg and corn 1146.3
238 mg/kg. Iron and manganese are mobile in plants and are found in both roots and aboveground parts of
239 corn and alfalfa. The highest concentration of manganese was registered in corn leaves (170.7 mg/kg).
240 This research shows higher values of manganese in corn grains at all three locations ranging between
241 7.24 and 20.1 mg/kg, probably due to increased concentrations of manganese in the soil. Zinc
242 accumulates in the leaf and grain of corn with the maximum measured value of 19.95 mg/kg. Nickel is
243 more abundant in alfalfa (roots 2.36 mg/kg and aboveground parts 2.43 mg/kg) in locations where the
244 pH is less than 6.5, while in all corn fractions content of nickel is the same and is less than 0.01 mg/kg.
245 Lead dominates in the aboveground part of alfalfa (4.92 mg/kg) in the Gradišće site, which is closest
246 to the metallurgical plants, due to the greater burden of this area with atmospheric depositions
247 containing heavy metals. The highest concentration of cadmium is in the roots of corn with 0.251
248 mg/kg at the Šerići location, which is related to soil acidity, while in other plant parts of corn and
249 alfalfa, low values were recorded and were less than 0.001 mg/kg. Corn roots accumulate higher
250 values of chromium (4.93 – 5.03 mg/kg), while alfalfa has a higher accumulated value in the
251 aboveground part (2.45 mg/kg). Copper mainly accumulates in the roots of the plant. The maximum
252 copper concentration was recorded in corn roots at 10.05 mg/kg, and in alfalfa roots at 7.47 mg/kg.
253 The plant fractions of corn and alfalfa (root, aerial part, and grain) contain the same amount of cobalt
254 and are less than 0.01 mg/kg. Vanadium (4.93 mg/kg), cadmium (0.251 mg/kg), chromium (4.93
255 mg/kg), and copper (10.05 mg/kg) are more accumulated in corn roots.

256 The presence of higher concentrations of metals in the aboveground parts of alfalfa makes it more
257 suitable for phytoremediation. Namely, by removing alfalfa cuttings from contaminated locations over
258 a longer period of time, one part of heavy metals whose values exceed permissible concentrations in

259 soil. In order to obtain better and more complete data on this issue, it is necessary to conduct tests with
260 a larger number of samples.

261

262 **References**

263 [1] Šerbula SM, Miljkovic DJD, Kovacevic MR, Ilic AA 2012. Assessment of airborne heavy metal
264 pollution using plant parts and topsoil. *Ecotoxicology and Environmental Safety*. 76: 209–214.

265 [2] Radojević AA 2017. Biomonitoring vazduha i fitoremedijacija zemljišta upotrebom hrasta, smreke
266 i lešnika. Doktorska disertacija. Univerzitet u Beogradu. Tehnički fakultet u Boru: 1-221.

267 [3] Kalinović JV, Šerbula SM, Kalinović TS, Radojević AA, Jordanović J.S 2024. Pollution indices as
268 useful tools for comprehensive evaluation of the soil contamination degree in the vicinity of
269 mining and metallurgical complexes. *Hemijska industrija*. 78 (3): 265-279.

270 [4] Nielsen M, Nielsen O, Hoffmann L 2013. Improved inventory for heavy metal emissions from
271 stationary combustion plants, 1990-2009, Scientific Report from DCE – Danish Centre for
272 Environment and Energy. No. 68: 9-111.

273 [5] Pravilnik o utvrđivanju dozvoljenih količina štetnih i opasnih materija u zemljištu i metode
274 njihovog ispitivanja („Službene novine Federacije BiH“ broj 72/09): 14-18.

275 [6] Federalni zavod za agropedologiju 2016. Izvještaj o monitoringu zemljišta na području Općine
276 Zenica 2011-2015. godina. Finalni izvještaj. Sarajevo:1-90

277 [7] Beganović S, Goletić Š, Prčanović H, Duraković M, Karić A 2018: Research of the relationship
278 between the content of heavy metals in sediment dust and agricultural soil in the area
279 surrounding the steelworks Zenica. *International Journal of Advanced Research (IJAR)*,
280 Article DOI: 10.21474/IJAR01/7447. 6 (7): 969-978

281 [8] Federalni zavod za agropedologiju septembar 2010. Prisustvo neorganskih polutanata u nekim
282 biljnim kulturama na području općine Zenice. Sarajevo: 3-22.

283 [9] Alagić SČ 2014. Strategije biljaka u borbi protiv fitotoksičnih koncentracija metala kao ključni
284 preduslov uspješne fitoremedijacije. Čelijski mehanizmi, deo I. Zaštita materijala. 55 (3): 314-
285 317.

286 [10] Ahmetović M, Šestan I, Odobašić A, Keran H, Junuzović H 2022. The influence of
287 physicochemical characteristics of the soil on the distribution of heavy metals in plant
288 cultures. *Food in Health and Disease, scientific-professional journal of nutrition and dietetics*.
289 11 (2): 41-49.

290 [11] Lazarević B, Milan Poljak M 2019. Fiziologija bilja. Sveučilište u Zagrebu. Agronomski fakultet
291 Zavod za ishranu bilja. Zagreb: 32-59.

292 [12] Sarić M 1983. Fiziologija biljaka. Naučna knjiga. Beograd: 1-538.

293 [13] Lazarević B, Milan Poljak M, Horvat T, Karažija T, Stepinac D 2013. Uticaj nedostatka željeza
294 na usvajanje mikrohraniva kod kukuruza. *Agroekologija, ekološka poljoprivreda i zaštita*
295 *okoliša*. Sveučilište u Zagrebu. 48. hrvatski i 8. međunarodni simpozij agronoma. Dubrovnik.
296 Hrvatska: 88-92.

297 [14] Marschner P 2012. Marschner's Mineral Nutrition of Higher Plants. Third Edition: pp 200

298 [15] Grbeša D 2016. Hranidbena svojstva kukuruza. Zagreb: 7-136.

299 [16] Rastija M, Rastija D, Šimić D, Mihaljević I Decembar 2009. Prinosi i koncentracije Mn i Zn u
300 znu samooplodnih linija i hibrida kukuruza. *Poljoprivreda. Osijek*. 15 (2):1-8.

301 [17] Kerovec D 2010. Određivanje koncentracije teških metala pomoću AAS-a i ICP-OES u uzorcima
302 tla i biljke. Diplomski rad. Sveučilište Josipa Jurja Strossmayera u Osijeku. Poljoprivredni
303 fakultet u Osijeku: 1-57.

304 [18] Kastori R 1998. Fiziologija biljaka. Feljton. Novi Sad: 1-527.

305 [19] Vukadinović V, Vukadinović V 2011. Ishrana bilja. Sveučilište J. J. Strossmayera. Poljoprivredni
306 fakultet. Osijek: 223-229.

- 307 [20] Varga I, Antunović M, Kristek A 2019. Sadržaj teških metala u zrnu uljarica, 4th International
308 Scientific and Professional Conference (CRODMA 2019). Section II. October 11th. Varaždin.
309 Croatia: 101-106.
- 310 [21] Weitzer G rujan 2023. Status teških metala i potencijalno onečišćujućih elemenata u plodu graha
311 mahunara na tržištu grada Zagreba. Diplomski rad. Agronomski fakultet Sveučilišta u
312 Zagrebu:1-66.
- 313 [22] Chhotu DJ, Fulekar MH 2008. Phytotoxicity and remediation of heavy metala by alfalfa
314 (*Medicago sativa*) in soil-vermicompost media,.Adv. Nat. Appl. Sci. 2: 141-151.
- 315 [23] Bašić F, Kisić I, Mesić M, Butorac A 1998. Studija stanja i projekt rekultivacije tla i isplačne
316 jame. Agronom. fak. Sveučilišta u Zagrebu:1-100.
- 317 [24] Kabata-Pendias A and Pendias H 2001.Trace Elements in Soils and Plants. 3rd Edition, CRC
318 Press, Boca Raton. 403 p: 1-403.
- 319 [25] Kumar V, Singh J and Kumar P 2019. Heavy metals accumulation in crop plants: Sources,
320 response mechanisms, stress tolerance and their effects. Chapter 4 in: Contaminants in
321 Agriculture and Environment. Health Risks and Remediation: 39-57.
- 322 [26] Aydın SD, Pakyürek M 2020: Heavy metal accumulation potential in pomegranate fruits and
323 leaves grown in roadside orchards. PeerJ, 8, e8990: 1-20.
- 324 [27] Vrsaljko A 2023. Dinamika nakupljanja olova u listu i dijelovima ploda bajama.Poljoprivreda 29.
325 35-42. (1): 15-30.
- 326 [28] Kachenko A, Singh B 2006. Heavy metals contamination in vegetables grown in urban and metal
327 smelter contaminated sites in Australia. Water, Air, and Soil Pollution 169: 101–123.
- 328 [29] Rajković M, Stojanović M, Glamočlija Đ, Tošković D, Miletić V, Stefanović V, Lačnjevac Č
329 2012. Pšenica i teški metali. Journal of Engineering & Processing Management. Volume 4,
330 No. 1: 85-126
- 331 [30] Benavides MP, Gallego SM, Tomaro ML 2005. Cadmium toxicity in plants. Brazilian Journal of
332 Plant Physiology Vol. 17 (1): 131-136.
- 333 [31] Alagić SČ, Šerbula SS, Tošić SB, Pavlović AN, Petrović JV 2013. Bioaccumulation of Arsenic
334 and Cadmium in Birch and Lime from the Bor Region. Arch Environ Contam Toxicol, 65:
335 671-682
- 336 [32] Mallick S, Sinam G, Mishra RK, Sinha S 2010. Interactive effects of Cr and Fe treatments on
337 plants growth, nutrition and oxidative status in *Zea mays* L..Ecotoxicol and Environmental
338 Safety, 73: 987-995.
- 339 [33] Hu X, Wei X, Ling J, Chen J 2021:Cobalt: an essential micronutrient for plant growth?. Frontiers
340 in Plant Science: Volume 12. Article 768523: 1-24.
- 341 [34] Beganović S, Prčanović H, Duraković M, Adrović A august 2024. Application of bioremediation
342 technologies in the remediation of agricultural land in the area of Zenica. International Journal
343 of Advanced Research (IJAR), Article DOI: 10.21474/IJAR01/19346. 12 (08): 1079-1085
- 344 [35] Alagić SČ, Nujkić MM, Dimitrijević MD 2014. Strategije biljaka u borbi protiv fitotoksičnih
345 koncentracija metala kao ključni preduslov uspješne fitoremedijacije: Ekskluderi i
346 hiperakumulatori, deo II. Zaštita materijala 55 (4): 435-440.

347 .

348

349

350