


Jana Publication & Research

DENTIGEROUS CYST ENCOMPASSING THE HEMI-MAXILLA: A CASE REPORT

 06

 BioTech

 Institut Seni Indonesia Surakarta

Document Details

Submission ID

trn:oid::1:3192534280

Submission Date

Mar 24, 2025, 11:57 AM GMT+7

Download Date

Mar 24, 2025, 12:11 PM GMT+7

File Name

IJAR-50762.docx

File Size

984.3 KB

5 Pages

1,649 Words

9,459 Characters





64% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.




Filtered from the Report

- ▶ Bibliography
- ▶ Quoted Text

Match Groups

-  **24 Not Cited or Quoted 64%**
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 62%  Internet sources
- 61%  Publications
- 25%  Submitted works (Student Papers)

Match Groups

- **24 Not Cited or Quoted 64%**
Matches with neither in-text citation nor quotation marks
- **0 Missing Quotations 0%**
Matches that are still very similar to source material
- **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 62% Internet sources
- 61% Publications
- 25% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Publication	Buyukkurt, M.C.. "Dentigerous cyst associated with an ectopic tooth in the maxill...	15%
2	Internet	www.ijdr.in	14%
3	Internet	ejo.springeropen.com	8%
4	Internet	pmc.ncbi.nlm.nih.gov	7%
5	Publication	Evren Ustuner, Suat Fitoz, Cetin Atasoy, Ilhan Erden, Serdar Akyar. "Bilateral maxi...	6%
6	Internet	www.ncbi.nlm.nih.gov	3%
7	Internet	www.thieme-connect.com	3%
8	Internet	datospdf.com	2%
9	Internet	storage.googleapis.com	1%
10	Internet	www.researchgate.net	1%

11	Internet	www.sciencegate.app	<1%
12	Internet	www.ajcr.com	<1%
13	Publication	Manuela Maltoni, Ivano Maltoni, Giorgia Santucci, Daniela Guiducci, Francesca Cr...	<1%
14	Internet	journals.lww.com	<1%

DENTIGEROUS CYST ENCOMPASSING THE HEMI-MAXILLA: A CASE REPORT.

Abstract.

Dentigerous cysts are the second most common odontogenic cysts after radicular cysts. Dentigerous cysts are benign odontogenic cysts associated with the crowns of permanent teeth. Cysts involve impacted, unerupted permanent teeth, supernumerary teeth, odontomas, and, rarely, deciduous teeth. The ectopic tooth embedded in a dentigerous cyst of the maxillary antrum is extremely rare in daily clinical practice. We are reporting a case of the ectopic left maxillary third molar tooth and the associated dentigerous cyst in the maxillary antrum

BACKGROUND.

Dentigerous cysts are the second most common odontogenic cysts after radicular cysts². Dentigerous cysts are benign odontogenic cysts associated with the crowns of permanent teeth. Cysts involve impacted, unerupted permanent teeth, supernumerary teeth, odontomas, and, rarely, deciduous teeth. In 75% of the cases, they are located in the mandible¹. The mandibular third molar and maxillary canine are involved most frequently. Dentigerous cysts are usually painless but may cause facial swelling and delayed tooth eruption. The usual presentation is in second or third decade of life. Extensive maxillary involvement and childhood presentation are rare⁷.

Introduction.

Displacement of the impacted tooth can occur due to a dentigerous cyst in an ectopic location like the mandibular condyle¹⁴, coronoid process¹⁴, nasal cavity, and maxillary antrum. There are possible etiological theories that explain the

pathogenesis of ectopic teeth, including developmental abnormalities, trauma, infection, and pathological problems like dentigerous cysts. However, the exact mechanism remains unclear. Dentigerous cysts are usually not associated with pain. However, if they are associated with an infection, there is a painful swelling⁶.

Case presentation.

A 40 year old female patient was referred to the oral and maxillofacial surgery department by a nearby dentist complaining of pain over the left infraorbital region. On primary evaluation tenderness over that region was noticed. Intraoral examination revealed the presence of distomolar and absence of third molar. On further evaluation with OPG (Fig 1) ectopically placed left upper third molar

was noticed and CBCT was advised.



Fig 1 .Pre- operative OPG

On Evaluation Single, unilateral, mostly well defined radiolucent lesion with variably corticated borders was noted in left body of maxilla Lesion extending antero – posteriorly from distal aspect of 25 to the pterygoid plates. Superiorly extending upto inferior border of orbit. Inferiorly lesion extends to 4 mm apical to the alveolar crest. Bucco- palatally causing buccal cortical plate expansion.

The internal structure was radiolucent with mucosal thickening. Buccal cortical plate expansion with multiple areas of perforation and extreme thinning and mild erosion of palatal cortical plate noted suggestive of a highly expansile lesion. Posteriorly lesion was causing perforation of posterior border of maxilla, involving pterygopalatine fossa, greater palatine foramen and pterygoid plates.



Fig 3 Perforation of inferior border of orbit




Fig 2 CBCT showing 28 in the medial wall of zygomatic bone

Complete obliteration of sinus noted with erosion of lateral wall of nasal cavity(fig 4), erosion of infraorbital canal and perforation of inferior border of orbit(fig 3). Vertically the impacted 28 was noted within sinus, in the medial border of zygoma.(Fig 2)




Fig 4

A Single fused root was noted. Root apex was in contact with and causing perforation of inferior border of the orbit. Lesion inferiorly extended to the apical third level of roots of 25, 26 and 27 and causing external root resorption.

Under GA a Caldwell-Luc incision was given over left posterior maxilla. Mucoperiosteal flap was raised. As the bone was too thin over the area, a natural window was opened into the maxillary sinus(fig 5)

(fig 5)



Complete removal of cyst along with the impacted 28 was done.(fig 6)Chemical cauterisation with carnoy's solution` was not done to prevent injury to orbital contents.



Fig 6

Sinus was packed with ribbon gauze soaked in betadine, which was removed after 48 hours. Biopsy report was indicative of dentigerous cyst.(fig 7)




Fig 7 .

The patient was given IV antibiotics, analgesics and steroids and discharged after 2 days as the postoperative period was uneventful.

Patient was reviewed after a week and satisfactory healing noticed .

After one month, healing was adequate with good function and patient satisfaction.

DISCUSSION.

Dentigerous cysts, also known as follicular cysts, are the second most frequent odontogenic cysts after those associated with the roots of the teeth⁸ (periapical cysts), originating from the weakened enamel epithelium during tooth crown development. They are most frequently found in the mandible (70%) than the maxilla (30%). The incidence is 14-20% and is slightly more prevalent among males (1.6:1) occurring in the second and third decades of life . Uncommon presentation of

4
1
2
third molar in the lateral wall (medial border of zygoma) of left maxilla prompted us to publish this case.

1
9
1
7
2
Tooth development results from a complicated multistep interaction between the oral epithelium and the underlying mesenchymal tissue. A series of complex tissue interactions result in the formation of mature teeth. Abnormal tissue interactions during tooth development may potentially result in ectopic tooth development and eruption. Ectopic eruption of a tooth into the dental environment is common whereas ectopic eruption of a tooth in other sites is rare. One such site for ectopic tooth eruption in a nondental location is the maxillary sinus. Due to its rarity, there is a dearth of literature discussing this entity. Ectopic eruption may result due to one of the three processes: developmental disturbance, pathological process and iatrogenic activity⁸. Tooth eruption into the maxillary sinus may cause sinusitis, the treatment of which (if infected) is surgical removal. We present a case of an ectopic maxillary third molar, which presented in the left maxillary sinus and was removed via a Caldwell-Luc procedure.

1
6
6
2
The treatment of an ectopic tooth in the maxillary sinus is usually removal, as if left untreated, it has the tendency to form a cyst or tumor. The differential diagnosis of a dentigerous cyst includes other odontogenic cysts, such as radicular cysts, odontogenic keratocysts, and odontogenic tumors, such as ameloblastoma, odontoma, odontogenic fibroma, and cementomas¹². However, mucoceles, retention cysts, and pseudocysts are also included in the differential diagnosis when a maxillary sinus cyst is visualized involving maxillary expansion. Histologically, dentigerous cysts are lined by a layer of nonkeratinized stratified squamous epithelium, with a

surrounding wall of thin connective tissue containing odontogenic epithelial rests¹⁰.

The standard treatment for a dentigerous cyst is enucleation and extraction of the cyst-associated impacted or unerupted tooth. Dentigerous cysts of the maxillary sinus, and the impacted tooth within, are often easily removed via a Caldwell-Luc procedure⁵. Removal of the entire cyst with the impacted tooth is a main treatment to prevent recurrence of the cyst. Marsupialization is another advisable treatment to preserve the cyst-associated tooth and promote its eruption. Large cysts that involve serious loss of bone and that thin the bone dangerously are often treated via marsupialization¹³. It was successfully used to minimize the amount of maxillary destruction and surgical morbidity that might have resulted from the immediate enucleation of the lesion for dentigerous cysts within the maxillary antrum.

The major disadvantage of marsupialization is recurrence or persistence of the lesion¹⁵. Although the traditional Caldwell-Luc procedure provides a direct view into the maxillary sinus, it is associated with more morbidity than transnasal endoscopy. Transnasal extradition of the tooth may be attempted if the tooth is small and sited near the maxillary ostium.

CONCLUSION.

The occurrence of an ectopic tooth in maxillary sinus and its association with dentigerous cyst is a rare phenomenon. Its presence is asymptomatic until it is infected. Routine X-rays and CBCT can confirm the diagnosis. The standard treatment for dentigerous cyst is enucleation and extraction of associated tooth via Caldwell Luc approach. In large cyst, an initial marsupialisation to diminish the size, followed by enucleation has been advocated.

REFERENCES.

1. Mamatha NS, KriShNaMoorthy B, Savitha JK, Bhai P. Diagnostic CBCT in dentigerous cyst with ectopic third molar in the maxillary sinus—a case report. *Journal of Clinical and Diagnostic Research: JCDR*. 2014 Jun;8(6):ZD07.
2. John T, Shekhar MG, Koshy M. Dentigerous cyst associated with supernumerary teeth: A report of three cases. *J Clin Diagn Res*. 2010 Jun;4:2601-6.
3. Girish G, Kumar M, Umashankar D, Sharma R, Veeresh M, Bhandari A. Case report dentigerous cyst in maxillary sinus: A Rare Occurrence. *Int J Oral Maxillofac Pathol*. 2011;2:19-23.
4. Alhashim FY, Almarhoon FS, Alhashim HY, Moumen A. Endonasal endoscopic management of different cases of dentigerous cysts and ectopic teeth. *Journal of Surgical Case Reports*. 2021 Apr;2021(4):rjab099.
5. Marino MJ, Luong A, Yao WC, Citardi MJ. Management of odontogenic cysts by endonasal endoscopic techniques: a systematic review and case series. *American Journal of Rhinology & Allergy*. 2018 Jan;32(1):40-5.
6. Amin ZA, Amran M, Khairudin A. Removal of extensive maxillary dentigerous cyst via a Caldwell-Luc procedure. *Archives of Orofacial Sciences*. 2008;3(2):48-51.
7. Prasad TS, Sujatha G, Niazi TM, Rajesh P. Dentigerous cyst associated with an ectopic third molar in the maxillary sinus: a rare entity. *Indian journal of dental research*. 2007 Jul 1;18(3):141-3.
8. MacDonald-Jankowski D, Chan KC. Clinical presentation of dentigerous cysts: systematic review. *Asian Journal of Oral and Maxillofacial Surgery*. 2005 Jun 1;17(2):109-20.
9. Austin RP, Nelson BL. Sine qua non: dentigerous cyst. *Head and neck pathology*. 2021 Dec;15(4):1261-4.
10. Nahajowski M, Hnitecka S, Antoszevska-Smith J, Rumin K, Dubowik M, Sarul M. Factors influencing an eruption of teeth associated with a dentigerous cyst: A systematic review and meta-analysis. *BMC Oral Health*. 2021 Dec;21:1-1.
11. Assael LA. Surgical management of odontogenic cysts and tumors. *Principles of oral and maxillofacial surgery*. 1992;2:685-8.
12. Regezi JA. Cyst and cystlike lesions. Regezi JA, Sciubba J, Pogrel M A. *Atlas of Oral and Maxillofacial Pathology*. Philadelphia: WB Saunders. 2000.
13. Shear M. *Cysts of the oral regions*. Wright Oxford, England. 1992.
14. Yusuf H, Quayle AA. Intracondylar tooth. *International journal of oral and maxillofacial surgery*. 1989 Dec 1;18(6):323.
15. Kasat VO, Karjodkar FR, Laddha RS. Dentigerous cyst associated with an ectopic third molar in the maxillary sinus: A case report and review of literature. *Contemporary clinical dentistry*. 2012 Jul 1;3(3):373-6.