

# Jana Publication & Research

## SURGICAL MANAGEMENT OF MALLEOLAR FRACTURES IN ADULTS

 18

 BioTech

 Institut Seni Indonesia Surakarta

---

### Document Details

Submission ID

trn:oid::1:3201395526

Submission Date

Apr 1, 2025, 11:58 AM GMT+7

Download Date

Apr 1, 2025, 12:52 PM GMT+7

File Name

IJAR-50844.docx

File Size

392.3 KB

10 Pages

2,030 Words

12,189 Characters

# 42% Overall Similarity





The combined total of all matches, including overlapping sources, for each database.

## Filtered from the Report




- ▶ Bibliography
- ▶ Quoted Text

---

## Match Groups

-  **52 Not Cited or Quoted 42%**  
Matches with neither in-text citation nor quotation marks
-  **1 Missing Quotations 1%**  
Matches that are still very similar to source material
-  **0 Missing Citation 0%**  
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

## Top Sources

- 37%  Internet sources
- 30%  Publications
- 11%  Submitted works (Student Papers)

### Match Groups

- **52 Not Cited or Quoted 42%**  
Matches with neither in-text citation nor quotation marks
- **1 Missing Quotations 1%**  
Matches that are still very similar to source material
- **0 Missing Citation 0%**  
Matches that have quotation marks, but no in-text citation
- **0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

### Top Sources

- 37% Internet sources
- 30% Publications
- 11% Submitted works (Student Papers)

### Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

|    |                |  |    |
|----|----------------|--|----|
| 1  | Internet       |  |    |
|    |                | www.researchgate.net                         | 5% |
| 2  | Internet       |  |    |
|    |                | repository-tnmgrmu.ac.in                     | 5% |
| 3  | Student papers |  |    |
|    |                | KLE Academy of Higher Education and Research | 5% |
| 4  | Internet       |  |    |
|    |                | www.orthopaper.com                           | 4% |
| 5  | Internet       |  |    |
|    |                | mafiadoc.com                                 | 3% |
| 6  | Internet       |  |    |
|    |                | orca.cardiff.ac.uk                           | 3% |
| 7  | Internet       |  |    |
|    |                | www.worldwidejournals.com                    | 2% |
| 8  | Internet       |  |    |
|    |                | iosrjournals.org                             | 1% |
| 9  | Internet       |  |    |
|    |                | www.scribd.com                               | 1% |
| 10 | Internet       |  |    |
|    |                | fdocuments.net                               | 1% |

|    |             |   |     |
|----|-------------|---|-----|
| 11 | Publication | Christian CMA Donken. "Surgical versus conservative interventions for treating a...   | <1% |
| 12 | Publication | Sankara Rao S, Chandra Sekaram Naidu M, Dharma Rao V, Arun Kumar V, Sravya ...        | <1% |
| 13 | Publication | Seçkin Özcan, Nizamettin Koçkara, Yalkın Camurcu, Hakan Yurten. "Magnetic Res...      | <1% |
| 14 | Publication | European Surgical Orthopaedics and Traumatology, 2014.                                | <1% |
| 15 | Internet    | gavinpublishers.com   | <1% |
| 16 | Publication | Bedri Karaismailoglu, Onur Yıldırım, Lercan Aslan, Necip Selcuk Yontar, Mehmet F...   | <1% |
| 17 | Internet    | mail.ijoro.org  | <1% |
| 18 | Publication | Chih-Wei Chang, Yen-Nien Chen, Guan-Heng Jhong, Kuo-Chih Su, Chun-Ting Li. "A ...     | <1% |
| 19 | Publication | Mario Herrera-Perez, Pablo Martín-Vélez, Diego Rendón-Díaz, José Luis Pais-Brito. ... | <1% |
| 20 | Internet    | www.medpulse.in   | <1% |
| 21 | Internet    | medicine.uiowa.edu  | <1% |
| 22 | Publication | Błażej Grzegorz Wójtowicz, Katarzyna Chawrylak, Jędrzej Lesman, Hubert Makow...       | <1% |
| 23 | Publication | Hasan Ulas Ogur, Firat Seyfettinoğlu, ÜmitX Ümit Tuhanioğlu, Hakan Cicek, Sefa ...    | <1% |
| 24 | Publication | Ram K. Alluri, Eric W. Tan. "Indications, Postoperative Management, and Advance...    | <1% |

|    |             |   |     |
|----|-------------|---|-----|
| 25 | Publication | S KOZIN. "Pediatric Distal Radius Fractures", Fractures and Injuries of the Distal R... | <1% |
| 26 | Internet    | www.msdlatinamerica.com   | <1% |
| 27 | Internet    | www.orthobullets.com  | <1% |
| 28 | Publication | "ECTES-Abstracts 2017", European Journal of Trauma and Emergency Surgery, 2017          | <1% |
| 29 | Internet    | www.ijoro.org   | <1% |
| 30 | Publication | Michael T. Archdeacon, Jeffrey O. Anglen, Robert F. Ostrum, Dolfi Herscovici. "Pre...   | <1% |

# SURGICAL MANAGEMENT OF MALLEOLAR FRACTURES IN ADULTS

## ABSTRACT

**Background and objectives:** Ankle fractures are one of the most common lower limb fractures. Acute injuries to ankle joint gain importance because body weight is transmitted through it. Conservative treatment can provide satisfactory results if anatomical reduction is maintained in many of the ankle fractures. The other unstable, displaced and open fractures require surgical treatment involving open reduction and internal fixation. Objectives were to study the efficacy, functional outcome and complications of surgical management of malleolar fractures of ankle.

**Methods:** A prospective study involving 30 patients admitted with malleolar fractures, which are classified according to Lauge-Hansen's classification and treated according to the AO system are presented. The follow-up period was for a minimum of 6 months. Assessment of the outcome was done using Olerud and Molander scoring system.

**RESULTS:** In this study, excellent results were found in 18(60%) cases, good results were found in 06(20%) cases, fair results were results were found in 04(13.33%) cases, poor results were found in 02(6.66%) cases according to Olerud-Molander ankle score.

**INTERPRETATION AND CONCLUSION:** The most prevalent cause of malleolar fractures was a vehicular accident (RTA). Supination external rotation injuries were the most common. Regardless of the kind of surgery, anatomical reduction is necessary for a successful functional result. The complications were minimal and rectified within two weeks.

## INTRODUCTION:

Ankle fractures are some of the most frequent fractures of the lower limb. They account for 10% of all fractures, accounting for a large amount of the trauma burden, although only around 2% of ankle fractures are open <sup>(1)</sup>. Twisting injuries, vehicular accidents, and falls are the most prevalent causes of ankle fractures, followed by sports injuries <sup>(2)</sup>. Diabetes mellitus and obesity are common comorbidities in adults above forty <sup>(3)</sup>. Ankle fractures are classified using the **Lauge-Hansen Classification** <sup>(4)</sup>.

1 Malleolar fracture treatment can be divided into three approaches: nonsurgical, staged, and surgical. In general, nonsurgical treatment for stable ankle fractures and surgical treatment for unstable ankle fractures are acceptable <sup>(5)</sup>.

5 Non-operative treatment with immobilization can give satisfactory results if anatomical reduction is maintained and closely followed. However, cast immobilization can also lead to muscle atrophy, cartilage degeneration, and a stiff, painful, and swollen joint <sup>(6)</sup>. Staged treatment with external fixation and delayed internal fixation is done in high-energy injuries or fracture dislocations with compromised surrounding soft tissue <sup>(7)</sup>. Open reduction and internal fixation of the fractured bone with devices such as metal plates, screws, tension bands, or external fixation are used in surgical treatment <sup>(8)</sup>. These surgical procedures are designed to give anatomical reduction and stability, allowing for quick mobilization. Complications from surgery include wound infection, pulmonary embolism, implant failure, amputation, and redo surgery <sup>(9)</sup>.

1 Although a non-operative approach is applicable in undisplaced fractures, surgical management has been favored for more than 40 years <sup>(10)</sup>. Malleolar fractures that are not properly treated might result in painful arthritis <sup>(11)</sup>. The importance of perfect anatomical reduction and firm fixation in malleolar fractures cannot be overstated <sup>(12)</sup>.

#### METHODOLOGY:

20 This is a prospective study involving 30 patients with fresh unimalleolar, bimalleolar and trimalleolar fractures, who were treated at Dr. PSIMS & RF, China Avutapally . AP and lateral view Xrays of the ankle joints of the patients, were taken. A detailed history of the mode of injury and any associated medical illness is recorded. The patient's general condition was assessed, and he or she were then thoroughly examined.

#### 23 SURGICAL TECHNIQUE:

25 All of the procedures were performed under spinal anesthesia without tourniquet. The fibula was the first bone to be exposed. After removing the hematoma from the fracture site, fracture is reduced and fixed. After clearing the soft tissue interposition, the fracture site was reduced. Tension bandwiring, cancellous screws, K-wire fixation, or semitubular plating were used for ORIF of malleolar fractures. Follow up was done for 4 months according to Olerud-Molander Ankle Score.

## RESULTS

### MODE OF INJURY

**Table 1:**

#### MODE OF INJURY

|                            |   |    |
|----------------------------|---|----|
| Road Traffic Accident      | - | 16 |
| Twisting injury/ self fall | - | 10 |
| Fall from height           | - | 4  |

### MALLEOLARITY

**Table 2:**

|                        |    |
|------------------------|----|
| Unimalleolar fractures | 10 |
| -                      |    |
| Bimalleolar fractures  | 18 |
| -                      |    |
| Trimalleolar fractures | 2  |
| -                      |    |

In present study 60% were bimalleolar fracture, 33.33% were unimalleolar fractures and 6.66% were trimalleolar fractures.

### INJURY PATTERN

Injury pattern according to Lauge-Hansen classification is as follows:

**Table 3:**

| Injury pattern               | Number of cases |
|------------------------------|-----------------|
| Supination-Adduction         | 3               |
| Supination-External rotation | 18              |
| Pronation-Abduction          | 2               |



|                             |   |
|-----------------------------|---|
| Pronation-External rotation | 7 |
|-----------------------------|---|

Supination-external rotation injuries are the commonest (60%) and pronation abduction injuries are the least common injuries in this study (6.66%).

## RESULTS

The results were analyzed based on Olerud-Molander ankle score.

**Table 4: Olerud-Molander ankle score**

| Parameters          |  | Score |
|---------------------|--|-------|
| 1. Pain             | Never                                  | 25    |
|                     | While walking on uneven surface        | 20    |
|                     | While walking on even surface outdoors | 10    |
|                     | While walking indoors                  | 05    |
|                     | Constant and severe                    | 0     |
| 2. Stiffness        | None                                   | 10    |
|                     | Present                                | 0     |
| 3. Swelling         | None                                   | 10    |
|                     | Only evenings                          | 05    |
|                     | Constant                               | 0     |
| 4. Stair climbing   | No problems                            | 10    |
|                     | Impaired                               | 05    |
|                     | Impossible                             | 0     |
| 5. Running          | Possible                               | 05    |
|                     | Impossible                             | 0     |
| 6. Jumping          | Possible                               | 05    |
|                     | Impossible                             | 0     |
| 7. Squating         | No problems                            | 05    |
|                     | Impossible                             | 0     |
| 8. Type of supports | None                                   | 10    |
|                     | Taping, wrapping                       | 05    |
|                     | Stick or crutch                        | 0     |

|                                      |                                      |    |
|--------------------------------------|--------------------------------------|----|
|                                      |                                      |    |
| 9. Work and Activities of daily life | Same as before injury                | 20 |
|                                      | Change to part time work/Simpler job | 15 |
|                                      | Severely impaired work capacity      | 0  |

Poor: 0 %-60 %, Fair: 61 %-80 %, Good: 81 %-90 %, Excellent: 91 %-100 %.

Follow up was done until fracture union. Results were analyzed clinically & radiographically. Union was seen in all the fractures at the end of 10 weeks. Overall functional outcome for our patients is as follows:

**Table 5:**

| Results          | No. of patients | Percentage |
|------------------|-----------------|------------|
| Excellent (>90%) | 18              | 60%        |
| Good (81%-90%)   | 06              | 20%        |
| Fair (61%-80%)   | 04              | 13.33%     |
| Poor (<60%)      | 02              | 6.66%      |

## COMPLICATIONS

The following complications were seen, postoperatively.

- Superficial infections with and without skin necrosis.



Fig 1 : Exposure of Fracture Site



Fig 2 : Fixation with C.C. Screw

**Fig1: cc screw fixation to medial malleolus**



Fig 1 : Draping of the Limb



Fig 2 : Lateral Skin Incision for Lateral Malleolus



Fig 3 : Exposure of Fracture in Lateral Malleolus



Fig 4 : Fracture Site Reduced



Fig : Fracture Site Reduced and Fixed with Semi Tubular Plate



Fig 6 : Fixation of Semi Tubular Plate with 3.5 mm Cortical Screw



Fig 7 : Semitubular Plate Closed with Soft Tissue



Fig 8 : Skin Closure

## DISCUSSION:

16 Ankle fractures are among the most prevalent lower limb fractures, comprising 10% of all fractures, with only a small proportion (2%) being open fractures [13]. The most common causes include twisting injuries, road traffic accidents, falls from height, and sports injuries [14]. In this study, road traffic accidents were the leading cause (53.33%), followed by twisting injuries or self-falls (33.33%) and falls from height (13.33%). These findings highlight the importance of preventive measures such as road safety awareness and protective gear for sports enthusiasts to reduce the incidence of ankle fractures [15].

9 27 Regarding fracture patterns, bimalleolar fractures (60%) were the most common type, followed by unimalleolar fractures (33.33%) and trimalleolar fractures (6.66%). The Lauge-Hansen classification revealed that supination-external rotation injuries were the most frequent (60%), while pronation-abduction injuries were the least common (6.66%) [4].



6 These findings align with previous studies indicating that rotational mechanisms, particularly supination-external rotation, are the most common causes of ankle fractures.

1 Treatment approaches varied based on fracture stability and soft tissue conditions. While non-surgical treatment with immobilization remains an option for stable fractures, its complications, such as muscle atrophy, cartilage degeneration, and joint stiffness, emphasize the need for careful patient selection and follow-up [16]. Staged treatment with external fixation and delayed internal fixation was utilized in high-energy injuries or cases with compromised soft tissue [3]. Surgical treatment remains the gold standard for unstable fractures, providing anatomical reduction and early mobilization [17]. However, complications such as wound infection and implant failure were noted, underscoring the importance of meticulous surgical technique and postoperative care [18].

22 9 The functional outcomes, assessed using the Olerud-Molander ankle score, revealed excellent results in 60% of patients, good outcomes in 20%, fair outcomes in 13.33%, and poor outcomes in 6.66% [19]. The high percentage of excellent and good results suggests that surgical intervention, when indicated, yields favorable functional outcomes. Nonetheless, the presence of fair and poor outcomes highlights the need for individualized treatment planning, rehabilitation, and patient education to optimize recovery and prevent long-term complications such as post-traumatic arthritis [20].

#### 18 SUMMARY:

21 1 7 7 15 Thirty cases of malleolar fractures, surgically repaired using various techniques, were followed in this study. Road traffic accidents were the leading cause of injury, accounting for 53.33 % of patients. Supination-external rotation (Lauge-Hansen's classification) type was the most common type of injury, comprising 60% of all the cases. Lateral malleolus was fixed with a semi-tubular plate / tension band wiring, and medial malleolus with malleolar screws / tension band wiring. The complication faced was postoperative skin infection, seen in 2 patients. After the study concluded, excellent to good results were seen in 24 (80%) cases, whereas 4 (13.33%) cases had fair results and 2 (6.66%) had poor results.

#### 10 2 CONCLUSION:

The most prevalent type of malleolar fracture is supination-external rotation injury, which is also the most frequently associated type with dislocations and complications. The Pronation-External rotation type

2 had minimal complications. A good functional outcome is achieved through perfect anatomical reduction and restoration of articular congruity, as well as early surgical fixation with appropriate implants. 2 By restoring sufficient stability and mobility to the ankle joint, a good 4 functional outcome was achieved. Understanding the mechanism of injury is critical for successful reduction and internal fixation. 26 When using one third tubular plate to fix the lateral malleolus, the lateral malleolar bend should be replicated. For lateral ankle stability, the length of fibula must be maintained. In all intraarticular fractures, 4 anatomical reduction is critical, especially if a weight-bearing joint such as the ankle is involved. Open reduction and internal fixation ensures a high level of reduction. Nearly 90% of patients had satisfactory outcomes. The more severe the injury, the less satisfactory the outcome. 30 Non-union of the malleoli because of soft tissue interposition should be avoided. Large percentage of complications were minor and resolved within two weeks. For small fracture fragments and osteoporotic bones, tension band wiring is preferred.

## REFERENCES

1. Court-Brown CM, McBirnie J. The epidemiology of tibial fractures. *J Bone Joint Surg Br.* 1995;77(3):417-21.
2. Kannus P, Palvanen M, Niemi S, Parkkari J, Järvinen M. Increasing number and incidence of low-trauma ankle fractures in elderly people: Finnish statistics during 1970–2000 and projections for the future. *Bone.* 2002;31(3):430-3.
3. Michelson JD. Ankle fractures resulting from rotational injuries. *J Am Acad Orthop Surg.* 2003;11(6):403-12.
4. Lauge-Hansen N. Fractures of the ankle. II. Combined experimental-surgical and experimental-roentgenologic investigations. *Arch Surg.* 1950;60(5):957-85.
5. Sanders DW, Tieszer C, Corbett B. Operative versus nonoperative treatment of unstable lateral malleolar fractures: a randomized multicenter trial. *J Orthop Trauma.* 2012;26(3):129-34.
6. Weber M, Simpson LA. Treatment of ankle fractures in the elderly: weight-bearing in a functional orthosis or immobilization in a cast? *J Bone Joint Surg Am.* 1984;66(6):847-62.
7. Bhandari M, Sprague S, Ayeni OR, Petrisor BA. Complications in the management of ankle fractures: a systematic review and meta-

- analysis of randomized controlled trials. *J Orthop Trauma*. 2012;26(2):47-51.
8. Herscovici D Jr, Scaduto JM, Infante A. Conservative treatment of isolated fractures of the medial malleolus. *J Bone Joint Surg Br*. 2007;89(1):89-93.
  9. Miller AN, Carroll EA, Parker RJ, Helfet DL, Lorich DG. Posterior malleolar stabilization of syndesmotic injuries is equivalent to screw fixation. *Clin Orthop Relat Res*. 2010;468(4):1129-35.
  10. Makwana NK, Bhowal B, Harper WM, Hui AW. Conservative versus operative treatment for displaced ankle fractures in patients over 55 years of age: a prospective, randomised study. *J Bone Joint Surg Br*. 2001;83(4):525-9.
  11. Stufkens SA, van den Bekerom MP, Kerkhoffs GM, Hintermann B, van Dijk CN. Long-term outcome after 1822 operatively treated ankle fractures: a systematic review of the literature. *Injury*. 2011;42(2):119-27.
  12. Schepers T, De Vries MR, Van Lieshout EM, Van der Elst M. The timing of ankle fracture surgery and the effect on infectious complications; a case series and systematic review of the literature. *Int Orthop*. 2013;37(3):489-95.
  13. Court-Brown CM, Caesar B. Epidemiology of adult fractures: A review. *Injury*. 2006;37(8):691-7.
  14. Lindsjö U. Operative treatment of ankle fractures. *Acta Orthop Scand*. 1985;56(2):201-6.
  15. Kannus P, et al. Epidemiology of osteoporotic ankle fractures in elderly persons. *Arch Intern Med*. 1999;159(8):878-85.
  16. Meena S, et al. Epidemiology of ankle fractures. *J Foot Ankle Surg*. 2015;54(3):432-8.
  17. Pakarinen HJ, et al. Operative treatment of ankle fractures in the elderly. *J Bone Joint Surg Am*. 2011;93(7):661-8.
  18. Lindsey RW, et al. Complications of ankle fracture management. *Clin Orthop Relat Res*. 1986;(206):152-8.
  19. Olerud C, Molander H. A scoring scale for symptom evaluation after ankle fracture. *Arch Orthop Trauma Surg*. 1984;103(3):190-4.
  20. Smeeing DP, et al. Weight-bearing versus non-weight-bearing after surgical treatment of ankle fractures. *J Bone Joint Surg Am*. 2018;100(13):1119-26.