

Jana Publication & Research

A rare case of cardiac metastasis from uterine leiomyosarcoma: A case report

 24 BioTech Institut Seni Indonesia Surakarta

Document Details

Submission ID

trn:oid:::1:3213753804

Submission Date

Apr 12, 2025, 12:19 PM GMT+7

Download Date

Apr 12, 2025, 2:11 PM GMT+7

File Name

IJAR-51027.docx

File Size

1.4 MB

6 Pages**1,456 Words****8,924 Characters**





35% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.




Filtered from the Report

- Bibliography
- Quoted Text

Match Groups

-  **25 Not Cited or Quoted 35%**
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 32%  Internet sources
- 26%  Publications
- 5%  Submitted works (Student Papers)

Match Groups

- 25 Not Cited or Quoted 35%**
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 32% Internet sources
- 26% Publications
- 5% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	jnsbm.com	6%
2	Internet	obgyn.onlinelibrary.wiley.com	5%
3	Publication	Aki Maebayashi, Masaji Nagaishi, Takahiro Nakajima, Mitsumasa Hata, Tang Xiao...	4%
4	Internet	www.hindawi.com	3%
5	Internet	journals.lww.com	2%
6	Internet	atm.amegroups.com	2%
7	Internet	coek.info	2%
8	Internet	crimsonsusa693.weebly.com	2%
9	Internet	auctoresonline.org	1%
10	Publication	Gabriel Stefan, Simona Stancu, Adrian Zugravu, Laura Predescu, Simona Cinca, G...	1%

11	Internet	www.ijpmonline.org	1%
12	Internet	jcp.bmj.com	1%
13	Internet	mafiadoc.com	1%
14	Publication	Girolamo Catapano-Minotti. "A case of metastatic melanoma in the heart: does e...	<1%
15	Internet	www.dovepress.com	<1%
16	Publication	Koji Numata, Nozomi Urata, Yuta Nakayama, Mihwa Ju et al. "Solitary Cardiac Me...	<1%
17	Publication	Ilaria Tortorelli, Federico Navarria, Antonio Di Maggio, Alberto Banzato et al. "Tra...	<1%
18	Publication	Junxiu Liu, Zijie Wang. "Advances in the Preoperative Identification of Uterine Sar...	<1%

A rare case of cardiac metastasis from uterine leiomyosarcoma: A case report

ABSTRACT:

Uterine leiomyosarcomas (ULMS) are rare malignant tumors of smooth muscle origin. Cardiac metastases from ULMS are extremely uncommon and remain underreported. Herein, we present the case of a 57-year-old woman with locally advanced uterine leiomyosarcoma who developed an intracardiac metastasis. Following chemotherapy, radiotherapy, and subsequent surgery for the primary tumor, postoperative imaging unexpectedly revealed a massive right atrial thrombus extending into the inferior vena cava, later confirmed to be metastatic. Despite surgical intervention, complete resection of the thrombus was not achieved, leading to the patient's demise from progressive heart failure. This case emphasizes the potential for cardiac involvement in ULMS and highlights the importance of considering metastatic cardiac lesions in the differential diagnosis of intracavitary masses.

Keywords: Uterine leiomyosarcoma, cardiac metastasis, malignant tumor

INTRODUCTION:

Uterine sarcomas are rare malignant tumors arising from the myometrial smooth muscle. (2) Cardiac metastases are among the least known and highly debated issues in oncology. The incidence of cardiac metastases reported in literature is highly variable, ranging from 2.3% and 18.3%. Endocardial metastases are usually the result of the invasion from the bloodstream through the heart's chambers with intracavitary lodging. (3).

Given the rarity of uterine sarcomas and intracardiac metastases, intracardiac metastases from uterine sarcomas have been very rarely reported in the literature. We report a case of 57-year-old woman with a locally advanced leiomyosarcoma of the uterus, that underwent radio and chemotherapy, followed by surgery. Post-operative investigation showed an intracardiac metastasis of the uterine leiomyosarcoma.

CASE REPORT:

A 57-year-old woman with no significant medical history, G3P3, postmenopausal for 9 years, being treated for a locally advanced uterine leiomyosarcoma, initial FIGO stage IIIC1, initially deemed unresectable. The patient underwent first-line chemotherapy with doxorubicin and dacarbazine, completing 4 cycles. Interim evaluation continued to indicate unresectability, prompting an additional 2 cycles of chemotherapy. In addition, the patient received radiotherapy to the pelvis with a boost to the parametrial area.

Radiological Findings:

Initial Pelvic MRI:

- Large uterine tumor causing a thrombotic tumor involvement of the left ovarian vein.
- Partial thrombosis of the left internal iliac vein.

Initial Thoraco-Abdomino-Pelvic CT (TAP):

- Large, locally advanced uterine tumor, FIGO stage IIIC1, associated with a thrombotic involvement of the left ovarian vein.

Post-Treatment Imaging Findings (MRI):

- Significant regression in the size of the large uterine cervico-isthmic tumor, predominantly on the left, still showing extension to the left parametrial region and ipsilateral fallopian tube, consistent with sarcomatous pathology.
- Extensive fibrotic changes in the pelvic region, particularly affecting the bladder and rectal walls, suggestive of post-radiation changes.

Clinical Examination after chemo and radiotherapy:**On Speculum Exam (SPC):**

- Cervix appears macroscopically normal with peri-orificial erythema.

On Vaginal Exam (TV):

- Cervix is supple. Fornices appear soft and free. A rounded mass is palpable through the left fornix, possibly indicative of tumor extension into the left ovary.

On Rectal Exam (TR):

- Right parametrium is supple. The left parametrium appears slightly infiltrated.

The patient underwent total hysterectomy with bilateral adnexectomy, resection of all visible tumor was made. Under general anesthesia, with prior spinal anesthesia by intrathecal morphine with 100 gamma of morphine in an analgesic purpose.

The procedure took place without incident except for a few episodes of hypotension corrected by phenylephrine.

The patient was then transferred ventilated intubated to the intensive care unit, then woken up and extubated 2h after warming and fully waking up.

Postoperative care mainly focuses on the hemodynamic and respiratory aspects and monitoring of drains.

The day after surgery, a transthoracic echocardiogram (TTE) was performed, complemented by a CT angiography, which revealed a massive thrombus in the right atrium extending into the inferior vena cava and reaching the renal veins.




Figure 2: The transthoracic ultrasound revealed the presence of a massive thrombus within the right atrium (RA).

Surgery was undertaken to extract the thrombus, but it was not possible to remove it entirely. Macroscopically, the thrombus had a tissue-like consistency, and the final histopathological analysis confirmed it to be a metastasis from the patient's leiomyosarcoma. The patient passed away 14 days after the surgery.



Figure 1: Macroscopic aspect the intra-cardiac metastasis after surgery

DISCUSSION:

Cardiac metastasis is highly reported in lung cancer, breast cancer, leukemia, malignant lymphoma, and malignant melanoma, only few cases have been reported on metastasis in uterine sarcoma (1, 4, 5).

Uterine leiomyosarcoma often metastasizes to the lungs, peritoneal cavity and liver, but metastasis to the heart is very rare. Our literature search yielded only 13 cases with cardiac metastasis, since the first report by Rosenblatt et al. in 1960, from 2000 through 2024.

Metastatic cancer cells from uterine leiomyosarcoma can reach the heart via lymphatic or hematogenous pathways; however, the lungs and breasts are more commonly the primary sites of metastasis. [8]

The patient can be totally asymptomatic, or can present signs of heart failure.

Cardiac metastases from uterine leiomyosarcoma most commonly present as atrial or ventricle masses detected on clinical presentation. [9]

In our case, the discovery of the intra-right atrial thrombus was incidental. A transthoracic echocardiography, performed almost systematically in the intensive care unit to assess volemia by measuring filling pressures, revealed the intra-right atrial thrombus.

Cardiac metastasis will generally produce clinical findings only when the heart is involved extensively. Findings are variable and include progressive heart failure, intracavitary obstruction, embolism and arrhythmia. (11)

Yazan Assaf et al. reviewed 12 patients with pulmonary infarcts caused by cardiac tumors and pulmonary tumor emboli. Approximately 10% of patients with metastatic cardiac

tumors develop cardiac dysfunction. The signs include tamponade, atrio-ventricular valve obstruction and pulmonary or systemic emboli. (10 - 11)

Appropriate candidates for cardiovascular surgery are those with low-grade malignancies and no evidence of metastases to other organs. However, urgent surgical intervention should be considered in cases of valvular insufficiency, obstruction of the outflow tract, or pulmonary infarction caused by the tumor.

Our patient underwent surgery due to the size of the thrombus, which caused an obstruction in the outflow tract. But complete extraction of the thrombus was not possible. Therefore, the patient suffered from a progressive heart failure which caused death.

Survival is largely dependent on metastasis, with a 5-year survival rate of 91.2% for nonmetastatic uterine leiomyosarcoma and 41.2% for metastatic uterine leiomyosarcoma. (7)

Given the aggressive nature of uterine leiomyosarcoma and the rarity of cardiac metastases, we present this case to provide additional evidence of the potential for cardiac involvement.

Conclusion:

Cardiac metastasis from uterine leiomyosarcoma is extremely rare and often detected incidentally, but it can cause severe complications, including heart failure and intracavitary obstruction. This case highlights the importance of early detection and multidisciplinary management to improve outcomes in patients with metastatic cardiac involvement.

Declaration of patient consent

The authors certify that they have obtained all appropriate the patient's legal representative consent forms.

Financial support and sponsorship

No.

Conflicts of interest

There are no conflicts of interest.

References:

- 1- Sasahara H, Yatsu S, Wada H, Sai E, Moriya S, Miyazaki S, Kajimoto K, Ogita M, Suwa S. Multiple Imaging and Surgical Characteristics in Cardiac Metastasis from Undifferentiated Uterine Sarcoma. Case Rep Cardiol. 2022 Nov 11;2022:6025354. doi: 10.1155/2022/6025354. PMID: 36407789; PMCID: PMC9674408.

- 2- Conklin C. M. and Longacre T. A., Endometrial stromal tumors, *Advances in Anatomic Pathology*. (2014) **21**, no. 6, 383–393, <https://doi.org/10.1097/PAP.0000000000000046>, 2-s2.0-84914144760.
- 3- Bussani R, De-Giorgio F, Abbate A, Silvestri F. Cardiac metastases. *J Clin Pathol*. 2007 Jan;60(1):27-34. doi: 10.1136/jcp.2005.035105. Epub 2006 Nov 10. PMID: 17098886; PMCID: PMC1860601.
- 4- Silva M. G., Banazol N., Coelho P., and Fragata J. I. G., Metastatic endometrial stromal sarcoma: a rare cause of right-sided intracardiac mass, *Journal of Cardiac Surgery*. (2021) **36**, no. 6, 2143–2145, <https://doi.org/10.1111/jocs.15463>, [33651415](https://doi.org/10.1111/jocs.15463).
- 5- Maebayashi A., Nagaishi M., Nakajima T., Hata M., Xiaoyan T., and Kawana K., Successful surgical treatment of cardiac metastasis from uterine leiomyosarcoma: a case report and literature review, *The Journal of Obstetrics and Gynaecology Research*. (2020) **46**, no. 5, 795–800, <https://doi.org/10.1111/jog.14231>, [32166826](https://doi.org/10.1111/jog.14231).
- 6- Rosenblatt WH, Featherston WP. Metastatic leiomyosarcoma of the heart. *Am J Cardiol* 1960; 6: 672–675.
- 7- Ahoubim, J., Sarcar, B., & Abdeen, Y. (2019). Cardiac metastasis of a uterine leiomyosarcoma. *Journal of Natural Science, Biology and Medicine*, 10(2), 217-219.
- 8- Liu T, Khan S, Behr S, Aparici CM. Diagnosis of cardiac metastasis from endometrial cancer by F-18 FDG-PET/CT. *Nucl Med Mol Imaging* 2014;48:237-40.
- 9- Peng YJ, Hueng GG, Lee HS. Acute heart failure as manifestation of metastatic uterine leiomyosarcoma to the heart and lung. *Heart Lung* 2004;33:46-9.
- 10- Assaf Y, Nasser M, Jneid H, Ott D. Pulmonary embolism following incomplete surgical resection of a right ventricular myxoma: A case report and review of the literature. *Cardiol Ther* 2018; 7: 107–117.
- 11- Martin JL, Boak JG. Cardiac metastasis from uterine leiomyosarcoma. *J Am Coll Cardiol*. 1983 Aug;2(2):383-6. doi: 10.1016/s0735-1097(83)80180-6. PMID: 6863772.