

A Silent Invader: Cutibacterium acnes Endocarditis Presenting with Obstructive Mitral Vegetation and Catastrophic Embolic Complications

Abstract:

Background: Infective endocarditis (IE) is a life-threatening condition, particularly in elderly patients with high mortality rates if not treated promptly. An uncommon but severe manifestation of IE is the development of obstructive vegetation on the native mitral valve, leading to systemic embolization including coronary embolism, splenic and renal infarction with hemorrhagic transformation of cerebral stroke. While cases of prosthetic mitral valve endocarditis with resultant stenosis have been reported such as cases involving Cutibacterium acnes infection, stenotic complications in native mitral valve endocarditis remain exceptionally rare. The management of such cases is particularly challenging due to the critical balance between the urgent need for surgical intervention and the heightened hemorrhagic risk that may contraindicate anesthesia, thus future guidelines should establish specific criteria to optimize surgical decision-making in these high-risk scenarios.

Case presentation: We present a case of infective endocarditis affecting the native mitral valve characterized by an obstructive vegetation caused by Cutibacterium acnes. The condition was complicated by multiple embolic events involving the coronary, cerebral, renal, and splenic circulation. The development of embolic phenomena in infective endocarditis is not uncommon, but their occurrence as a result of a vegetation obstructing the mitral valve is an atypical presentation, requiring immediate intervention.

Conclusion: While embolic complications are a well-recognized consequence of infective endocarditis, their occurrence secondary to a mitral valve obstruction remains an unusual presentation, necessitating urgent intervention.

Keywords: Infective endocarditis, Mitral stenosis, Systemic embolization, Cutibacterium acnes, Native valve endocarditis

Introduction:

Infective endocarditis (IE) is a severe condition associated with high in-hospital mortality rates (15-20%) [1], with stroke being the second most frequent cause of death in affected patients [2]. While IE primarily predominantly involves the aortic and mitral valves, its clinical presentation can be highly variable, posing significant diagnostic and therapeutic challenges. While IE is most commonly caused

by *Staphylococcus aureus* and *Streptococcus* species, *Cutibacterium acnes* is a rare causative pathogen, typically associated with prosthetic valve infections and postsurgical endocarditis [3,4]. Native mitral valve involvement by *Cutibacterium acnes* is exceptionally rare, and its indolent course can delay diagnosis, increasing the risk of severe complications [5] with a mortality rate of 16 % [6].

We present a rare case of obstructive vegetation on a native mitral valve caused by *Cutibacterium acnes*, an infrequent etiology in clinical practice. The resulting vegetation led to significant functional mitral stenosis, complicated by coronary, cerebral, renal, and splenic embolism, as well as hemorrhagic transformation of an ischemic stroke.

Case report:

A 56-year-old patient with no known cardiovascular risk factors or notable medical history presented to the emergency department with a two-week history of fever, asthenia, and progressive weakness associated with febrile low back pain.

On examination, the patient was conscious, hemodynamically stable, and able to tolerate the supine position. He was eupneic with a respiratory rate of 14 cycles per minute and a peripheral oxygen saturation SpO₂ of 94%. Notably, he was tachycardic (heart rate: 140 beats per minute) but normotensive (blood pressure: 120/55 mmHg). The patient was febrile at 39°C, and oral examination revealed poor dental hygiene. There were no signs of congestive heart failure, and pulmonary auscultation was unremarkable. Cardiovascular examination revealed a murmur suggestive of mitral stenosis. Neurological assessment was unremarkable, and the neck was supple.

The initial electrocardiogram (Figure 1) demonstrated sinus tachycardia at 100 bpm with a normal axis and signs of left ventricular hypertrophy (LVH) with ST segment depression in lateral and inferior leads. A skin examination revealed track marks over the left antecubital fossa and Janeway lesions on the fingertips. A chest X-ray was unremarkable. Given the high clinical suspicion of infective endocarditis (IE), two sets of blood cultures were obtained.

A transesophageal echocardiography (TEE) identified an 11 mm obstructive vegetation attached to the distal third of the anterior mitral valve leaflet, with fine added elements but no evidence of intra-atrial or intraventricular thrombus. Left ventricular ejection fraction was preserved (Figure 2). Given the potential risk of atrioventricular (AV) block progression, the patient was transferred to the cardiac intensive care unit for continuous monitoring.

Laboratory assessment showed a significant infectious syndrome with a C-reactive protein CRP of 369mg/dl, procalcitonin of 115 ng/mL, leukocytosis (17,000 cells/mm³), and thrombocytopenia (platelet count: 80,000/mm³), the troponin level was elevated to 5000 ng/ml (Table 1). A thoracoabdominal computed tomography (CT) scan was performed as part of the embolic workup,

revealing multiple splenic and renal infarctions (Figure 3), which explained the patient's low back pain.

On the second day of hospitalization, the patient developed a sensorimotor deficit characterized by dysarthria and left-sided hemiparesis. An urgent cerebral CT scan identified an acute ischemic stroke in the left frontoparietal region. Twenty-four hours later, the patient exhibited a decline in Glasgow Coma Scale (GCS) score, prompting an emergent follow-up neuroimaging, which revealed a large hemorrhagic transformation involving the right frontoparieto-occipital region (fig 4).

The clinical course rapidly deteriorated with the onset of bilateral mydriasis and signs of brain herniation, necessitating urgent endotracheal intubation. Unfortunately, the patient suffered a sudden cardiorespiratory arrest on the spot, which remained refractory to resuscitative measures. Blood cultures grew *Cutibacterium acnes* ten days after they were drawn.

Variable	Results
Sodium (mmol/L)	139
Potassium (mmol/L)	4.4
Chloride (mmol/L)	98
Blood glucose (g/L)	1.20
Serum bicarbonate (mmol/L)	27
Urea (g/L)	0.32
Creatinine (mg/L)	10.3
ASAT (U/L)	16
ALAT (U/L)	19
GGT (U/L)	23
CRP (mg/dL)	369
Procalcitonin (ng/mL)	115
Troponin (ng/ml)	5000
White blood cells (/mCL)	17000
Neutrophils	12000

76	Hemoglobin (g/dL)	12.9
77	Platelets (/mcL)	80000

Table 1 :Biological parameters on admission to emergency room




Figure 1 :Electrocardiogram demonstrating sinus tachycardia at 100 bpm with a normal axis and signs of left ventricular hypertrophy (LVH) with ST segment depression in lateral and inferior leads




Figure 2 : Transesophageal echocardiography (TEE) demonstrating an 11 mm obstructive vegetation attached to the distal third of the anterior mitral valve leaflet, with fine added elements (blue arrow)




Figure 3: Thoracoabdominal computed tomography (CT) scan revealing multiple splenic and renal infarctions




Figure 4: Cerebral tomography scan revealing a large hemorrhagic transformation involving the right frontoparieto-occipital region

130

131 **Discussion:**

132 Infective endocarditis (IE) presenting primarily as valvular stenosis is an exceptionally rare
133 phenomenon. The estimated annual incidence of IE in developed countries ranges between 3 and 10
134 cases per 100,000 individuals, a figure that has remained relatively stable over the past two decades
135 [1]. While valvular regurgitation is the most frequently observed hemodynamic consequence of IE,
136 mitral stenosis patent hemodynamically remains an unusual finding [2]. Few cases have been
137 documented in which mitral valve obstruction resulted from extensive vegetation, and only two cases
138 have described diffuse thickening of a prosthetic valve leaflet leading to stenosis [2]. To date, only 15
139 cases of bacterial endocarditis affecting a mitral prosthetic valve and causing stenosis by obstruction
140 have been reported in the literature [3]. Despite its rarity, IE should remain a differential diagnosis in
141 patients presenting with mitral valve obstruction, particularly in the presence of predisposing factors.

142 Infective endocarditis represents a clinical challenge, particularly when associated with large mitral
143 valve vegetations that carry a significant risk of hemodynamic deterioration and embolic
144 complications. The case presented herein describes an extensive vegetation leading to severe mitral
145 stenosis and a cascade of systemic embolization, including ischemic stroke, renal infarction, and
146 hepatic embolization.

147 The causative pathogen, *Cutibacterium acnes*, poses a significant diagnostic challenge due to its
148 ubiquitous presence as a skin commensal [9], frequently leading to its misinterpretation as a
149 contaminant when isolated from a single blood culture [10]. Unlike more common IE pathogens, *C.*
150 *acnes* requires both aerobic and anaerobic culture conditions, with prolonged incubation to ensure
151 reliable detection [11]. Several studies recommend extending blood culture incubation to 10 to 14 days
152 to reduce the risk of false-negative results [12]. There have been reported cases of *Cutibacterium acnes*
153 infective endocarditis in which patients remained afebrile and exhibited normal C-reactive protein
154 levels, thereby complicating the clinical diagnosis [13]. This delay presents a critical therapeutic
155 dilemma, as illustrated in the present case, where native mitral valve endocarditis progressed rapidly,
156 culminating in multi-organ embolization and fatal neurological deterioration within 24 hours of
157 diagnosis on transesophageal echocardiography, before the initiation of targeted antimicrobial therapy.
158 This highlights the need for heightened clinical suspicion and revised microbiological protocols to
159 enhance the timely detection and management of *C. acnes* endocarditis.

160 The diagnosis of IE remains primarily based on the modified Duke criteria [14], which emphasize key
161 echocardiographic and microbiological findings such as vegetations, abscesses, prosthetic valve
162 dehiscence, and new valvular regurgitation. However, valvular stenosis is not currently recognized as a
163 diagnostic criterion, even in the most recent updates to the European Society of Cardiology (ESC)
164 guidelines [15]. While these guidelines provide essential diagnostic and therapeutic frameworks,

certain cases such as the present one challenge existing classification paradigms and call for a more nuanced approach to defining IE-related complications.

Early cardiac surgery is recommended in cases complicated by acute heart failure, uncontrolled infection, or large vegetations [16]. However, the presence of concomitant intracranial hemorrhage introduces a significant management challenge, necessitating a careful risk-benefit analysis. In patients with an unstable clinical course due to heart failure, refractory infection, or a persistent high embolic burden, urgent or emergent surgery should be considered, provided there is a reasonable expectation of neurological recovery [17]. Current ESC guidelines have become more restrictive regarding the indication for surgery based on vegetation size. In native aortic or mitral valve endocarditis, urgent surgery should be considered in patients with vegetations exceeding 10 mm when associated with severe valve stenosis or regurgitation, particularly in cases of low operative risk. This raises the question of whether surgical intervention should be prioritized in all cases where the vegetation is mitral, even before embolic complications ensue. The case described herein illustrates the devastating consequences of delayed intervention, as the occurrence of hemorrhagic transformation, manifesting as meningeal hemorrhage, ultimately precluded surgery and led to a fatal outcome.

This underscores the urgent need for refined surgical criteria in IE, particularly in cases of mitral valve involvement, to optimize outcomes and prevent irreversible complications.

Conclusion:

In conclusion, this case of infective endocarditis on a native mitral valve, leading to mitral stenosis and cerebral and splenic embolizations with hemorrhagic transformation, highlights the complexities and challenges of managing such severe cases. The contraindication to surgery and the poor prognosis underlines the importance of early diagnosis and intervention. Future perspectives should focus on improving diagnostic techniques, optimizing medical therapies to control infection and prevent complications, and developing more effective surgical approaches for high-risk patients. These advancements could potentially improve outcomes and reduce mortality associated with infective endocarditis.

Abbreviations

EI	Infective endocarditis
C. acnes	Cutibacterium acnes
LVH	Left ventricular hypertrophy
TEE	Transesophageal echocardiography
CRP	C-Reactive protein
CT	Computed tomography
GCS	Glasgow coma scale

Funding

None to declare

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Written informed consent was obtained from the patient for publication of this case report and accompanying images

References:

1. Nappi, F., Singh, S. S. A., Spadaccio, C., Acar, C., & Chello, M. (2022). Infective Endocarditis in High-Income Countries. *Journal of Clinical Medicine*, 11(17), 4981.
2. Hart, M.A., Shroff, G.R. Infective endocarditis causing mitral valve stenosis – a rare but deadly complication: a case report. *J Med Case Reports* 11, 44 (2017). <https://doi.org/10.1186/s13256-017-1197-3>
3. Gakuto Bando, Taiji Okada, Hideki Tsubota, Yutaka Furukawa, Prosthetic valve infective endocarditis with severe mitral stenosis caused by *Cutibacterium acnes*: a case report, *European Heart Journal - Case Reports*, Volume 8, Issue 5, May 2024, ytae205, <https://doi.org/10.1093/ehjcr/ytae205>
4. Hassan, T, Mahfooz, F, Tandon, S. et al. culprit in the shadows: cutibacterium acnes endocarditis unveiling prosthetic valve dysfunction. *JACC*. 2024 Apr, 83 (13_Supplement) 3052. [https://doi.org/10.1016/S0735-1097\(24\)05042-3](https://doi.org/10.1016/S0735-1097(24)05042-3)
5. Perry A, Lambert P. *Propionibacterium acnes*: infection beyond the skin. *Expert Rev Anti Infect Ther* 2011;9:1149–1156.
6. Boyle M, Tennyson C, Guleri A, Walker A. Infective endocarditis: acne to zoonoses on the valve, an A to Z perspective. *Br J Cardiol*. 2022;29(2):112–116. doi:10.5837/bjc.2022.025
7. Sohail MR, Gray AL, Baddour LM, Tleyjeh IM, Virk A. Infective endocarditis due to *Propionibacterium* species. *Clin Microbiol Infect* 2009;15:387–94. <https://doi.org/10.1111/j.1469-0691.2009.02703.x>
8. Li, Mingfang et al. Infective endocarditis. *The Lancet*, Volume 404, Issue 10450, 377 – 392.
9. Tattevin P, Watt G, Revest M, Arvieux C, Fournier P-E. Update on blood culture- negative endocarditis. *Med Mal Infect* 2015;45:1–8.

10. Lodhi SH, Abbasi A, Ahmed T, Chan A. Acne on the Valve: Two Intriguing Cases of Cutibacterium Acnes Endocarditis. *Cureus*. 2020 Jun 9;12(6):e8532. doi: 10.7759/cureus.8532. PMID: 32665879; PMCID: PMC7352884.
11. Achermann, Y., Goldstein, E. J., Coenye, T., & Shirtliff, M. E. (2014). *Propionibacterium acnes*: from commensal to opportunistic biofilm-associated implant pathogen. *Clinical microbiology reviews*, 27(3), 419-440.
12. Verkaik NJ, Schurink CAM, Melles DC. Letter to the editor, Antibiotic treatment of *Propionibacterium acnes* endocarditis. *Clin Microbiol Infect* 2018;24:209.
<https://doi.org/10.1016/j.cmi.2017.07.024>
13. Heinen FJ, Arregle F, van den Brink FS, Marsan NA, Bernts L, Houthuizen P, et al. Clinical characteristics and outcomes of patients with *Cutibacterium acnes* endocarditis. *JAMA Netw Open* 2023;6:e2323112.
14. Li JS, Sexton DJ, Mick N, Nettles R, Fowler VG Jr, Ryan T, et al. Proposed modifications to the Duke criteria for the diagnosis of infective endocarditis. *Clin Infect Dis* 2000;30:633–638.
15. Delgado V, Ajmone Marsan N, de Waha S, Bonaros N, Brida M, Burri H, et al. 2023 ESC guidelines for the management of endocarditis. *Eur Heart J* 2023;44:3948–4042.
16. Richard Rezar, Michael Lichtenauer, Markus Haar, Georg Hödl, Jan Marco Kern, Zhichao Zhou, Thomas Wuppinger, Johannes Kraus, Bernhard Strohmmer, Uta C. Hoppe, Bernhard Wernly. Infective endocarditis – A review of current therapy and future challenges. *Hellenic Journal of Cardiology*, Volume 62, Issue 3, 2021, Pages 190-200, ISSN 1109-9666, <https://doi.org/10.1016/j.hjc.2020.10.007>
17. Victoria Delgado, Nina Ajmone Marsan, Suzanne de Waha, Nikolaos Bonaros, Margarita Brida, Haran Burri, Stefano Caselli, Torsten Doenst, Stephane Ederhy, Paola Anna Erba, Dan Foldager, Emil L Fosbøl, Jan Kovac, Carlos A Mestres, Owen I Miller, Jose M Miro, Michal Pazdernik, Maria Nazarena Pizzi, Eduard Quintana, Trine Bernholdt Rasmussen, Arsen D Ristić, Josep Rodés-Cabau, Alessandro Sionis, Liesl Joanna Zühlke, Michael A Borger, ESC Scientific Document Group, 2023 ESC Guidelines for the management of endocarditis: Developed by the task force on the management of endocarditis of the European Society of Cardiology (ESC) Endorsed by the European Association for Cardio-Thoracic Surgery (EACTS) and the European Association of Nuclear Medicine (EANM), *European Heart Journal*, Volume 44, Issue 39, 14 October 2023, Pages 3948–4042,
<https://doi.org/10.1093/eurheartj/ehad193>