

Efficacy of Transcutaneous Electrical Nerve Stimulation and Soft Splint Therapy in the Treatment of Patients with Temporomandibular Joint Disc Displacement with Reduction – A Comparative Study

by Jana Publication & Research

Submission date: 24-May-2025 12:26PM (UTC+0700)

Submission ID: 2664988079

File name: IJAR-51815.doc (1.26M)

Word count: 3869

Character count: 23351

Efficacy of Transcutaneous Electrical ⁴Nerve Stimulation and Soft Splint Therapy in the Treatment of Patients with Temporomandibular Joint Disc Displacement with Reduction – A Comparative Study

Abstract:

Background: The variability in treatment approaches ¹⁷for temporomandibular disc displacement with reduction underscores the pressing need for this study. With disparate clinical practices, there is a critical gap in establishing the most effective treatment. Addressing this gap will optimize patient care and minimize the risk of suboptimal treatment outcomes.

Aims: The study aimed ¹to compare the effectiveness of Transcutaneous Electrical Nerve Stimulation and splint therapy in managing temporomandibular joint disc displacement with reduction by evaluating ⁶pain intensity, mouth opening, and masticatory muscle tenderness post-treatment.

Settings and Design: A randomized controlled trial was conducted ¹in patients with temporomandibular joint disc displacement with reduction. ¹Patients were randomly assigned to receive either TENS therapy or splint therapy.


Methods and Material: Baseline assessments ²of pain intensity, maximum mouth opening, and masticatory muscle tenderness were recorded preoperatively. Treatment was administered over four weeks, with weekly evaluations to monitor efficacy.

Statistical analysis used: ²⁷Inter-group analysis was done using independent t-tests, and intra-group analysis was done using ANOVA tests to evaluate the efficacy of each treatment.

Results: In the TENS group, there were notable reductions in pain intensity and masticatory muscle tenderness, accompanied by an improvement in mouth opening over the four weeks. Conversely, the splint therapy group showed minimal changes in these parameters.

Conclusions: TENS therapy shows promise as ²⁴an effective treatment modality for temporomandibular joint disc displacement with reduction. The significant improvements observed in pain intensity, muscle tenderness, and mouth opening highlight TENS therapy's potential to alleviate symptoms of DDWR.

Key-words: Temporomandibular joint disc displacement with reduction, Trans electronic nerve stimulation, Soft Splint therapy, Temporomandibular disorders



Introduction:

Orofacial pain commonly stems from temporomandibular disorders (TMDs), which typically have origins beyond dental concerns. They encompass various conditions affecting the temporomandibular joint and jaw muscles. Symptoms include pain in the auricular, temporomandibular joint, or masticatory muscles, restricted jaw movement, and clicking sounds during jaw movement.¹ Interestingly, only a quarter of individuals with TMD recognize or report their symptoms, despite 70% of the general population exhibits signs of the disorder.²

Research indicates a TMD prevalence of 40% to 75%. For instance, Solberg et al. found 76% of individuals aged 18 to 25 displayed TMD signs, with 26% experiencing associated symptoms. While half of TMD patients may exhibit jaw sounds or deviation upon opening, only a tiny fraction seek treatment.³ The most TMDs involve anterior disc displacement, which can be classified into disc displacement with and without reduction.⁴

Treatment modalities range from non-invasive options like physical, exercise, and thermal therapies, acupuncture, occlusal splint, and Transcutaneous Electric Nerve Stimulation therapy (TENS), to invasive options like arthrocentesis, arthroscopy, joint reconstructive surgery, and condylotomy. Among these, TENS and occlusal splint therapy are most commonly used. TENS delivers pulsed biphasic electrical waves through skin-surface electrodes to manage pain by promoting muscle relaxation and neuromuscular stimulation. Occlusal splint therapy, involving bite guards, protects teeth from excessive forces during clenching or grinding, reducing TMJ stress and muscle pain.²

Due to the varied treatment options for temporomandibular disc displacement with reduction, consensus on the most effective approach is lacking, leading to varied clinical practices and potential suboptimal outcomes.

This study aims to assess and compare the effectiveness of TENS therapy versus soft splint therapy in managing temporomandibular disc displacement with reduction by evaluating the intensity of pain, maximum mouth opening, and masticatory muscle tenderness post-treatment.

Subjects and Methods:

Study Population:

¹⁴ The study was conducted in the Outpatient Department of Oral and Maxillofacial Surgery at GITAM Dental College and Hospital, Visakhapatnam. ³⁹ Patients diagnosed with temporomandibular disc displacement with reduction were enrolled. Patients were randomly selected and voluntarily participated by signing informed consent forms. ⁵ The study was approved by the institutional review board, and ethical clearance was obtained from the institutional ethical committee.

The inclusion criteria were patients aged 16 to 50 years, no gender predisposition, patients no predisposing medical history, class I occlusion and no requirement for orthodontic intervention, and diagnosed with temporomandibular disc displacement with reduction. Diagnosis was made solely through clinical examination, without radiographs. Selected patients exhibited a clicking sound, masticatory muscle tenderness, temporomandibular joint pain upon palpation.

The exclusion criteria were patients with systemic diseases or dentofacial anomalies, ongoing medical or psychological treatments, history of temporomandibular joint surgery or trauma, parafunctional habits like bruxism and/or clenching, refusal to participate, edentulous or partially edentulous patients, those with cardiac pacemakers, and epileptic patients.

A randomized controlled study involving 40 patients diagnosed with temporomandibular disc displacement with reduction was conducted. Patients were informed about clinical research's nature and purpose, and their informed consent was obtained. ⁵ Ethical clearance was diligently obtained from the Institutional Ethical Committee, ensuring compliance with ethical standards. A double-blinding protocol was implemented. Participant Blinding involved not informing patients of their assigned treatment group (TENS or Splint), and researcher blinding ensured that the researchers were unaware of each participant's treatment allocation.

The study parameters included the intensity of pain, mouth opening, and masticatory muscle tenderness, evaluated initially pre-treatment as baseline data and then weekly for four weeks post-treatment. ² Pain intensity was measured using the Visual Analogue Scale (VAS), where patients marked their pain level on a ⁹ 10 cm line ranging from 0 (no pain) to 10 (worst possible pain). ¹³ Mouth opening was measured by the distance between the incisal edges of the upper and lower central incisors using a divider. Masticatory muscle tenderness was evaluated by manually palpating the temporalis, masseter, lateral and medial pterygoid muscles bilaterally with firm yet gentle and constant pressure (Figure 1). The tenderness assessment also utilized the Visual Analogue Scale,

with patients marking their pain level on a 10 cm line from 0 (no pain) to 10 (worst possible pain). These assessments were conducted weekly for four consecutive weeks following treatment initiation.

In the TENS Group, patients received treatment with transcutaneous electric nerve stimulation therapy using a TENS device (PHYSIO TENS, manufactured by Scientific Medical Systems). The active electrode was positioned between the temporomandibular joint (TMJ) and coronoid process to ensure the stimulus reached the trigeminal and facial nerves, while the control electrode was placed at the back of the neck. The therapy was administered at a frequency of 50 Hz, a pulse width of 0.5 msec, and an intensity ranging from 0 to 60 mA for 20 minutes per session, conducted weekly intervals for four consecutive weeks. (Figure 2)

In the Splint Group, patients underwent soft splint therapy. Impressions were made, and a 2mm thick rubber sheet was vacuum-formed using a BIOSTAR vacuum former to fabricate the soft splint. After cooling, the splint was trimmed to shape and checked for retention and occlusal interferences. Patients were instructed to wear the splint continuously for 24 hours daily. Follow-up assessments were conducted for four weeks. Pain score, maximum mouth opening, and masticatory muscle tenderness were reassessed and recorded at each follow-up. (Figure 3) Depending on the patients' tolerance levels, analgesics were prescribed as needed for pain management. During each visit, patients were asked to report if their pain was intolerable and if they required medication. Patients who required additional analgesics were excluded from the study. Only those not taking additional pain medication were included in the final analysis.

In the TENS Group (A) and Splint group (B), baseline data for pain (PQ), mouth opening (OQ), and muscle tenderness (TQ) were recorded, followed by weekly evaluations pain (P1- P4), mouth opening (O1- O4), and muscle tenderness (T1 - T4). This systematic approach ensured comprehensive monitoring of treatment outcomes.

Statistical analysis:

Data analysis was conducted using SPSS version 15.0. The Independent t-test compared data between the TENS and SPLINT groups weekly and ANOVA tests analyzed data across consecutive weeks for both groups.

Results:

The mean and standard deviation scores for pain intensity, mouth opening, and muscle tenderness were recorded weekly for both groups.

Pain intensity and masticatory muscle tenderness were consistently higher in the SPLINT group compared to the TENS group. In the SPLINT Group, the mean pain score slightly decreased from 6.15 in the first week and 5.15 at the end of the fourth week. In contrast, the TENS group showed a significant reduction in pain scores, from 5.80 in the first week to 1.20 at the end of the fourth week.

(Graph 1)

Similarly, the mean masticatory muscle tenderness scores in the SPLINT group showed only 1.1 variation from 6.90 in the first week to 5.80 at the end of the fourth week. However, the TENS group experienced a significant decrease, from 6.65 in the first week to 1.50 at the end of the fourth week.

(Graph 2)

Regarding mouth opening, the SPLINT group showed a slight increase the measurements showed an increase from 38.85 mm in the first week to 39.69 mm at the end of the fourth week. In contrast, the TENS group saw a substantial increase from 39.90 mm in the first week to 50.22 mm at the end of the fourth week. (Graph 3)

Inter-group and intra-group analyses were conducted to evaluate the efficacy of each treatment. Independent t-test evaluations revealed significant differences ($p\text{-value} < 0.05$) in all three parameters. The intensity of pain and masticatory muscle tenderness significantly decreased in the TENS group, while mouth opening significantly increased compared to the SPLINT group (Table 1).

Furthermore, the parameters were compared to baseline data each week. In the TENS group, there was a notable decrease in the intensity of pain and muscle tenderness and a marked increase in mouth opening with each consecutive measurement. These differences were statistically significant ($p\text{-value} < 0.05$) across all three parameters, as analyzed by the ANOVA test.

In contrast, the SPLINT group showed only minor variations from baseline data for all parameters, with no statistical significance in the scores between baseline and subsequent weeks as revealed by the ANOVA test (Table 2).

Discussion:

² The temporomandibular joint (TMJ) is often affected by pain-related and intraarticular conditions, with ²⁵ anterior disc displacement being the most common type of TMJ arthropathy. Anterior disc displacement with reduction (DDWR) accounts 41% of clinical TMJ disorder diagnoses, as highlighted Poluha et al.⁵ In DDWR, the articular disc displaces ⁴¹ when the mouth is closed and repositions between the condyle and ⁴² articular tubercle upon opening. Misalignment of the condyle-disc complex ³⁴ is attributed to factors like elongation of discal collateral ligaments, thinning of the posterior disc border, and condyle resting on posterior disc regions, leading to abnormal condyle movement over the disc's posterior border during the opening, defining the DDWR stage.⁵

Disc displacement involves abnormal disc positioning ¹ to the mandibular condyle and articular eminence. Mild disc displacement may initially manifest as occasional painful clicking, intermittent locking, and headaches. As the displacement worsens, symptoms often include ⁷ joint pain and tenderness, locking, and a restricted mandibular motion range. Severe displacement can lead to disc deformation, degeneration of osseous components, ⁷ deterioration of articular cartilage and disc surface, and bone remodelling. In this study, the diagnosis of DDWR was made solely through clinical findings of patients with clicking sounds, TMJ pain, and tenderness of masticatory muscles on palpation. Radiographs or MRI scans were not used due to ethical concerns regarding radiation exposure and cost-effectiveness.

The three most common symptoms of DDWR, i.e., ⁶ pain intensity, mouth opening, and masticatory muscle tenderness were selected for evaluation in this study. Treatment options for TMJ internal derangement are categorized into non-invasive, minimally invasive, and invasive management. The initial approach prioritizes the least invasive and reversible treatments, reserving more invasive options for cases where initial treatments fail to alleviate the symptoms. Non-invasive treatment modalities include pharmacotherapy, physical therapy, exercise therapy, thermal therapy, acupuncture, occlusal splint therapy, and TENS therapy. Disorders of articular or muscular origin, typically respond well to non-invasive interventions like TENS and occlusal splint therapy.^{6,7} Therefore, this study compared these two treatment modalities for effectiveness. Additionally, analgesics were prescribed as needed, but patients requiring additional medication were excluded, ensuring only those not taking extra pain medication were included in the final analysis.

Occlusal splint therapy aims to establish neuromuscular harmony in the masticatory system by using removable appliances to create a mechanical disadvantage against parafunctional forces. These

splints serve various purposes, including muscle relaxation, condylar positioning, diagnostic aid, protection against bruxism, and reduce cellular hypoxia levels.⁷ TENS therapy, a non-invasive and cost-effective modality, effectively controls chronic and acute pain by activating the descending inhibitory system. According to the gate control theory, TENS modulates pain perception by recruiting A-beta afferent fibers and inhibiting pain impulse transmission. TENS units consist of a TENS unit, lead wires, and electrodes, and are classified into high-frequency (>50 Hz) and low-frequency (<10 Hz), with advantages including safety, self-administration, and muscle relaxation.^{8, 9, 10}

In our study, the TENS group showed significantly lower pain intensity and masticatory muscle tenderness scores along significant increase in mouth opening, compared to the splint group. These findings align with other studies by Nunez et al. and Kota et al., which demonstrated significant improvements in mouth opening with TENS therapy.^{11, 12} Contradictory findings from Farheen Jahan et al. suggest higher post-intervention pain scores in the TENS group compared to ultrasound and splint groups when observed for four weeks.⁹ Therefore, the observation period in this study was standardized to four weeks for consistency in the outcomes assessment. TENS therapy yielded a more pronounced reduction in pain scores over one month than splint interventions.

Blinding procedures were rigorously employed in this study, to mitigate potential sources of bias, particularly regarding subjective assessment of treatment outcomes. A double-blinding protocol was implemented, as both the participants and researchers were unaware of the treatment allocation for each participant. This ensured participants' expectations or perceptions did not influence reported outcomes researchers' judgments and interpretations remained unbiased throughout the study.

A notable reduction in masticatory muscle tenderness was observed in the TENS group compared to the Splint group, with statistical significance ($p < 0.05$). This aligns with the findings by A. Monaco et al., who demonstrated that a single 60-minute TENS session application effectively diminished masticatory muscle tenderness, while increasing interocclusal distance.¹³ Furthermore, the results from Siefi et al.'s study further support our findings. Their investigation revealed a substantial 53.88% decrease in masticatory muscle tenderness among TENS therapy participants.¹⁴ In a randomized controlled study by Ana Paula de Lima Ferreira et al., it was found that transcutaneous electrical nerve stimulation (TENS) could effectively increase the pressure pain threshold (PPT) of masticatory muscles.¹⁵ Moreover, Remi Escassan et al. suggested that an optimal application of ultra-low frequency TENS (ULF-TENS) for 40 minutes is necessary to achieve adequate muscle relaxation.

This recommendation aligns with our study's findings regarding the efficacy of TENS in reducing masticatory muscle tenderness.¹⁶ These collective findings emphasize the efficacy of TENS in alleviating muscle tenderness associated with TMDs, underscoring its potential as a valuable treatment modality.

In the TENS group, consecutive weekly treatment led to a significant decrease in pain intensity, ¹ masticatory muscle tenderness, and a notable increase in mouth opening. In contrast, such improvements were not observed in the SPLINT group. This finding aligns with previous research by Farheen Jahan et al., which also demonstrated significant reductions in muscle tenderness and temporomandibular joint (TMJ) pain with TENS therapy over consecutive weeks.⁹ Conversely, the SPLINT group did not show significant improvements in pain intensity, mouth opening, or muscle tenderness over consecutive weeks, likely due to the longer time required for clinical symptom reduction with splint therapy. ¹⁵ Studies by Tecco et al. and Malgorzata Pihut et al. support this, indicating that patients treated with splints experienced reduced pain levels over months.^{17,18}

TENS therapy offers additional advantages, including its applicability for needle-phobic patients, absence of post-operative anaesthesia, and the ability for patients to self-administer treatment and adjust dosage as needed, leading to positive patient acceptance. Furthermore, TENS helps relax hyperactive muscles and acts as a neuromuscular stimulator.¹⁹ On the other hand, occlusal splint therapy has limitations, including dependence on patient compliance for wearing the splint, interference with finding the patient's usual occlusion, and longer treatment duration required for therapeutic outcomes.²⁰

²¹ The study's limitations include a relatively small sample size, a brief four-week treatment duration, and a lack of long-term follow-up. The short treatment duration in this study may not fully capture the benefits of occlusal splint, which typically requires longer application periods.

Despite these limitations, our study indicates that TENS therapy is more effective than occlusal splint therapy for managing TMJ-DDWR. Over the 4-week period, TENS therapy significantly reduced ⁶ pain intensity and masticatory muscle tenderness and improved mouth opening compared to occlusal splint therapy. ³⁷ Further research with larger sample sizes and extended treatment durations is needed for more comprehensive results.

References:


1. Cairns BE. Pathophysiology of TMD pain--basic mechanisms and their implications for pharmacotherapy. *J Oral Rehabil.* 2010 May;37(6):391-410.
2. Dimitroulis, G. (2018) Management of temporomandibular joint disorders: a surgeon's perspective. *Australian Dental Journal*, 63: S1 S79–S90.
3. Jyoti Devi, Mahesh Verma, Rekha Gupta. Assessment of treatment response to splint therapy and evaluation of TMJ function using joint vibration analysis in patients exhibiting TMJ disc displacement with reduction. A clinical study. *Indian J Dent Res.* 2017; 28:33-43.
4. U. Siva Kalyan, Kishore Moturi, K. Padma Rayalu. The Role of Ultrasound in Diagnosis of Temporomandibular Joint Disc Displacement: A Case–Control Study. *J. Maxillofac. Oral Surg.* 2018 Sep;17(3):383-388.
5. Poluha RL, Canales GT, Costa YM, Grossmann E, Bonjardim LR, Conti PCR. Temporomandibular joint disc displacement with reduction: a review of mechanisms and clinical presentation. *J Appl Oral Sci.* 2019 Feb 21;27
6. Melissa Thiemi Kato, Evelyn Mikaela Kogawa, Carlos Neanes Santos, Paula Cesar Rodrigues Conti. TENS and Low-Level Laser therapy in the management of temporomandibular disorders. *J Appl Oral Sci.* 2006;14(2): 130-5.
7. I-Yueh Huang, Ju-Hui Wu, Yu-Hsun Kao, Chao-Ming Chen, Chun-Ming Chen, Yi-Hsin Yang. Splint therapy for disc displacement with reduction of the temporomandibular joint. Part I: Modified mandibular splint therapy. *Kaohsiung Journal of Medical Sciences.* 2011; 27: 323-329.
8. Fahimeh Razazadeh, Khadijeh Hajian, Shoaleh Shahidi, Soraya Piroozi. Comparison of the effects of transcutaneous electric nerve stimulation and low-level therapy on drug-resistant temporomandibular disorders. *J Dent Shiraz Univ Medi Sci.* 2017; 18(3): 187-192.
9. Farheen Jahan, Vinod V. C, Rashmi Sapkal. Efficacy of TENS therapy, Therapeutic Ultrasound and Stabilization Splint as an Adjuvant to pharmacotherapy for temporomandibular disorders. *Innovative science and Research International Journal of Technology.* 2020; 5(3).

10. Vikrant kasat, Aditi Gupta, Ruchi Ladda, Mitesh Kathariya, Harish Saluja, Anjum- Ara Farooqui. Transcutaneous electric nerve stimulation (TENS) in dentistry –A review. *J Clin Exp Dent*. 2014; 6(5): 562-8.
11. Silvia Cristina Nunez, Aguinaldo Silva Garcez, Selly Sayuri Suzuki and Martha Simoes Riberio. Management of mouth opening in patients with temporomandibular disorders through low-level laser therapy and transcutaneous electrical neural stimulation. *Photomedicine and Laser surgery*. 2006; 24: 1.
12. Melissa Thiemi Kato, Evelyn Mikaela Kogawa, Carlos Neanes Santos, Paula Cesar Rodrigues Conti. TENS and Low-Level Laser therapy in the management of temporomandibular disorders. *J Appl Oral Sci*. 2006;14(2): 130-5.
13. Annalisa Monaco, Fabrizio Sgolastra, Irma Ciarrochi, Ruggero Cattaneo. Effects of transcutaneous electric nerve stimulation on electromyographic and kinesiographic activity of patients with temporomandibular disorders: A placebo-controlled study. *Journal of Electromyography and Kinesiology*. 2012; 22: 463-468.
14. Massoud Seifi, Asghar Ebadifar, Sattar Kabiri, Mohammad Reza Badiie, Zahra Abdolazimi, Parisa Amdjadi. Comparative effectiveness of Low-Level Laser Therapy and Transcutaneous Electric Nerve Stimulation on Temporomandibular Joint Disorders. *J Lasers Medi Sci*. 2017; 8 (1): S27-S31
15. Ana Paula de Lima Ferreira, Dayse Regina Alves da Costa, Ana Izabela Sobral de Oliveria.et al. Short term transcutaneous electric nerve stimulation reduces pain and improves the masticatory muscle activity in temporomandibular disorder patients: A randomized controlled trial. *J Appl Oral Sci*. 2017; 25(2): 112-20.
16. Remi Esclassan, Anaïs Rumerio, Paul Monsarrat, Jean Claude Combadazou. et al. Optimal duration of ultra low frequency-transcutaneous electric nerve stimulation (ULF-TENS) therapy for muscular relaxation in neuromuscular occlusion: A preliminary clinical study. *The Journal of Craniomandibular & Sleep Practice*. 2016.
17. Simona Tecco, Felice Festa, Vincenzo Salini, Ettore Epifania, Michele D' Attilio. Treatment of Joint pain noises associated with a recent TMJ internal derangement: a comparison of an Anterior Repositioning Splint, a Full-arch Maxillary Stabilization Splint, and an Untreated Control group. *The Journal of Craniomandibular Practice*. 2004; 22(3).

18. Malgorzata Pihut, Malgorzata Gorecka Piotr Ceranowicz and Mieszko Wieckiewicz. The efficacy of anterior repositioning splints in the management of pain related to temporomandibular joint disc displacement with reduction. *Hindawi Pain Research and Management*. 2018; 6.
19. Vikrant kasat, Aditi Gupta, Ruchi Ladda, Mitesh Kathariya, Harish Saluja, Anjum- Ara Farooqui. Transcutaneous electric nerve stimulation (TENS) in dentistry –A review. *J Clin Exp Dent*. 2014; 6(5): 562-8.
20. Ré, J-P & Pérez Ureña, Carolain & Darmouni, L. & Carlier, J. & J.D, Orthlieb. (2009). The occlusal splint therapy. *international journal of stomatology & occlusion medicine*. 2. 82-86. 10.1007/s12548-009-0015-y


Graphs:

Graph 1: Comparison of mean value of pain between TENS group and SPLINT group at initial




assessment and after treatment during the 1st, 2nd, 3rd, 4th consecutive weeks.

Graph 2: Comparison of mean value of masticatory muscle tenderness between TENS group and SPLINT group at initial assessment and after treatment during the 1st, 2nd, 3rd, 4th consecutive weeks.



Graph 3: Comparison of mean value of mouth opening between TENS group and SPLINT group at initial assessment and after treatment during the 1st, 2nd, 3rd, 4th consecutive weeks.




Tables:

Table 1: Inter-group analysis of three parameters between TENS and Soft Splint therapy by independent t-test

Parameters	Examined at	TENS (A)		SPLINT (B)		p-value (t-test)
		Mean	SD	Mean	SD	
Pain	Initial (P ₀)	5.80	1.24	6.15	1.09	0.35
	1st week (P ₁)	4.50	1.15	6.15	1.09	0.000*
	2nd week (P ₂)	3.35	1.18	5.80	1.06	0.000*
	3rd week (P ₃)	2.35	1.09	5.35	0.99	0.000*
	4th week (P ₄)	1.20	1.11	5.15	1.09	0.000*
Mouth Opening	Initial (O ₀)	39.90	3.93	38.85	3.36	0.369
	1st week (O ₁)	42.89	4.05	39.07	3.36	0.002*
	2nd week (O ₂)	45.55	3.80	39.29	3.34	0.000*
	3rd week (O ₃)	48.14	3.16	39.48	3.34	0.000*
	4th week (O ₄)	50.22	2.98	39.69	3.34	0.000*
Masticatory Muscle Tenderness	Initial (T ₀)	6.65	1.18	6.90	1.12	0.496
	1st week (T ₁)	5.20	1.40	6.90	1.12	0.000*
	2nd week (T ₂)	3.80	1.28	6.35	1.04	0.000*
	3rd week (T ₃)	2.40	1.31	5.95	1.10	0.000*
	4th week (T ₄)	1.50	0.76	5.80	1.01	0.000*

Table 2: Intra-group analysis of three parameters among TENS and Soft Splint therapy at consecutive weeks by ANOVA test



Measurement of parameters (A): Mouth opening, (B) Temporalis muscle tenderness, (C): Masseter muscle tenderness

Figure 2: Application of Transcutaneous electric nerve stimulation (TENS)

Figure 3: Soft splint adaptation in the patient

Ethical Clearance:




GITAM DENTAL COLLEGE AND HOSPITAL
RUSHIKONDA, VISAKHAPATNAM - 530045


ETHICAL COMMITTEE CLEARANCE CERTIFICATE

This is to certify that Dr. P.VIJAY ANIL KUMAR is a bonafide Post Graduate student in the speciality of Oral & Maxillofacial Surgery in GITAM DENTAL COLLEGE & HOSPITAL, Visakhapatnam.

The Ethical Committee gives its consent for the study on
"TRANSCUTANEOUS ELECTRIC NERVE STIMULATION AND
SOFT SPLINT THERAPY IN THE TREATMENT OF PATIENTS
WITH TEMPOROMANDIBULAR JOINT DISC
DISPLACEMENT WITH REDUCTION- A COMPARATIVE
STUDY".


Dr. B. RAMESH

(CHAIR PERSON)
Chairperson, Ethical Committee
GITAM Dental College & Hospital
Rushikonda - 530 045


Dr. K. GANGADHARA PRASAD

(PRINCIPAL)
PRINCIPAL
GITAM Dental College & Hospital
Rushikonda, Visakhapatnam - 45

Efficacy of Transcutaneous Electrical Nerve Stimulation and Soft Splint Therapy in the Treatment of Patients with Temporomandibular Joint Disc Displacement with Reduction – A Comparative Study

ORIGINALITY REPORT

22%

SIMILARITY INDEX

17%

INTERNET SOURCES

17%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | www.researchgate.net
Internet Source | 3% |
| 2 | www.science.gov
Internet Source | 2% |
| 3 | seer.ufrgs.br
Internet Source | 1% |
| 4 | gdch.edu.in
Internet Source | 1% |
| 5 | www.arcjournals.org
Internet Source | 1% |
| 6 | Norazlin Mohamad, Ana Izabela Sobral de Oliveira-Souza, Ester Moreira de Castro-Carletti, Frauke Muggenborg et al. "The effectiveness of different types of acupuncture to reduce symptoms and disability for patients with orofacial pain. A systematic review and meta-analysis", Disability and Rehabilitation, 2024
Publication | 1% |
| 7 | B. E. CAIRNS. "Pathophysiology of TMD pain - basic mechanisms and their implications for pharmacotherapy : PATHOPHYSIOLOGY OF TMD PAIN", Journal of Oral Rehabilitation, 05/11/2010 | 1% |

8	Melike Donmez, Nese Yorguner, Kaan Kora, Volkan Topcuoglu. "Efficacy of bright light therapy in perinatal depression: A randomized, double-blind, placebo-controlled study", Journal of Psychiatric Research, 2022 Publication	1 %
9	cyberleninka.org Internet Source	1 %
10	www.neliti.com Internet Source	1 %
11	Submitted to University of Malaya Student Paper	1 %
12	www.edp-open.org Internet Source	1 %
13	turkjorthod.org Internet Source	1 %
14	Gokkulakrishnan Sadhasivam, Satya Bhushan, Kho Chai Chiang, Neha Agarwal, P. L. Vasundhar. "Clinical Trial Evaluating the Risk of Thromboembolic Events During Dental Extractions", Journal of Maxillofacial and Oral Surgery, 2016 Publication	<1 %
15	www.tandfonline.com Internet Source	<1 %
16	Akansha Budakoti, Nidhi Puri, Manu Dhillon, Upasana Sethi Ahuja, Akshay Rathore, Anuridhi Choudhary, Manmeet Kour. "A comparative evaluation of the effectiveness of low-level laser therapy, ultrasound therapy, and transcutaneous electric nerve stimulation in the treatment of patients with TMDs: a	<1 %

-
- 17 Shigemitsu Sakuma, Yoshihiro Yamaguchi, Kei Taguchi, Rina Kobayashi, Shinya Takagi, Nozomu Taguchi. "Short-term effects of range-of-motion exercise on temporomandibular joints of patients who undergo disc displacement with reduction of temporomandibular joint", Journal of Physical Therapy Science, 2021
- Publication

-
- 18 Ester Moreira de Castro-Carletti, Frauke Müggenborg, Liz Dennett, Ana Izabela Sobral de Oliveira-Souza et al. "Effectiveness of electrotherapy for the treatment of orofacial pain: A systematic review and meta-analysis", Clinical Rehabilitation, 2023
- Publication

-
- 19 Submitted to University of Birmingham
- Student Paper

-
- 20 www.oraljournal.com
- Internet Source

-
- 21 beonbrand.getbynder.com
- Internet Source

-
- 22 www.cambridge.org
- Internet Source

-
- 23 www.ijss-sn.com
- Internet Source

-
- 24 Alpaslan, C.. "Does the use of soft or hard splints affect the short-term outcome of temporomandibular joint arthrocentesis?",

25 Anne L. Harrison, Jacob N. Thorp, Pamela D. Ritzline. "A Proposed Diagnostic Classification of Patients With Temporomandibular Disorders: Implications for Physical Therapists", Journal of Orthopaedic & Sports Physical Therapy, 2014

Publication

26 worldwidescience.org

Internet Source

27 www.jcdr.net

Internet Source

28 dentjods.sums.ac.ir

Internet Source

29 libmast.utm.my

Internet Source

30 www.bmj.com

Internet Source

31 www.journalcra.com

Internet Source

32 Ömer Ekici, Ümit Dünder, Murat Büyükbosna. "Comparison of the Efficiency of High-Intensity Laser Therapy and Transcutaneous Electrical Nerve Stimulation Therapy in Patients With Symptomatic Temporomandibular Joint Disc Displacement With Reduction", Journal of Oral and Maxillofacial Surgery, 2021

Publication

33 Melissa Thiemi Kato, Evelyn Mikaela Kogawa, Carlos Neanes Santos, Paulo César Rodrigues

Conti. "TENS and low-level laser therapy in the management of temporomandibular disorders", Journal of Applied Oral Science, 2006

Publication

34

[ebin.pub](#)

Internet Source

<1 %

35

[healthdocbox.com](#)

Internet Source

<1 %

36

[thejournalofheadacheandpain.biomedcentral.com](#)

Internet Source

<1 %

37

[tnsroindia.org.in](#)

Internet Source

<1 %

38

[www5.bahiana.edu.br](#)

Internet Source

<1 %

39

Afaf A. M. Shaheen, Hajar Alhajri, Noura Alrajeeb, Ruba Almoammar, Arwa Alyousef, Syamala Buragadda, Ganeswararao Melam. "Level of awareness of dentists about the role of physiotherapy in temporomandibular disorders: a pilot study in Riyadh, Saudi Arabia", Bulletin of Faculty of Physical Therapy, 2020

Publication

<1 %

40

Dorrit Nitzan, Rafael Benoliel, Gary Heir, Franklin Dolwick. "Pain and dysfunction of the temporomandibular joint", Elsevier BV, 2008

Publication

<1 %

41

Rodrigo Lorenzi POLUHA, Giancarlo De la Torre CANALES, Yuri Martins COSTA, Eduardo GROSSMANN et al. "Temporomandibular joint disc displacement with reduction: a review of

<1 %

mechanisms and clinical presentation",
Journal of Applied Oral Science, 2019

Publication

42

"MRI of the Temporomandibular Joint",
Springer Science and Business Media LLC,
2020

Publication

<1 %

43

Abdalwhab Zwiri, Manal Ahmad Alrawashdeh,
Mohammad Khan, Wan Muhamad Amir W.
Ahmad et al. "Effectiveness of the Laser
Application in Temporomandibular Joint
Disorder: A Systematic Review of 1172
Patients", Pain Research and Management,
2020

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On