



International Journal of Advanced Research

Publisher's Name: Jana Publication and Research LLP

www.journalijar.com

REVIEWER'S REPORT

Manuscript No.: IJAR-51821 Date: 5/23/2025

Title: Misuse of Anxiolytics Among Women with Psychiatric Disorders

Recommendation:

Accept as it is					
Accept after minor revision	Rating	Excel.	Good	Fair	P
Accept after major revision	Originality		$\sqrt{}$		
Do not accept (Reasons below)	Techn.		ما		
	Quality		V		
	Clarity				

Significance

Reviewer Name: Dr Azadeh Mousavi Date: 5/23/2025

Reviewer's Comment for Publication.

(To be published with the manuscript in the journal)

The reviewer is requested to provide a brief comment (3-4 lines) highlighting the significance, strengths, or key insights of the manuscript. This comment will be Displayed in the journal publication alongside with the reviewers name.

This study provides valuable insights into the misuse of anxiolytics among women with psychiatric disorders, highlighting key sociodemographic and clinical factors. The findings underscore the need for targeted interventions to mitigate dependence risks. The comparative analysis with international studies adds depth, though clarity in methodology presentation could be improved.

Detailed Reviewer's Report Strengths:

The study addresses a critical public health issue with well-defined objectives.

Robust use of the Severity of Dependence Scale (SDS) and comparative analysis with global data.

Clear implications for clinical practice, including preventive measures (e.g., prescription limits, patient education).

Areas for Improvement:

<u>Clarity:</u> The methodology section lacks detail on sample selection criteria (e.g., exclusion/inclusion criteria).

<u>Data Presentation:</u> Tables and figures (e.g., SDS results) could be better integrated into the text for readability.

ISSN: 2320-5407

International Journal of Advanced Research

Publisher's Name: Jana Publication and Research LLP

www.journalijar.com

REVIEWER'S REPORT

Discussion: Elaborate on why Alprazolam misuse is prevalent compared to other benzodiazepines.