

Hypo-fractionated radiotherapy: a technical and societal innovation in the treatment of breast cancer.

by Jana Publication & Research

Submission date: 27-May-2025 01:32PM (UTC+0700)

Submission ID: 2664989466

File name: IJAR-51882.docx (18.18K)

Word count: 1479

Character count: 8520

Hypo-fractionated radiotherapy: a technical and societal innovation in the treatment of breast cancer.

Abstract: Adjuvant radiotherapy is a key element in the treatment of early breast cancer.

In recent years, there has been a growing trend towards hypofractionated external radiotherapy regimens.

Many have been shown to be equivalent to the standard regimen in terms of tumour control and cosmetic results, while improving patients' quality of life.

This paper provides an update on the main rationales for the use of hypofractionated radiotherapy in breast cancer, and reviews the bulk of the literature that has concluded that hypofractionated radiotherapy is effective and safe.

We examine various hypofractionation regimens used in our Mohammed VI Centre for the treatment of cancers, aiming at care centred on the quality of life of our patients while improving their productivity at home and at work in order to facilitate their social reintegration in the face of the disease.

We retrospectively analysed all patients treated in our department using the Fast Forward protocol and other hypofractionated regimens after the original publication was published and after an institutional consensus had been reached regarding the selection of patients with breast cancer without indication for lymph node irradiation.

Patients were selected between January 2021 and December 2024 at the Mohammed VI Cancer Treatment Centre.

The median age of the patients was 53 years. Invasive ductal carcinoma was the most common histological type (90%). Lumpectomy was performed in 19 patients. The preferred site was the left breast.

All patients were treated using modern conformal techniques with coverage of the predicted target volume of between 95 and 100%.

Treatment toxicity was assessed in the short, medium and long term in accordance with world standards. Today, 70% of our patients are still active in their social and professional lives, and 80% of them are mothers.

At present, we are joining the growing body of scientific evidence in making hypo-fractionated radiotherapy the therapeutic standard for the irradiation of early breast cancer.

With fewer fractions of treatment and more fractions of life for our patients, hypofractionated radiotherapy is an example of a care pathway that aims to deploy effective strategies for de-escalating the therapeutic burden and reducing the after-effects of cancer treatment.

keywords: External beam radiotherapy ,Hypofractionated radiotherapy, breast cancer, patient-centred care, quality of life.

Introduction:

Adjuvant radiotherapy is a key element in ⁶the treatment of early breast cancer.

¹In recent years, there has been a growing trend towards hypofractionated external beam radiotherapy regimens.

⁴Many have been shown to be equivalent to the standard regimen in terms of tumour control and cosmetic results, while improving patients' quality of life.

Materials and methods:

This is a retrospective analytical study: We are examining various hypofractionation regimens in the treatment of early breast cancer, used in our Mohammed VI Center for Cancer Treatment, aiming at care focused on the quality of life of our patients while improving their productivity at home, at work in order to facilitate their social reintegration in the face of the disease.

Two questionnaires were carried out with the 25 patients treated and the care team: 30 doctors and radiotherapy machine technicians.

Patient files were selected between January 2021 and December 2024 at the Mohammed VI Cancer Treatment Center.

Results:

⁵The median age of the patients was 53 years. Invasive ductal carcinoma was ³the most common histological type (90%), with stages limited to pT1-T2 N0 M0.

³Tumorectomy, or any other conservative surgery was performed in 19 patients.

The preferred site was the left breast.

¹²All patients received radiation to the breast ⁷alone at a total dose of 26 Gy in five fractions of 5.2 Gray, ¹¹in accordance with the Fast Forward protocol. All patients were seen in consultation and treated for the first fraction on Monday.

The blocked inspiration technique was systematically proposed. The end-of-treatment consultation took place on Friday (after five sessions).

All patients were treated using modern intensity-modulated conformal techniques, with coverage of the predicted target volume of between 95 and 100%.

Treatment toxicity was assessed in the short, medium and long term in accordance with world standards. There was only one case of Grade I radiodermatitis in the short term.

Today, 70% of our patients are still active in their social and professional environment, and 80% of them are mothers. The adapted hypofractionated treatment regime has enabled our patients to remain independent both at home and at work.

For patients who live a long way from the hospital, it has reduced travel costs and the cost of a long stay compared with the conventional regime.

We also found that One Week Radiotherapy was very well tolerated, marking the end of a long battle after the heavy impact of chemotherapy and surgery.

The thirty professionals questioned, including radiotherapists and specialised technicians, all agreed on the numerous advantages of hypofractionated radiotherapy, reducing the problems of managing the reproducibility of treatment positions, increasing the number of patients recruited to the machine, with better tolerance of the treatment and fewer undesirable effects, with local control and efficacy superimposable on the conventional regime.

Discussion:

Doctors in the UK's FAST group have shown that it is possible, in patients with no risk of recurrence, to reduce the protocol to 5 weeks, with one session per week.

explains the radiotherapy oncologist. She and her colleagues studied this approach and further concentrated treatment time with a schedule of 5 sessions over 1 week. In their 'FAST Forward' study, published in 2020, the Curie researchers presented data comparable to the 15-session, 3-week protocol, with a 5-year follow-up(1).

In practice, we administer a slightly higher dose at each session than with the conventional hypo-fractionated regimen, but the total dose of radiation, accumulated over the 5 sessions, is lower.

Explains Youlia Kirova. With 5 years' hindsight, this irradiation regimen, known as 'FAST Forward', showed equal efficacy to hypo-fractionation in 15 fractions and 3 weeks in a controlled phase III study involving more than 4,000 patients (1). The aim of this approach is to maintain the same therapeutic result while reducing the impact of treatment on patients' lives.

For the moment, this protocol is aimed at patients aged over 70, with no high risk of recurrence. They lead active lives, look after their grandchildren... It's a real benefit for them to be able to be treated in just one week.

8

These patients are treated with radiotherapy after breast-conserving surgery. Only patients without lymph node involvement can benefit from this extreme hypo-fractionation regime.

The Institut Curie is a pioneer in this field, and the results of the first patients to be treated have already been published (2).

1

Breast cancer is one of the most prevalent diseases in women. Prevention and treatments have lowered mortality; nevertheless, the impact of the diagnosis and treatment continue to impact all aspects of patients' lives (physical, emotional, cognitive, social, and spiritual) (3).

1

Supporting and accompanying breast cancer patients throughout this process requires a deep understanding of their experiences. To describe the patient's experiences, including thoughts, emotions, feelings, worries, and concerns, the phrase "patient voice" has been used, which is becoming increasingly common in healthcare (4).

1

The women diagnosed with breast cancer describe the journey as a process tremendously affected by the different medical stages. Each stage has its own characteristics that condition the experiences, unleashing specific physical, emotional, cognitive, and social processes. Additionally, the patients perceive this entire process as pre-established journey they must undertake to save their life, with its protocols based on the type and stage of cancer⁽³⁾.

Our work is in line with all this scientific evidence, and its aim is to highlight the benefits of reducing the burden of radiotherapy or any other stage in this long process, while ensuring its therapeutic effectiveness.

Conclusion:

We are joining the growing body of scientific evidence in making hypofractionated radiotherapy the therapeutic standard for the irradiation of early breast cancer: **Fewer treatment fractions, more fractions of life for our patients.**

Hypofractionated radiotherapy is an example of a care pathway designed to deploy effective strategies for de-escalating the therapeutic burden for our patients.


References:

(1) Brunt AM, Haviland JS, Wheatley DA, Sydenham MA, Alhasso A, Bloomfield DJ, et al. Hypofractionated breast radiotherapy for 1 week versus 3 weeks (FAST-Forward): 5-year efficacy and late normal tissue effects results from a multicentre, non-inferiority, randomised, phase 3 trial. The Lancet [Internet]. 23 mai 2020

(2) Chaffai I, Cao K, Kissel M, El Ayachy R, Aboudaram A, Fourquet A, Kirova Y. : Patient selection and early tolerance of whole breast irradiation according to the "Fast Forward" protocol: Preliminary results]. Cancer Radiother. 2021:S1278-3218(21)00192-X. doi: 10.1016/j.canrad.2021.07.036. Online ahead of print. PMID: 34454837 French.

(3) Ciria-Suarez L, Jiménez-Fonseca P, Palacín-Lois M, Antoñanzas-Basa M, Fernández-Montes A, Manzano-Fernández A, et al. (2021) Breast cancer patient experiences through a journey map: A qualitative study. PLoS ONE 16(9): e0257680. <https://doi.org/10.1371/journal.pone.0257680>

(4) Hall LK, Kunz BF, Davis EV., Dawson RI, Powers RS. The cancer experience map: An approach to including the patient voice in supportive care solutions. Journal of Medical Internet Research. 2015. <https://doi.org/10.2196/jmir.3652> PMID: 26022846



Hypo-fractionated radiotherapy: a technical and societal innovation in the treatment of breast cancer.

ORIGINALITY REPORT

22%

SIMILARITY INDEX

19%

INTERNET SOURCES

22%

PUBLICATIONS

11%

STUDENT PAPERS

PRIMARY SOURCES

1

hdl.handle.net

Internet Source

11%

2

pubmed.ncbi.nlm.nih.gov

Internet Source

3%

3

&NA;, . "15th Biennial Meeting of the International Gynecologic Cancer Society :", International Journal of Gynecological Cancer, 2014.

Publication

1%

4

api.repository.cam.ac.uk

Internet Source

1%

5

covid19dataportal.es

Internet Source

1%

6

pmc.ncbi.nlm.nih.gov

Internet Source

1%

7

I. Chaffai, K. Cao, M. Kissel, R. El Ayachy, A. Aboudaram, A. Fourquet, Y. Kirova. "Sélection de patientes et tolérance précoce de l'irradiation mammaire selon le protocole «Fast Forward»: résultats préliminaires", Cancer/Radiothérapie, 2021

Publication

1%

8

www.actaoncologicturcica.com

Internet Source

1%

9 S.K. Vinod, R. Merie, S. Harden. "Quality of Decision Making in Radiation Oncology", Clinical Oncology, 2024 1 %
Publication

10 www.researchsquare.com 1 %
Internet Source

11 Jihane Bouziane, Pierre Loap, Sofiane Allali, Kim Cao, Alain Fourquet, Youlia Kirova. "Ultra hypofractionated breast radiotherapy according to the Fast-Forward scheme: Excellent early tolerance or delayed skin reactions? Real life experience", Cancer/Radiothérapie, 2025 1 %
Publication

12 S. Haldar, B. Sarkar, A. Dixit. "Dose to Organ at Risk and its Characteristic Variation with the Clinically Used Different Prescription Levels for Early-stage Left-sided Breast Cancer", Clinical Oncology, 2023 <1 %
Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On