

# Evaluation of Biochemical and Hematological Alterations in Patients with Cholelithiasis: A Hospital-Based Cross-Sectional Study

*by* Jana Publication & Research

---

**Submission date:** 28-May-2025 01:07PM (UTC+0700)

**Submission ID:** 2664984353

**File name:** IJAR-51920.docx (58.01K)

**Word count:** 2660

**Character count:** 16953

**Evaluation of Biochemical and Hematological Alterations in Patients with  
Cholelithiasis: A Hospital-Based Cross-Sectional Study**

## Abstract:

**Background:** Cholelithiasis, or gallstone disease, is a common gastrointestinal condition marked by the formation of gallstones within the gallbladder, often leading to biliary obstruction, hepatic dysfunction, and inflammation. It is particularly prevalent in northern India, including Uttar Pradesh. This study was conducted to assess the biochemical and hematological changes in cholelithiasis patients.

**Methods:** A hospital-based cross-sectional study was conducted at Teerthanker Mahaveer Hospital and Research Centre, Moradabad, Uttar Pradesh, over a duration of 6 months. A total of 240 participants were enrolled, including 118 confirmed cholelithiasis patients and 122 age- and sex-matched healthy controls. Blood samples were analyzed for total, direct, and indirect bilirubin; liver enzymes (AST, ALT, ALP); pancreatic enzymes (amylase, lipase); and inflammatory markers (CRP, TLC). Data analysis was done using SPSS software and statistical comparisons were performed using the Mann-Whitney U test. Ethical approval was obtained from the Institutional Ethical Committee, College of Paramedical Sciences (Ref. No.: PM/ETHICAL/COPS/2024/018).

**Results:** Cholelithiasis patients showed significantly elevated levels of total bilirubin ( $2.60 \pm 0.11$  mg/dL), direct bilirubin ( $0.64 \pm 0.06$  mg/dL), and indirect bilirubin ( $1.96 \pm 0.10$  mg/dL). Liver enzymes including AST ( $33.26 \pm 1.24$  U/L), ALT ( $31.96 \pm 2.20$  U/L), and ALP ( $312.66 \pm 17.38$  U/L) were significantly increased in the patient group. Inflammatory markers such as CRP ( $7.38 \pm 0.48$  mg/dL) and TLC ( $6729.17 \pm 409.84 \times 10^3/\mu\text{L}$ ) also showed significant elevations. Pancreatic enzymes (amylase and lipase) showed mild but not statistically significant changes.

**Conclusion:** The study demonstrates that cholelithiasis is associated with notable biochemical and hematological alterations, particularly in liver function and inflammatory markers. These parameters can be effectively used in the clinical assessment and early detection of gallstone-related complications.

**Keywords:** Cholelithiasis, Liver Enzymes, Bilirubin, Inflammatory Markers.

## Introduction

<sup>19</sup>Cholelithiasis, commonly known as gallstone disease, is a significant global health concern <sup>25</sup>characterized by the formation of gallstones <sup>12</sup>within the gallbladder due to bile composition imbalances (1). It is a prevalent gastrointestinal disorder affecting millions worldwide, with <sup>23</sup>variations in prevalence based on geographical, genetic, and lifestyle factors (2). The global prevalence of gallstones ranges between 5% and 25%, with higher rates observed in developed countries. In the United States, approximately 15% of the population is affected, while in Europe, prevalence varies from 9% to 21%. In contrast, regions such as Africa and Southeast Asia report significantly lower prevalence rates (3). In India, gallstone disease affects an estimated 4% of the population, contributing to nearly 1 million new cases annually. Notably, the northern states, including Uttar Pradesh, exhibit a higher prevalence than southern regions, with North Indians being nearly seven times more at risk than their South Indian counterparts (4). Uttar Pradesh, in particular, has emerged as a high-burden region for gallstone disease, making it crucial to understand its biochemical and hematological implications in this population. Gallstones primarily develop due to an imbalance in bile constituents, including cholesterol, bilirubin, and bile salts, leading to their precipitation and stone formation (5). While many individuals remain asymptomatic, others experience severe <sup>29</sup>complications such as acute cholecystitis, obstructive jaundice, and pancreatitis (6). Timely diagnosis and management are essential to prevent disease progression and associated morbidity (7). Biochemical and hematological alterations play a pivotal role in diagnosing and monitoring gallstone disease (8,9). <sup>9</sup>Liver function tests (LFTs), including total, direct, and indirect bilirubin, aspartate aminotransferase (AST), alanine aminotransferase (ALT), and alkaline phosphatase (ALP), provide insight into hepatobiliary dysfunction. Additionally, serum amylase and lipase are assessed to rule out pancreatic involvement, while <sup>11</sup>inflammatory markers like total leukocyte count (TLC) and C-reactive protein (CRP) help evaluate systemic inflammation. Identifying variations in these parameters can enhance diagnostic accuracy and improve clinical management strategies (10). This study aims to evaluate and compare biochemical and hematological alterations in cholelithiasis patients and healthy controls at a tertiary care hospital in Moradabad, a region within Uttar Pradesh with a high burden of gallstone disease.

## Material and Methodology

This hospital-based cross-sectional study was conducted at Teerthanker Mahaveer Hospital and Research Center, Moradabad, over six months. A total of 236 participants were enrolled,

comprising 118 USG-confirmed cholelithiasis patients and 118 age- and sex-matched healthy controls on the basis of inclusion and exclusion Blood samples were collected from all participants under aseptic conditions from the antecubital vein using sterile syringes and transferred into EDTA and plain vacutainers. After getting informed consent form from all participants. The collected samples were analyzed for total, direct, and indirect bilirubin, AST, ALT, ALP, serum amylase, serum lipase, TLC, and CRP using standard laboratory protocols. Data were recorded in an Excel sheet and statistically analyzed using SPSS software. Ethical approval for this study was obtained from the Institutional Ethical Committee, College of Paramedical Sciences, (PM/ETHICAL/COPS/2024/018).

Result

Gender Distribution Among Study Populations

The study included a total of 240 participants, comprising 148 females (61.7%) and 92 males (38.3%), indicating a higher prevalence of gallstone disease among females as show in Figure 1 (a). Participants were categorized into two groups: cases (cholelithiasis patients) and controls (healthy individuals). Among the 120 cholelithiasis cases, 80 (66.7%) were female, while 40 (33.3%) were male as show in Figure 1 (b).




Figure 1. Show the Gender distribution in study populations overall and group wise.

Mean Age Among Study Populations

The mean age of cholelithiasis patients (cases) was  $37.23 \pm 11.48$  years, whereas the mean age of the control group was  $39.18 \pm 12.34$  years as show in Figure 2 (a). Gender-based analysis revealed that the mean age for females in the case group was  $36.02 \pm 11.45$  years,

while males had a mean age of  $39.65 \pm 11.31$  years. Among the control group, females had a mean age of  $38.91 \pm 12.40$  years, and males had a mean age of  $39.51 \pm 12.37$  years as show in Figure 2 (b).




Figure 2. Show the mean age of study populations overall populations and group wise.

### Biochemical and Hematological Biomarkers in Study Populations

Significant differences were observed in biochemical and hematological markers between cholelithiasis patients and healthy controls as show in Table 1. Total bilirubin ( $2.60 \pm 0.11$  mg/dL vs.  $0.91 \pm 0.04$  mg/dL,  $p < 0.01$ ), direct bilirubin ( $0.64 \pm 0.06$  mg/dL vs.  $0.23 \pm 0.02$  mg/dL,  $p < 0.01$ ), and indirect bilirubin ( $1.96 \pm 0.10$  mg/dL vs.  $0.68 \pm 0.04$  mg/dL,  $p < 0.01$ ) were markedly elevated in cholelithiasis patients. Liver enzymes, including AST ( $33.26 \pm 1.24$  U/L) and ALT ( $31.96 \pm 2.20$  U/L), were significantly higher in cases compared to controls ( $p < 0.01$ ). ALP, a marker of biliary obstruction, was significantly increased ( $312.66 \pm 17.38$  U/L vs.  $143.31 \pm 2.78$  U/L,  $p < 0.01$ ).

Inflammatory markers, CRP ( $7.38 \pm 0.48$  mg/dL) and TLC ( $6729.17 \pm 409.84$  ( $10^3/\mu\text{L}$ )), were also elevated in cholelithiasis patients ( $p < 0.01$ ), indicating systemic inflammation. Amylase levels were slightly higher in cases ( $46.94 \pm 1.07$  U/L) than in controls ( $44.60 \pm 1.44$  U/L,  $p < 0.01$ ), while lipase levels showed no significant difference ( $p = 0.107$ ). These findings confirm biochemical alterations associated with gallstone disease, highlighting liver dysfunction and inflammation as show in Table 1.

**Table 1. Comparison of Study Parameters in Study Populations According to Groups**

Parameters	Control	Case	Z-Vale	P-Value
<b>Total Bilirubin</b> (mg/dl)	0.91 ± 0.04	2.60 ± 0.11	-13.95	<0.00**
<b>Direct Bilirubin</b> (mg/dL)	0.23 ± 0.02	0.64 ± 0.06	-13.40	<0.00**
<b>Indirect Bilirubin</b> (mg/dL)	0.68 ± 0.04	1.96 ± 0.10	-13.39	<0.00**
<b>AST (U/L)</b>	26.74 ± 3.43	33.26 ± 1.24	-13.03	<0.00**
<b>ALT (U/L)</b>	26.42 ± 4.23	31.96 ± 2.20	-9.99	<0.00**
<b>ALP (U/L)</b>	143.31 ± 2.78	312.66 ± 17.38	-13.38	<0.00**
<b>Amylase (U/L)</b>	44.60 ± 1.44	46.94 ± 1.07	-10.61	<0.00**
<b>Lipase (U/L)</b>	76.78 ± 18.81	80.74 ± 16.29	-1.61	0.107
<b>CRP (mg/dL)</b>	4.39 ± 0.70	7.38 ± 0.48	-13.38	<0.00**
<b>TLC ((10<sup>3</sup>/μL))</b>	6444.17 ± 467.39	6729.17 ± 409.84	-4.66	<0.00**

All values were mean and standard deviation (SD), AST (Aspartate Aminotransferase); ALT (Alanine Aminotransferase); ALP (Alkaline Phosphatase); Amylase; Lipase; CRP (C-Reactive Protein); TLC (Total Leukocyte Count), mg/dL (milligrams per deciliter); U/L (units per liter); (10<sup>3</sup>/μL) is thousands per microliter. The statistical analysis was performed using the Mann-Whitney U test (a non-parametric test), as indicated by the Z-values. Asterisks (\*\*) indicate highly significant p-values (p < 0.01).

## Discussion

The present study evaluated biochemical and hematological parameters in cholelithiasis patients, revealing significant alterations compared to healthy controls. These findings align with existing literature and provide insights into the pathophysiological mechanisms underlying gallstone disease. The present study evaluated biochemical and hematological alterations in patients with cholelithiasis, revealing significant differences compared to healthy controls. Elevated levels of total bilirubin (2.60 ± 0.11 mg/dL), direct bilirubin (0.64 ± 0.06 mg/dL), and indirect bilirubin (1.96 ± 0.10 mg/dL) were observed in the case group, which is indicative of biliary obstruction. Similar findings have been reported by Aslam et al.

(2013), who found a strong correlation between hyperbilirubinemia and the presence of gallstones. Elevated bilirubin levels suggest impaired hepatic function due to obstructed bile flow, which is commonly seen in cholelithiasis patients (11). Moreover, our study showed a significant increase in alkaline phosphatase (ALP) levels ( $312.66 \pm 17.38$  U/L) in cases compared to controls ( $143.31 \pm 2.78$  U/L), supporting the findings of **L et al. (2009)**, who suggested that ALP serves as a key biomarker for biliary obstruction (12). Liver enzymes were significantly elevated in cholelithiasis patients, with AST ( $33.26 \pm 1.24$  U/L) and ALT ( $31.96 \pm 2.20$  U/L) levels higher than in controls ( $26.74 \pm 3.43$  U/L and  $26.42 \pm 4.23$  U/L, respectively). These results align with the findings of **Hawar and Lor (2022)**, who reported a significant rise in transaminase levels in cholelithiasis patients, particularly in those with obstructive pathology (13). Elevated AST and ALT levels are likely due to hepatocellular injury caused by bile stasis and inflammation, as reported by **Rangaswamy et al. (2017)** (14). This indicates the need for routine liver function testing in suspected gallstone cases for early detection and intervention. The inflammatory markers in our study showed a considerable rise in CRP ( $7.38 \pm 0.48$  mg/dL) and total leukocyte count (TLC) ( $6729.17 \pm 409.84$  ( $10^3/\mu\text{L}$ )) in cholelithiasis patients compared to controls ( $4.39 \pm 0.70$  mg/dL and  $6444.17 \pm 467.39$  ( $10^3/\mu\text{L}$ ), respectively). These findings agree with **Rajab et al. (2020)**, who reported that elevated CRP levels in gallstone patients indicate an ongoing inflammatory response, which may be due to recurrent biliary irritation (15). Likewise, **Napolitano et al. (2021)** demonstrated that higher leukocyte counts in cholelithiasis patients are associated with an increased risk of gallstone-related complications, such as acute cholecystitis (16). Elevated inflammatory markers suggest that gallstone disease is not merely a structural disorder but also involves an active inflammatory component. Our study also assessed pancreatic enzymes, showing a slight increase in amylase levels ( $46.94 \pm 1.07$  U/L) in cholelithiasis patients compared to controls ( $44.60 \pm 1.44$  U/L), while lipase levels remained statistically insignificant. Similar observations were made by **DM et al. (2017)**, who noted that isolated amylase elevation without lipase involvement is not uncommon in gallstone disease unless pancreatitis is present (17). This suggests that routine assessment of pancreatic enzymes in uncomplicated gallstone cases may not provide significant diagnostic value, as also reported by **Napolitano et al. (2021)** (16). Overall, our findings align with multiple studies emphasizing the diagnostic value of biochemical and hematological markers in gallstone disease. The significant alterations in bilirubin, liver enzymes, and inflammatory markers highlight their clinical relevance in evaluating cholelithiasis patients. However, our study's cross-sectional design limits causal interpretation, and larger longitudinal studies are



needed to confirm these associations. Future research should focus on correlating these biochemical markers with disease severity and treatment outcomes to improve clinical decision-making in cholelithiasis management.

### Conclusion

Cholelithiasis is associated with significant biochemical and hematological alterations, reflecting hepatic dysfunction, biliary obstruction, and systemic inflammation. Elevated bilirubin levels indicate impaired bile flow, while increased liver enzymes suggest hepatocellular damage. The rise in inflammatory markers, including CRP and TLC, highlights the role of inflammation in disease progression. These findings support the clinical relevance of routine biochemical assessments for early diagnosis and effective management of gallstone disease. **Further studies are needed to explore the** underlying pathophysiological mechanisms **and** potential therapeutic targets.

### Reference

1. Carey MC. Pathogenesis of gallstones. *Am J Surg* [Internet]. 1993 Apr 1 [cited 2025 Mar 11];165(4):410–9. Available from: <https://www.sciencedirect.com/science/article/pii/S0002961005809328>
2. Miquel JF, Covarrubias C, Villaroel L, Mingrone G, Greco AV, Puglielli L, et al. Genetic epidemiology of cholesterol cholelithiasis among Chilean Hispanics, Amerindians, and Maoris. *Gastroenterology* [Internet]. 1998 Oct 1 [cited 2025 Mar 11];115(4):937–46. Available from: <https://www.sciencedirect.com/science/article/pii/S0016508598702665>
3. Acalovschi M, Lammert F. World Gastroenterology Organisation (WGO). 2012 [cited 2025 Apr 5]. The Growing Global Burden of Gallstone Disease. Available from: <https://www.worldgastroenterology.org>
4. He M, Gu R, Huang X, Zhao A, Liu F, Zheng Y. Age-period-cohort analysis of gallbladder and biliary diseases epidemiological trends from 1990 to 2021 and forecasts for 2035: a systematic analysis from the global burden of disease study 2021. *BMC Gastroenterol* [Internet]. 2025 Jan 8 [cited 2025 Apr 5];25(1):7. Available from: <https://doi.org/10.1186/s12876-025-03592-w>
5. Kratzer W, Mason RA, Kächele V. Prevalence of gallstones in sonographic surveys worldwide. *J Clin Ultrasound* [Internet]. 1999 [cited 2025 Mar 11];27(1):1–7. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/%28SICI%291097-0096%28199901%2927%3A1%3C1%3A%3AAID-JCU1%3E3.0.CO%3B2-H>
6. Shabanzadeh DM, Sørensen LT, Jørgensen T. A Prediction Rule for Risk Stratification of Incidentally Discovered Gallstones: Results From a Large Cohort Study. *Gastroenterology* [Internet]. 2016 Jan 1 [cited 2025 Mar 11];150(1):156–167.e1. Available from: <https://www.sciencedirect.com/science/article/pii/S0016508515013086>

7. Tenner S, Dubner H, Steinberg W. Predicting gallstone pancreatitis with laboratory parameters: a meta-analysis. *Am J Gastroenterol*. 1994 Oct;89(10):1863–6.
8. Pratt DS, Kaplan MM. Evaluation of Abnormal Liver-Enzyme Results in Asymptomatic Patients. *N Engl J Med* [Internet]. 2000 Apr 27 [cited 2025 Mar 11];342(17):1266–71. Available from: <https://www.nejm.org/doi/full/10.1056/NEJM200004273421707>
9. Wang DQH, Cohen DE, Carey MC. Biliary lipids and cholesterol gallstone disease. *J Lipid Res* [Internet]. 2009 Apr 1 [cited 2025 Mar 11];50:S406–11. Available from: [https://www.jlr.org/article/S0022-2275\(20\)30647-7/abstract](https://www.jlr.org/article/S0022-2275(20)30647-7/abstract)
10. Shaffer EA. Epidemiology of gallbladder stone disease. *Best Pract Res Clin Gastroenterol* [Internet]. 2006 Jan 1 [cited 2025 Mar 11];20(6):981–96. Available from: <https://www.sciencedirect.com/science/article/pii/S1521691806000540>
11. Aslam HM, Saleem S, Edhi MM, Shaikh HA, Khan JD, Hafiz M, et al. Assessment of gallstone predictor: comparative analysis of ultrasonographic and biochemical parameters. *Int Arch Med* [Internet]. 2013 Apr 24 [cited 2025 Apr 4];6(1):17. Available from: <https://doi.org/10.1186/1755-7682-6-17>
12. L H, Mr M, M AC, Wh W. Role of liver function tests in symptomatic cholelithiasis. *J Ayub Med Coll Abbottabad JAMC* [Internet]. 2009 Jun [cited 2025 Apr 4];21(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/20524486/>
13. Taban S, Hawar, Diyari A. Lor. ResearchGate. 2024 [cited 2025 Apr 4]. (PDF) Comparative Study of Liver Function Test in Cholelithiasis Patient Pre and Post Cholecystectomy. Available from: [https://www.researchgate.net/publication/380889352\\_Comparative\\_Study\\_of\\_Liver\\_Function\\_Test\\_in\\_Cholelithiasis\\_Patient\\_Pre\\_and\\_Post\\_Cholecystectomy](https://www.researchgate.net/publication/380889352_Comparative_Study_of_Liver_Function_Test_in_Cholelithiasis_Patient_Pre_and_Post_Cholecystectomy)
14. Rangaswamy R, Singh CG, Singh HM, Punyabati P, Nyuwi KT. Impact of Biliary Calculi on the Liver. *J Clin Diagn Res JCDR* [Internet]. 2017 Apr [cited 2025 Apr 4];11(4):PC04–7. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5449850/>
15. Rajab IM, Majerczyk D, Olson ME, Addams JMB, Choe ML, Nelson MS, et al. C-reactive protein in gallbladder diseases: diagnostic and therapeutic insights. *Biophys Rep* [Internet]. 2020 Jun 1 [cited 2025 Apr 4];6(2):49–67. Available from: <https://doi.org/10.1007/s41048-020-00108-9>
16. Napolitano LM, Alam HB, Biesterveld BE, Gemmete JJ, Kronick SL, Pomerantz B, et al. Evaluation and Management of Gallstone-Related Diseases in Non-Pregnant Adults [Internet]. Ann Arbor (MI): Michigan Medicine University of Michigan; 2021 [cited 2025 Apr 4]. (Michigan Medicine Clinical Care Guidelines). Available from: <http://www.ncbi.nlm.nih.gov/books/NBK569245/>
17. Dm KN, B K, Karthick R. A Cross Sectional Study on Gallstone Disease and its Risk Factors in a Tertiary Care Hospital of South India. *JSM Gastroenterol Hepatol* [Internet]. 2017 Dec 29 [cited 2025 Apr 4]; Available from: <https://www.jscimedcentral.com/article/A-Cross-Sectional-Study-on-Gallstone-Disease-and-its-Risk-Factors-in-a-Tertiary-Care-Hospital-of-South-India>

**Acknowledgment:** I sincerely thank the Department of Medical Laboratory Techniques, College of Paramedical Sciences, Teerthanker Mahaveer University, Moradabad, Uttar Pradesh, India, for their continuous support. I extend my gratitude to the Principal, Head of Department, my Guide, esteemed faculty members, and dedicated laboratory staff for their valuable guidance and encouragement throughout this research. I am also deeply grateful to the Teerthanker Mahaveer Hospital and Research Center and the patients for their kind cooperation and contribution.

**Conflict of Interest:** No

# Evaluation of Biochemical and Hematological Alterations in Patients with Cholelithiasis: A Hospital-Based Cross-Sectional Study

## ORIGINALITY REPORT

17%	14%	10%	4%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

## PRIMARY SOURCES

1	Submitted to Teerthanker Mahaveer University Student Paper	1%
2	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet Source	1%
3	<a href="http://journals.squ.edu.om">journals.squ.edu.om</a> Internet Source	1%
4	Submitted to South College Student Paper	1%
5	<a href="http://pmc.ncbi.nlm.nih.gov">pmc.ncbi.nlm.nih.gov</a> Internet Source	1%
6	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	1%
7	<a href="http://gsconlinepress.com">gsconlinepress.com</a> Internet Source	1%
8	<a href="http://link.springer.com">link.springer.com</a> Internet Source	1%
9	Jami Jackson, Melissa Miller, Mary Moffatt, Shannon Carpenter, Ashley Sherman, James Anderst. "Bruising in Children", Clinical Pediatrics, 2014 Publication	1%

10	Sujin Lee, Jongyoon Kim, Bongjin Lee. "Dose-Response Analysis of Nafamostat Mesylate in Pediatric Critical Care: Implications for Anticoagulation Monitoring During Continuous Venovenous Hemodiafiltration", Springer Science and Business Media LLC, 2025 Publication	1 %
11	<a href="https://pesquisa.bvsalud.org">pesquisa.bvsalud.org</a> Internet Source	1 %
12	<a href="https://f1000research.com">f1000research.com</a> Internet Source	1 %
13	<a href="https://iris.univr.it">iris.univr.it</a> Internet Source	1 %
14	Weingartner, O.. "Vascular Effects of Diet Supplementation With Plant Sterols", Journal of the American College of Cardiology, 20080422 Publication	<1 %
15	<a href="https://www.medrxiv.org">www.medrxiv.org</a> Internet Source	<1 %
16	<a href="https://www.researchgate.net">www.researchgate.net</a> Internet Source	<1 %
17	Ahmet Usta, Emin Gemcioglu, Salih Baser, Osman Ersoy, Yunus Halil Polat, Abdülsamet Erden, Seval Izdes. "Comparison of APACHE II scores and mortality with CRP/albumin, neutrophil/lymphocyte and thrombocyte/lymphocyte ratios in patients admitted to internal medicine and anesthesia reanimation intensive care unit", Journal of Laboratory Medicine, 2021 Publication	<1 %

18	Rasoul Goli, Navid Faraji, Himan Maroofi, Amireh Hassanpour. "Effect of spiritual care on the quality of life in patients who underwent intracranial hemorrhage surgery: a randomized controlled trial", International Journal of Surgery, 2024 Publication	<1 %
19	Submitted to Sefako Makgatho Health Science University Student Paper	<1 %
20	healthdocbox.com Internet Source	<1 %
21	Mohammed Elmowafy, Khaled Shalaby, Mohammed Elkomy, Nabil K. Alruwaili et al. "Impact of highly phospholipid-containing lipid nanocarriers on oral bioavailability and pharmacodynamics performance of genistein", Pharmaceutical Development and Technology, 2022 Publication	<1 %
22	courses.washington.edu Internet Source	<1 %
23	epdf.pub Internet Source	<1 %
24	jssm.org Internet Source	<1 %
25	www.ijmse.com Internet Source	<1 %
26	www.mdedge.com Internet Source	<1 %
27	www.scielo.br Internet Source	<1 %

---

28	Anna Mueller, Roberto Warman. "Low-Dose Atropine for High Myopia: An Observational Cohort Study", AJO International, 2024	<1 %
Publication		

---

29	Gottfried Novacek. "Gender und Gallensteine", Wiener Medizinische Wochenschrift, 2006	<1 %
Publication		

---

30	boneandjoint.org.uk	<1 %
Internet Source		

---

31	www.science.gov	<1 %
Internet Source		

---

---

Exclude quotes	On	Exclude matches	Off
Exclude bibliography	On		