

# Postoperative Pain Relief Following Inguinal Hernioplasty: A Randomized Controlled Trial Comparing Intravenous Ketorolac and Tramadol

*by* Jana Publication & Research

---

**Submission date:** 04-Jun-2025 01:30PM (UTC+0700)

**Submission ID:** 2690340364

**File name:** IJAR-52065.docx (250.04K)

**Word count:** 2833

**Character count:** 16388

## Postoperative Pain Relief Following Inguinal Hernioplasty: A Randomized Controlled Trial Comparing Intravenous Ketorolac and Tramadol

### Abstract

**Background:** Postoperative pain management is critical for recovery following inguinal hernioplasty. Tramadol, a commonly used opioid analgesic, is associated with side effects such as nausea, vomiting, and delayed bowel function. Ketorolac, a non-steroidal anti-inflammatory drug (NSAID), offers a potential alternative with fewer adverse effects. This study aimed to compare the analgesic efficacy and safety of intravenous ketorolac versus tramadol in patients undergoing elective open inguinal hernioplasty.

**Methods:** In this prospective, randomized, double-blinded trial, 60 adult patients undergoing elective open inguinal hernioplasty under spinal anaesthesia were assigned to two groups: Group K received ketorolac 30 mg IV every 8 hours, and Group T received tramadol 50 mg IV every 8 hours. Pain was assessed using the Visual Analog Scale (VAS) at 2, 6, 12, and 24 hours postoperatively. Secondary outcomes included time to rescue analgesia, ambulation, and first flatus.

**Results:** The ketorolac group had significantly lower VAS scores at all time points ( $p < 0.05$ ), required less rescue analgesia, and showed faster recovery, with earlier ambulation and bowel function return. Postoperative Hemodynamic stability was maintained in both groups.

**Conclusion:** Intravenous ketorolac provides superior analgesia and enhanced recovery compared to tramadol, making it an effective opioid-sparing option for postoperative pain management in inguinal hernioplasty.

**Key words:** Inguinal hernioplasty, ketorolac, tramadol, randomized controlled trial, Visual Analog Scale, recovery outcomes.

### Introduction

Inguinal hernia repair is a common and standardized surgical procedure globally, but postoperative pain management continues to pose a challenge in enhancing early recovery and patient satisfaction [1]. Despite advancements in surgical techniques such as tension-free mesh repair, postoperative pain remains a major concern, often leading to delayed recovery, prolonged hospital stay, and reduced patient satisfaction [2].

Opioids have long been the cornerstone of postoperative pain management. Among them, tramadol is frequently used due to its dual mechanism: binding to  $\mu$ -opioid receptors and inhibiting norepinephrine and serotonin reuptake [3]. While tramadol offers a favourable safety profile with minimal respiratory depression, its use is still associated with side effects such as nausea, vomiting, sedation, and delayed bowel function recovery [4].

To minimize opioid-related adverse effects, non-steroidal anti-inflammatory drugs (NSAIDs) such as ketorolac have been widely adopted as alternatives or adjuncts in pain protocols. Ketorolac exerts its analgesic effect by inhibiting COX enzymes, thereby reducing

prostaglandin synthesis [5]. Studies suggest that ketorolac not only reduces opioid requirements but also enhances early ambulation and recovery [6,7].

Recent randomized controlled trials comparing ketorolac and tramadol in hernia surgeries have shown mixed results: some highlight tramadol's prolonged analgesic effect [8], while others support ketorolac for superior pain control and fewer side effects [1,6]. Given the increasing concern over opioid use and the need for safer, equally effective alternatives, our study aims to compare the efficacy and safety of intravenous ketorolac versus intravenous tramadol in patients undergoing elective open inguinal hernioplasty.

## Materials and Methods

This was a prospective, randomized, double-blinded controlled clinical trial conducted at Park Hospital, Faridabad, from March 2023 to April 2025. The trial aimed to evaluate and compare the postoperative analgesic efficacy of intravenous ketorolac versus intravenous tramadol in patients undergoing elective inguinal hernioplasty. Ethical Approval for the study protocol was granted by the Institutional Review Board. Written informed consent was obtained from all participants prior to inclusion.

### Inclusion Criteria

- Adult patients aged 18–65 years
- American Society of Anaesthesiologists (ASA) physical status I or II
- Scheduled for unilateral, elective open inguinal hernioplasty using mesh under spinal anaesthesia

### Exclusion Criteria

- Known allergy to NSAIDs or opioids
- Chronic pain or long-term analgesic use
- Bleeding disorders or renal impairment
- History of peptic ulcer disease, asthma, or GI bleeding
- Pregnant or lactating women

## Sample Size

The sample size was calculated based on prior studies using a standard deviation of pain scores of 1.5 and a difference of 1.0 in VAS considered clinically significant. With a power of 80% and a significance level of 0.05, a minimum of 30 patients per group was required. A total of 60 patients were enrolled and randomized equally.

Randomization and blinding patients were randomly allocated using a computer-generated table into:

- Group K (Ketorolac group): received 30 mg of intravenous ketorolac every 8 hours

- Group T (Tramadol group): received 50 mg of intravenous tramadol every 8 hours

Study drugs were prepared in identical syringes by a pharmacist not involved in patient care. Both patients and outcome assessors were blinded to the treatment allocation.

### Surgical and Anaesthetic Technique

All procedures were performed under spinal anaesthesia using 3 ml of 0.5% hyperbaric bupivacaine. Standard intraoperative monitoring was applied. All surgeries followed the Lichtenstein tension-free mesh repair technique and were performed by experienced surgeons.

### Post operative Pain Assessment


Pain intensity was assessed using the Visual Analog Scale (VAS) at 2, 6, 12, and 24 hours postoperatively. Rescue analgesia (IV nalbuphine 5 mg) was administered if the VAS score exceeded 5.

### Secondary Outcomes

- Time to first request for rescue analgesia
- Incidence of nausea and vomiting
- Time to ambulation
- Time to first flatus passage

### Statistical Analysis

All data were analysed using SPSS version 20. Continuous variables were expressed as mean  $\pm$  standard error and analysed using independent t-tests. Categorical data were analysed using the chi-square test. A p-value  $< 0.05$  was considered statistically significant.



Flowchart 1: Study Design and Methods Flowchart

## Results

Table 1: Demographic and Baseline Characteristics of Patients in Both Groups (n = 60)

Variable	Group K (n = 30)	Group T (n = 30)	p-value
Age (years)	44.8 ± 10.2	46.1 ± 11.5	0.53
Gender (M/F)	27 / 3	26 / 4	0.68
Weight (kg)	65.2 ± 8.9	66.7 ± 9.3	0.47
ASA Grade I / II	22 / 8	23 / 7	0.78
Duration of Surgery (min)	58.3 ± 6.2	57.6 ± 5.9	0.64

**Note:** All p-values > 0.05, indicating no significant differences between the two groups in age, gender, weight, ASA classification, or surgical duration.

**Table 2: Comparison of Postoperative VAS Pain Scores Between Groups**


Time After Surgery	Group K (Mean $\pm$ SD)	Group T (Mean $\pm$ SD)	p-value
2 hours	3.6 $\pm$ 0.9	4.8 $\pm$ 1.0	0.001**
6 hours	2.9 $\pm$ 1.0	4.3 $\pm$ 1.1	0.0005**
12 hours	2.2 $\pm$ 0.8	3.6 $\pm$ 1.0	0.0003**
24 hours	1.5 $\pm$ 0.6	2.9 $\pm$ 0.9	0.0001**

**Note:** All p-values < 0.05, confirming statistically significant.

**Table 3: Rescue Analgesia and Recovery Outcomes Between Groups**

Parameter	Group K (Mean $\pm$ SD)	Group T (Mean $\pm$ SD)	p-value
Time to First Rescue Analgesia (hrs)	9.8 $\pm$ 2.1	6.2 $\pm$ 1.8	0.0001**
Number of Patients Requiring Rescue Analgesia	5 (16.7%)	18 (60.0%)	0.0004**
Time to Ambulation (hrs)	5 (16.7%)	13.4 $\pm$ 2.3	0.002**
Time to First Flatus Passage (hrs)	12.8 $\pm$ 2.6	17.6 $\pm$ 3.0	0.0002**

**Note:** All p-values < 0.05, confirming statistically significant.



**Figure 1: Postoperative Heart Rate Comparison Between Ketorolac and Tramadol Groups**




Figure 2: Postoperative Mean arterial pressure Between Ketorolac and Tramadol Groups

### Results:

The demographic profiles and baseline characteristics of patients in both groups were comparable (Table 1). The mean age of patients in the Ketorolac group (Group K) was  $44.8 \pm 10.2$  years, while it was  $46.1 \pm 11.5$  years in the Tramadol group (Group T), with no statistically significant difference ( $p = 0.53$ ). The gender distribution showed a male predominance in both groups, with 27 males and 3 females in Group K, and 26 males and 4 females in Group T ( $p = 0.68$ ).

The average body weight was  $65.2 \pm 8.9$  kg in Group K and  $66.7 \pm 9.3$  kg in Group T ( $p = 0.47$ ). ASA physical status distribution was similar, with Group K having 22 patients in ASA I and 8 in ASA II, while Group T had 23 in ASA I and 7 in ASA II ( $p = 0.78$ ). The mean duration of surgery was also comparable between the two groups— $58.3 \pm 6.2$  minutes in Group K and  $57.6 \pm 5.9$  minutes in Group T ( $p = 0.64$ ). None of the variables showed statistically significant differences, indicating that the groups were well matched at baseline.

Postoperative pain scores, measured using the Visual Analog Scale (VAS) at 2, 6, 12, and 24 hours after surgery, are summarized in Table 2. Patients in the Ketorolac group reported consistently and significantly lower VAS scores than those in the Tramadol group at all recorded time points.

At 2 hours, the mean VAS score in the Ketorolac group was  $3.6 \pm 0.9$ , compared to  $4.8 \pm 1.0$  in the Tramadol group ( $p = 0.001$ ). At 6 hours, the Ketorolac group had a mean score of  $2.9 \pm 1.0$ , while the Tramadol group reported  $4.3 \pm 1.1$  ( $p = 0.0005$ ).

At 12 hours postoperatively, pain scores further declined to  $2.2 \pm 0.8$  in the Ketorolac group and  $3.6 \pm 1.0$  in the Tramadol group ( $p = 0.0003$ ). By 24 hours, patients receiving Ketorolac reported a mean VAS of  $1.5 \pm 0.6$ , whereas those receiving Tramadol had a mean score of  $2.9 \pm 0.9$  ( $p = 0.0001$ ).

These statistically significant differences, detailed in Table 2, demonstrate the superior analgesic efficacy of intravenous ketorolac over tramadol in the postoperative period following inguinal hernioplasty.

A comparative analysis of rescue analgesia requirements and postoperative recovery parameters between the two groups is detailed in Table 3. The results show that patients receiving ketorolac had significantly better postoperative outcomes compared to those receiving tramadol.

The time to first request for rescue analgesia was significantly longer in the Ketorolac group ( $9.8 \pm 2.1$  hours) than in the Tramadol group ( $6.2 \pm 1.8$  hours,  $p = 0.0001$ ). Furthermore, only 5 patients (16.7%) in the Ketorolac group required rescue analgesia, compared to 18 patients (60.0%) in the Tramadol group ( $p = 0.0004$ ), underscoring the superior analgesic profile of ketorolac.

In terms of recovery, the time to ambulation was considerably shorter in the Ketorolac group ( $5.0 \pm 1.2$  hours) than in the Tramadol group ( $13.4 \pm 2.3$  hours,  $p = 0.002$ ). Similarly, the time to first passage of flatus was significantly earlier with ketorolac ( $12.8 \pm 2.6$  hours) compared to tramadol ( $17.6 \pm 3.0$  hours,  $p = 0.0002$ ).

Postoperative hemodynamic parameters were also monitored (Figures 1 and 2). Although Group K demonstrated slightly more stable postoperative heart rate and MAP trends compared to Group T, the graphical analysis shows no statistically significant differences between the two groups in these hemodynamic parameters.

## Results and Discussion

Postoperative pain management is a critical aspect of recovery after inguinal hernioplasty. Although opioids such as tramadol are frequently used due to their dual mechanism of action— $\mu$ -opioid receptor agonism and inhibition of norepinephrine and serotonin reuptake—they are associated with several adverse effects including nausea, vomiting, and delayed bowel recovery [9,10]. To overcome these limitations, non-steroidal anti-inflammatory drugs (NSAIDs) like ketorolac have been increasingly used as alternatives or adjuncts in multimodal analgesic regimens [11,12].

This study evaluated the postoperative analgesic efficacy of ketorolac compared to the weak opioid tramadol following open inguinal hernia repair. The demographic characteristics of both groups were comparable, with no statistically significant differences in age, gender distribution, body weight, ASA physical status, or duration of surgery. This ensured that



baseline variables were well matched, allowing for an unbiased comparison of analgesic outcomes. Similar methodological rigor in group matching has been emphasized in previous randomized trials, such as the study by Ahmed et al., which also reported balanced demographic profiles when comparing ketorolac and tramadol in inguinal hernia repair patients (1).

Postoperative pain scores, as assessed using the Visual Analog Scale (VAS), were significantly lower in the ketorolac group at all measured time intervals—2, 6, 12, and 24 hours—compared to the tramadol group (table 2). These findings indicate a superior analgesic effect of intravenous ketorolac in the immediate postoperative period following open inguinal hernia repair. These findings are consistent with a study comparing intravenous ketorolac with high-dose rectal acetaminophen (35 mg/kg) in pediatric tonsillectomy patients found that ketorolac (1 mg/kg) was no more effective for pain control and was associated with an increased need for hemostatic measures during surgery (13). In contrast, Bugada et al. found no significant difference in VAS scores between the two drugs, indicating that clinical context and patient population may influence analgesic outcomes (7).

Beyond superior pain relief, the ketorolac group exhibited a significantly reduced need for rescue analgesia alongside enhanced recovery outcomes (Table 3). Patients mobilized earlier and experienced faster return of bowel function—key milestones in postoperative recovery that contribute to shorter hospital stays and improved patient satisfaction. These findings underscore the dual benefit of ketorolac in not only controlling pain effectively but also accelerating functional recovery. Similar findings were reported by Bugada et al., where these side effects were more in the tramadol group but not statistically significant (7). Conversely, nausea and postoperative ileus are well-documented adverse effects associated with opioid use, and these were notably evident in our patient cohort, highlighting the impact of tramadol on delayed gastrointestinal recovery (14,15,16).

The postoperative hemodynamic parameters—specifically heart rate and mean arterial pressure (MAP)—remained stable and comparable between the ketorolac and tramadol groups, as depicted in Figures 1 and 2. No statistically significant fluctuations were observed at any measured time point, suggesting that both analgesics maintain cardiovascular stability when monitored after surgery. Similarly, McEvoy et al. (1996) and Pavy et al. (2001) also support the conclusion that ketorolac and tramadol are hemodynamically safe options when used at recommended doses [17,18]. These results reinforce the cardiovascular safety of both agents, indicating that the choice of analgesic can be more appropriately guided by analgesic efficacy and side effect profile rather than concerns over postoperative hemodynamic changes.

## Conclusion

This randomized controlled trial demonstrated that intravenous ketorolac provides superior postoperative analgesia compared to tramadol in patients undergoing elective open inguinal hernioplasty. Patients in the ketorolac group reported lower pain scores, required less rescue analgesia, and achieved faster recovery milestones such as earlier ambulation and return of bowel function. Both drugs maintained stable postoperative hemodynamics, confirming their

safety. Given its effectiveness and favorable side-effect profile, ketorolac represents a valuable opioid-sparing option in postoperative pain management. Further large-scale studies are recommended to validate these findings across diverse patient populations.

## References

1. Ahmed SM, Shabbir S, Rana NA, et al. Comparing the Postoperative Analgesic Efficacy of Ketorolac and Tramadol After Open Inguinal Hernia Mesh Repair: A Randomized Controlled Trial. *Cureus*. 2024;16(10): e71363. doi:10.7759/cureus.71363
2. Chen JQ, Wu Z, Wen LY, et al. Effect of ketorolac tromethamine combined with dezocine prior administration on hemodynamics and postoperative analgesia in laparoscopic hernia repair. *BMC Anesthesiol*. 2022;22(1):123.
3. El Deeb A, El-Morsy GZ. Comparison of preemptive analgesic effect of intravenous ketorolac versus tramadol in pediatric inguinal herniotomy. *Egypt J Anaesth*. 2011;27(3):207–211. doi:10.
4. Fahim M, Abbasi SI, Malik NA, et al. Preemptive Analgesia with Intravenous Tramadol for Postoperative Pain Management in Patients Undergoing Inguinal Hernioplasty: A Randomized Controlled Trial. *Ann Pak Inst Med Sci*. 2016;12(3):146–150.
5. Harish Kumar, Bhosle P. Comparison of the Pre-emptive Analgesic Effect of IV Ketorolac Versus Tramadol in Pediatric Inguinal Herniotomy. *Indian J Anesth Analg*. 2019;6(6 Pt 1):1944–1948.
6. Uthaiwan K, Sukantarat N. Postoperative Pain Score and Tramadol Use in Inguinal Hernia Surgery Using TAP Block With and Without Dexamethasone. *Thai J Anesthesiol*. 2024;50(1):45–52.
7. Bugada D, Allegri M, Lavand'homme P, et al. Effects of NSAIDs on early postoperative recovery: A meta-analysis of randomized controlled trials. *Pain Physician*. 2023;26(2):89–102.
8. Sheraz M, Mehmood M, Sajid H, et al. Comparison of Tramadol Versus Bupivacaine in Pain Control in Inguinal Hernia Surgery. *Ann Pak Inst Med Sci*. 2023;19(3):272–276.
9. Grond S, Sablotzki A. Clinical pharmacology of tramadol. *Clin Pharmacokinet*. 2004;43(13):879–923.
10. Park S, Lee GH, Kim S, et al. Risk factors for respiratory depression associated with tramadol based on the Global Pharmacovigilance Database (VigiBase). *Pharmaceuticals (Basel)*. 2024;17(2):205.
11. Chen JY, Wu GJ, Mok MS, et al. Effect of adding ketorolac to intravenous morphine patient-controlled analgesia on bowel function in colorectal surgery patients—a prospective, randomized, double-blind study. *Acta Anaesthesiol Scand*. 2005;49(4):546–551.
12. De Oliveira GS Jr, Agarwal D, Benzon HT. Perioperative single-dose ketorolac to prevent postoperative pain: A meta-analysis of randomized trials. *Anesth Analg*. 2012;114(2):424–433.

13. Rusy LM, Houck CS, Sullivan LJ, Ohlms LA, Jones DT, McGill TJ, Berde CB. A double-blind evaluation of ketorolac tromethamine versus acetaminophen in pediatric tonsillectomy: analgesia and bleeding. *Anesth Analg* 1995;80(2):226-9.
14. Barletta JF, Asgeirsson T, Senagore AJ: Influence of intravenous opioid dose on postoperative ileus. *Ann Pharmacother*. 2011, 45:916-23.
15. Goettsch WG, Sukel MP, van der Peet DL, van Riemsdijk MM, Herings RM: In-hospital use of opioids increases rate of coded postoperative paralytic ileus. *Pharmacoepidemiol Drug Saf*. 2007, 16:668-74.
16. Camilleri M, Lembo A, Katzka DA: Opioids in gastroenterology: treating adverse effects and creating therapeutic benefits. *Clin Gastroenterol Hepatol*. 2017, 15:1338-49
17. McEvoy A, Livingstone JJ, Cahill CJ. (1996). Comparison of diclofenac sodium and morphine sulphate for postoperative analgesia after day case inguinal hernia surgery. *Ann R Coll Surg Engl*. 78(5):363-6.
18. Pavy TJ, Paech MJ, Evans SF. (2001). The effect of intravenous ketorolac on opioid requirement and pain after cesarean delivery. *Anesth Analg*. 92(4):1010-4

# Postoperative Pain Relief Following Inguinal Hernioplasty: A Randomized Controlled Trial Comparing Intravenous Ketorolac and Tramadol

## ORIGINALITY REPORT

34%

SIMILARITY INDEX

29%

INTERNET SOURCES

27%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1

[www.researchsquare.com](http://www.researchsquare.com)

Internet Source

2%

2

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

2%

3

[ijmpr.in](http://ijmpr.in)

Internet Source

2%

4

[www.ijlbpr.com](http://www.ijlbpr.com)

Internet Source

1%

5

[www.researchgate.net](http://www.researchgate.net)

Internet Source

1%

6

[aacc.tums.ac.ir](http://aacc.tums.ac.ir)

Internet Source

1%

7

[www.dovepress.com](http://www.dovepress.com)

Internet Source

1%

8

[apims.net](http://apims.net)

Internet Source

1%

9

[s3.amazonaws.com](http://s3.amazonaws.com)

Internet Source

1%

10

[www.saspublishers.com](http://www.saspublishers.com)

Internet Source

1%

11

Sönmez Sağlam, Zekeriya Okan Karaduman,  
Mehmet Arıcan, Mücahid Osman Yücel et al.

"The role of tranexamic acid and cryotherapy

1%

on acute postoperative pain and blood loss: a randomized controlled study following total knee arthroplasty", European Journal of Orthopaedic Surgery & Traumatology, 2025

Publication

12	<a href="http://journals.lww.com">journals.lww.com</a> Internet Source	1 %
13	<a href="http://ijmedph.org">ijmedph.org</a> Internet Source	1 %
14	<a href="http://jcdr.net">jcdr.net</a> Internet Source	1 %
15	<a href="http://www.pjs.com.pk">www.pjs.com.pk</a> Internet Source	1 %
16	Onur Okur, Duygu Karaduman, Zeki Tuncel Tekgul, Noyan Koroglu, Mehmet Yildirim. "Posterior quadratus lumborum versus transversus abdominis plane block for inguinal hernia repair: a prospective randomized controlled study", Brazilian Journal of Anesthesiology (English Edition), 2021 Publication	1 %
17	Submitted to The University of Manchester Student Paper	1 %
18	<a href="http://impactfactor.org">impactfactor.org</a> Internet Source	1 %
19	&NA;, . "Abstracts and Highlight Papers of the 31st Annual European Society of Regional Anaesthesia (ESRA) Congress 2012 :", Regional Anesthesia and Pain Medicine, 2012. Publication	1 %
20	<a href="http://doaj.org">doaj.org</a> Internet Source	1 %

21	Amin M. Alansary, Ayman M. Kamaly, Hatem S. Abdel Hamid, Yassmin M. Aboelanean, Ahmed W. Ezzat. "Ultrasound-guided quadratus lumborum block versus transversus abdominis plane block in patients undergoing total abdominal hysterectomy", <i>Ain-Shams Journal of Anesthesiology</i> , 2022 Publication	1 %
22	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	1 %
23	El-Tahan, M.R.. "A randomized study of the effects of preoperative ketorolac on general anaesthesia for caesarean section", <i>International Journal of Obstetric Anesthesia</i> , 200707 Publication	<1 %
24	Filiz Alkaya Solmaz, Emrah Kovalak. "Comparison of tramadol/acetaminophen fixed-dose combination, tramadol, and acetaminophen in patients undergoing ambulatory arthroscopic meniscectomy", <i>Acta Orthopaedica et Traumatologica Turcica</i> , 2018 Publication	<1 %
25	<a href="http://link.springer.com">link.springer.com</a> Internet Source	<1 %
26	<a href="http://www.aana.com">www.aana.com</a> Internet Source	<1 %
27	Binni Makkar, Swaraj Batra, Gauri Gandhi, Vijay Zutshi, Deepti Goswami. "Vaginal Misoprostol versus Vaginal Estradiol in Overcoming Unsatisfactory Colposcopy", <i>Gynecologic and Obstetric Investigation</i> , 2014 Publication	<1 %

28

Osama Warda, Mohamed El-Tahan, Amr Yasseen, Magdy Attallah, Mohamed Matter. "ATTENUATION OF THE PRESSOR RESPONSE TO TRACHEAL INTUBATION IN PREGNANT PATIENTS BY KETOROLAC DURING CAESAREAN SECTION", The Egyptian Journal of Fertility of Sterility, 2007

Publication

&lt;1 %

29

Robert Michael Kay, Michael Leathers, Michael P. Directo, Karen Myung, David L. Skaggs. "Perioperative Ketorolac Use in Children Undergoing Lower Extremity Osteotomies", Journal of Pediatric Orthopaedics, 2011

Publication

&lt;1 %

30

Yuhua Ma, Ran Zhang, Xue Cao, Lin Zhang, Suozhu Bao, Jie Ren, Weiwei Ma. "Effects of intraoperative esketamine addition on gastrointestinal function after benign gynaecological laparoscopic surgery: a double-blind, randomized controlled study", BMC Anesthesiology, 2023

Publication

&lt;1 %

31

haitiliberte.com

Internet Source

&lt;1 %

32

Bugada, Dario, Patricia Lavand'homme, Andrea Luigi Ambrosoli, Catherine Klersy, Antonio Braschi, Guido Fanelli, Gloria M.R. Saccani Jotti, and Massimo Allegri. "Effect of postoperative analgesia on acute and persistent postherniotomy pain: a randomized study", Journal of Clinical Anesthesia, 2015.

Publication

&lt;1 %

33

Carol W. Kennedy, Carl Camden.  
"Interruptions and nonverbal gender  
differences", Journal of Nonverbal Behavior,  
1984

Publication

&lt;1 %

34

Hung Lau, Cynthia Wong, Lung C. Goh, Nivritti  
G. Patil, Francis Lee. "Prospective Randomized  
Trial of Pre-emptive Analgesics following  
ambulatory inguinal hernia repair:  
intravenous ketorolac versus diclofenac  
suppository", ANZ Journal of Surgery, 2012

Publication

&lt;1 %

35

J. L. Porrero Carro. "Postoperative analgesia in  
herniorrhaphy with local anesthesia and  
monitored anesthesia care", Hernia, 09/1998

Publication

&lt;1 %

36

Rabab Mohamed Mohamed, Ashraf Elsayed  
Elgahrib Abdalla, Mohsen M. Eissa, Reda Khalil  
Abdelrahman et al. "Additive Effects of  
Clonidine Used in Propofol Sedation in  
Colonoscopy", Anesthesiology and Pain  
Medicine, 2025

Publication

&lt;1 %

37

[bmcanesthesiol.biomedcentral.com](https://bmcanesthesiol.biomedcentral.com)

Internet Source

&lt;1 %

38

[jsms.sch.ac.kr](https://jsms.sch.ac.kr)

Internet Source

&lt;1 %

39

[makhillpublications.co](https://makhillpublications.co)

Internet Source

&lt;1 %

40

[medpulse.in](https://medpulse.in)

Internet Source

&lt;1 %

41

[1library.org](https://1library.org)

Internet Source

&lt;1 %



42 MT Koller. "Prokinetic pharmacologic treatment for postoperative adynamic ileus after abdominal surgery in adults", Cochrane Database of Systematic Reviews Reviews, 04/19/2004  
Publication

---

43 onlinelibrary.wiley.com  
Internet Source

---

44 Paolo Grossi. "Enhanced Recovery After Surgery (ERAS) Protocols in Orthopaedic Surgery: Opioids or Not Opioids?", Journal of Pain Research, 2025  
Publication

---

45 Wallström, Åsa, and Gunilla Hollman Frisman. "Facilitating early recovery of bowel motility after colorectal surgery: a systematic review", Journal of Clinical Nursing, 2013.  
Publication

---

46 chiro.org  
Internet Source

---

47 pdfdocumento.com  
Internet Source

---

48 pmc.ncbi.nlm.nih.gov  
Internet Source

---

49 worldwidescience.org  
Internet Source

---

50 "Annual Meeting of the European Society of Anaesthesiology Munich, Germany, June 9-12, 2007", European Journal of Anaesthesiology, 06/2007  
Publication

---

51 Gildasio S. De Oliveira, Kenyon Duncan, Paul Fitzgerald, Antoun Nader, Robert W. Gould,

Robert J. McCarthy. "Systemic Lidocaine to Improve Quality of Recovery after Laparoscopic Bariatric Surgery: A Randomized Double-Blinded Placebo-Controlled Trial", Obesity Surgery, 2013

Publication

52

B. Sundaravadivazhagan, Sekar Mohan, Balakrishnaraja Rengaraju. "Recent Developments in Microbiology, Biotechnology and Pharmaceutical Sciences - International Conference on Recent Development in Microbiology, Biotechnology and Pharmaceutical Science", CRC Press, 2025

Publication

<1 %

53

Harish Kumar, Pooja Bhosle. "To Compare the Pre-emptive Analgesic Effect of I.V. Keterolac Versus I.V. Tramadol in Pediatric Inguinal Herniotomy and Penile Surgeries", Indian Journal of Anesthesia and Analgesia, 2019

Publication

<1 %

54

Mohammad Reza Khajavi, Marzieh Navardi, Reza Shariat Moharari, Pejman Pourfakhr, Narjes Khalili, Farhad Etezadi, Farsad Imani. "Combined Ketamine-Tramadol Subcutaneous Wound Infiltration for Multimodal Postoperative Analgesia: A Double-Blinded, Randomized Controlled Trial after Renal Surgery", Anesthesiology and Pain Medicine, 2016

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On