

Comparative Assessment of Oral Health Status between Orphanage raised and Family raised Children of Lucknow City: A Cross- Sectional Study

by Jana Publication & Research

Submission date: 06-Jun-2025 03:18PM (UTC+0700)

Submission ID: 2690355736

File name: IJAR-52117.docx (122.23K)

Word count: 3954

Character count: 22591

Comparative Assessment of Oral Health Status between Orphanage raised and Family raised Children of Lucknow City: A Cross- Sectional Study

Running Title: Oral Health of Orphanage and Family- Raised Children

Abstract

Background: Despite sufficient support, lack of parental care make orphanage raised children vulnerable to various health and psychological problems. Health problems like nutritional deficiencies, compromised oral health and various psychological conditions are the common problems in these children. Considering the various problems faced by orphanage raised children the study is designed to better assess the facts and status of oral health and quality of life in these children.

Aims & Objective: 1. To assess and compare the oral health status in orphanage raised and family raised children of Lucknow City. 2.To compare BMI and correlate with oral health among these children.

Methodology: Eight Hundred Seventy children (342 Family raised and 528 Orphanage raised) were included according to inclusion and exclusion criteria in the study, after obtaining written consent and institutional ethical approval. Comprehensive oral examination was done according to WHO criteria.

Results: Results showed statistically significant poor oral hygiene, higher DMFT score and lower BMI in orphanages raised children. There was no correlation between BMI and DMFT /OHI scores in both the groups.

Conclusion: In conclusion, our study highlights the compromised oral health status of orphanage children compared to family-raised children.

Key Words: Orphan, Oral Health, Family raised children, BMI

Introduction

Based on definition, orphans are children under the age of 18 years old who have lost their mother or father or both. ^[1] Many children lose their families due to issues such as financial poverty, cultural problems, mental health, or other factors and encounter difficult situations as family deprivation and poor living conditions. ^[2,3] Though the children get protection, food, and shelter in orphanages, they meet with problematic hygiene, nutritional deficiencies, compromised oral health and psychological conditions due to lack of parental support and care. ^[4,5,6] Dental caries, periodontal and gingival problems are the most common oral conditions. ^[4,7]

Considering this present ⁶ study was conducted to assess and compare the oral health status and BMI (Body mass index) in family raised children and children living in orphanages of Lucknow city.

Materials And Methods

The cross-sectional study was conducted on 870 participants aged 3-16 years, who were selected in the camps that were organised at both orphanages and primary schools. The study was conducted from July 2023 to July 2024. ¹ Written informed consent was taken from the parents, school principal and orphanages authorities prior to start observations or examinations. Institutional ethical approval (BBDCODS/IEC/09/2022 dated 16/9/22) was obtained prior to start the study.

¹ The sample size for the present study was calculated on the basis of proportion of unfavorable oral health among orphan and non-orphan groups using the formula:

$$n = d \left(\frac{z_{\alpha} + z_{\beta}}{\ln(1-e)} \right)^2 \left[\frac{1-p_1}{p_1} + \frac{1-p_2}{p_2} \right]$$

Where $p_1 = 0.88$ (88%) proportion of unfavorable oral health among orphan group

$p_2 = 0.33$ (33%) proportion of unfavorable oral health among non-orphan group ^[8]

$e = 0.6(p_1 - p_2)$.

The proportion ¹⁴ difference considered to be clinically significant

$d = 2.0$, the design effect

Type I error $\alpha=5\%$, for the significance level of 95%.

⁴ Type II error $\beta=10\%$, for detecting the results with 90% power of study,

The minimum sample size required $n = 340$ each group

The study sample consisted of two groups based on the upbringing environment. Both orphanage and family raised ¹ children were similar in gender distribution and mean age. The exclusion criteria were age less than three years and more than 16 years, lack of cooperation in the examination, systemic disorders, mental or other behavioural disorders. According to inclusion criteria all children of both sex who were in good health, Children between ²¹ the age of 3-16 years and Children ² who were cooperative were included in the study. Children who were not physically participating, had enamel/dental defects other than dental caries, and were suspected of having a chronic medical condition (constant use of sugar-containing drugs is mandatory) were excluded.

¹ Intraoral examinations were performed by a trained examiner (postgraduate student) using a mouth mirror and personal protective barriers (gloves and masks). The data were recorded on a standard format. ³⁶ Decayed, Missing, Filled Teeth/Surfaces (DMFT/DMFS) score

¹ of teeth was recorded as per World Health Organization (WHO) criteria including the indices ¹ of for the permanent teeth and the decayed, missing and filled teeth/surfaces (dmft/dmfs) for the primary teeth. ¹ T-test was used to analyze DMFT/DMFS, dmft/dmfs and OHI-S scores.

Other soft tissue/hard tissue findings of the oral cavity were recorded according to the pre-set format. ² As per the World Health Organization (WHO), the anthropometric measurement ² method of BMI calculation was used for body weight and height measurements using a digital ² scale of 150 kg and 200 cm of tape. The height of the body is measured with subjects wearing ² no shoes and their heads contacting the ruler with their line of sight horizontally aligned. ² As per WHO, the BMI scores are as follows: underweight (18.5), average weight (18.5-24.99), ² overweight (>25), and obese (>30). ¹⁹ The BMI and oral health status assessments were analysed according to aims and objectives of the study. Results were calculated and analysed with suitable statistical tests.

Results were calculated using chi square test for group distributions and associations, ANOVA for correlation between BMI, DMFT and OHI scores and MANN Whitney test to compare BMI in the groups.

Results

1.Sample Distribution: A total 870 children were included in the study. 528 individuals (60.7%) were raised in an orphanage, while 342 individuals (39.3%) were family-raised. (Table 1, Figure 1)

2. Age distribution of the Study Groups

Among children aged three 3-9 years, 372 (70.5%) were from orphanages, while only 138 (40.4%) were family-raised. In contrast, in the 10–16 years age group, 156 (29.5%) were from orphanages compared to 204 (59.6%) from family settings. The chi-square analysis indicated

²⁶ a statistically significant difference in age distribution between the two groups ($\chi^2 = 77.54$, ⁴¹ $p < 0.001$) (Table 2, Figure 2)

3. Sample distribution according to Sex

The sex distribution between orphanage and family-raised groups showed a significant difference. Among those in orphanages, 248 (47.0%) were male and 280 (53.0%) were female. In contrast, within the family-raised group, 189 (55.3%) were male and 153 (44.7%) were female. The ²⁰ chi-square test revealed a statistically significant association between sex and upbringing environment ($\chi^2 = 5.71$, $p = 0.017$), indicating that males were more prevalent in the family-raised group, while females were more predominant in the orphanage group. (Table 3, Figure 3)

4. Oral Hygiene Status

In the orphanage group, 236 (44.7%) individuals had good oral hygiene, 180 (34.1%) had fair hygiene, and 112 (21.2%) had poor hygiene. Conversely, in the family-raised group, 156 (45.6%) had good oral hygiene, 163 (47.7%) had fair hygiene, and only 23 (6.7%) had poor hygiene. ²⁷ The chi-square test showed a highly significant association ($\chi^2 = 37.81$, $p < 0.001$).

(Table 4, Figure 4)

5. Intergroup Comparison of Orthodontic Problems

The occurrence of orthodontic problems showed no significant difference between orphanage and family-raised groups. Among the orphanage group, 50 (9.5%) individuals had orthodontic problems, while 478 (90.5%) did not. In the family-raised group, 28 (8.2%) had orthodontic issues, and 314 (91.8%) did not. The chi-square test result ($\chi^2 = 0.42$, $p = 0.518$) indicates ¹² that

there was no statistically significant association between living environment and the presence of orthodontic problems. (Table 5, Figure 5)

6. Intergroup Comparison of Dental Carries

The orphanage group exhibited higher DMFT score for both primary and permanent teeth with mean score (2.3 ± 1.5 and 2.7) than the family-raised group (2.0 and 1.5 ± 0.9), with a z-value of 5.924 and a p-value of <0.001 . (Table 6, Figure 6)

7. Intergroup Comparison of BMI

The comparison of Body Mass Index (BMI) between orphanage and family-raised children, analysed using the Mann-Whitney test, showed a significant difference. The mean BMI for children in the orphanage group was 15.67 ± 1.71 , whereas the family-raised group had a higher mean BMI of 16.24 ± 2.71 . The test yielded a z-value of 3.783 with a p-value of 0.001, indicating a statistically significant difference. (Table 7, Figure 7)

8. Correlation of DMFT Score with BMI

The correlation analysis between Body Mass Index (BMI) and dental caries, measured by DMFT scores (Decayed, Missing, and Filled Teeth), revealed weak and non-significant associations in both orphanage and family-raised groups. For permanent teeth, the overall correlation showed an r-value of -0.055 with a p-value of 0.181, indicating no significant relationship. Similarly, in the orphanage group, the r-value was -0.007 ($p = 0.910$), and in the family-raised group, it was -0.069 ($p = 0.230$), both showing no significant correlation. For primary teeth, the overall correlation had an r-value of -0.040 ($p = 0.391$), with the orphanage

group showing an almost negligible r-value of 0.002 ($p = 0.977$), and the family-raised group also exhibiting no significant relationship ($r = -0.002$, $p = 0.968$). (Table 8)

9. Correlation of BMI with OHIS

The analysis of the relationship between oral hygiene status and Body Mass Index (BMI) using ANOVA revealed no significant differences across the groups. Overall, individuals with good oral hygiene had a mean BMI of 15.81 ± 2.22 , those with fair hygiene had a mean BMI of 16.03 ± 2.16 , and those with poor hygiene had a mean BMI of 15.81 ± 2.07 . The overall F-value was 1.03 with a p-value of 0.359, indicating no statistically significant difference.

In the orphanage group, the BMI for individuals with good, fair, and poor oral hygiene was 15.59 ± 1.66 , 15.78 ± 1.64 , and 15.67 ± 1.92 respectively, with an F-value of 0.59 and a p-value of 0.556, showing no significant variation.

Similarly, in the family-raised group, those with good oral hygiene had a BMI of 16.14 ± 2.85 , fair hygiene had 16.30 ± 2.60 , and poor hygiene had 16.50 ± 2.65 . The F-value of 0.26 and p-value of 0.775 also indicated no significant differences. (Table 9, Figure8)

Discussion

Oral health is an essential component of general health and well-being, particularly in children, as it influences their nutrition, growth, self-esteem, and quality of life. The World Health Organization (WHO) defines oral health as a state of being free from chronic mouth and facial pain, oral and throat cancer, oral infection and sores, periodontal disease, tooth decay, tooth loss, and other diseases and disorders that limit an individual's capacity in biting, chewing, smiling, speaking, and psychosocial well-being (WHO, 2012).^[10] In children, oral

diseases such as dental caries, gingivitis, and malocclusion are common and, if left untreated, can have detrimental effects on physical and psychological development.

Multiple factors⁴⁶ contribute to the oral health status of children, including genetic predisposition, socioeconomic status, dietary habits, oral hygiene practices, access to dental services, and parental or caregiver involvement (Petersen & Kwan, 2010).^[11] In the absence of a stable family environment, orphaned children are often deprived of consistent supervision and care regarding their oral health practices, exacerbating their risk of developing dental problems. Children in orphanages often face many barriers to oral healthcare, including financial constraints of the institution, absence of routine dental check-ups, and limited availability of dental care facilities, particularly in rural or underdeveloped areas. Additionally, orphanage staff may lack adequate training in paediatric oral health management, resulting in inconsistent or insufficient supervision³⁷ of oral hygiene practices among the children.

Present study was attempted to identify the Oral health status in orphanage children as compare to family raised children. Our study corroborates earlier research by demonstrating that children living in orphanages²⁵ have a higher prevalence of dental caries, poor oral hygiene⁴⁷ status, and lower body mass index (BMI) compared to family-raised children.

Total 872 children were included in the study. This distribution of sample subjects highlighted a higher representation of participants from orphanage settings compared to those raised in a family environment, the distribution was justified statistically as the calculated sample size was 340 subjects(minimum) in each group. The age distribution of participants revealed significant differences between the orphanage and family-raised groups. There were more elder age group (7-16) children in family raised group whereas the orphanage children group consisted more younger group children.

We found a highly significant association between OHI score and both the study groups indicating that poor oral hygiene was notably more prevalent among those in orphanages compared to family-raised children. Vinay Suresan et al 2021¹ also found poor oral hygiene status among orphans in their study.^[12]

The DMFT score comparison suggested that orphanage children⁵ have a significantly higher prevalence of dental caries in both permanent and primary teeth compared to those raised in family environments. Similarly Nagwa Mohamed Ali Khattab⁶ and Mennat Allah Ashraf Abd-Elsabour (2023) also reported significantly high caries index in the governmental orphanage in their study.^[13]

A study by Rao et al. (2016)¹⁷ similarly observed a higher prevalence of dental caries and gingival diseases in institutionalized children than in children living with their families⁴⁵ in India. The lack of individualized attention in maintaining oral hygiene routines, irregular access to dental care services, and inadequate oral health education programs within orphanage settings contribute substantially to this disparity.^[14]

In both the groups there was no significant difference⁸ in orthodontic problems. Few studies had suggested high prevalence of oral habits and malocclusion in orphanage children.⁴⁴

Our study showed that orphanage children¹ had significantly lower BMI values compared to family raised group. This finding was in accordance with studies by Naveen et al (2015) and Dosumu et al. (2013),^[15, 16] which reported a significant association between low BMI and poor oral health status in institutionalized children. In present study there was no significant correlation found between BMI and DMFT and OHI scores in either group. Contradictory Kunal Jha (2022) et al found that there was a significant correlation between the OHI and DC in their study.^[17] Few studies showed mixed correlation between BMI and oral health status.^[18, 19]

The findings of our study clearly showed that children residing in orphanages exhibit significantly compromised oral health status as compared to their counterparts raised in family environments. This disparity may ⁷ be attributed to multiple interrelated factors, including lack of oral health awareness, nutritional deficiencies, and insufficient psychological and emotional support that typically characterizes a familial setting. Psychological and emotional well-being ³⁵ has been identified as a significant determinant of oral health behaviour and outcomes in children. Family environments typically provide emotional support, affection, and positive reinforcement that encourage the adoption of healthy habits, including proper oral hygiene practices (Locker, 2000). ^[20] Conversely, orphanage children often experience emotional neglect, stress, and low self-esteem, which may manifest as neglect of personal hygiene, including oral care.

Lack of details regarding comprehensive diet history, analysis of psychological component ⁵ and oral health knowledge among children and their parents/ caretakers can be considered as drawback of our study, yet the study reflects ⁵ light on the demand of orphan children for regular professional dental care.

Oral health awareness is a key modifiable determinant of oral health status. In family settings, parents and guardians ¹¹ play a pivotal role in educating children about proper oral hygiene practices, dietary habits, and the importance of regular dental visits. In contrast, children in orphanages often lack access to consistent oral health education programs. For instance, a study by Petersen et al. (2004) in Kenya demonstrated that regular oral health education significantly reduced the ⁷ prevalence of dental caries and improved oral hygiene among school children. ^[21] Implementing similar programs in orphanage settings could potentially mitigate the disparities in oral health outcomes observed in our study.

Timely access to professional dental care is essential ⁴⁰ for the prevention, early detection, and management of oral diseases in children. Family-raised children typically benefit from parental supervision and facilitation of dental visits. In contrast, orphanage children often lack such facilitation due to financial constraints, logistical difficulties, and the prioritization of other health concerns over dental care within institutional settings. Studies by Ramazani (2014) and Kumar et al. (2016) have reported low utilization rates of dental services among institutionalized children, contributing to the accumulation of untreated dental caries and other oral health problems. ^[22,23] Integrating dental services into routine healthcare programs for orphanages and establishing partnerships with dental schools and community health centres could enhance access to preventive and curative oral healthcare for these children.

The findings of our study emphasize the urgent need for policy interventions and programmatic strategies aimed at improving the oral health status of orphanage children. Key recommendations may include: Implementation of Institution-based Regular Oral Health Screening Programs and Nutritional Support Programs, and training of Caregivers and Orphanage Staff.

Conclusion

³² In conclusion, our study highlights the compromised oral health status of orphanage children compared to family-raised children, attributed primarily to lack of awareness, nutritional deficiencies, and the absence of familial psychological support. Addressing these challenges requires a multifaceted approach encompassing government policies and recommendations for oral health education, improved access to dental care, nutritional support, and psychosocial interventions. Future research should ⁴³ explore the long-term impact of targeted oral health interventions in institutional/orphanages settings and develop evidence-based models for promoting oral health equity among vulnerable child populations.

References

1. UNICEF. The State of the World's Children 2021: On My Mind – Promoting, protecting and caring for children's mental health. New York: United Nations Children's Fund (UNICEF); 2021.
2. Nair M, Brennan E, Bhattacharya S. A comparison of oral health status between institutionalized and non-institutionalized children: A systematic review. *Children*. 2020;7(12):248.
3. Sujatha R, Ravi GR. Oral health status of orphan children in Chennai. *J Indian Soc Pedod Prev Dent*. 2014;32(4):274-9.
4. Kumar S, Goyal A, Tadakamadla J, Tibdewal H, Duraiswamy P, Kulkarni S. Oral health status and practices of 6- to 12-year-old children in orphanages in Jaipur, India. *Int J Paediatr Dent*. 2010;20(3):167–73.
5. Patil S, Shigli AL. Oral health in children residing in orphanages in India: A review. *J Clin Pediatr Dent*. 2012;36(4):385-90.
6. Indian Council of Medical Research (ICMR). Nutrient Requirements and Recommended Dietary Allowances for Indians. New Delhi: ICMR; 2018.

7. World Health Organization (WHO). Oral health. Geneva: WHO; 2020. Available from: <https://www.who.int/news-room/fact-sheets/detail/oral-health>
8. Meshki R, Basir L, Motaghi S, Kazempour M. Oral health status among orphan and non-orphan children in Mashhad: a case-control study. *J Med Life*. 2022 Sep;15(9):1198-1201
9. World Health Organization. BMI classification. Geneva: WHO; 2020. Available from: <https://www.who.int/data/gho/indicator-metadata-registry/imr-details/26>
10. World Health Organization. Oral health. Geneva: WHO; 2012. Available from: <https://www.who.int/news-room/fact-sheets/detail/oral-health>
11. Petersen PE, Kwan S. The burden of oral disease: challenges to improving oral health in the 21st century. *Bull World Health Organ*. 2005;83(1):3–5.
12. Suresan V, Muthu MS, Mohan A, Rathna PV, Kirubakaran R, Selvaraju G. Oral health status of children in orphanages in Chennai, India. *J Indian Soc Pedod Prev Dent*. 2021;39(2):123–7
13. Khattab NMA, Abd-Elsabour MAA. Oral health status and treatment needs among institutionalized orphan children in Cairo, Egypt: a cross-sectional study. *F1000Research*. 2023;12:95.
14. Rao A, Sequeira SP, Peter S. Prevalence of dental caries and gingivitis among 6–13-year-old children in orphanages in Mangalore City. *J Indian Soc Pedod Prev Dent*. 2016;34(2):133–7.
15. Naveen N, Reddy CVK, Ramesh T, Kiranmayi M. Association between body mass index and dental caries among institutionalized orphan children. *J Indian Soc Pedod Prev Dent*. 2015;33(4):292–6.

16. Dosumu EB, Ayanbadejo PO, Nwhator SO, Umezudike KA, Oginni AO. Oral health status of orphanage children in Lagos State, Nigeria. *Niger Postgrad Med J.* 2013;20(3):188–93.
17. Jha K, Bhatia HP, Thakur S, Jha M. Correlation between body mass index and oral hygiene index among school children in Patna, India. *Int J Contemp Med Res.* 2022;9(2):B1–B4.
18. Alm A, Wendt LK, Koch G, Birkhed D. Body adiposity status in teenagers and snacking habits in early childhood in relation to approximal caries at 15 years of age. *Int J Paediatr Dent.* 2008;18(3):189–96.
19. Willershausen B, Haas G, Krummenauer F, Hohenfellner K. Relationship between high weight and caries frequency in German school children. *Eur J Med Res.* 2004;9(8):400–4.
20. Locker D. Self-esteem and socioeconomic disparities in self-perceived oral health. *J Public Health Dent.* 2000;60(2):78–84.
21. Petersen PE, Hobo B, Yayas A, Kikwilu E. Oral health status and oral health behavior of school children in Kenya, Tanzania, and Uganda. *Bull World Health Organ.* 2004;82(9):637–44.
22. Ramazani N. Child dental neglect: a short review. *Int J High Risk Behav Addict.* 2014;3(4):e21861.
23. Kumar S, Goyal A, Tadakamadla J, Tibdewal H, Duraiswamy P, Kulkarni S. Oral health status and practices of 6- to 12-year-old children in orphanages in Jaipur, India. *Int J Paediatr Dent.* 2016;26(5):373–9.

Tables

Group	No.	%
Group A Orphanage	528	60.7%
Group B Family raised	342	39.3%

42

Table 1: Distribution of Cases according to Group

Age	Orphanage		Family raised		Significance	
	No.	%	No.	%	chi sq	p-value
3 - 9 years	372	70.5%	138	40.4%	77.54	<0.001

10 - 16 years	156	29.5%	204	59.6%		
---------------	-----	-------	-----	-------	--	--

Table 2: Age Distribution of the Groups

Sex	Orphanage		Family raised		Significance	
	No.	%	No.	%	chi sq	p-value
Male	248	47.0%	189	55.3%	5.71	0.017
	280	53.0%	153	44.7%		

Table 3: Groupwise Distribution of Sex

Oral Hygiene Status	Orphanage		Family raised		Significance	
	No.	%	No.	%	chi sq	p-value
Good	236	44.7%	156	45.6%	37.81	<0.001

Fair	180	34.1%	163	47.7%		
Poor	112	21.2%	23	6.7%		

Table 4: Intergroup Comparison of OHI

Orthodontic problem	Orphanage		Family raised		Significance	
	No.	%	No.	%	chi sq	p-value
Present	50	9.5%	28	8.2%	0.42	0.518
Not Present	478	90.5%	314	91.8%		

Table 5: Intergroup Comparison of Orthodontic Problem

Dental Caries	Orphanage		Family raised		Mann Whitney test	
	Mean	SD	Mean	SD	z-value	p-value
DMFT (Permanent)	2.7	2.1	2.0	1.2	4.993	<0.001
DMFT (Primary)	2.3	1.5	1.5	.9	5.924	<0.001

Table 6: Intergroup Comparison of Dental Caries

Dental Caries	BMI		Mann Whitney test	
	Mean	SD	z-value	p-value
Orphanage	15.67	1.71	3.783	0.001
Family raised	16.24	2.71		

Table 7: Intergroup Comparison of BMI

Correlation with BMI	Overall		Orphanage		Family Raised		
	r-value	p-value	r-value	p-value	r-value	p-value	N
DMFT (PERMANENT)	-.055	.181	-.007	.910	-.069	.230	308
DMFT (primary)	-.040	.391	.002	.977	-.002	.968	303

Table 8: Correlation of DMFT Score with BMI

Group	Oral Hygiene Status	BMI		ANOVA	
		Mean	SD	F-value	p-value
Overall	Good	15.81	2.22	1.03	0.359
	Fair	16.03	2.16		
	Poor	15.81	2.07		
Orphanage	Good	15.59	1.66	0.59	0.556
	Fair	15.78	1.64		
	Poor	15.67	1.92		
	Good	16.14	2.85	0.26	0.775
	Fair	16.30	2.60		
	Poor	16.50	2.65		

Table 9: Correlation of BMI with OHIS

Figures




Figure 1: Distribution of Cases according to Group




Figure 2: Age Distribution of the Groups





Figure 3: Groupwise Distribution of Sex



8
Figure 4: Intergroup Comparison of OHI




Figure 5: Intergroup Comparison of Orthodontic Problem




Figure 6: Intergroup Comparison of Dental Caries




Figure 7: Intergroup Comparison of BMI




Figure 8: Correlation of BMI with OHIS

List of Abbreviations:

29	Abbreviation	Definition
1.	BMI	Body Mass Index
2.	DMFT	Decayed, Missing, Filled Teeth
3.	WHO	World Health Organisation
4.	BBDCODS	Babu Banarasi Das College of dental sciences
5.	DMFS	Decayed, Missing, Filled Surfaces
6.	IEC	Institutional Ethical Committee
30	OHI	Oral hygiene index
8.	OHI-S	Oral hygiene index-Simplified
9.	ANOVA	Analysis of Variance

Acknowledgements: We acknowledge the orphanage and primary schools of Lucknow city where the oral health camps were organized.

34
Source(s) of support: Nil

Conflicting Interest : None

Contribution Details:

10	Role (Concepts, Design, Definition of intellectual content, investigation, manuscript writing, etc.)	Contributor 1	Contributor 2	Contributor 3	Contributor 4	Contributor 5	Contributor 6
	1. Concepts & Design	Dr. Vandana Singh					
	2. Definition of intellectual content		Dr. Subash Singh				
	3. Manuscript writing	Dr. Vandana Singh		Dr. Sonalika Srivastava			
	4. Investigation				Dr. Ayushi Bhardwaj		

5. Data Collection			Dr. Sonalika Srivastava		Dr. Rajlakshmi Mitra	Dr. Karina Hyanki
--------------------	--	--	----------------------------	--	----------------------------	----------------------

24

Ethical policy and Institutional Review board statement: Institutional ethical approval

(BBDCODS/IEC/09/2022 dated 16/9/22) was obtained prior to start the study.

Patient declaration of consent statement: Written informed consent was taken from the parents, school principal and orphanages authorities prior to start observations or examinations.

Data Availability statement: N/A

Comparative Assessment of Oral Health Status between Orphanage raised and Family raised Children of Lucknow City: A Cross- Sectional Study

ORIGINALITY REPORT

26%

SIMILARITY INDEX

22%

INTERNET SOURCES

16%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1

www.ncbi.nlm.nih.gov

Internet Source

6%

2

journals.lww.com

Internet Source

3%

3

ir.mu.ac.ke:8080

Internet Source

2%

4

www.iosrjournals.org

Internet Source

1%

5

go.gale.com

Internet Source

1%

6

www.researchgate.net

Internet Source

1%

7

bmcoralhealth.biomedcentral.com

Internet Source

1%

8

Arunima Saikia, Raju Prasad Tayung, Maumita Saha. "ATTENUATION OF HAEMODYNAMIC RESPONSE TO LARYNGOSCOPY & INTUBATION- A COMPARATIVE STUDY BETWEEN ORAL GABAPENTIN AND ORAL CLONIDINE", Journal of Evolution of Medical and Dental Sciences, 2019

Publication

1%

9

Salwa Salsabiela, Metana Puspitasari, Safari Wahyu Jatmiko. "Blood Parameters and Length of Hospitalization in Pediatric Dengue Virus Infection Patients", Proceeding ISETH

1%

(International Summit on Science,
Technology, and Humanity), 2025

Publication

10	www.jispcd.org Internet Source	1 %
11	www.prospersmilestudio.com Internet Source	1 %
12	www.scielo.br Internet Source	<1 %
13	Miseon Yoon. "A cinematographic analysis of English-dubbed Korean films", Humanities and Social Sciences Communications, 2025 Publication	<1 %
14	Submitted to Sharda University Student Paper	<1 %
15	Arsany Yassa, Olivia Ayad, Aman Patel, Ved Vengsarkar, Mina Mikhael. "Artificial Intelligence in Patient Care: A Study of Artificial Intelligence's Responses to Blepharoplasty Concerns", Research Square Platform LLC, 2024 Publication	<1 %
16	Submitted to University of College Cork Student Paper	<1 %
17	Submitted to University of Mosul Student Paper	<1 %
18	www.coursehero.com Internet Source	<1 %
19	Shah Sumaya Jan. "Anatomic differences in patellar dimensions: A comparative study of left and right sides", Indian Journal of Clinical Anatomy and Physiology, 2023 Publication	<1 %
20	ir.lib.uwo.ca Internet Source	<1 %

21	Delfin Lovelina Francis, Kumara Raja, Srudhy Raghavan, Chithraa R Chandran. "Relationship between obesity and dental caries experience in patients attending Tagore Dental College, Chennai, India: A cross-sectional survey", Journal of Global Oral Health, 2022 Publication	<1 %
22	Submitted to Rutgers University, New Brunswick Student Paper	<1 %
23	Shalli Rani, Ayush Dogra, Ashu Taneja. "Smart Computing and Communication for Sustainable Convergence", CRC Press, 2025 Publication	<1 %
24	herniasurgeryjournal.org Internet Source	<1 %
25	ijcpd.com Internet Source	<1 %
26	www.bioinformation.net Internet Source	<1 %
27	www.health2joy.com Internet Source	<1 %
28	GISELLE D'MELLO. "Childhood obesity and dental caries among paediatric dental clinic attenders : Childhood obesity and dental caries", International Journal of Paediatric Dentistry, 05/2011 Publication	<1 %
29	ijpm.mui.ac.ir Internet Source	<1 %
30	repositorio.ufpe.br Internet Source	<1 %

31 "Abstracts of The 51st Annual Conference of Research Society for the Study of Diabetes in India", International Journal of Diabetes in Developing Countries, 2023
Publication

32 André Hajek, Hans-Helmut König. "Oral health-related quality of life, probable depression and probable anxiety: evidence from a representative survey in Germany", BMC Oral Health, 2022
Publication

33 Ezekiel Ijaopo, Ruth Ijaopo. "A Review of Oral Health in Older Adults: Key to Improving Nutrition and Quality of Life", OBM Geriatrics, 2018
Publication

34 Saruchi Saxena, Neerja Mahajan, Vineet Vinay. "Reliability and validity of a Hindi version of the Oral Health Impact Profile (OHIP-EDENT-H) for edentulous subjects: A Cross-Sectional study", Journal of Oral Biology and Craniofacial Research, 2024
Publication

35 digital.library.adelaide.edu.au
Internet Source

36 journals.sagepub.com
Internet Source

37 mansapublishers.com
Internet Source

38 pmc.ncbi.nlm.nih.gov
Internet Source

39 repository-tnmgrmu.ac.in
Internet Source

40 revmedicaelectronica.sld.cu
Internet Source

41	wajradiology.org Internet Source	<1 %
42	www.longdom.org Internet Source	<1 %
43	Elisenda Tarrats-Pons, Marc Mussons-Torras, Yirsa Jiménez-Pérez. "Efficacy of a Positive Psychology Intervention in Enhancing Optimism and Reducing Depression Among University Students: A Quasi-Experimental Study", Behavioral Sciences, 2025 Publication	<1 %
44	Nagwa Mohamed Ali Khattab, Mennat Allah Ashraf Abd-Elsabour, Ola Moustafa Omar. "Parent-perceived oral habits among a group of school children: prevalence and predictors", BDJ Open, 2024 Publication	<1 %
45	Ayat Gamal-AbdelNaser, Mennat Allah Ashraf A.Elsabour, Nagwa Mohamed Ali Khattab. "Caries in orphan children: prevalence and determinants—a systematic review and meta-analysis", BMC Oral Health, 2024 Publication	<1 %
46	Duangporn Duangthip, Sherry Shiqian Gao, Edward Chin Man Lo, Chun Hung Chu. "Early childhood caries among 5- to 6-year-old children in Southeast Asia", International Dental Journal, 2017 Publication	<1 %
47	eprints.ugd.edu.mk Internet Source	<1 %