

Clinical utility of Shock Index in the Early Detection of Adverse Outcomes In Postpartum Hemorrhage.

by Jana Publication & Research

Submission date: 11-Jun-2025 01:55PM (UTC+0700)

Submission ID: 2690333996

File name: IJAR-52168.docx (170.8K)

Word count: 5069

Character count: 28473

Clinical utility of Shock Index in the Early Detection of Adverse Outcomes In Postpartum Hemorrhage.

6 ABSTRACT

Background: Postpartum hemorrhage is a major cause of maternal morbidity and mortality. Shock Index (SI), defined as heart rate (HR) divided by systolic blood pressure (SBP), is emerging as a valuable early predictor of hemodynamic instability.

Objective: This study evaluates the role of SI in predicting adverse maternal outcomes, including ICU admission, need for transfusion, and surgical intervention.

Materials and Methods: A prospective cohort study was conducted from March 2023 to March 2024 at Dr.B.R.A.M Hospital, Raipur, involving 65 patients diagnosed with PPH. SI was measured at 15-minute intervals for 1 hour post-delivery. The primary outcomes included ICU admission, blood transfusion, and surgical intervention.

Results: SI > 1.1 was significantly associated with increased ICU admissions (40%), need for massive transfusion (68%), and surgical interventions (55%). ROC curve analysis demonstrated an area under the curve (AUC) of 0.80, indicating strong predictive value of SI.

Conclusion: SI is an effective tool for early detection of hemodynamic instability in PPH and should be integrated into obstetric early warning systems for better maternal outcomes.

Keywords: Shock Index, Postpartum Hemorrhage, Maternal Mortality, Hemodynamic Instability, Obstetric Emergencies.

INTRODUCTION

The maternal mortality ratio (MMR) in India has declined to 97 per 100,000 live births for the period 2018–2020, down from 130 in 2014–2016, according to the SRS report.[1]

However, this remains above the Sustainable Development Goal (SDG) 3.1 target of reducing global MMR to below 70 per 100,000 live births by 2030.[2]

Postpartum hemorrhage (PPH), defined as blood loss >500 mL after vaginal delivery or >1000 mL after cesarean section, remains a leading cause of maternal mortality and morbidity globally.[3,4] The World Health Organization reports that PPH affects approximately 14 million women annually, leading to around 70,000 deaths worldwide.[5]

The causes of PPH are classified into the “Four Ts”: tone (uterine atony), trauma, tissue (retained placenta), and thrombin (coagulopathies), with uterine atony being the most common etiology.[6]

Prompt recognition and management of PPH are crucial. However, conventional methods of estimating blood loss, such as visual assessment, are often inaccurate, leading to diagnostic delays and suboptimal intervention.[7,8] In response, there has been increasing interest in objective tools like the Shock Index (SI), calculated as the ratio of heart rate to systolic blood pressure, to assess hemodynamic instability.[9]

In healthy adults, normal SI ranges from 0.5 to 0.7, while in pregnant women, due to physiological changes, it ranges from 0.7 to 0.9.[10,11] Elevated SI values have been shown to correlate with greater blood loss, hemodynamic compromise, and increased risk of adverse maternal outcomes in PPH.[12,13] Unlike individual vital signs that may remain deceptively normal, SI offers a more sensitive marker of early decompensation.[9,14]

This study aims to evaluate the role of shock index in assessing adverse maternal outcomes in postpartum hemorrhage and to determine its clinical utility in comparison to traditional assessment methods. By identifying SI thresholds predictive of poor outcomes, this research seeks to support more timely and effective interventions in PPH, thereby contributing to improved maternal health outcomes.

OBJECTIVE

Primary objective - To study the role of shock index in assessing the adverse maternal outcomes in postpartum hemorrhage.

Secondary objective – To correlate shock index with visual estimation of blood loss in women with postpartum hemorrhage.

MATERIALS AND METHODOLOGY

Study Design & Setting

A prospective cohort study was conducted in the Department of Obstetrics and Gynaecology at Dr. B.R.A.M Hospital, Raipur, from March 2023- March 2024. The study included pregnant women diagnosed with postpartum hemorrhage (PPH), and their hemodynamic parameters were continuously monitored to evaluate the predictive utility of Shock Index in determining adverse maternal outcomes.

Inclusion criteria

Women who delivered after 28 weeks of gestation.

Patients diagnosed with PPH based on visual blood loss ≥ 500 mL in vaginal delivery and ≥ 1000 mL in LSCS.

Patients with normal baseline hemodynamic parameters before labor.

Exclusion criteria

Antepartum hemorrhage

Pre-existing maternal heart disease or severe anaemia (<7 gm/dL).

Pregnancy induced hypertension, preeclampsia, eclampsia

Patients with pre-existing coagulopathies.

Methodology

Immediately after delivery blood loss estimation was done using blood collected in drapes, fixed size mops of 45*45cm, swabs of 10*10cm, perianal pads which when fully soaked amounted to a blood loss of approximately 350ml, 60ml and 100ml respectively. Blood loss estimation in case of caesarean section was done

using fixed size mops, kidney tray and suction machine. A full kidney tray amounted to a blood loss of approximately 500ml whereas a partially filled tray amounted to a blood loss of approximately 250ml.

Baseline vitals were recorded at the time of admission. Thereafter as soon as postpartum haemorrhage was anticipated by visual estimation of blood loss study participants were subjected to BP and HR measurement every 15 minutes for 1 hour postpartum. Shock index was evaluated by dividing heart rate by systolic blood pressure. The highest SI that was recorded was selected for further analysis. Active management of third stage of labor was routinely performed.

Pre specified potential confounding factors included age, gestational age at delivery, height, weight, BMI, parity, mode of delivery, type of anaesthesia and use of oxytocin for AMTSL.

The following outcome measures were recorded: need for ICU care, need for blood and blood products transfusion, need for operative intervention, acute renal failure, surgical site infection and maternal mortality.

RESULT

The present study aimed to assess the role of Shock Index (SI) in predicting adverse maternal outcomes in postpartum hemorrhage (PPH). The average age of participants was 24.97 years, with most being unbooked cases. The mean gestational age was 38.52 weeks, and the average BMI was 24.65. Most deliveries were vaginal (58.46%), followed by LSCS (38.46%) and VBAC (3.08%). The primary cause of PPH was uterine atony (60%).

The mean shock index was 1.26 (range 1.0–1.81).

A strong positive correlation was found between SI and blood loss ($r = 0.88$), with an average loss of 902.92 ± 340 mL.

Table 1: Association of Shock Index and Blood Loss

Shock Index Range	Mean Blood Loss (mL)	Median (25th–75th percentile)	Range	p-value
0.9 to <1.2	643.07 ± 148.62	600 (550–800)	500–1050	<0.0001
1.2 to <1.5	940.34 ± 197.08	900 (850–1000)	600–1550	
1.5 to <1.7	1408.33 ± 316.89	1300 (1262.5–1450)	1100–2000	
≥1.7	1587.5 ± 184.28	1600 (1537.5–1650)	1350–1800	

Significant adverse outcomes included ICU admission (52.31%), transfusion (64.62%), operative intervention (69.23%), acute renal failure (23.08%), severe anemia (50.77%), and maternal mortality (6.15%). The mean SI associated with ICU admission was 1.4 ± 0.18 , ventilatory support 1.63 ± 0.16 , inotropic support 1.55 ± 0.17 , and mortality 1.73 ± 0.06 .

Table2: Association of Shock Index with Adverse Maternal Outcomes

Outcome	Shock Index 29 (Mean \pm SD)	Median (25th– 75th percentile)	Range	P-value
Need for ICU Admission - No	1.11 \pm 0.09	1.08 (1.045–1.165)	1–1.4	<0.0001
Need for ICU Admission - Yes	1.4 \pm 0.18	1.33 (1.282–1.5)	1.19–1.81	
Ventilatory Support - No	1.21 \pm 0.15	1.2 (1.08–1.31)	1–1.61	<0.0001
Ventilatory Support - Yes	1.63 \pm 0.16	1.69 (1.48–1.73)	1.42–1.81	
Inotropic Support - No	1.19 \pm 0.14	1.19 (1.08–1.292)	1–1.6	<0.0001
Inotropic Support - Yes	1.55 \pm 0.17	1.5 (1.42–1.72)	1.28–1.81	
Transfusion Required - No	1.07 \pm 0.06	1.06 (1.03–1.1)	1–1.26	<0.0001
Transfusion Required - Yes	1.37 \pm 0.18	1.31 (1.22–1.468)	1.1–1.81	
Operative Intervention - No	1.09 \pm 0.09	1.08 (1.04–1.105)	1–1.4	<0.0001
Operative Intervention - Yes	1.34 \pm 0.2	1.31 (1.21–1.46)	1–1.81	

ARF - No	1.2 ± 0.17	1.18 (1.08–1.288)	1–1.81	<0.0001
ARF - Yes	1.46 ± 0.19	1.42 (1.32–1.64)	1.2–1.76	
Dialysis - No	1.25 ± 0.19	1.22 (1.09–1.325)	1–1.81	0.0006
Dialysis - Yes	1.74 ± 0.03	1.74 (1.73–1.75)	1.72–1.76	
Surgical Site Infection - No	1.25 ± 0.21	1.21 (1.08–1.33)	1–1.81	0.167
Surgical Site Infection - Yes	1.4 ± 0.17	1.39 (1.295–1.498)	1.22–1.61	
Severe Anemia - No	1.11 ± 0.09	1.09 (1.048–1.185)	1–1.28	<0.0001
Severe Anemia - Yes	1.41 ± 0.18	1.34 (1.3–1.5)	1.1–1.81	
Maternal Mortality - Alive	1.23 ± 0.17	1.21 (1.08–1.32)	1–1.76	<0.0001
Maternal Mortality - Died	1.73 ± 0.06	1.72 (1.708–1.742)	1.67–1.81	

ROC analysis showed excellent predictive power: AUC values were 0.958 for ICU admission, 0.978 for transfusion, 0.896 for operative intervention, 0.864 for acute renal failure, and 0.988 for maternal mortality (all $p < 0.05$).

Table3:ROC Curve Summary Table

Outcome	AUC	Cut-off Value	Interpretation
ICU Admission	0.958	>1.18	Excellent discrimination. High accuracy for identifying patients needing ICU care.
Ventilatory Support	0.966	>1.4	Outstanding discrimination. Very high sensitivity and specificity.
Inotropic Support	0.944	>1.4	Excellent discrimination. Strong predictive value.
Transfusion Requirement	0.978	>1.18	Outstanding discrimination. Most accurate among all outcomes evaluated.
Operative Intervention	0.896	>1.12	Very good discrimination. Slightly lower but still reliable.

Acute Renal Failure (ARF)	0.864	>1.3	Good discrimination. Moderate predictive capacity.
Need for Dialysis	0.98	>1.67	Outstanding discrimination. Very high accuracy despite small sample size.
Surgical Site Infection	0.752	>1.21	Fair discrimination. Predictive power is weaker here.




Figure1:Compiled ROC curves of Shock Index for Adverse Maternal Outcomes

SI was categorized into four ranges (0.9–<1.2, 1.2–<1.5, 1.5–<1.7, ≥ 1.7), with a stepwise increase in the frequency and severity of adverse outcomes across higher SI ranges. ICU admissions, transfusion needs, operative interventions, renal complications, mortality, and hospital stay duration increased with rising SI value

Table 4: Association of Maternal Outcomes with Shock Index Ranges

Maternal Outcome	0.9 to <1.2 (n=26)	1.2 to <1.5 (n=29)	1.5 to <1.7 (n=6)	≥ 1.7 (n=4)	P-value
Need for ICU Admission	1 (3.85%)	23 (79.31%)	6 (100%)	4 (100%)	<0.0001
Ventilatory Support	0 (0%)	2 (6.90%)	2 (33.33%)	4 (100%)	<0.0001
Inotropic Support	0 (0%)	5 (17.24%)	4 (66.67%)	4 (100%)	<0.0001
Operative Intervention	8 (30.77%)	27 (93.10%)	6 (100%)	4 (100%)	<0.0001
ARF	0 (0%)	9 (31.03%)	3 (50%)	3 (75%)	<0.0001
Dialysis	0 (0%)	0 (0%)	0 (0%)	2 (50%)	0.003

Surgical Site Infection	0 (0%)	3 (10.34%)	1 (16.67%)	0 (0%)	0.223
Maternal Mortality	0 (0%)	0 (0%)	1 (16.67%)	3 (75%)	<0.0001

Table5:Correlation of Shock Index with SBP,DBP,PR and MAP

Variable	²⁴ Correlation Coefficient	P-value
Systolic Blood Pressure (SBP)	-0.404	.001
Diastolic Blood Pressure (DBP)	-0.270	0.030
Pulse Rate (PR)	0.380	.002
Mean Arterial Pressure	-0.372	.002

¹¹ A weak positive correlation was observed between SI and pulse rate ($r = 0.38$), and negative correlations were noted with ³⁵ systolic BP ($r = -0.404$), diastolic BP ($r = -0.27$), and MAP ($r = -0.372$).

DISCUSSION

The study titled “Clinical Utility of Shock Index in the Early Detection of Adverse Outcomes in Postpartum Hemorrhage” was conducted at Pt. J.N.M Medical College Raipur (C.G) from March 2023 to March 2024. ²⁰

Postpartum hemorrhage (PPH) remains a major cause of maternal mortality, especially in low-resource settings where early recognition is challenging. Shock Index (SI)—the ratio

of heart rate to systolic blood pressure—is emerging as a simple, cost-effective tool for early identification of hemodynamic instability in PPH.

This study aimed to assess the role of SI in predicting adverse maternal outcomes in PPH patients, evaluating its correlation with clinical parameters such as ICU admission, need for transfusion, operative interventions, and maternal mortality.

Descriptive statistics of Shock Index

The shock index (SI) in our study demonstrated a mean value of 1.26, ranging from 1 to 1.81, indicating variability in patients' physiological responses to shock. El Ayadi et al. (2016) reported a comparable median SI of 1.3. The primary utility of SI lies in its ability to detect hemodynamic instability earlier than conventional vital signs, identifying significant blood loss and hypovolemia before overt hypotension develops.

Association between Shock Index and Blood Loss

In non-pregnant individuals, an SI of 1.0 typically corresponds to a blood loss of 750–1500 mL.^[51] In obstetric settings, massive hemorrhage is defined as blood loss >2000 mL or >30% of blood volume.⁽¹⁶⁾ SI values differ between pregnant and non-pregnant women due to physiological changes. Literature indicates that a 10–30% blood loss in pregnant women correlates with an SI of ~1.0, while in non-pregnant women, a similar SI reflects a 15–20% loss.^(17,18) In our study, SI strongly correlated with blood loss ($r = 0.88$, $p < 0.0001$), with the highest blood loss observed in the $SI \geq 1.7$ group (1587.5 ± 184.28 mL). These findings are consistent with studies by Dziadosz et al. (2020), Sanchez et al. (2023), and Talbot et al. (2023), all of which reported a positive association between SI and hemorrhage severity. However, contrasting studies by Huang et al. (2022) and Ushida et al. (2021) found weaker correlations, suggesting that SI alone may not always reliably quantify blood loss.

Association of Shock Index with Adverse Maternal Outcomes

Operative Intervention

In our study, the mean SI among patients requiring operative intervention was significantly elevated at 1.34 ± 0.2 , compared to 1.09 ± 0.09 in those managed medically. Among surgical procedures, the mean SI progressively increased with the severity of intervention: vaginal tear repair (1.13), uterine artery ligation (1.32), uterine compression sutures (1.44), uterine artery embolisation (1.51), and hysterectomy (1.63).

An SI cut-off of >1.12 predicted the need for operative intervention with 91.11% sensitivity, 85% specificity. Similar findings were reported by Nathan et al. (2019), where SI measured after PPH diagnosis predicted emergency hysterectomy risk. SI <0.9 indicated low risk (2% underwent hysterectomy), 0.9–1.69 moderate risk (14.7%), and ≥ 1.7 high risk (28.6%). El Ayadi et al. (2016) reported an SI of 1.35 (95% CI; 60% specificity) for hysterectomy, which was lower than our study's mean SI of 1.63 ± 0.12 . Chaudhary et al. (2020) reported a slightly higher mean SI (1.58 ± 0.51) for surgical intervention.

Our findings on SI cut-off values were consistent with Agarwal et al. (2021), who reported thresholds >1 for interventions: hysterectomy >1.32 (90.91% sensitivity, 89.74% specificity), vaginal/cervical tear repair >1.32 (75%, 78.41%), internal artery ligation >1.3 (90%, 77.78%), and compression sutures >1.24 (100%, 58.76%). Studies by Sakshi Agarwal et al. (2023) and Kohn et al. (2017) also reported comparable SI values (≥ 1.1 and ≥ 1.14) with moderate sensitivity and specificity.

ICU Admission

The mean SI among patients requiring ICU admission was 1.32 ± 0.15 , significantly higher than 1.11 ± 0.09 in non-ICU patients. A cut-off value of >1.18 demonstrated 100% sensitivity, 80.65% specificity. These values are consistent with previous reports. El Ayadi et al. identified a threshold of 1.35, Chaudhary et al. noted 1.23, and Sakshi Agarwal et al. reported 1.32 as predictive of ICU admission. Our findings reaffirm that SI is a strong predictor of ICU-level care in PPH.

Nathan et al. (2019) demonstrated increasing ICU admission rates with rising SI: 25.5% (SI <0.9), 48.3% (SI 0.9–1.69), and 78.6% (SI ≥ 1.7). El Ayadi et al. (2016) reported a higher mean SI of 1.35 for ICU admission, while Chaudhary et al. (2020) found a mean SI of 1.23 ± 0.35 —both comparable to our findings. Sakshi Agarwal et al. (2023) reported a cut-off SI ≥ 1.1 (sensitivity 97.62%, specificity 93.41%), similar to Koch et al. (2019) who found SI >1 predictive of ICU admission. Agarwal et al. (2021) and El Ayadi et al. (2016) reported higher thresholds of >1.3 and ≥ 1.4 , with the latter showing 70.5% sensitivity and 74.8% specificity. Nathan et al. (2015) suggested an even lower threshold of SI ≥ 0.9 .

Inotropic and Ventilatory Support

Patients who required ventilatory support had a mean SI of 1.63 ± 0.16 . An SI cut-off of >1.4 predicted ventilatory requirement with 100% sensitivity, 89.47% specificity

Sakshi Agarwal et al. similarly found high SI (mean 1.34) in ventilated patients, with high sensitivity and specificity. Our results further confirm that elevated SI is a reliable marker for identifying patients who may require respiratory support. Inotropic Support

The mean SI in patients needing inotropic support was 1.55 ± 0.17 . A cut-off of >1.4 yielded 84.62% sensitivity, 94.23% specificity. Similar to the findings of the current study Agarwal et al. 2021 established cut-off thresholds of shock index (SI) to predict the need for ICU admission with ventilatory support and ICU admission with inotropic support. The values were >1.34 (sensitivity: 95.45%; specificity: 92.31%) and >1.446 (sensitivity: 91.67%; specificity: 93.18%) respectively.

The close alignment across studies enhances the external validity of our results.

Blood and Blood Products Transfusion

The mean SI among those receiving transfusions was 1.37 ± 0.18 , compared to 1.07 ± 0.06 in non-transfused patients. Among women with Hb <7 g/dL, the mean SI rose to 1.41 ± 0.18 . An SI cut-off of >1.18 predicted transfusion need with 92.86% sensitivity, 95.65% specificity.

Nathan et al. (2019) reported rising transfusion requirements with increasing SI: 25.5% (SI <0.9), 37.1% (SI $0.9-1.69$), and 71.4% (SI ≥ 1.7). In contrast, our study showed fewer transfusions in the SI range $0.9-1.2$, with a mean of 2.61 ± 2.31 units. El Ayadi et al. (2016) and Kwon H et al. (2024) found mean SIs of 1.35 and 1.22, respectively, for predicting massive transfusion—findings comparable to ours.

Chaudhary et al. (2020) reported a lower mean SI (1.15 ± 0.41), possibly due to inclusion of hypertensive and anaemic patients. Studies by Le Bas et al. (2013), Agarwal et al. (2023), and Kwon H et al. (2024) consistently identified SI $>1-1.1$ as predictive of massive transfusion, with high sensitivity and specificity. Koch et al. (2019) also supported SI >1 as a marker of morbidity.

Higher thresholds were reported by Guerrero-De León et al. (2018), Kohn et al. (2017), and Agarwal et al. (2021), with SI $>1.32-1.4$ predicting ≥ 4 to ≥ 10 units

of transfusion with strong diagnostic accuracy. Despite variations, most studies affirm SI >1 as a reliable predictor of transfusion need.

Maternal Morbidity

Patients who experienced significant maternal morbidity had a mean SI of 1.34 ± 0.21 , significantly higher than 1.10 ± 0.09 in those without morbidity. An SI cut-off of >1.14 predicted maternal morbidity with 88% sensitivity, 88.57% specificity.

El Ayadi et al. (2016) reported an SI of 1.57 (95% CI; 80% specificity) for predicting severe end-organ damage, aligning with our findings. Chaudhary et al. (2020) also found a comparable mean SI of 1.47 ± 0.84 in patients with MODS. Similarly, Agarwal et al. (2021) identified an SI >1.3 as predictive of MODS (sensitivity 95%, specificity 88.75%), and El Ayadi et al. (2016) noted a cut-off of ≥ 1.4 for end-organ damage with 80.6% sensitivity and 71.4% specificity.

Maternal Mortality

Among the four maternal deaths in our cohort, the mean SI was markedly elevated at 1.71 ± 0.11 . While our sample size for mortality is small, the high SI reinforces previous evidence linking very high SI to fatal outcomes.

Nathan et al. (2019) reported increasing mortality with rising SI: 0% for SI <0.9, 4.3% for SI 0.9–1.69, and 7.1% for SI ≥ 1.7 . El Ayadi et al. (2016) found a mean SI of 1.58 (95% CI; 80% specificity) for maternal mortality, aligning with our study. Chaudhary et al. (2020) and Liu et al. (2012) reported lower SI values of 1.39 ± 0.85 and 1.3 respectively. Agarwal et al. (2021) and El Ayadi et al. (2016) reported mortality cut-off SI values of >1.65 and ≥ 1.7 , both comparable with our findings.

7

Area under the curve value of Shock Index to predict adverse maternal outcome

43

In our study, the Shock Index (SI) demonstrated excellent predictive performance for multiple adverse outcomes, with AUC values of 0.958 for ICU admission, 0.978 for blood product transfusion, and 0.896 for operative intervention. Additionally, AUC values for surgical site infection (0.752), acute renal failure (0.864), and maternal mortality (0.988) were statistically significant, reinforcing SI as a robust predictor in postpartum hemorrhage.

Our results are consistent with Agarwal et al. (2021), who reported AUROC values of 0.95 and 0.98 for ICU admissions requiring inotropic and ventilatory support, respectively, 0.91 for blood product transfusion and operative intervention, and 0.99 for maternal mortality. Similarly, Lee et al. (2019) and Kwon H et al. (2024) reported AUCs of 0.815 and 0.829, respectively, for predicting massive transfusion, which align closely with our findings.

M. Chaudhary et al. (2020) found lower AUC values: ICU admission (0.8), operative intervention (0.8), maternal death (0.9), and blood transfusion (0.68). The variation may stem from their inclusion of patients with pregnancy-induced hypertension and severe anaemia— conditions that can distort SI interpretation due to altered hemodynamics.

Nathan et al. (2019) also reported lower AUCs: ICU admission (0.68), hysterectomy (0.79), transfusion ≥ 4 units (0.65), and maternal mortality (0.86). Despite the variability, a consistent trend is evident across all studies—an elevated SI is strongly associated with adverse maternal outcomes.

Clinical significance of Shock Index thresholds

The normal Shock Index (SI) range in healthy pregnant women is 0.7–0.9. An SI >0.9 has been associated with adverse outcomes including ICU admission, significant blood loss, surgical intervention, and increased morbidity and mortality (21, 33). Nathan et al. (2019)

found that $SI < 0.9$ offered reassurance, while $SI \geq 1.7$ indicated urgent need for intervention. Similarly, El-Ayadi et al. (2016) suggested $SI > 0.9$ for referral, ≥ 1.4 for urgent tertiary care, and ≥ 1.7 for high risk of maternal complications. Our study identified a slightly lower SI threshold of ≥ 1.1 to predict adverse outcomes, which may be attributed to population differences, anaemia prevalence, and study design.

Comparable findings were reported by Kohn et al. (2017), where $SI \geq 1.14$ predicted PPH with 93% specificity. Guerrero-De León et al. (2018) also found $SI \geq 1.0$ to be predictive of severe outcomes, recommending care at tertiary centers for such patients.

We stratified SI into four categories: $0.9 < 1.2$, $1.2 < 1.5$, $1.5 < 1.7$, and ≥ 1.7 . A stepwise increase in adverse outcomes was noted across these groups. For instance, transfusion was needed in 15.38% of patients with $SI 0.9 < 1.2$, versus 96.55%, 100%, and 100% in the higher ranges, respectively. Operative intervention rose from 30.77% to 100% across these SI brackets. No acute renal failure was noted below SI 1.2, but increased substantially in higher groups—up to 75% in $SI \geq 1.7$. Maternal mortality occurred exclusively in the ≥ 1.7 SI group.

Length of hospital stay also correlated with SI: the longest durations were observed in the ≥ 1.5 groups (mean 7.5 days), versus 2.77 days in $SI < 1.2$. These findings mirror those of Nathan et al. (2019), who reported rising rates of transfusion, ICU admission, and hysterectomy with increasing SI. In their study, no mortality occurred in $SI < 0.9$, while 7.1% mortality was reported for $SI \geq 1.7$. Nathan et al. (2015) also established $SI \geq 0.9$ as a reliable threshold for ICU admission and ≥ 1.7 as a critical alert trigger.

Collectively, these results reaffirm SI as a sensitive early marker of hypovolemia. Unlike conventional vital signs, which may initially remain stable due to compensatory mechanisms, SI captures the critical rise in heart rate alongside stable or declining SBP, providing a more reliable early indicator of clinical deterioration.

Overall, rising SI is a clear marker of worsening clinical status in PPH. Sustained elevation reflects ongoing hypovolemia, tissue hypoperfusion, and risk of MODS. Vital organs such as the kidneys and brain are particularly susceptible to ischemic injury in this state. Hemorrhage-induced coagulopathy further complicates management and increases the likelihood of adverse maternal outcomes.

Correlation of Shock Index with vital signs

Our study categorized patients based on SI ranges and found the following mean values:

SI 0.9–<1.2: SBP 98.48 mmHg, DBP 59 mmHg, PR 106 bpm, MAP 72.64 mmHg

SI 1.2–<1.5: SBP 92 mmHg, DBP 58 mmHg, PR 120 bpm, MAP 65.6 mmHg

SI 1.5–<1.7: SBP 84 mmHg, DBP 56.3 mmHg, PR 135 bpm, MAP 65 mmHg

SI ≥ 1.7 : SBP 78 mmHg, DBP 52 mmHg, PR 138 bpm, MAP 61 mmHg

We observed that while pulse rate and diastolic blood pressure changed in accordance with SI, other parameters such as SBP and MAP remained relatively stable until SI reached ≥ 1.5 .

These findings align with ⁴¹El Ayadi et al. (2019), who noted that at SI ≥ 1.4 , PR increased to 112 bpm and SBP dropped to 80 mmHg, and at SI 1.7, PR reached 130 bpm with SBP 70 mmHg.

CLINICAL IMPLICATIONS

Our findings support integrating SI into obstetric early warning systems (EWS) for PPH management. Key applications include:

Early Recognition & Triage:

Women with SI >1.1 should receive immediate hemodynamic monitoring and blood cross matching.

SI >1.3 should prompt early ICU transfer consideration.

Blood Transfusion Protocols:

SI >1.1 correlates strongly with the need for transfusion, suggesting SI can be used to guide blood product administration before overt hypovolemia develops.

Surgical Preparedness:

SI >1.3 may predict surgical intervention, allowing teams to mobilize resources for emergency hysterectomy or B-Lynch suture placement.

Strengths

Prospective design reduced recall bias and improved data accuracy.

Objective blood loss estimation was also incorporated along with visual estimation of blood loss.

Standardized SI measurements at multiple time points, ensuring dynamic monitoring of hemodynamic changes.

Limitations

Single center study: Findings may not be generalizable to different populations.

Small sample size (n=65): A larger multicenter study would improve statistical power.

CONCLUSION

The Shock Index is a valuable, cost-effective, and early predictor of adverse maternal outcomes in postpartum hemorrhage. By incorporating SI into standard clinical protocols, healthcare providers can improve early detection, reduce delays in intervention, and ultimately enhance maternal survival. Further multicenter studies are warranted to establish universal SI thresholds tailored to diverse populations.

ACKNOWLEDGMENT

³⁶ I would like to express my heartfelt gratitude to the Department of Obstetrics and Gynaecology, Dr. B.R.A.M Hospital, Raipur, for their support throughout the study period. I am deeply thankful to my guide and faculty members for their continuous encouragement, guidance, and valuable suggestions during the research. I also extend my sincere appreciation to all the patients who participated in this study, without whom this work would not have been possible.

REFERENCES

1. Trends in maternal mortality 2000 to 2020: estimates by WHO, UNICEF, UNFPA, World Bank Group and UNDESA/Population Division. Geneva: World Health Organization; 2023 (<https://iris.who.int/handle/10665/366225>)

2. New maternal mortality estimates: United Nations Maternal Mortality Estimation Inter-Agency Group (UN MMEIG)

3. Say L, Chou D, Gemmill A, Tunçalp Ö, Moller A-B, Daniels J et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Glob Health*. 2014;2(6):e323–33. doi: 10.1016/S2214-109X(14)70227-X.

4. Pacagnella RC, Souza JP, Durocher J et al. A systematic review of the relationship between blood loss and clinical signs. *PLoS One* 2013;8(3):e57594. DOI: 10.1371/journal.pone.00575

5. A roadmap to combat postpartum haemorrhage between 2023 and 2030. Geneva: World Health Organization; 2023 (<https://iris.who.int/handle/10665/373221>)

6. Bienstock JL, Eke AC, Hueppchen NA. Postpartum Hemorrhage. *N Engl J Med*. 2021 Apr 29;384(17):1635-1645. doi: 10.1056/NEJMra1513247. PMID: 33913640; PMCID: PMC10181876.

7. Lee SY, Kim HY, Cho GJ, Hong SC, Oh MJ, Kim HJ. Use of the shock index to predict maternal outcomes in women referred for postpartum hemorrhage. *Int J Gynaecol Obstet*. 2019 Feb;144(2):221-224. doi: 10.1002/ijgo.12714. Epub 2018 Dec 10. PMID: 30447073.

8. Schorn MN. Measurement of blood loss: review of the literature. *J Midwifery Womens Health* 2010;55:20-7

9. Allgower M, Burri C. Shock index. *Dtsch Med Wochenschr*. 1967;92 (43):1947–1950. doi:10.1055/s-0028-1106070

10.Rady MY, Smithline HA, Blake H, Nowak R, Rivers E. A comparison of the shock index and conventional vital signs to identify acute, critical illness in the emergency department. *Ann Emerg Med* 1994; **24**: 685–90.

11.Rady MY, Rivers EP, Martin GB, Smithline H, Appelton T, Nowak RM. Continuous central venous oximetry and shock index in the emergency department: use in the evaluation of clinical shock. *Am J Emerg Med* 1992; **10**: 538–41.

12.Vandromme MJ, Griffin RL, Kerby JD, McGwin G Jr, Rue LW 3rd, Weinberg JA. Identifying risk for massive transfusion in the relatively normotensive patient: utility of the prehospital shock index. *J Trauma* 2011; **70**: 384–8;discussion 8–90.


13.Cannon CM, Braxton CC, Kling-Smith M, Mahnken JD, Carlton E, Moncure M. Utility of the shock index in predicting mortality in traumatically injured patients. *J Trauma* 2009; **67**: 1426–30.

14..Krishna H, Chava M, Jasmine N, et al. Patients with postpartum hemorrhage admitted in intensive care unit: patient condition, interventions, and outcome. *J Anaesthesiol Clin Pharmacol* 2011;27(2):192–194. DOI: 10.4103/0970-9185.81826.

15.El Ayadi AM, Nathan HL, Seed PT, Butrick EA, Hezelgrave NL, Shennan AH, Miller S. Vital Sign Prediction of Adverse Maternal Outcomes in Women with Hypovolemic Shock: The Role of Shock Index. *PLoS One*. 2016 Feb 22;11(2):e0148729. doi: 10.1371/journal.pone.0148729. PMID: 26901161; PMCID: PMC4762936.

16. Mavrides E, et al. Prevention and management of postpartum hemorrhage. BJOG 2016;124:e106–e149. DOI: [10.1111/1471-0528.14178](https://doi.org/10.1111/1471-0528.14178)
17. Obstetrics Subgroup CSO, Gynecology CiMA, Association OSCSoO, Medical GC. [Guideline of prevention and treatment about postpartum haemorrhage (2014)]. Zhonghua Fu Chan Ke Za Zhi. 2014;49(9):6
18. Liu LM, Bai XJ, Li T, Zhou XY, Yang GM, Gao W, et al. Early treatment specifications for traumatic haemorrhagic shock. Journal of Traumatic Surgery. 2012; 19 (12): (881–883, 891).41–6.
19. Nathan HL, Seed PT, Hezelgrave NL, De Greeff A, Lawley E, Anthony J, Steyn W, Hall DR, Chappell LC, Shennan AH. Shock index thresholds to predict adverse outcomes in maternal hemorrhage and sepsis: A prospective cohort study. Acta Obstet Gynecol Scand. 2019 Sep;98(9):1178-1186. doi: 10.1111/aogs.13626. Epub 2019 May 14. PMID: 31001814; PMCID: PMC6767575.
20. Chaudhary M, Maitra N, Sheth T, Vaishnav P. Shock Index in the Prediction of Adverse Maternal Outcome. J Obstet Gynaecol India. 2020 Oct;70(5):355-359. doi: 10.1007/s13224-020-01355-z. Epub 2020 Jul 25. PMID: 33041552; PMCID: PMC7515997.
21. Agarwal V, Suri J, Agarwal P, et al. Shock Index as a Predictor of Maternal Outcome in Postpartum Hemorrhage. J South Asian Feder Obst Gynae 2021;13(3):131–136.
22. Agarwal, S. and Pandey, U. (2023) 'Shock index as a predictor of maternal outcome in postpartum hemorrhage: An experience from a tertiary care centre in Northern India', *International Journal of Medical Reviews and Case Reports*, (0), p. 1. doi:10.5455/ijmrcr.172-1669772235.

- 23.Jaden R. Kohn, Gary A. Dildy & Catherine S. Eppes (2017): Shock index and delta-shock index are superior to existing maternal early warning criteria to identify postpartum hemorrhage and need for intervention, *The Journal of Maternal-Fetal & Neonatal Medicine*, DOI: 10.1080/14767058.2017.1402882
- 24.Koch E, Lovett S, Nghiem T, Riggs RA, Rech MA. Shock index in the emergency department: utility and limitations. *Open Access Emerg Med*. 2019 Aug 14;11:179-199. doi: 10.2147/OAEM.S178358. PMID: 31616192; PMCID: PMC6698590.
- 25.Nathan HL, El Ayadi A, Hezelgrave N, Seed P, Butrick E, Miller S, et al. Shock index: an effective predictor of outcome in postpartum haemorrhage? *BJOG*. 2015;122:268-75. Medline:25546050 doi:10.1111/1471-0528.13206
- 26.Le Bas A, Chandrachan E, Addei A, Arulkumaran S. Use of the "obstetric shock index" as an adjunct in identifying significant blood loss in patients with massive postpartum hemorrhage. *Int J Gynaecol Obstet*. 2014 Mar;124(3):253-5. doi: 10.1016/j.ijgo.2013.08.020. Epub 2013 Dec 4. PMID: 24373705
- 27.Guerrero-De Leon MC, Escarcega-Ramos LR, Gonzalez-Dias OA. Utility of the shock index in obstetric hemorrhage as a predictive value for the transfusion requirement. *Ginecol Obstet Mex*. 2018;86(10):665-75.
- 28.Lee SY, Kim HY, Cho GJ, Hong SC, Oh MJ, Kim HJ. Use of the shock index to predict maternal outcomes in women referred for postpartum hemorrhage. *Int J Gynaecol Obstet*. 2019 Feb;144(2):221-224. doi: 10.1002/ijgo.12714. Epub 2018 Dec 10. PMID: 30447073.



Clinical utility of Shock Index in the Early Detection of Adverse Outcomes In Postpartum Hemorrhage.

ORIGINALITY REPORT

15%

SIMILARITY INDEX

11%

INTERNET SOURCES

13%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	www.researchgate.net Internet Source	2%
2	www.jsp.org.pk Internet Source	1%
3	link.springer.com Internet Source	1%
4	ijrcog.org Internet Source	1%
5	Takafumi Ushida, Tomomi Kotani, Kenji Imai, Tomoko Nakano-Kobayashi et al. "Shock Index and Postpartum Hemorrhage in Vaginal Deliveries: A Multicenter Retrospective Study", Shock, 2021 Publication	1%
6	www.ncbi.nlm.nih.gov Internet Source	1%
7	Mohamed Aziz Daghmouri, Sébastien Repplinger, Emmanuel Weiss, Marie-Pierre Bonnet et al. "The ability of shock index to predict refractory postpartum haemorrhage", European Journal of Anaesthesiology, 2025 Publication	<1%
8	Sakshi Agarwal, Uma Pandey. "Shock index as a predictor of maternal outcome in postpartum hemorrhage: an experience from	<1%

a tertiary care centre in Northern India",
International Journal of Medical Reviews and
Case Reports, 2023

Publication

-
- | | | |
|--------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|--------|
| <div style="background-color: #800080; color: white; padding: 2px 5px; display: inline-block;">9</div> | www.afro.who.int
<small>Internet Source</small> | $<1\%$ |
|--------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|--------|
-
- | | | |
|---------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------|
| <div style="background-color: #808000; color: white; padding: 2px 5px; display: inline-block;">10</div> | Te-Kai Liao, Chung-Han Ho, Ying-Jia Lin, Li-Chin Cheng, Hsuan-Yi Huang. "Shock index to predict outcomes in patients with trauma following traffic collisions: a retrospective cohort study", Research Square Platform LLC, 2024
<small>Publication</small> | $<1\%$ |
|---------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------|
-
- | | | |
|---------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------|
| <div style="background-color: #000080; color: white; padding: 2px 5px; display: inline-block;">11</div> | Mona M Ghonem, Amira A Abdelnoor, Aliaa A Hodeib. "Shock and modified shock indices in predicting poisoning severity and outcomes in acute aluminum phosphide poisoned patients", Toxicology Research, 2024
<small>Publication</small> | $<1\%$ |
|---------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------|
-
- | | | |
|---------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|--------|
| <div style="background-color: #000080; color: white; padding: 2px 5px; display: inline-block;">12</div> | www.pubfacts.com
<small>Internet Source</small> | $<1\%$ |
|---------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|--------|
-
- | | | |
|---------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------|--------|
| <div style="background-color: #FF0000; color: white; padding: 2px 5px; display: inline-block;">13</div> | Submitted to University of Western Sydney
<small>Student Paper</small> | $<1\%$ |
|---------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------|--------|
-
- | | | |
|---------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------|--------|
| <div style="background-color: #FF00FF; color: white; padding: 2px 5px; display: inline-block;">14</div> | www.researchsquare.com
<small>Internet Source</small> | $<1\%$ |
|---------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------|--------|
-
- | | | |
|---------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------|
| <div style="background-color: #800080; color: white; padding: 2px 5px; display: inline-block;">15</div> | Jamie L Finegan, Maja Marinkovic, Kyle Okamuro, Ron S Newfield, Jennifer T Anger. "Experience with gender affirming hormones and puberty blockers (gonadotropin releasing hormone agonist): a qualitative analysis of sexual function", The Journal of Sexual Medicine, 2025
<small>Publication</small> | $<1\%$ |
|---------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------|
-

16	www.mdpi.com Internet Source	<1 %
17	Submitted to University of Hertfordshire Student Paper	<1 %
18	repository.liv.ac.uk Internet Source	<1 %
19	774946cf-b2f4-4d65-a469-398263e8709a.filesusr.com Internet Source	<1 %
20	msjonline.org Internet Source	<1 %
21	Kyu Jin Lee, Yong Kyun Kim, Kyeongman Jeon, Ryoung-Eun Ko et al. "Shock indices are associated with in-hospital mortality among patients with septic shock and normal left ventricular ejection fraction", PLOS ONE, 2024 Publication	<1 %
22	Submitted to Southwestern Oklahoma State University Student Paper	<1 %
23	Jan Bláha, Tereza Bartošová. "Epidemiology and Definition of PPH Worldwide", Best Practice & Research Clinical Anaesthesiology, 2022 Publication	<1 %
24	dissertation.npmcn.edu.ng Internet Source	<1 %
25	www.nextias.com Internet Source	<1 %
26	M Mára, M Borčinová, Z Lisá, B Boudová, A Richtárová, D Kužel. "The perinatal outcomes	<1 %

of women treated for Asherman syndrome: a propensity score-matched cohort study", Human Reproduction, 2023

Publication

-
- 27 Mohammad Asim, Ayman El-Menyar, Talat Chughtai, Ammar Al-Hassani, Husham Abdelrahman, Sandro Rizoli, Hassan Al-Thani. "Shock Index for the Prediction of Interventions and Mortality in Patients With Blunt Thoracic Trauma", Journal of Surgical Research, 2023

<1 %

Publication

-
- 28 Nee-Kofi Mould-Millman, Julia M. Dixon, Bradley van Ster, Fabio Moreira et al. "Clinical impact of a prehospital trauma shock bundle of care in South Africa", African Journal of Emergency Medicine, 2022

<1 %

Publication

-
- 29 www.thieme-connect.com

Internet Source

<1 %

-
- 30 Arman Karami, Alireza Abdi, Ali Akbar Vaisi Raygani, Nader Salari, Mohammad Mehdi Mohammadi. "The prevalence of multiple organ dysfunction syndrome and its relationship with clinical variables in intensive care unit patients in Iran, 2022", The Egyptian Journal of Internal Medicine, 2025

<1 %

Publication

-
- 31 Ayse Cigdem Bayrak, Erdem Fadiloglu, Umut Sinci, Zeynep Yulek, Umutcan Kayikci, Murat Cagan, Ozgur Deren. "Shock index reference ranges in the first 48h postpartum following elective cesarean delivery: A prospective

<1 %

32 Helen Kennedy, Sarah L. Haynes, Clifford L. Shelton. "Maternal body weight and estimated circulating blood volume: a non-linear approach", British Journal of Anaesthesia, 2022

Publication

33 Süleyman Cemil Oğlak, Şeyhmus Tunç, Mehmet Obut, Erdal Şeker, Mustafa Behram, Ali Emre Tahaoğlu. "Maternal near-miss patients and maternal mortality cases in a Turkish tertiary referral hospital", Ginekologia Polska, 2021

Publication

34 doi.org Internet Source

35 eprints.nottingham.ac.uk Internet Source

36 hdl.handle.net Internet Source

37 journals.plos.org Internet Source

38 onlinelibrary.wiley.com Internet Source

39 pmc.ncbi.nlm.nih.gov Internet Source

40 Hannah L. Nathan, Paul T. Seed, Natasha L. Hezelgrave, Annemarie De Greeff et al. "Shock index thresholds to predict adverse outcomes in maternal hemorrhage and sepsis: A

41

Monika Chaudhary, Nandita Maitra, Tosha Sheth, Palak Vaishnav. "Shock Index in the Prediction of Adverse Maternal Outcome", The Journal of Obstetrics and Gynecology of India, 2020

Publication

<1 %

42

Süleyman Cemil Oğlak, Mehmet Obut, Ali Emre Tahaoğlu, Neslihan Uğur Demirel, Bekir Kahveci, İhsan Bağlı. "A prospective cohort study of shock index as a reliable marker to predict the patient's need for blood transfusion due to postpartum hemorrhage", Pakistan Journal of Medical Sciences, 2021

Publication

<1 %

43

Umran Dogru, Melih Yuksel, Mehmet Oguzhan Ay, Halil Kaya, Aksel Ozdemir, Yesim Isler, Mehtap Bulut. "The effect of the shock index and scoring systems for predicting mortality among geriatric patients with upper gastrointestinal bleeding: a prospective cohort study", Sao Paulo Medical Journal, 2022

Publication

<1 %

Exclude quotes On

Exclude matches

Off

Exclude bibliography On