

Obstructive Laryngeal Dyspnea Revealing a Large Supraglottic Fibrolipoma: A Case report

Introduction :

Laryngeal fibrolipoma is an uncommon benign tumor of the larynx, histologically composed of mature adipose tissue interspersed with fibrous connective elements. It is classified as a rare mesenchymal tumor and accounts for approximately 0.6% of all benign laryngeal lesions. These tumors most frequently occur in males during their sixth decade of life.[1]

The clinical diagnosis of laryngeal fibrolipoma can be particularly challenging due to its often nonspecific or absent symptomatology. Patients may present with a range of manifestations, including exertional dyspnea, paroxysmal coughing, dysphagia, or features of obstructive sleep apnea. In certain cases, progressive tumor growth may lead to partial or complete upper airway obstruction, necessitating prompt evaluation and management.[2]

This case report aims to present the clinical course of a rare laryngeal fibrolipoma, with emphasis on the diagnostic process and a review of current strategies for both medical and surgical intervention.

Case report :

We report the case of a 56-year-old male patient, a chronic smoker with a 20 pack-year history, referred for the management of a laryngeal fibrolipoma.

The patient reported a nine-month history of symptoms, initially presenting with dysphonia. Over time, progressive dysphagia and inspiratory dyspnea developed, prompting a nasofibrosopic examination. This revealed a supraglottic, exophytic mass arising from the larynx.

A cervical CT scan demonstrated a hypodense, fat-density lesion consistent with a supraglottic lipoma, measuring up to 4.77 cm in maximum diameter.

The patient underwent a tracheotomy under local anesthesia due to the obstructive nature of the mass, which contraindicated orotracheal intubation, followed by a suspension microlaryngoscopy under general anesthesia. Intraoperative examination revealed a large, pinkish, pear-shaped tumor attached by a pedicle to the left aryepiglottic fold. The vocal cords were visualized by gently retracting the mass to the left, and no involvement of the vocal folds was noted. Complete resection of the lesion was achieved via blunt dissection using standard laryngeal instruments. The excised specimen was sent for histopathological analysis.

Postoperative management included a six-day course of systemic corticosteroids and antibiotics. The patient's clinical condition improved progressively, and the tracheotomy site was closed on postoperative day 10 following endoscopic reassessment.

At the one-month follow-up visit, the patient reported complete resolution of symptoms and restoration of a normal voice. Follow-up pharyngolaryngoscopy confirmed excellent mucosal healing and a normal postoperative appearance.

42 Discussion

43 Laryngeal fibrolipomas are rare benign tumors, representing a histological variant of lipomas that
44 contain both mature adipose tissue and fibrous connective tissue. Lipomas represent approximately
45 0.6% of all benign laryngeal tumors, with the supraglottic region, especially the aryepiglottic fold.
46 They are the most commonly involved site due to its relative richness in adipose tissue.[1] These
47 tumors are more frequently observed in middle-aged men and tend to grow slowly, often remaining
48 asymptomatic until they reach a size that causes functional impairment.

49 Although lipomas are uncommon in the head and neck region, they can develop in a variety of
50 anatomical sub-sites, including the tonsils, tongue, parotid gland, palate, and larynx.

51 Laryngeal lipomas are categorized into two main types: extrinsic and intrinsic.
52 Extrinsic lipomas typically involve the posterior aspect of the larynx, the piriform sinus, and the
53 lingual surface of the epiglottis, while intrinsic lipomas are most often located in the false vocal
54 cords, the laryngeal surface of the epiglottis, the aryepiglottic folds, and the subglottic larynx.[2]

55 The clinical presentation of laryngeal fibrolipoma is often nonspecific and may overlap with more
56 common laryngeal pathologies and lead to diagnosis delays. Symptoms can include dysphonia,
57 progressive dysphagia, inspiratory stridor, and signs of upper airway obstruction such as exertional
58 dyspnea or sleep apnea. In some cases, sudden respiratory distress may occur, where the mass
59 intermittently occludes the glottic inlet.

60 Imaging studies, particularly CT scan and MRI, are critical for preoperative evaluation. On CT,
61 lipomas typically appear as well-circumscribed, hypodense lesions with fat attenuation. MRI offers
62 superior soft-tissue resolution and can help differentiate lipomas from liposarcomas based on signal
63 characteristics and the presence of internal septations or nodularity.

64 Histopathological examination remains the gold standard for definitive diagnosis. Fibrolipomas are
65 characterized by lobulated masses of mature adipocytes interspersed with fibrous septa, lacking
66 cellular atypia, mitotic activity, or necrosis, these features help distinguish them from
67 liposarcomas[3]. In infiltrative forms, malignant transformation remains a potential concern.

68
69 Given the rarity of fibrolipomas in the supraglottic larynx, establishing a correct diagnosis requires
70 careful distinction from a range of benign and malignant lesions that may present with similar
71 clinical and endoscopic findings.

72 The most relevant differential diagnoses, in addition to liposarcomas, include laryngocele, which is
73 one of the most common benign mimics. It represents an air-filled dilation of the laryngeal saccule,
74 often presenting as a cystic neck mass or intralaryngeal swelling. On CT imaging, it appears as an
75 air- or fluid-filled lesion without solid components, in contrast to the fat-density seen in
76 fibrolipomas.[4]

77 Another important consideration is hemangioma, a benign vascular tumor that enhances with
78 contrast and may appear bluish on endoscopy. It is composed histologically of proliferating blood
79 vessels, distinguishing it clearly from fibroadipose tissue.

80 Chondromas are also part of the differential diagnosis. These cartilaginous tumors are typically firm
81 and may show calcifications on imaging. [3][5]

82 Finally, neurogenic tumors, including schwannomas and neurofibromas, although uncommon, can
83 also occur in the supraglottic region and should be taken into account during diagnostic
84 evaluation.[5]

85 Complete surgical excision is the treatment of choice, with the approach tailored to tumor size,
86 location, and airway status. Endoscopic resection via suspension microlaryngoscopy is feasible for
87 most lesions, offering excellent visualization and preservation of surrounding structures. In cases of
88 large or obstructive tumors, a preliminary tracheotomy may be required to secure the airway, as in
89 the present case. Or an external surgical approach is preferred for tumors exceeding 2cm
90 diameter.[6]

91 Postoperative outcomes are typically excellent, as observed in our case, with low recurrence rates
92 and restoration of normal voice and swallowing function. However, regular follow-up is advisable
93 to monitor for recurrence or delayed complications such as scarring or granuloma formation.[7]

94
95

96 **Conclusion**

97 The evaluation of a supraglottic mass should include fibrolipoma in the differential diagnosis,
98 especially when imaging demonstrates a well-defined, homogeneous, fat-density lesion. Accurate
99 diagnosis relies on a multidisciplinary approach combining clinical examination, endoscopic
100 visualization, cross-sectional imaging, and confirmatory histopathological analysis. Distinguishing
101 fibrolipoma from other entities such as laryngoceles, liposarcomas, or vascular and neurogenic
102 tumors is essential for determining the appropriate surgical and therapeutic strategy.

103
104

105

106

107

108

109

110

111


112

- 113 **1. Azar SS, Buen F, Chia JJ, Ma Y, Caron J, Dry S, Bhuta S, Abemayor E.** Spindle Cell
114 Lipoma Arising from the Supraglottis: A Case Report and Review of the Literature. Head
115 Neck Pathol. 2021;15(4):1299–1302.
- 116 **2. J.-R. Lechien, S. Hans, M. Remacle.** Tumeurs bénignes du larynx .
- 117 **3. Kumar S, Ghosh A, Suri V, et al.** Large Laryngeal Lipoma with Extra Laryngeal
118 Component Mimics Mixed Form Laryngocele: A Case Report. Ear Nose Throat J.
119 2019;98(7):E20–E23.
- 120 **4. López-González MA, Rodríguez-Campo FJ, González-Moles MA, et al.** Laryngeal
121 Fibrolipoma: A Case Report. Acta Otorrinolaringol Esp. 2010;61(4):292–294.


- 122 **5. Kumar S, Ghosh A, Suri V, et al.** Dyspnea and Dysphagia Associated with
123 Hypopharyngeal Fibrolipoma: A Case Report. Case Rep Otolaryngol. 2017;2017:1–4.
- 124 **6. Santos-Silva AR, Oliveira-Neto HH, Lima Júnior LC, et al.** An Atypical Lipomatous
125 Tumor of the Hypopharynx, a Unique Entity? A Rare Case Report. Case Rep Otolaryngol.
126 2021.
- 127 **7. Joyson Kodiyan, B.S, Jason R. Rudman, M.D., David E. Rosow, M.D., Giovana R.**
128 **Thomas, M.D.** Lipoma and liposarcoma of the larynx: case reports and literature review.

129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144

UNDER PEER REVIEW IN IJAR



Axial views showing the supraglottic laryngeal mass without evidence of cartilaginous erosion.



Axial (left) and sagittal (right) CT scan of the neck showing a supraglottic fibrolipoma measuring $4.77 \times 3.66 \times 3.61$ cm.



Resected fibrolipoma.