

Pancreatico-pleural fistula-A rare sequelae of pancreatitis

by Jana Publication & Research

Submission date: 30-Jun-2025 01:57PM (UTC+0700)

Submission ID: 2690319141

File name: IJAR-52511.docx (65.91K)

Word count: 2352

Character count: 14720

15 ABSTRACT

Pancreatico-pleural fistula-A rare sequelae of pancreatitis

INTRODUCTION: Pancreaticopleural fistula is an uncommon complication of pancreatitis, presenting both diagnostic and therapeutic challenges. The following case report highlights the distinctive difficulties associated with pancreaticopleural fistula. The objective is to provide valuable insights to the scientific literature by detailing the case of a middle-aged gentleman suffering from sequelae of acute pancreatitis and left sided pleural effusion.

MATERIALS AND METHOD: A 41-year-old gentleman with a history of acute pancreatitis and chronic alcohol use presented with breathlessness, fever, chest pain, and abdominal pain. Computed tomography showed a pseudocyst involving the tail of pancreas & Spleen, a large left-sided pleural effusion and a moderate right-sided pleural effusion, along with signs of pneumonitis. Analysis of the pleural fluid from left pleural cavity revealed markedly elevated levels of Amylase. A MRCP (Abbrev) confirmed the findings & suggested a pancreaticopleural fistula.

RESULTS: Following a multidisciplinary consensus, an intercostal drainage tube was inserted into the left pleural cavity preoperatively. He was started on Octreotide therapy, and underwent nutritional & pulmonary optimization for elective surgery. He underwent a distal pancreato-splenectomy. Postoperative period was uneventful. He is doing well on 6 months of follow up.

CONCLUSION: The management of pancreaticopleural fistula demands a comprehensive and individualized approach. Recognition guided by high clinical suspicion coupled with appropriate investigations and a careful balance between medical and surgical interventions is crucial for achieving favorable outcomes. This case report adds to the scientific literature by providing insights into the complexities of sequelae of pancreatitis and emphasizes the importance of individualized strategies in their management.

Key points

1. Pancreatico pleural fistula being rare complication of pancreatitis
2. Clinical features mimics signs of breathlessness and fever which may cause delay in accurate diagnosis and their appropriate management

Key words

Pancreatitis, pancreatico pleural fistula, pleural effusion breathlessness psuedocyst

⁴INTRODUCTION

- Pancreaticopleural fistula has been recognized as a clinical entity since case reports were published in late 1960s [1]. Since that time, pancreaticopleural fistulae and pancreatic ascites have been termed as internal pancreatic fistulae which share common pathogenesis which includes the disruption of main pancreatic duct, resulting in leakage of pancreatic fluid [2–6]. This rare entity may be seen in patients with acute and chronic pancreatitis or may follow traumatic and surgical disruption of the pancreatic duct [2–6]. Pancreaticopleural fistula is a⁶ rare complication of chronic pancreatitis consequent to posterior disruption of the pancreatic duct. The fistulous track ascends into the pleural cavity and gives rise to large volumes of pleural fluid.⁶ To further complicate the matter, abdominal pain is seldom the presenting or significant feature.⁴ It is characterized by massive pleural fluid and has a tendency to recur following treatment. While conservative management with pancreatic duct stenting and inhibition of pancreatic secretion with octreotide may achieve closure of fistula in 31 to 45% of cases, surgery leads to healing in 80 to 90% of cases but carries a mortality up to 10% [2–6]. Pleural effusion resulting from a pancreaticopleural fistula is⁸ exceptionally rare, occurring in less than 1% of cases [7]. It is observed in approximately 3% to 7% of patients with pancreatitis [8]. The combined incidence of internal pancreatic fistulae—which includes both pancreatic ascites and pancreaticopleural fistulae—ranges from 0.4% to¹² 7% in individuals with chronic pancreatitis, and from 6% to 14% in

those with pancreatic pseudocysts. Among these, pancreaticopleural fistulae are less common than pancreatic ascites. Clinically, they often manifest as large, recurrent pleural effusions, typically affecting one side of the chest, with the left side being more frequently involved—reported in about 76% of cases.

Pancreaticopleural fistula typically arises from a leak associated with a partially formed or ruptured pancreatic pseudocyst, though in a smaller subset of cases, it results from a direct leak of the pancreatic duct. The fistulous tract may traverse the diaphragm via the aortic or esophageal hiatus or penetrate directly through the diaphragm. The underlying pathophysiology and etiology are similar for both pancreatic ascites and pancreatic pleural effusion. When the disruption of the pancreatic duct occurs anteriorly without being contained, it can lead to the formation of a pancreaticoperitoneal fistula, manifesting as ascites. Conversely, if the ductal disruption occurs posteriorly, pancreatic secretions may track into the retroperitoneum and ascend through the aortic or esophageal hiatus into the mediastinum. This can result in a pleural fistula directly or through the formation of a mediastinal pseudocyst, which may subsequently rupture into the pleural cavity, establishing a pleural fistula.

Pancreaticopleural fistula most commonly occurs in middle-aged men between the ages of 40 and 50, particularly those with a history of chronic alcohol use and pancreatitis. However, approximately 50% of affected individuals do not have a documented history of pancreatitis. Trauma is a much less frequent cause, accounting for only about 0.5% of cases. Pancreatic pseudocysts are present in 69% to 77% of patients who develop a pancreaticopleural fistula.

The clinical presentation can be misleading, as symptoms are typically related to a large pleural effusion. Common complaints include dyspnea, cough, chest pain, fever, and occasionally sepsis. Pulmonary symptoms are more prominent than abdominal ones and are often the initial

presenting features, with dyspnea being the most frequently reported symptom. Classic abdominal pain associated with acute pancreatitis is rarely reported. On average, symptoms persist for about 5.6 weeks before diagnosis.

Pleural effusion are predominately left sided; however, right-sided and bilateral effusion occurs in 19% and 14% of patient's, respectively.

Pleural effusion of this nature tends to be large and recurrent despite repeated thoracentesis. Many patients go through extensive pulmonary evaluation before pancreas is identified as the site of primary pathology. The pleural effusion is associated with ascites in 20% and pericarditis in 4% [9].

CASE REPORT


- A 41-year-old male farmer presented to our hospital with a 2-week history of progressively worsening shortness of breath, a dry cough, and left-sided chest pain. He denied experiencing recent abdominal pain, distension, nausea, or vomiting. The patient reported having undergone two thoracentesis procedures for left-sided pleural effusion at an outside facility over the past four months, which provided symptomatic relief.
- On physical examination, decreased breath sounds and dullness to percussion were noted on the left side of the chest, indicating the presence of a pleural effusion. On admission, his vital signs were: temperature 38.0°C, blood pressure 130/80 mmHg, pulse rate 95 bpm, respiratory rate 22 breaths per minute, and oxygen saturation of 92% on room air. The patient was tachypneic. He has history of smoking. He had a history of regular alcohol consumption (approximately 120 ml twice weekly for 20 years), which he discontinued 2 months prior to presentation. Six months ago, he experienced an acute episode of abdominal pain and was diagnosed with pancreatitis, which was managed conservatively.
- Computed tomography showed a pseudocyst involving the tail of pancreas & Spleen, a large left-sided pleural effusion along with signs of pneumonitis.

- Analysis of the pleural fluid from left pleural cavity following insertion of a intercostal drainage tube revealed markedly elevated levels of Amylase. A Magnetic Resonance Cholangio-Pancreaticography confirmed the findings & suggested a Left sided pancreaticopleural fistula

¹MANAGEMENT

Initial treatment focused on symptomatic relief and reduction of pancreatic secretions. An intercostal chest tube drainage was placed in the right pleural cavity to alleviate the pleural effusion. Octreotide injection was administered to inhibit pancreatic secretions after the cause of pleural effusion was attributed to PPF. The patient was provided total parenteral nutrition (TPN) to allow the pancreas to rest. A pancreatic duct stent was placed on ERCP to promote fistula closure¹. In addition to these interventions, the patient was given prophylactic antibiotic cover to prevent secondary infections.

- Following a multidisciplinary consensus underwent nutritional & pulmonary optimization for elective surgery. He underwent a distal pancreatectomy because of large number of collaterals which had developed between the tail of pancreas and the spleen
- Postoperative period was uneventful. He is doing well on 6 months of followup.



²DISCUSSION

Pleural fistulas develop due to abnormal communication of the pleural space with various structures in its vicinity, often due to iatrogenic causes, malignancy, or necrotizing infective processes(10). The different forms of pleural fistula include relatively common bronchopleural fistula, esophagopleural fistula and gastro pleural fistula, uncommon pancreaticopleural fistula, and hepatopleural fistula, and rarely encountered duropleural (subarachnoid-pleural) and nephropleural fistulas. Pleural space has negative pressure and deep recesses with large areas of surface contact with various structures, making it vulnerable to fistula formation. Pancreaticopleural fistulas (PPF) are uncommon and are usually linked to pancreatic trauma, pancreatitis, pancreatic surgery (such as pancreaticoduodenectomy), or ruptured pancreatic pseudocysts (11). Most patients with PPF are

middle-aged men, typically between 40 and 50 years old, with a history of chronic alcohol consumption and pancreatitis. PPF is usually seen in patients with necrotizing and recurrent pancreatitis; however, PPF may have insidious development in cases of pancreatitis with persistent inflammation. The fistulous tract may pass directly through the diaphragm—though this is uncommon—or through diaphragmatic openings such as the aortic or esophageal hiatus. Anterior ²disruption of the pancreatic duct (PD), when uncontained, typically results in ²a pancreaticoperitoneal fistula, often presenting as ascites. In contrast, posterior PD disruption can allow ¹pancreatic secretions to spread into the retroperitoneal space, potentially reaching the mediastinum via the esophageal or aortic hiatus and leading to a pancreaticopleural fistula (PPF). Another possible pathway involves the formation of a ¹mediastinal pseudocyst that eventually ruptures into the pleural cavity, creating a PPF(12). ¹Serum amylase and lipase levels are also often elevated in patients with pancreatic pseudocysts, as was seen in our case. Serum amylase and ¹lipase levels are also often elevated in patients with pancreatic pseudocysts, as was seen in our case. Clinical signs linked to a recurrent large pleural effusion typically include dyspnea, cough, fever, chest discomfort, and septicemia. ¹Dyspnea is the most common presentation of pleuropulmonary symptoms in PPF, which are more common than abdominal symptoms. Patients rarely report the characteristic abdominal pain associated with pancreatitis, and many patients undergo an extensive pulmonary evaluation with delayed diagnosis of PPF. A plain chest radiograph is typically the first diagnostic tool used to identify pleural effusion, which may appear unilaterally—most often on the left—or bilaterally. Thoracocentesis is performed to relieve symptoms and to obtain pleural fluid for analysis. In cases of pancreaticopleural fistula (PPF), ⁷pancreatic enzymes leak into the pleural space, resulting in pleural effusions rich in amylase and lipase, with ⁷pleural fluid amylase levels often exceeding 1,000 U/L. Elevated pleural fluid amylase can also be seen in other conditions, such as ¹acute pancreatitis, gastropneural and esophagopleural fistulas, pulmonary tuberculosis, liver cirrhosis, and certain malignancies. ⁹However, markedly elevated amylase levels—above 10,000 U/L—are more characteristic of PPF or ruptured pancreatic pseudocysts, with levels surpassing 50,000 U/L typically indicating PPF (12). In the current case, persistent and rapid re-accumulation

of pleural fluid—despite 2 previous thoracentesis draining over a liter each time—prompted further investigation into rarer causes. The markedly elevated ²pleural fluid amylase and lipase levels ¹were critical diagnostic clues pointing toward a pancreatic origin. ¹The management of PPF typically involves a combination of medical, endoscopic, and sometimes surgical interventions. Initial treatment includes chest tube drainage of the pleural effusion, octreotide administration to reduce pancreatic secretions, and nutritional support through total parenteral nutrition (TPN) to rest the pancreas. ¹Surgical management is curative, with a success rate of up to 90%, and is typically done in patients who fail to respond to conservative treatment.

CONCLUSION

- This case highlights the necessity ¹of considering pancreaticopleural fistula (PPF) in the differential diagnosis of recurrent pleural effusion, particularly in patients with a history of pancreatic disease or relevant risk factors. Early diagnosis and appropriate therapeutic intervention are crucial for improving clinical outcomes.
- ³The management of pancreaticopleural fistula demands a comprehensive and individualized approach. Recognition guided by high clinical suspicion coupled with appropriate investigations and a careful balance between medical and surgical interventions is crucial for achieving favorable outcomes.
- This case report adds to the scientific literature by providing insights into the complexities of sequelae of pancreatitis and emphasizes the importance of individualized strategies in their management. It also emphasizes the essential role of advanced imaging and endoscopic


techniques in the accurate diagnosis and effective management of this uncommon but significant clinical entity

REFERENCES

- 1.Anderson WJ, Skinner DB, Zuidema GD, Cameron JL. Chronic pancreatic pleural effusions. Surgery Gynecology and Obstetrics. 1973;137(5):827–830. [[PubMed](#)] [[Google Scholar](#)]
- 2.King JC, Reber HA, Shiraga S, Hines OJ. Pancreatic-pleural fistula is best managed by early operative intervention. Surgery. 2010;147(1):154–159. doi: 10.1016/j.surg.2009.03.024. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
- 3.Sut M, Gray R, Ramachandran M, Diamond T. Pancreaticopleural fistula: a rare complication of ERCP-induced pancreatitis. Ulster Medical Journal. 2009;78(3):185–186. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- 4.Dhebri AR, Ferran N. Nonsurgical management of pancreaticopleural fistula. Journal of the Pancreas. 2005;6(2):152–161. [[PubMed](#)] [[Google Scholar](#)]
- 5.Vyas S, Gogoi D, Sinha SK, Singh P, Yadav TD, Khandelwal N. Pancreaticopleural fistula: an unusual complication of pancreatitis diagnosed with magnetic resonance cholangiopancreatography. Journal of the Pancreas. 2009;10(6):671–673. [[PubMed](#)] [[Google Scholar](#)]
- 6.Safadi BY, Marks JM. Pancreatic-pleural fistula: the role of ERCP in diagnosis and treatment. Gastrointestinal Endoscopy. 2000;51(2):213–215. doi: 10.1016/s0016-5107(00)70422-6. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
- 7.Burgess NA, Moore HE, Williams JO, Lewis MH. A review of pancreatico-pleural fistula in pancreatitis and its management. HPB Surgery. 1991;5(2):79–86. doi: 10.1155/1992/90415. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

8. Materne R, Vranckx P, Pauls C, Coche EE, Deprez P, Van Beers BE. Pancreaticopleural fistula: diagnosis with magnetic resonance pancreatography. *Chest*. 2000;117(3):912–914. doi: 10.1378/chest.117.3.912. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
9. Hastier P, Rouquier P, Buckley M, Simler JM, Dumas R, Delmont JP. Endoscopic treatment of wirsungo-cysto-pleural fistula. *European Journal of Gastroenterology and Hepatology*. 1998;10(6):527–529. doi: 10.1097/00042737-199806000-00017. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
10. Kapoor H, Gulati V, Gulati A, Donuru A, Parekh M (2022) Comprehensive imaging review of pleural fistulas from diagnosis to management. *Radiographics* 42(7):1940–1955. <https://doi.org/10.1148/rg.220083>. (Epub 2022 Oct 21 PMID: 36269669)
11. Koussayer B, Kattih M, Nester M, Peterson P, DuCoin CG. (2023) Gastropleural fistula presenting as a complication of gastric sleeve surgery: a case report. *Cureus* 4;15(4):e37133. <https://doi.org/10.7759/cureus.37133>. PMID: 37162785; PMCID: PMC10163983
12. Tay CM, Chang SK (2013) Diagnosis and management of pancreaticopleural fistula. *Singapore Med J* 54(4):190–4. <https://doi.org/10.11622/smedj.2013071>. (PMID: 23624444)





Pancreatico-pleural fistula-A rare sequelae of pancreatitis

ORIGINALITY REPORT

55%
SIMILARITY INDEX

51%
INTERNET SOURCES

54%
PUBLICATIONS

1%
STUDENT PAPERS

PRIMARY SOURCES

1 ejb.springeropen.com 18%
Internet Source

2 Mithun Nilgiri K., Varsha P. Rangankar, Shaily K. P., Tushar R. Sahasrabudhe, Rishi Govind Orakkan. "Pancreaticopleural fistula: a rare cause of recurrent right-sided pleural effusion", The Egyptian Journal of Bronchology, 2025 8%
Publication

3 jmedicalcasereports.biomedcentral.com 8%
Internet Source

4 www.hindawi.com 7%
Internet Source

5 pancreapedia.org 3%
Internet Source

6 pancreas.imedpub.com 3%
Internet Source

7 CM Tay, S Change. "Diagnosis and management of pancreaticopleural fistula", Singapore Medical Journal, 2013 2%
Publication

8	www.ncbi.nlm.nih.gov Internet Source	1 %
9	Rajesh Kumar, Harikrishnan Premdeep, Saurabh Dawra, Roshan Ray. "Uncommon Connections—Pancreaticopleural Fistula as a Cause of Recurrent Pleural Effusion: A Case Report", Journal of Postgraduate Medicine, Education and Research, 2025 Publication	1 %
10	www.thieme-connect.com Internet Source	1 %
11	pubmed.ncbi.nlm.nih.gov Internet Source	1 %
12	Hakim Elkaoui, Fouad Atoini, Sidi Mohamed Bouchentouf, Fatima El Omari et al. "Pancreatic-pleural fistula in chronic pancreatitis", Arab Journal of Gastroenterology, 2012 Publication	1 %
13	Choe, In Soo. "Acute mediastinitis arising from pancreatic mediastinal fistula in recurrent pancreatitis", World Journal of Gastroenterology, 2014. Publication	<1 %
14	jmas.in Internet Source	<1 %
15	Manoj Munirathinam, Pugazhendhi Thangavelu, Ratnakar Kini. "Pancreatico-pleural Fistula: Case Series", Journal of Digestive Endoscopy, 2019 Publication	<1 %
16	Zehra Dhanani, Stephen Dachert. "Pancreatopleural Fistula: A Rare Cause of Recurrent Pleural Effusion", Respiratory Medicine Case Reports, 2025	<1 %

17	Kosmas, Epaminondas, and Vlasis Polychronopoulos. "Pleural Effusions in Gastrointestinal Tract Diseases", Lung Biology in Health and Disease, 2004. <small>Publication</small>	<1 %
18	Milan Khadka, Suzit Bhusal, Binod Pantha, Rabin Gautam, Kapil Gautam, Ashlesha Chaudhary. "Pancreaticopleural fistula causing pleural effusion: a case report and review of the literature", Journal of Medical Case Reports, 2024 <small>Publication</small>	<1 %
19	Piyush Ranjan, Rinkesh Bansal, Munish Sachdeva, Mandhir Kumar. "Pancreaticopleural fistula: Report of two cases and review of literature", Journal of Digestive Endoscopy, 2019 <small>Publication</small>	<1 %

Exclude quotes

On

Exclude matches

Off

Exclude bibliography

On