

Profile fatty acid of Chironomidae larvae produced from rabbit manures fish farms in the Guinean forested region

by Jana Publication & Research

Submission date: 12-Jul-2025 05:35PM (UTC+0700)

Submission ID: 2690332272

File name: IJAR-52757.docx (29.57K)

Word count: 1924

Character count: 11062

Profile fatty acid of Chironomidae larvae produced from rabbit manures fish farms in the Guinean forested region

Abstract

This study confirms the richness in fatty acids of Chironomidae larvae produced from rabbit droppings. These Chironomidae larvae can be used in aquaculture for more environmentally friendly and less costly larval production for fry rearing in fish farms. This simple technique for producing Chironomidae larvae would reduce environmental impact while strengthening the resilience of aquaculture to climate change.

Key words: Aquaculture, Chironomidae Larvae, Rabbit Manure, Fatty Acids

Introduction

Use live feed in hatcheries in Africa general and in Guinea particular is a promising alternative following the various failures of using dry feed during larval and fry rearing, as well as its unavailability and high cost (Adande et al. 2025). Live feeds contain essential compounds beneficial to fish growth and survival (Iraj et al., 2024). Similarly, the effectiveness of live prey depends on the supply sufficient quantities amino acids (AA) and unsaturated fatty acids (UFA) for larval growth and development (Samat et al., 2020). Similarly, according to Arts et al. (2001), benthic macroinvertebrates are one the most important components freshwater ecosystems and a primary food source for fish. Through the process energy transfer (Eddy et al., 2021), the fatty acid richness of fish could play an important role in men's need for essential fatty acids. Improving the nutritional quality of live fatty acid foods is therefore essential. The aim this study was to evaluate the different proportions essential fatty acids in Chironomidae produced from rabbit manure.

Materials and methods

Producing Chironomidae from rabbit manure

Chironomidae larvae were produced using an optimal dose of 600 g/m³ or 25 g/dm² of rabbit droppings, according to Adandé et al. (2018). These rabbit manure consist of 15%, 1.3%, and 0.8% N, P, and K, respectively. The characteristics are presented in Table 1. The first step in this production technique involves setting up a medium composed of sixteen (16) liters of borehole water and a dose of 600 g/m³ or 25 g/dm² of rabbit. The second step consists of adding four (04) liters of pond water, previously filtered using a 200µm metal sieve, three

(03) days after fertilization, in order to eliminate any macroinvertebrates and allow phytoplankton, a source of food, to develop. The third step, three (03) days after fertilization, involves seeding the media with macroinvertebrates (Chironomidae larvae) at an initial density of at least 6 to 12 individuals per 20 liters. The production areas are covered with mosquito netting to prevent potential predators of macroinvertebrates from entering the culture environment.

Chemical analyses

Physicochemical parameters (temperature, pH, conductivity, salinity (from rabbit manure), TDS, dissolved oxygen) were measured in situ every 7 days (at 8 a.m. and 5 p.m.) using the CALYPSO multi-parameter probe (softer version/2015 2138 SN-ODEON CALYPSO; $\pm 0.1^\circ\text{C}$ sensitivity). Turbidity measurements were taken in the various production environments using a turbidimeter (EUTECH TN-100). Five hundred milliliters of water from the production environment were sampled every seven days and stored in a refrigerator (at 4°C) for the measurement of nitrates, nitrites, orthophosphates, and ammonium using a molecular absorption spectrophotometer (HACH DR/2800) according to Rodier & Merlet (2012).

Fatty acid analysis.

To assess the fatty acid profile, Chironomidae (bloodworms) produced from rabbit droppings were harvested from the production buckets and sorted using a $200\ \mu\text{m}$ sieve. Harvested Chironomidae worms were placed in microtubes and rapidly frozen in liquid nitrogen. These samples were then immediately transferred to the Aquatic Animal Nutrition Laboratory at the Regional Laboratory for Health Safety Expertise and Analysis (LSSEA/IRGIB) and stored at -80°C until analysis.

Measurement fatty acid composition involved two steps: a) fat extraction and esterification according to Folch et al. (1957) and Firston (1998). Fat extraction was performed using the methanol-chloroform extraction method, followed by fat esterification using 2% methanolic sodium and BF_3 (boron trifluoride). b) Fatty acid samples were then analyzed using a gas chromatograph (Philips, Sussex, England) equipped with an SGE BPX70 capillary column (ID: $0.25\ \text{mm} \times 0.22\ \mu\text{m} \times 30\ \text{m}$). We used a flame ionization detector at a temperature of 300°C and an injector set at 250°C . A volume of $0.2\ \mu\text{L}$ of the ester sample was injected into the gas chromatograph for analysis. The initial column temperature was set at 160°C and gradually adjusted up to 230°C and maintained for 5 minutes until all compounds were eluted. In addition, helium was used as carrier gas, hydrogen as fuel, nitrogen as auxiliary gas and synthetic air. By comparing the inhibition times of chromatograms from unknown samples with those obtained from a standard

solution, the fatty
produced from rabbit
results expressed as a

Parameters	Averages
Temp (°C)	31.51±0.63
pH	7.12±0.10
Cond (μs/cm)	621.06±14.53

Results and
Physicochemical
Temperature is an
optimizing the
Chironomidae
value recorded
to those obtained by
on the drift of

acids present in Chironomidae
manure were identified and the
percentage.

discussion
parameters
important parameter for
production of freshwater
larvae (Table 1). The average
during this study is comparable
Williams (1985) during his work
Chironomidae egg masses. The

neutral pH obtained in this study is similar to those obtained by Sotaro et al. (2023) during the conservation of Chironomidae larvae in the laboratory for DNA sequencing. The production of Chironomidae larvae at a neutral pH would allow for the conservation of their genetic material (Bisthoven et al., 2003). Conductivity, salinity, turbidity, and transparency (Table 1) are values selected for the successful production of Chironomidae larvae (Adandé et al., 2018). The average values for ammonia, nitrite, and nitrate are close to those obtained by Poulsen et al. (2014). Chironomidae are capable of mobilizing and denitrifying ammonium (Samuiloviene et al., 2019). The average orthophosphate level obtained during this study is higher than those obtained by Biswas et al. (2009), which ranged from 0.03 to 0.189 mg/l. Chironomidae larvae are capable of increasing orthophosphate levels during the bioturbation process (Biswas et al., 2009). All of these average values for the various physical and chemical parameters would allow for the production of good-quality Chironomidae larvae for fish larvae.

Table 1: Production medium

OD (mg/l)	3.28±0.46
TDS	318.46±8.58
Sal (mg/l)	0.34±0.00
Transp (cm)	16.85±1.51
Turb (NTU)	84.77±4.01
¹³ N-NH ₃ (mg/l)	0.98±0.09
N-NO ² (mg/l)	0.29±0.05
N-NO ³ (mg/l)	0.19±0.01
P-PO ₄ (mg/l)	1.21±0.11

⁷
T: temperature, pH: Hydrogen potential, OD: dissolved oxygen, TDS: Total dissolved solids, Cond: conductivity and sal: salinity, Turb: turbidity, Transp: transparency

Quantity and quality of Chironomidae fatty acids produced from rabbit manure

Chironomidae larvae produced from rabbit manure have the right fatty acid proportions in terms of quantity and quality. Such larvae are likely to meet the needs of fish larvae. Indeed, ¹² the ratio of total long-chain polysaturated fatty acids ω6 to polysaturated fatty acids ω3 (TAGP) ω6/TAGP ω3 is equal to 1.33 (Table 2). This ratio highlights the richness of larvae in long-chain polysaturated fatty acids, and perfectly meets the needs of fish larvae and fry. However, this ratio is lower than those obtained by Iraj et al. (2024), which is 2.81 in the Chironomidae. This difference may be due to the production environments and species used. On the other hand, these results corroborate those obtained by Kiyashko et al. (2004) in the larvae *Stictochironomus pitulus* (Diptera: Chironomidae). One the most important factors is

Fatty Acid	Names	%
------------	-------	---

the nutritional quality of the live feed for rearing larvae and fry, including the content⁹ essential fatty acids, in particular eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA).

They have values 5.79 and 1.52%²¹ respectively and are referred to as highly unsaturated fatty acids (HUFA) (Table 2). The long-chain polyunsaturated fatty acids in the (ω3) series are 3.03% higher than those obtained in the fish larvae feed formulation by Zambonino (2000). According to Michaïl et al (2013), these fatty acids are physiologically indispensable to animals, including humans.

Table 2 : Fatty acid profile Chironomidae larvae produced from rabbit manure (% of total proportions)

10 C14 :0	Myristic acid	1,65
C15 :0	Pentadecanoic acid	1,08
C16 :0	Palmitic acid	19,67
6 C17 :0	Heptadecanoic acid	1,85
C18 :0	Stearic acid	5 ,14
C20 :0	Arachidic acid	0 ,23
C21 :0	Heneicosanoic acid	0 ,78
C24 :0	Tetracosanoic acid	0 ,18
Total saturated fatty acids	TOTAL SFA	30,58
5 C14 :1	Myristoleic acid	0,17
C15 :1	Pentadecanoic acid	0,39
C16 :1	Palmitoleic acid	13,49
C17 :1	Heptadecanoic acid	0,53
C18 :1	Oleic acid	22,64
C20 :1	Gondoic acid	1,82
C24 :1	Nervonic acid	0,51
Total monosaturated fatty acids	TOTAL MsFA	39,55
14 C18 :2 ω6	Linoleic acid	14,86
C18 :3 ω6	α-linolenic acid	0,52
C20 :3 ω6	Dihomolinolenic acid	0,54
C20 :4 ω6	Arachidonic acid	4,97
Total polysaturated fatty acids ω6 (PFGA ω6)	TOTAL : PsFGA ω6	20,89
C18 :3 ω3	γ-Linolenic Acid	3,03
C18 :4 ω3	Stearidonic Acid	4,29
C20 :5 ω3	Eicosapentaenoic Acid (EPA)	5,79
C22 :5 ω3	Docosapentaenoic Acid	1,06

C22 :6 ω3	Docosanoic Acid	1,52
Total polyunsaturated fatty acids ω3	Total : PsFGA ω3	15,69
PsFGA ω6/ PsFGA ω3		1,33

Conclusion

This study of the fatty acid profile Chironomidae freshwater produced from rabbit droppings, shows that these fatty acids can be used in hatchery larval production for more environmentally friendly, efficient, and less costly production for rural fish farmers. However, an optimization technique in a healthily controlled environment is needed.

Conflict of interest: The authors declare that they have no conflict of interest.

Author contributions: ADANDE Richard: Conceived the study, directed the research writing, editing and interpretation of the results. KEITA Oumar and BAKAYOKO Ibrahima: Conceived the study, directed the research writing, editing and interpretation of the results. SANGARE Aboubacar: Editing and directed the research.

Références

1. Adande, R., G. Djidohokpin, A.S.M. Djissou, P. Bilivogui, J-C. Micha. 2025. "Bromatological value of Chironomidae produced from organic fertilizers and their effects on the growth of *Clarias gariepinus* fry in the Guinean forest region". Blue Biotechnology 2:14. <https://doi.org/10.1186/s44315-025-00024-y>
2. Adandé, R., Liady, M.N.D., H.K.J. Bokossa, G. Djidohokpin, F. Zouhir, G.A. Mensah, E.D. Fiogbe .2018. "Utilisation rationnelle de fertilisants organiques pour la production de macroinvertébrés benthiques d'eau douce en pisciculture". BASE. 208-219. <https://doi.org/10.25518/1780-4507.16660>
3. Arts M.T., R.G. Ackman and B.J. Holub. 2001. "Essential fatty acids" in aquatic ecosystems: a crucial link between diet and human health and evolution". Can. J. Fish. Aquat. Sci., 58, 122. <https://doi.org/10.1139/f00-224>
4. Eddy, T.D., J.R. Bernhardt, J.L. Blanchard, W.W.L. Cheung, M. Colléter et al. 2021. "Energy Flow Through Marine Ecosystems: Confronting Transfer Efficiency". Tends, in Ecology and Evolution 36 (1):76-86. <https://doi.org/10.1016/j.tree.2020.09.006>
5. Firstone, D. 1998. Official Methods and Recommended Practices of the American Oil Chemists Society, Vol. I-II, AOCS, Champaign, (5 ed.) edition.
6. Folch, J., M. Lees, and G. H. Sloane-Stanley. 1957. "A simple method for the isolation and purification of total lipides from animal tissues". Journal of Biological Chemistry, 226 (1) : 497-509. <https://doi.org/10.1016/j.prostaglandins.2013.03.002>
7. Iraj, E., Bahram F., Mir M.S. and M. M. Shokri. 2024. "The Effect of Feeding with Chironomid and Artemia on Fatty Acids and Amino Acids Profiles in Persian Sturgeon (*Acipenser persicus*) Larvae. Aquaculture Nutrition, 13. <https://doi.org/10.1155/2024/6975546>
8. Kiyashkoa, S.I., A.B. Imbs, T.Narita, V. I. Svetashev, W. Eitaro .2004. "Fatty acid composition of aquatic insect larvae *Stictochironomus pictulus* (Diptera: Chironomidae): evidence of feeding upon methanotrophic bacteria". Comparative Biochemistry and Physiology, Part B 139 :705-711.
9. Michaïl, G. Ier., N. Souchtchik Nadezhda, N. Makhutova Olesia. 2013. "Prostaglandines et autres médiateurs lipidiques". 107 :117-126.
10. Rodier, J.B.L, N. Merlet, coll. 2012. "Analyse de l'eau". 9e éd. Paris: Dunod.
11. Samat, N.A., F.M. Yusoff, N.W. Rasdi, and M. Karim. 2020. "Enhancement of live food nutritional status with essential nutrients for improving aquatic animal health: a review," Animals, 10 (12) : 1-27.

12. Zambonino, J. 2000. "Brevet WO 00/64273 Aliment complet pour larves de poissons et procédé pour sa préparation". PCT/FR00/01068.

Profile fatty acid of Chironomidae larvae produced from rabbit manures fish farms in the Guinean forested region

ORIGINALITY REPORT

26%

SIMILARITY INDEX

18%

INTERNET SOURCES

24%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|----|
| 1 | downloads.hindawi.com
Internet Source | 7% |
| 2 | Richard Adande, Gildas Djidohokpin, Asm Djissou, Papa Bilivogui, Micha Jean-Claude. "Bromatological value of Chironomidae produced from organic fertilizers and their effects on the growth of <i>Clarias gariepinus</i> fry in the Guinean forest region", Blue Biotechnology, 2025
Publication | 3% |
| 3 | www.hindawi.com
Internet Source | 2% |
| 4 | Pablo Intriago, Andrés Medina, Nicole Cercado, Kelly Arteaga et al. "Passive surveillance for shrimp pathogens in <i>Penaeus vannamei</i> submitted from 3 regions of Latin America", Aquaculture Reports, 2024
Publication | 2% |
| 5 | Michael D. Moths, John A. Dellinger, Bruce Holub, Michael P. Ripley, Joseph E. McGraw, Ronald E. Kinnunen. "Omega-3 Fatty Acids in Fish from the Laurentian Great Lakes Tribal Fisheries", Human and Ecological Risk Assessment: An International Journal, 2012
Publication | 1% |
| 6 | B. Kierończyk, S. A. Kaczmarek, M. Hejdysz, P. Szymkowiak et al. "Implementation of a | 1% |

metabolizable energy regression model for black soldier fly larvae fat in broiler chicken diets: effect on growth performance, nutrient digestibility, and selected physiological indices", Journal of Animal and Feed Sciences, 2024

Publication

-
- | | | |
|----------|---|------------|
| 7 | link.springer.com
<small>Internet Source</small> | 1 % |
|----------|---|------------|
-
- | | | |
|----------|---|------------|
| 8 | publinestorage.blob.core.windows.net
<small>Internet Source</small> | 1 % |
|----------|---|------------|
-
- | | | |
|----------|---|------------|
| 9 | www.raingarden.us
<small>Internet Source</small> | 1 % |
|----------|---|------------|
-
- | | | |
|-----------|---|------------|
| 10 | Antti Aro. "Inverse Association Between Dietary and Serum Conjugated Linoleic Acid and Risk of Breast Cancer in Postmenopausal Women", Nutrition and Cancer, 01/01/2000
<small>Publication</small> | 1 % |
|-----------|---|------------|
-
- | | | |
|-----------|---|------------|
| 11 | N.N. Sushchik, M.I. Gladyshev, A.V. Moskvichova, O.N. Makhutova, G.S. Kalachova. "Comparison of fatty acid composition in major lipid classes of the dominant benthic invertebrates of the Yenisei river", Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology, 2003
<small>Publication</small> | 1 % |
|-----------|---|------------|
-
- | | | |
|-----------|---|------------|
| 12 | research.library.mun.ca
<small>Internet Source</small> | 1 % |
|-----------|---|------------|
-
- | | | |
|-----------|---|------------|
| 13 | L Yang. "Biofilter treatment of aquaculture water for reuse applications", Water Research, 2001
<small>Publication</small> | 1 % |
|-----------|---|------------|
-

14	garri.overcastcdn.com Internet Source	1 %
15	www.science.gov Internet Source	1 %
16	Shohei Hayashi, Mai Naka, Kenshin Ikeuchi, Makoto Ohtsuka et al. "Control Mechanism for Carbon-Chain Length in Polyunsaturated Fatty-Acid Synthases", Angewandte Chemie, 2019 Publication	<1 %
17	d-nb.info Internet Source	<1 %
18	www.fisheries.go.th Internet Source	<1 %
19	Hidayu Suhaimi, Amirah Yuslan, Nizalmie Azani, Ahasan Habib, Hon Jung Liew, Nadiah W. Rasdi. "Effect of dietary enhanced Moina macrocopa (Straus, 1820) on the growth, survival and nutritional profiles of hybrid Nile tilapia fry", The Egyptian Journal of Aquatic Research, 2021 Publication	<1 %
20	Richard Adande, Hervé Kouessivi Janvier Bokossa, Mouhamadou Nourou Dine Liady, Emile Didier Fiogbé. "Valorization of various sources of rabbit manure in agro-piscicultural system in Benin (West Africa): dynamics and effect of mineralization upon quality of fresh water", International Journal of Recycling of Organic Waste in Agriculture, 2017 Publication	<1 %
21	orbi.uliege.be Internet Source	<1 %

Exclude quotes	On	Exclude matches	Off
Exclude bibliography	On		