

“Antimicrobial and Antioxidant Activity of Triphala: An Ayurvedic Formulation”

by Jana Publication & Research

Submission date: 12-Jul-2025 05:47PM (UTC+0700)

Submission ID: 2690345585

File name: IJAR-52771.docx (263.6K)

Word count: 2281

Character count: 14110

“Antimicrobial and Antioxidant Activity of Triphala: An Ayurvedic Formulation”

Abstract

Triphala, an ancient Ayurvedic formulation composed of the dried fruits of *Emblica officinalis* (Amla), *Terminalia chebula* (Haritaki), and *Terminalia bellerica* (Bahera), is widely recognized for its broad-spectrum therapeutic applications. This study aimed to scientifically evaluate its **phytochemical composition**, **antimicrobial efficacy**, and **antioxidant potential** using different solvent extracts. Preliminary phytochemical screening revealed that the **ethanolic extract** was the most bioactive, containing alkaloids, tannins, glycosides, saponins, terpenoids, and phlobatannins. Antimicrobial testing against *Escherichia coli* and *Staphylococcus aureus* using the **disc diffusion method** demonstrated significant activity in ethanolic and aqueous extracts, with the ethanolic extract showing the highest inhibition zones. Furthermore, **DPPH radical scavenging assays** confirmed the antioxidant capacity of both ethanolic and aqueous extracts in a dose-dependent manner, although their efficacy was lower than that of standard ascorbic acid. These results substantiate the traditional use of Triphala and highlight its potential as a natural source of antimicrobial and antioxidant agents. Further phytochemical characterization and in vivo studies are recommended for clinical validation.

Keywords

Triphala, Antioxidant Activity, Antimicrobial Activity, Phytochemical Screening, Ayurvedic Medicine, Ethanolic Extract, DPPH Assay, Herbal Formulation, Natural Therapeutics, *Escherichia coli*, *Staphylococcus aureus*.

1. INTRODUCTION

Triphala, a traditional Ayurvedic formulation, has been extensively used for centuries in Indian medicine owing to its broad spectrum of therapeutic properties. Composed of the dried fruits of three medicinal plants—*Emblica officinalis* (Amla), *Terminalia chebula* (Haritaki), and *Terminalia bellerica* (Bahera)—Triphala is revered for its ability to promote digestive health, enhance immunity, and delay the aging process. Its multifaceted medicinal applications are increasingly being validated by modern scientific investigations, particularly for its antimicrobial and antioxidant potential.

The biological activities of Triphala are largely attributed to the presence of diverse phytochemicals. Phytochemicals are naturally occurring, biologically active compounds found in plants, excluding essential nutrients such as vitamins and minerals. These compounds act as the plant's defense mechanisms against pathogens and environmental stress, and they offer similar protective benefits when consumed by humans. Rich in flavonoids, tannins, phenolic acids, and other bioactive constituents, phytochemicals exhibit a variety of therapeutic properties, including antioxidant, anti-inflammatory, antibacterial, antiviral, and immune-boosting effects.

Each component of Triphala contributes uniquely to its overall pharmacological efficacy. *Embl*¹⁵*ca officinalis* (Amla) is a potent source of ascorbic acid (vitamin C), along with tannins, flavonoids, and polyphenols such as gallic acid and ellagic acid, all known for their antioxidant activity. *Terminalia chebula* (Haritaki) is rich in tannic acid, chebulinic acid, and betulinic acid, which possess significant antimicrobial and free radical scavenging properties. *Terminalia bellerica* (Bahera), on the other hand, contains ellagitannins, gallotannins, and a spectrum of polyphenolic compounds that further enhance the formulation's antioxidant and antimicrobial activities.

Given the rising global concern regarding antibiotic resistance and oxidative stress-related disorders, there is a pressing need to explore plant-based alternatives with minimal side effects. Triphala, with its rich phytochemical profile, presents a promising candidate for such therapeutic exploration. This study aims to evaluate the antimicrobial and antioxidant activities of Triphala, thereby providing scientific validation for its traditional usage and exploring its potential as a natural alternative for managing microbial infections and oxidative stress.

2. MATERIAL & METHOD

2.1 Materials

Triphala powder (100 g) was procured from Vindhya Herbals, Sanjeevani Ayurveda, Bhopal, and stored for further use. Extraction solvents included chloroform, petroleum ether, and ethanol, all of analytical grade. Microbial activity was assessed using Nutrient Agar Media (NAM) and Nutrient Broth, both with standard compositions and pH adjusted to 7.0.

2.2 Instruments

Key instruments used included a Soxhlet extractor for phytochemical extraction, rotary vacuum evaporator for solvent recovery, digital pH meter, vertical autoclave, laminar airflow chamber, incubator, and antibiotic zone scale for zone measurement.

2.3 Extraction Procedure

Soxhlet extraction was carried out using 25 g of Triphala powder and 250 mL of solvent at 50–60°C until clear extract was obtained. Solvent was then recovered via rotary evaporation under vacuum, and the residue was mixed with chloroform water and refrigerated at 4°C for further analysis.

2.4 Phytochemical Screening

Preliminary phytochemical screening of Triphala extracts was conducted following standard protocols (Brain & Turner, 1975; Evans, 1996) to identify key bioactive constituents. Various qualitative tests were performed using specific reagents to detect the presence of the following phytochemicals:

- **Alkaloids:** Detected using Mayer's reagent, indicated by a cream-colored precipitate.
- **Carbohydrates & Glycosides:** Identified by a red precipitate upon reaction with Fehling's solution.
- **Phenolic Compounds & Tannins:** Confirmed by a bluish-black coloration with ferric chloride.

- **Proteins & Amino Acids:** Indicated by a purplish-pink color upon addition of ninhydrin.
- **Terpenoids:** Detected by the formation of a red-brown interface with chloroform and concentrated H₂SO₄.
- **Phlobatannins:** Identified through the deposition of red precipitate after boiling with 1% HCl.
- **Saponins:** Confirmed by persistent frothing upon boiling with distilled water.
- **Flavonoids:** Detected by a yellow color with NaOH, which disappears after adding dilute HCl.

2.4 Antimicrobial Activity (Disc Diffusion Method)

²¹ *Escherichia coli* and *Staphylococcus aureus* were ²⁴ used to evaluate antimicrobial activity via the disc diffusion method. Inocula were prepared in sterile nutrient broth and incubated at 37°C. Nutrient Agar Media (NAM) was prepared and sterilized. Using the pour plate method, 1 mL of inoculum was added to Petri dishes, followed by 15 mL of NAM. ⁸ Sterile filter paper discs loaded with 20 µL of Triphala extract were placed on the solidified media and incubated at 37°C for 24 hours. Zones of inhibition were measured in millimeters to assess antimicrobial efficacy.

¹¹

2.5 Antioxidant Activity (DPPH Method)

Antioxidant activity was determined using the DPPH free radical scavenging assay. A 0.004% DPPH methanolic solution and ascorbic acid (standard) were prepared in varying concentrations (20–100 µg/mL). Triphala extract was similarly diluted. Each sample (2 mL) was mixed with 0.5 mL DPPH solution and incubated in the dark for 10 minutes. Absorbance was measured at 517 nm.

Calculation of Antioxidant Activity

⁶

The percentage of DPPH radical scavenging activity was calculated using the following formula:

$$\% \text{ Inhibition} = \frac{(A_0 - A_t)}{A_0} \times 100$$

- A_0 = absorbance of the control
- A_t = absorbance in the presence of sample or standard

⁵

All tests were performed in triplicate, and the results were expressed as mean ± standard deviation.

3. RESULT

3.1 Phytochemical Screening

⁴ Preliminary phytochemical screening of Triphala extract ²⁵ using various solvents revealed the presence of several bioactive compounds (Table 1). The ethanolic extract exhibited the presence of alkaloids, carbohydrates and glycosides, tannins, saponins, terpenoids, and phlobatannins.

Chloroform extract showed positive results for alkaloids and terpenoids, while the aqueous extract confirmed the presence of tannins, saponins, and terpenoids. No phytochemicals were detected in the petroleum ether extract.

These findings affirm the presence of secondary metabolites in Triphala, which are known for their therapeutic potential. The ethanolic extract, being the richest in phytochemicals, was subjected to further antimicrobial and antioxidant testing.

Phytochemical tests	Triphala			
	Chloroform	Ethanol	Petroleum ether	Aqueous
Alkaloids (Mayers Reagent)	+	+	-	-
Carbohydrates & glycosides (Fehling Solution)	-	+	-	-
Phenolic compounds & tannins (Ferric Chloride solution)	-	+	-	+
Proteins/Aminoacids (Ninhydrin test)	-	-	-	-
Saponins	-	+	-	+
Terpenoid	+	+	-	+
Phlobatannins	-	+	-	-
Flavanoïdes	-	-	-	-

3.2 Antimicrobial Activity

The antimicrobial efficacy of Triphala extracts was evaluated using the disc diffusion method against *Escherichia coli* and *Staphylococcus aureus*. The ethanolic extract exhibited the highest antimicrobial activity, with zones of inhibition measuring 10 mm against *E. coli* and 9 mm against *S. aureus*. The aqueous extract showed moderate activity with inhibition zones of 9 mm

and 7 mm, respectively. The chloroform and petroleum ether extracts displayed negligible or no activity.

When compared with the standard antibiotic streptomycin, which showed inhibition zones of 18 mm (*E. coli*) and 13 mm (*S. aureus*), the ethanolic extract demonstrated moderate but significant antimicrobial potential. This activity may be attributed to the presence of phenolic compounds and other bioactive constituents in the extract.

Extract Type	Microorganism	Zone of Inhibition (mm)
Ethanolic	<i>E. coli</i>	10
	<i>S. aureus</i>	9
Aqueous	<i>E. coli</i>	9
	<i>S. aureus</i>	7
Streptomycin	<i>E. coli</i>	18
	<i>S. aureus</i>	13




Fig: Zone of inhibition of triphala for *S. aureus*

3.3 Antioxidant Activity

The antioxidant potential of Triphala was evaluated using the DPPH free radical scavenging assay. Both ethanolic and aqueous extracts demonstrated dose-dependent radical scavenging activity, although lower than the standard antioxidant, ascorbic acid.


At the highest tested concentration (100 $\mu\text{g/mL}$), the aqueous extract showed a scavenging activity of 53.6%, followed by the ethanolic extract with 44.2%, compared to 67.6% for ascorbic acid. This suggests a significant antioxidant potential, likely due to the presence of phenolic compounds, flavonoids, and other phytochemicals.



Fig: Antioxidant activity of extract of triphala

16

Concentration ($\mu\text{g/mL}$)	Ethanolic Extract (%)	Aqueous Extract (%)	Ascorbic Acid (%)
20	12.3	16.2	19.2
40	19.5	24.15	28.15
60	26.3	31.3	39.6
80	37.3	42.3	51.7
100	44.2	53.6	67.6



2

These findings highlight Triphala as a promising source of natural antioxidants with potential applications in food and pharmaceutical industries. The presence of flavonoids, tannins, alkaloids, glycosides, and phenolics may contribute to its free radical scavenging activity. Further isolation and characterization of the active compounds are warranted to understand their mechanisms and enhance the formulation of polyherbal therapeutics.

4. DISCUSSION

This study validated the traditional medicinal use of *Triphala* by confirming its phytochemical richness, antimicrobial, and antioxidant properties through various solvent extractions. The ethanolic extract exhibited the highest concentration of bioactive compounds—including alkaloids, tannins, glycosides, and phenolics—correlating with its superior antimicrobial and antioxidant effects. Antimicrobial activity was strongest against *E. coli* and *S. aureus* with the ethanolic extract, while antioxidant potential was observed in both ethanolic and aqueous extracts, though less than that of ascorbic acid. These findings highlight *Triphala's* therapeutic potential and suggest its suitability as a natural source for antimicrobial and antioxidant agents.

5. CONCLUSION

The present study affirms the pharmacological potential of **Triphala**, validating its traditional Ayurvedic applications with modern scientific evidence. Among the various solvent extracts analyzed, the **ethanolic extract** demonstrated the most potent antimicrobial and antioxidant activities, likely due to its rich phytochemical content, especially phenolic compounds, tannins, and glycosides. The findings support the role of *Triphala* as a promising **natural alternative to synthetic antimicrobial and antioxidant agents**, with possible applications in the pharmaceutical and nutraceutical industries. However, **further chemical characterization and in vivo investigations** are essential to isolate active constituents and determine their mechanisms of action and clinical safety profiles.

6. REFERENCES

- [1] Bhowmik D, Tiwari P, Tripathi K, Kumar S. Traditional Indian memory enhancer herbs and their medicinal importance. *Ann Biol Res*. 2010;1(1):41–46.
- [2] Biradar Y, Sharma S, Khendalwal KR. Preparation method of optimization and physicochemical evaluation of traditional formulation *Triphala*. *Indian J Tradit Knowl*. 2006;6(2):292–297.
- [3] Chaitali H, Nikhil S, Bhamicee S, Bhamicee SB. Cytotoxicity screening of selected Indian medicinal plants using Brine-Shrimp lethality bioassay. *Adv Nat Appl Sci*. 2010;4(3):389–395.
- [4] Dhanalakshmi S, Kumar R, Sundaramahalingam P, Naraganaperumal Jeya. Antioxidant property of *Triphala* on cold stress induced oxidative stress in experimental mice. *J Health Sci*. 2006;52(6):843–842.
- [5] Girish C, Koner B, Jayanthi S, Rao KR, Rajesh B, Pradhan SC. Hepatoprotective activity of six polyherbal formulations in paracetamol induced liver toxicity in mice. *Indian J Med Res*. 2009;129:569–578.
- [6] Gupta D, Mradu R. Therapeutic uses of the polyherbal drug *Triphala* in geriatric diseases. *Int J Pharm Bio Sci*. 2010;1(2):145–158.
- [7] Gupta M. Therapeutic uses of the polyherbal drug *Triphala* in geriatric diseases. *Int J Pharm Bio Sci*. 2010;1(2):267–285.
- [8] Hussin A. Adverse effects of herbs and drug-herbal interactions. *Malays J Pharm*. 2001;1(2):39–44.
- [9] Kaur S, Jaggi RK. Antinociceptive activity of chronic administration of different extracts of

Terminalia bellirica Roxb and *Terminalia chebula* Retz. fruits. Indian J Exp Biol. 2010;48:925–930.

- [10] Khan KH. *Terminalia chebula* reduces the oxidative stress induced by *Salmonella typhimurium* in mice and may reduce the risk of getting typhoid. Adv Biol Res. 2009;3:1–2.
- [11] Khan K, Hayat. The effect of regular intake of *Terminalia chebula* on oxidative stress in mice originated from *Salmonella typhimurium*. EurAsian J Biosci. 2009;3:113–121.
- [12] Lagae L, Buyse G, Ceulemans B, Claeys P, Dedeurwaerdere S, De Meirleir L, Hauman R, Janssen A. Anti-epileptogenesis research: the clinical relevance. Acta Neurol Belg. 2003;103:78–82.
- [13] Masola SN, Mosha RD, Wambura PN. Assessment of antimicrobial activity of crude extracts of stem and root barks from *Adansonia digitata* (Bombacaceae) (African baobab). Afr J Biotechnol. 2009;8:5076–5083.
- [14] Mahajan A, Pai D, Nandini R. Development and validation of HPLC method for quantification of phytoconstituents in Haritaki Churna. Int J ChemTech Res. 2011;3:329–336.
- [15] Mahesh R, Ramalingam B, Bhuvana S, Begum HV, Vava M. Effect of *Terminalia chebula* aqueous extract on oxidative stress and antioxidant status in the liver and kidney of young and aged mice. Cell Biochem Funct. 2009;27:358–363.
- [16] Modak M, Dixit P, Londhe J, Ghaskadbi S, Devasagayam TPA. Indian herbs and herbal drugs used for the treatment of diabetes. J Clin Biochem Nutr. 2007;40:163–173.
- [17] Preethi R, Vimal V, Devanathana V, Loganathan M. Antimicrobial and antioxidant efficiency of some medicinal plants against food borne pathogens. Adv Biol Res. 2010;4(2):122–125.
- [18] Rahman S, Akbar M, Howlader A, Jabber A. Antimicrobial and cytotoxic activity of the alkaloids of Amlaki (*Emblica officinalis*). Pak J Biol Sci. 2009;12(16):1152–1155.
- [19] Singh M, Sharma CS. Wound healing activity of *Terminalia chebula* in experimentally induced diabetic mice. Int J PharmTech Res. 2009;1(4):1267–1270.
- [20] Sharma US, Sharma U, Kumar U, Singh A, Singh N, Puspak J. Screening of *Terminalia bellirica* fruits extracts for its analgesic and antipyretic activities. Int J PharmTech Res. 2010;3(3):121–124.
- [21] Tambekar DH, Dahikar SB, Lahare MD. Antibacterial potentials of some herbal preparations available in India. Res J Med Med Sci. 2009;4(2):224–227.
- [22] Tambekar DH, Dahikar SB. Exploring antibacterial potential of some Ayurvedic preparations to control bacterial enteric infections. J Chem Pharm Res. 2010;2(5):494–501.

"Antimicrobial and Antioxidant Activity of Triphala: An Ayurvedic Formulation"

ORIGINALITY REPORT

21%

SIMILARITY INDEX

18%

INTERNET SOURCES

16%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	mail.journalcra.com Internet Source	3%
2	www.researchgate.net Internet Source	1%
3	www.mdpi.com Internet Source	1%
4	"Medicinal Plants and their Bioactive Compounds in Human Health: Volume 1", Springer Science and Business Media LLC, 2024 Publication	1%
5	www2.mdpi.com Internet Source	1%
6	www.xjcistanche.com Internet Source	1%
7	Shahid Akbar. "Handbook of 200 Medicinal Plants", Springer Science and Business Media LLC, 2020 Publication	1%
8	Submitted to Community College System of New Hampshire Student Paper	1%
9	Submitted to Sharda University Student Paper	1%
10	Submitted to University of Newcastle Student Paper	1%

11	Submitted to Higher Education Commission Pakistan Student Paper	1 %
12	Nayeema Akther, A.S. Shawl, Sarwat Sultana, B.K. Chandan, Mymoona Akhter. "Hepatoprotective activity of Marrubium vulgare against paracetamol induced toxicity", Journal of Pharmacy Research, 2013 Publication	1 %
13	repository-tnmgrmu.ac.in Internet Source	1 %
14	repository.sustech.edu Internet Source	1 %
15	Baoru Yang, Pengzhan Liu. "Composition and Biological Activities of Hydrolyzable Tannins of Fruits of Phyllanthus emblica", Journal of Agricultural and Food Chemistry, 2014 Publication	1 %
16	pharmacologyonline.silae.it Internet Source	1 %
17	www.jetir.org Internet Source	1 %
18	www.yumpu.com Internet Source	1 %
19	Bukar, AM, MA Isa, HS Bello, and AS Abdullahi. "Antibacterial Activity of Aqueous And Ethanol Leaf Extracts Of Vernonia Amygdalina On Selected Species Of Gram Positive And Gram Negative Bacteria", International Journal of Environment, 2013. Publication	<1 %
20	phcogj.com Internet Source	<1 %

21	updatepublishing.com Internet Source	<1 %
22	www.frontiersin.org Internet Source	<1 %
23	www.jabonline.in Internet Source	<1 %
24	www.jocpr.com Internet Source	<1 %
25	www.phcogj.com Internet Source	<1 %
26	www.thepharmajournal.com Internet Source	<1 %
27	Ameya Joshi, Akshay Baheti, Manish Wani, Ranjeet Nimbalkar. "A review through therapeutic attributes of Ayurvedic formulation mashi", Journal of Ayurveda and Integrative Medicine, 2021 Publication	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On