

Wilkie's Syndrome: Case Report and Surgical Approach in a Patient with High Intestinal Obstruction

by Jana Publication & Research

Submission date: 17-Jul-2025 12:21PM (UTC+0700)

Submission ID: 2690332272

File name: IJAR-52833.docx (33.19K)

Word count: 854

Character count: 5320

Wilkie's Syndrome: Case Report and Surgical Approach in a Patient with High Intestinal Obstruction²¹

Abstract

² Superior mesenteric artery syndrome (SMAS), also known as Wilkie's syndrome, is a rare cause of upper intestinal obstruction due to duodenal compression between the abdominal aorta and the superior mesenteric artery. This condition arises from a decreased aortomesenteric angle and distance, typically following significant weight loss. We report the case of a 38-year-old male with upper intestinal obstruction symptoms and unintentional weight loss. A diagnosis of SMAS was confirmed via contrast-enhanced CT revealing a 17° aortomesenteric angle and a 7 mm distance. ⁵ The patient underwent a laparoscopic duodenojejunostomy in a Roux-en-Y configuration with an uneventful recovery. ⁴ This report highlights the importance of early recognition and surgical intervention in SMAS to avoid complications.

Keywords

¹⁹ Wilkie's syndrome; intestinal obstruction; superior mesenteric artery; duodenojejunostomy; laparoscopic surgery

¹¹ Introduction

Superior mesenteric artery syndrome (SMAS) is an uncommon but significant condition characterized by the external compression of the third portion of the duodenum between the superior mesenteric artery and the aorta. This narrowing is primarily due to rapid loss of retroperitoneal fat, reducing the normal aortomesenteric angle (38–65°) and distance (10–33 mm). ⁸ SMAS has an estimated incidence of 0.013–0.3% in the general population [1]. The clinical presentation is often nonspecific, with symptoms including early satiety, postprandial fullness, nausea, vomiting, and weight loss, which delays diagnosis [2,3]. ¹⁷ Imaging studies, especially contrast-enhanced CT, are essential for identifying anatomical criteria of SMAS. This condition may be secondary to rapid weight loss due to trauma, burns, anorexia nervosa, or chronic illness [4]. ¹³

Case Presentation

⁶ A 38-year-old male presented with a 7-day history of intense abdominal pain, nausea, vomiting, and absence of bowel movements. His history included tobacco, marijuana, and methamphetamine use, with a reported weight loss of approximately 20 kg over the previous months. Initial laboratory tests revealed leukocytosis (20.8 x10⁹/L), serum creatinine of 1.34 mg/dL, CRP 1.69 mg/dL,

and procalcitonin of 1.28 ng/mL. Contrast-enhanced abdominal CT showed gastric and proximal duodenal distension with abrupt caliber change at the third portion, and an aortomesenteric angle of 17°, along with a 7 mm distance, confirming SMAS. Initial conservative management with nasogastric decompression and nutritional support was attempted for 5 days without sufficient improvement. The patient subsequently underwent elective laparoscopic duodenojejunostomy with Roux-en-Y reconstruction. The third portion of the duodenum was mobilized (Kocher maneuver), and an anastomosis was created using a 45 mm linear stapler and reinforced with V-Loc 3-0 suture and serosal Lambert stitches. Estimated intraoperative blood loss was 10 mL. Postoperative recovery was uneventful: the contrast study at 24 hours showed no leak, oral intake resumed at 72 hours, and discharge occurred on postoperative day 5.

Discussion

SMAS is a rare condition, frequently underdiagnosed due to the vague nature of gastrointestinal symptoms. In many cases, weight loss precedes the onset of symptoms, as in our patient, highlighting the role of nutritional depletion in pathogenesis [5,6]. CT and MRI are valuable tools for measuring the aortomesenteric angle and distance. An angle <22° and distance <8 mm are diagnostic indicators [7]. Although conservative treatment is recommended initially (nutritional support, positional therapy), surgery becomes necessary when symptoms persist or complications arise. Laparoscopic duodenojejunostomy is considered the gold standard due to lower morbidity, quicker recovery, and favorable cosmetic outcomes [8,9]. Studies have demonstrated high success rates with laparoscopic approaches, with resolution of symptoms and early return to oral intake [10]. Our case supports these findings and emphasizes the importance of timely surgical referral in nonresponding patients.

Conclusion

SMAS should be considered in patients with upper gastrointestinal obstruction symptoms and a history of rapid weight loss. Timely diagnosis and treatment, including surgical intervention when indicated, can result in excellent outcomes. Laparoscopic duodenojejunostomy remains the preferred approach for definitive management in selected patients.

Informed Consent

Written informed consent was obtained from the patient for publication of this case and accompanying images.

Conflict of Interest

The authors declare no conflict of interest.

Funding

This study received no external funding.

References

1. Welsch T, Büchler MW, Kienle P. Recalling superior mesenteric artery syndrome. *Dig Surg*. 2007;24(3):149-156.
2. Merrett ND, Wilson RB, Cosman P, Biankin AV. Superior mesenteric artery syndrome: diagnosis and treatment strategies. *J Gastrointest Surg*. 2009;13(2):287-292.
3. Biank V, Werlin S. Superior mesenteric artery syndrome in children: a 20-year experience. *J Pediatr Gastroenterol Nutr*. 2006;42(5):522-525.
4. Neri S, Signorelli SS, Mondati E, et al. Ultrasound imaging in diagnosis of superior mesenteric artery syndrome. *J Intern Med*. 2005;257(4):346-351.
5. Welsch T, Kienle P, Schäfer M. SMAS: current diagnostic and therapeutic approach. *Langenbecks Arch Surg*. 2010;395(6):681-688.
6. Shiu JR, Chao HC, Luo CC, et al. Clinical and nutritional outcomes in children with SMA syndrome. *Pediatr Neonatol*. 2012;53(4):252-257.
7. Unal B, Aktas A, Kemal G, et al. Superior mesenteric artery syndrome: CT and ultrasonography findings. *Diagn Interv Radiol*. 2005;11(2):90-95.
8. Gersin KS, Heniford BT. Laparoscopic duodenojejunostomy for SMAS. *J Laparoendosc Adv Surg Tech A*. 1998;8(6):367-373.
9. Lee TH, Lee JS, Jo Y, et al. Laparoscopic management of SMAS: case series and review. *Ann Surg Treat Res*. 2014;86(3):139-142.
10. Palanivelu C, Rangarajan M, Senthilkumar R, Parthasarathi R. Laparoscopic duodenojejunostomy for SMAS: review and case report. *JSLS*. 2006;10(4):531-534.

Wilkie's Syndrome: Case Report and Surgical Approach in a Patient with High Intestinal Obstruction

ORIGINALITY REPORT

37%

SIMILARITY INDEX

22%

INTERNET SOURCES

33%

PUBLICATIONS

13%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to Teerthanker Mahaveer University

Student Paper

4%

2

Jair Díaz-Martínez, Francisco Tomás Hizojo-Aloé, Renata Jimena Rivera-Chávez, Nidia Alejandra González-Hernández.

"Misdiagnosed superior mesenteric artery syndrome due to diabetic gastroparesis. Case report and literature review", International Journal of Surgery Case Reports, 2024

Publication

3%

3

jfmo-dz.net

Internet Source

3%

4

"Clinical Vignettes/Case Reports - Esophagus", The American Journal of Gastroenterology, 2014.

Publication

2%

5

Mohammad Maraqaq, Mohammad Farid Al Tamimi, Sara Abu Aram, Shukran Hmedat, Maimana Abueisha, Yazan Hmeedat.

"Possible familial predisposition in superior mesenteric artery syndrome with surgical management: A case report", Medicine, 2025

Publication

2%

6	Carl Vaughan, Cornelius C Cronin, Elizabeth Kenny Walsh, Michael Whelton. "The Jarisch-Herxheimer reaction in leptospirosis", Postgraduate Medical Journal, 1994 Publication	2%
7	Gabriela Lozano, Eleanor Dunlap, Khanjan H. Nagarsheth, Daniel Shouhed. "Chapter 8 Gastrointestinal Compression Associated with Median Arcuate Ligament Syndrome: Superior Mesenteric Artery Syndrome", Springer Science and Business Media LLC, 2025 Publication	2%
8	acamedicine.org Internet Source	2%
9	surgrad.com Internet Source	1%
10	Gautam A. Agrawal. "Multidetector Row CT of Superior Mesenteric Artery Syndrome", Journal of Clinical Gastroenterology, 01/2007 Publication	1%
11	M. E. C. Bakker, R. van Delft, N. A. L. Vaessens, P. R. R. Rosias. "Superior mesenteric artery syndrome in a 15-year-old boy during Ramadan", European Journal of Pediatrics, 2013 Publication	1%
12	rrtjournal.biomedcentral.com Internet Source	1%
13	Booka, Eisuke, Hirofumi Kawakubo, Ken Ishii, Tomohiro Hikata, Kazumasa Fukuda, Rieko Nakamura, Tsunehiro Takahashi, Norihito	1%

Wada, Yoshiyuki Ishii, Hiroya Takeuchi, and Yuko Kitagawa. "Superior Mesenteric Artery Syndrome Caused by Massive Lumbar Osteophytes : A Case Report", Spine, 2015.

Publication

14

Jeune, Florence. "A rare cause of obstructive jaundice and gastric outlet obstruction", World Journal of Gastrointestinal Surgery, 2013.

Publication

1 %

15

Nariman Abol Oyou, Muayad Kadhim, John P. Dormans. "Late-onset superior mesenteric artery syndrome four years following scoliosis surgery – a case report", SICOT-J, 2015

Publication

1 %

16

Raluca Cristina Apostu, Lucian Chira, Doina Colcear, Andrei Lebovici, Georgiana Nagy, Radu Razvan Scurtu, Radu Drasovean. "Wilkie's syndrome as a cause of anxiety-depressive disorder: A case report and review of literature", World Journal of Clinical Cases, 2022

Publication

1 %

17

link.springer.com

Internet Source

1 %

18

www.karger.com

Internet Source

1 %

19

www.mdpi.com

Internet Source

1 %

20

Jihui Li. "Laparoscopic Roux-en-Y Duodenojejunal Bypass for Superior Mesenteric Artery Syndrome : Case Reports

1 %

and Review of the Literature", Surgical
Laparoscopy Endoscopy & Percutaneous
Techniques, 12/2011

Publication

21

Luis A Chirinos, Danitza Lukac, Mateo E
Garland, Jorge André Céspedes, Victor M
Salcedo. "Weight Loss-associated Superior
Mesenteric Artery Syndrome, an Unusual
Condition in a Peruvian Patient: A Case
Report", Cureus, 2018

Publication

1 %

22

Myung Jin Oh. "Superior Mesenteric Artery
Syndrome Combined with Renal Nutcracker
Syndrome in a Young Male: A Case Report",
The Korean Journal of Gastroenterology, 2017

Publication

1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On