

The Effect of the Educational Program Based on Roy's Adaptation Model on the Quality of Life for

by Jana Publication & Research

Submission date: 28-Jul-2025 12:33PM (UTC+0700)

Submission ID: 2692518556

File name: IJAR-53022.docx (93.82K)

Word count: 6699

Character count: 37903

¹⁵The Effect of the Educational Program Based on Roy's Adaptation Model on the Quality of Life for Patients with Thalassemia

ABSTRACT

Thalassemia is an autosomal recessive blood disorder. ⁶⁵The chronic nature of the disease, along with its complications and treatment, causes multiple ⁶¹physical, psychological, and social problems that affect ³¹the quality of life in these patients. **Aim of the study:** Evaluate the effect of provided educational program based on Roy's adaption model on quality of life for patients with thalassemia. ²**Research design:** A quasi experimental design. **Setting:** General medicine department and the blood transfusion unit in Fayoum university hospital. **Methods:** A purposive sample of 80 adult patients from previously mentioned setting, allocated randomly into two equal groups (40 patients in each). **Tools:** four tools were utilized, included: Interview patient's assessment questionnaire, patients knowledge assessment questionnaire, ²⁰World Health Organization Quality of Life-BREF (WHOQOL-BREF) questionnaire ⁷and Roy's Adaptation Model questionnaire. **Results:** There was a statistical significant difference between both groups as regarding knowledge, quality of life and adaptation level (P-value <0.001). **Conclusion:** Application of the educational program based on Roy's adaptation model had a positive effect on the outcomes of patients with thalassemia. ³**Recommendations:** Nurses should apply the educational instrument that would help patients improve their adaptation to reach positive outcomes.

⁵⁴**Key words:** Adaptation, Quality of life, Thalassemia.

Introduction

Thalassemia ⁴is an autosomal recessive blood disorder and is characterized by abnormal hemoglobin production. Owing to genetic abnormalities, the production of one or more types of globin peptide

chains is diminished or entirely lacking, which results in hemolytic anemia,²⁵ fatigue, and a range of complications related to poor oxygen delivery to tissues (Thein & Rees, 2025).

⁴Thalassemia is one of the most common genetic diseases in the world. Worldwide, it is estimated that 7 % of the population carries the thalassemia gene.⁴ The World Health Organization (WHO) has listed thalassemia as a major public health problem. Thalassemia has clear regional characteristics and group specificity, mainly occurring in the Mediterranean region, Southeast Asia, India, the Middle East, Africa, and southern China. With the flow of migrants, thalassemia has spread from the Mediterranean region, Africa, Asia to Europe, America, and Australia, becoming a serious global public health problem (Su, et al., 2025).

⁴Individuals with mild forms of thalassemia might not exhibit any symptoms, whereas those with severe cases require frequent blood transfusions and iron chelation therapy to sustain life or potentially a hematopoietic stem cell transplant for a cure. The thalassemia treatment and monitoring process is cumbersome, expensive, seriously affects the quality of life of patients, and has a heavy disease burden. These impose substantial financial and social burdens on patients, families and society. Therefore, early, timely, and accurate screening is crucial (Setiawan, et al., 2025).

Patients with thalassemia often experience various²⁸ physical, psychological, and social challenges that can significantly impact their²⁰ overall quality of life. Improving the quality of life for these patients requires comprehensive and multidimensional care that extends beyond pharmacological treatment (Wangi, et al., 2025).

The Roy⁶ Adaptation Model Roy's adaption model developed by Sister Callista Roy, offers a holistic nursing framework that focuses on⁴² promoting adaptation in four key domains: physiological, self-concept, role function, and interdependence.³³ This model views individuals as adaptive systems responding to internal and external stimuli and it emphasizes the role of nursing interventions in facilitating effective coping mechanisms and enhancing adaptive responses (El Arab, et al., 2025).

Educational programs based on the Roy's adaption model have shown promise in promoting patient-centered care by empowering individuals with knowledge, skills, physical, psychosocial and environmental support. For patients with thalassemia, such programs may help improve physical practices, psychological resilience, social integration, environmental adaptation and ultimately, quality of life. So, this study aims to evaluate the effect of provided educational program¹⁵ based on Roy's adaption model on quality of life for patients with thalassemia. By addressing the multidimensional needs of these patients⁶⁴ through a structured and theory-based educational approach, the program seeks to foster improved adaptation and well-being of patients with thalassemia (Hassan, et al., 2025).

Significance of Study:

Thalassemia is a lifelong hereditary disorder that imposes considerable physical, emotional, and social burdens on patients, often leading to a diminished quality of life. Despite advances in

medical management, many patients continue to struggle with complications arising from their condition, including frequent hospital visits, dependency on transfusions, and psychological stress. Therefore, addressing the holistic needs of patients is essential (Al_Hamadiny, 2024)

³⁴ **From the researcher point of view and experience.** This study is significant as it applies ⁶⁹ the Roy Adaptation Model, a well-established ⁴¹ nursing framework that emphasizes adaptive responses in four domains: physiological, self-concept, role function, and interdependence. By integrating Roy Adaptation Model into an educational program tailored for thalassemia patients. The findings can provide evidence-based support for implementing theory-driven nursing interventions that empower patients to adapt to their chronic condition. Moreover, the outcomes may inform healthcare professionals and policymakers about the value of structured educational programs ⁷³ in improving quality of life, especially in resource-limited settings such as Egypt where thalassemia remains a public health challenge.

⁷² **Aim of study:**

The present study aimed to evaluate ¹⁵ the effect of an education program based on Roy's adaptation model on the quality of life for patients with thalassemia through:

1. Assess level of quality of life for patients with thalassemia.

2. Design ¹²educational program based on Roy's adaption model and patients' basic quality of life assessment needs for patients with thalassemia.
3. Implement the ¹²educational program based on Roy's adaption model and patients' basic quality of life assessment needs for patients with thalassemia.
4. Evaluate ¹²the effect of provided educational program based on Roy's adaption model on quality of life for patients with thalassemia.

Research Hypothesis:

H0: Patients who will receive educational program based on Roy's adaption model will not exhibit better physical, psychological, social and environmental well-being more than those who will not receive the program.


H1: Patients who will receive educational program based on Roy's adaption model will be exhibit better physical, psychological, social and environmental well-being more than those who will not receive the program as measure by **tool (III)**.

Theoretical Framework

This study is grounded in ⁶Roy's Adaptation Model (RAM), which views the individual as a biopsychosocial ²²being in constant interaction with a changing environment. According to RAM, health is

defined as a state and process of being and becoming integrated and whole. The goal of nursing is to promote adaptation in four adaptive modes: **physiological, self-concept, role function, and interdependence**. (Kumar, 2022).

This conceptual framework illustrates the cyclical process of Roy's Adaptation Model, where ⁴⁷ the nurse continuously assesses, diagnoses, plans, implements, and evaluates the patient's adaptation to promote optimal health outcomes. The Roy Adaptation Model (RAM) can be seamlessly integrated with the nursing process to provide a comprehensive and holistic approach for patient care. The model emphasizes the importance of assessing the individual's adaptation levels, identifying adaptation problems, and developing interventions to enhance adaptation. This nursing process based on the RAM model focuses on promoting adaptation and improving health outcomes by addressing the patient's physiological, emotional, and social needs (Majeed, et al., 2020).






Figure (1): Nursing process according to Roy's theoretical model for patients with thalassemia designated by the researcher.

Subject and Methods

Research Design:

A ¹⁰Quasi-experimental design identifies a comparison group that is as similar as possible to the treatment group in terms of baseline (pre-intervention) characteristics. The comparison group captures what would have been the outcomes if the program/policy had not been implemented. Hence, the program or policy can be said to have caused any difference in outcomes between the treatment and comparison groups.

Setting:

This ²³study was conducted in the General Medicine Department and the Blood Transfusion Unit at Fayoum University Hospital. The General Medicine Department is located on the 3rd floor and consists of twenty rooms—ten ¹¹for male patients and ten for female patients—with each room containing two beds. The total capacity of the department is forty beds. The Blood Transfusion ⁵⁰Unit is located on the ground floor of the hospital. It consists of a large room that includes a healthcare provider's desk and eight beds for patients

Subject:

⁵A purposive sample of 80 adult patients from both genders. They were recruited from the previously mentioned setting and divided randomly into two equal groups ^{study} and ^{control} (40 patients for each group).

Inclusion criteria:

- Patients from both genders agree to participate in the study.
- Patients diagnosed with thalassemia.
- Patient able to communicate verbally..

1

Exclusion criteria:

- Psychiatric patients.
- Patients suffering from other types of anemia..

60

Sample size:

Based on sample size equation, ¹¹ the sample size was calculated by adjusting the power of the test to 80% and the confidence interval to 95% with margin of error accepted adjusted to 5% and a known total population of 80 patients using the following equation: (Chow & Wang, 2007).

$$n = \frac{N \times p(1-p)}{\left[\left(N-1 \right) \times \left(d^2 \div z^2 \right) \right] + p(1-p)}$$

$$N \times p(1-p) = (180 \times (0.20 \times (1 - 0.20))) /$$

$$N-1 = (180-1) \times$$

$$= 0.0025 / 3.8416 +$$

$$d^2 / z^2$$

$$\frac{30}{p(1-p)} = 0.20 \times (1 - 0.20)$$

$$N = 80$$

- P= 0.5
- N= Total population
- Z= Z value “1.96”
- D= Standard Error
- n= sample size

24

Tools of Data Collection:

Four tools were used to collect the data according to the following :

Tool I: Interview Patient's assessment Questionnaire:

This tool was developed by the researcher in English and Arabic language after reviewing relevant recent literatures (Haq, et al., 2017) to collect baseline data pertinent to the current study. It was consisting of two parts as follow:

Part I: Patients' Socio-demographic data: This part concerned with assesses patients' socio-demographic characteristics which include patients' age, sex, marital status, educational level and occupation, residence and income. It contains seven questions in form of multiple-choice questions.

Part II: Patients' present, past and family history: This part aimed to assess patients' medical history that includes information about the present history, duration of the disease, past history, previous hospitalization and family history of thalassemia. It contains 12 questions in form of closed, open ended and multiple-choice questions.

Tool II: patients' knowledge assessment questionnaire.

It was developed by the researcher in the light of related literatures (Haq, et al., 2017; Olwi, et al., 2018; Man, et al., 2019) to assess knowledge of patient regarding thalassemia. It includes questions about definition, causes, signs and symptoms, diagnostic test, prognosis, complications, treatment, iron therapy, diet and activity. It consisted of 40 questions in form of multiple-choice questions.

Scoring system: Patient's knowledge assessment questionnaire consisted of 40 questions, the correct answers were predetermined according to literature review, a correct answer was scored 1 point and incorrect answer was scored 0 point, and satisfactory level was detected based on statistical analysis as following:

- ☐ Satisfactory knowledge level $\geq 75\%$

- Unsatisfactory knowledge level < 75 %

Tool (III): World Health Organization Quality of Life-BREF (WHOQOL-BREF) questionnaire:

The WHOQOL-BREF was developed by GENEVA, 1996. It was used to assess quality of life for patient with thalassemia. It was translated into an Arabic language and back translated into English. The tool included (26) questions. The items of the WHOQOL-BREF questionnaire are divided into 4 domains (Physical health domain, psychological domain, social relationships domain and environmental domain).

Scoring system:

WHOQoL-BREF composed of 26 items, which assess the following broad domains: physical health (7 items), psychological health (6 items), social relationships (3 items), and environmental factors (8 items), as well as the overall quality of life and general health (2 items). Each item is rated on a five-point likert scale that ranged from 1 to 5 where 1 is the lowest score and 5 are the highest score.

The score were categorized according Skevington et al., (2004) into:

- Good quality of life: $\geq 75\%$
- Average quality of life: $50 < 75\%$
- Poor quality of life: $< 50\%$

Tool (IV): Roy's Adaptation Model questionnaire:

It was adapted from (DeSanto& Fawcett, 2009).It is consisted of four modes physiological, self-concept, role function, and interdependence 10 items for each mode. Each mode composed of 10 questions which are used for assessment level of adaptation among studied patients. The answers was measured and given a score based on five-point Likert scale as the following: Strongly Disagree (1) Disagree (2), Neither Agree or Disagree (3), Agree (4), and Strongly Agree (5).The

¹total grade for this questionnaire is 200 grades that represent 100%. The total ⁷score ranged from 40- 200 with the higher score indicated more adaptive behavior for thalassemia patients.

The studied patients'adaptation can be classified according Abdelrahman et al.. (2023) as:

- High adaption: $\geq 75\%$
- Moderate adaption: $50 < 75\%$
- Low adaptation: $< 50\%$

¹**Validity:**

The validity of the developed tools was tested by a panel of five experts from adult health nursing department, faculty of nursing, Helwan University. The panel of experts was from different academic categories (four assist professors and one lecture). ¹The experts reviewed the tools for clarity, relevance, comprehensiveness, understanding, applicability and easiness for administration. ²Suggestions were given and modifications were done.

Reliability:

Reliability of the tool was tested to determine the extent to which the questionnaire items are related to each other. The Cronbach's alpha model, which is a model of internal consistency, was used in the analysis. ²Statistical equation of Cronbach's alpha reliability coefficient normally ranges between 0 and 1. Higher values of Cronbach's alpha (more than 0.7) denote acceptable reliability.

Ethical consideration:

An ethical approval to conduct the proposed study was obtained from the Scientific Research, Ethical Committee of the faculty of Nursing, Helwan University. An official permission was obtained from the administrative authority of the selected setting for the current study.

The researcher obtained consent from the studied patients, explaining the purpose and nature of the study, stating the possibility to withdraw at any time, confidentiality of data assured by the researcher by using codes to identify participants instead of names or any other personal identifiers.

Pilot study:

A Pilot study was carried out with 10% (not less than 10 patients) of the sample under study⁸ to test the applicability, clarity and efficiency of the tools, then the tools modified according to the results of the pilot study. Modifications included: rephrasing and rearrangement of some questions. After modification, the final form of the tools was developed. Patients who shared in pilot study excluded from the study sample.

Field Work:

Study was conducted within nine months from beginning of July² 2024 to the end of March 2025 and carried out through four phases according to nursing process: ⁷assessment, planning, implementing and evaluation.

I-Assessment phase:

During this phase, ⁸the researcher visited the selected setting regularly, four days per week, selected patient according to inclusion criteria, and then assigned them randomly to either a study or control group. ⁹Initial assessment was done by the researcher for all study subjects in study and control groups regarding to participants' age, gender, educational level, occupation, and health history. Determine whether the deficit is due to lack of needs, lack of skills, or limited ability. ³Data collection was held through structured interviews and medical record chart. During this phase each patient was assessed individually during follow up period, and data collection was filled by the researcher, by

using tools (I), tool (II), tool (III) and tool (IV) for study and control groups as follows:

Tool I: Was utilized to assess patients' socio-demographic characteristics and medical history data that filled for the study and control groups, it took around 10 minutes.

Tool II: To assess patient about knowledge regarding patients with thalassemia; it took around 15-20 minutes.

Tool III: To assess quality of life specific to patient with thalassemia, it took around 15-20 minutes.

Tool III: To assess level of adaptation among patient with thalassemia it took around 15-20 minutes.

⁵ **II- Planning and Design phase:**

This phase included analysis of the pre-test findings; where goals and outcomes are formulated that directly impact patient care. The researcher plan intervention, design the educational section's content according to the patient's needs. Detected needs, requirements and deficiencies were translated into the aim and objectives of the educational program sections in the form of guidelines booklet.

² **III- Implementation phase:**

Based on the results obtained from the interviewing and observational sheets, as well as literature review, the educational program was developed by the researcher. It was implemented immediately after the pre-test. This includes ³implementing the designated nursing guidelines for the study group (40) patient, in term of educational sessions, instructions and follow up. The program was divided into 5 sessions (2 theoretical and 3 practical sessions) each session was implemented in one day. The duration of each session varied, according to its contents as well as the clients' response.

IV- Evaluation phase:

Evaluation phase aimed to reassess patients after implementation of nursing program to identify progress in term of differences in patients' level of response from baseline. Evaluation was done by using the posttest questionnaire which was the same format of pre-test in order to compare the changes in patients' need, and practices, using assessment tools (II, III, IV) post 2 weeks and follow up 3month from program.

Results:

Table (1): Frequency and percentage distribution of Socio-demographic data for the studied patients. (n=80).

Socio-demographic data	Control group (n=40)		Study group (n=40)		x2	P-value
	N	%	N	%		
Age group						
18-24 Years	21	52.5	22	55	0.753	0.686
25-34 Years	9	22.5	11	27.5		
35-45 Years	10	25	7	17.5		
Mean±SD	27.8±8.83		27.3±7.07			
Gender						
Male	24	60	21	52.5	0.457	0.499
Female	16	40	19	47.5		
Social Status						
Single	29	72.5	29	72.5	2.39	0.495
Married	9	22.5	8	20		
Widowed	2	5	1	2.5		
Divorced	0	0	2	5		
Level of Education						
Can't read and write	20	50	21	52.5	1.66	0.312
Primary education	10	25	8	20		
Secondary education	7	17.5	8	20		
University education	3	7.5	3	7.5		
Occupation						
Work	15	25	13	27.5	1.16	0.2799

Doesn't work	25	75	27	72.5		
Place of residence						
Urban	10	25	11	27.5	0.065	0.799
Rural	30	75	29	72.5		
Income						
Sufficient	11	27.5	13	32.5	0.238	0.626
Insufficient	29	72.5	27	67.5		

*Significant at $P \leq 0.05$.

³ **Table (1)** illustrates that there was no statistical significant difference between study and control groups with p-value >0.05 , as regarding socio-demographic characteristics like; age, gender, social status, education level, occupation, place of residence, and income which indicated proper matching between groups in these variables.

Table (2): Frequency and percentage distribution of past health and family history for the studied patients. (n=80).

Past health history and family history	Control group (n=40)		Study group (n=40)		Test	P-value
	N	%	N	%		
Suffering from Chronic Disease:						
No	1	4.3	2	8.6	3.85	0.696
Diabetes	9	23.07	8	21.05		
High blood pressure	10	25.6	11	28.9		
Heart problems	15	38.4	17	44.7		
Chest sensitivity	0	0	1	2.6		
Diabetes, High blood pressure and Heart problems	2	5.1	2	5.2		
Diabetes and Heart problems	3	7.6	2	5.2		
Family history for thalassemia:						
Yes	30	75	29	72.5	0.065	0.799
No	10	25	11	27.5		
Degree of relationship:						
First	20	66.6	19	65.5	0.037	0.901
Second	10	33.3	10	34.4		

*Significant at $P \leq 0.05$.

Table (2) illustrated that there was no statistical significant difference between study and control groups as regarding past health history and family history chronic disease, previous surgery and family history.

Table (3): Frequency and Percentage Distribution of the patient's total knowledge for the two groups within pre, post, and follow up test.

Variable	Total Patients' Knowledge												χ^2	P-Value
	Pre				Post				Follow Up					
	Control group (n=40)		Study group (n=40)		Control group (n=40)		Study group (n=40)		Control group (n=40)		Study group (n=40)			
	N	%	N	%	N	%	N	%	N	%	N	%		
Unsatisfactory	25	62.5	23	57.5	22	55	0	0	20	50	1	2.5	32.72	0.00*
Satisfactory	15	37.5	17	42.5	18	45	40	100	20	50	39	97.5		
Test & P-Value	$\chi^2 = 0.208$ (0.648)				$\chi^2 = 30.34$ (0.00*)				$\chi^2 = 23.30$ (0.00*)					

*Significant at $P \leq 0.05$.

Table (3) showed that; there was no statistically significant difference in knowledge between the control and study groups at the pre-test, while there was a statistically significant difference in knowledge between the control and study groups at the post and follow up test.

Table (4): Frequency and Percentage Distribution of the patient's total quality of life for the two groups within pre, post, and follow up test.

Variable	Total Patient's Quality of Life												χ^2	P-Value
	Pre				Post				Follow Up					
	Control group (n=40)		Study group (n=40)		Control group (n=40)		Study group (n=40)		Control group (n=40)		Study group (n=40)			
	N	%	N	%	N	%	N	%	N	%	N	%		
Poor QoL	6	15	5	12.5	6	15	2	5	6	15	1	2.5	10.68	0.00*
Average QoL	32	80	30	75	33	82.5	32	80	32	80	34	85		
Good QoL	2	5	5	12.5	1	2.5	6	15	2	5	5	12.5		
Test & P-Value	$\chi^2 = 1.82$ (0.799)				$\chi^2 = 6.212$ (0.02*)				$\chi^2 = 8.21$ (0.01*)					

* Significant at $P \leq 0.05$.

Table (4): illustrated that there was a statistically significant difference between the two groups within post and follow up test. The control group remained unchanged across all phases.

Table (5): Frequency and percentage distribution of total patient's Roy Adaptation for two groups Modes during pre, post, and follow up phase (n=80).

Total patient’s Roy Adaptation Modes	Studied patients (n = 80)				χ^2	P- Value
	Control group (n=40)		Study group (n=40)			
	N	%	N	%		
Pretest:					0.057	0.972
Low adaptation.	14	35	13	32.5		
Moderate adaptation.	24	60	25	62.5		
High adaptation.	2	5	2	5		
Posttest:					23.69	0.00*
Low adaptation.	14	35	3	7.5		
Moderate adaptation.	24	60	21	52.5		
High adaptation.	2	5	16	40		
Follow up test:					45.35	0.00*
Low adaptation.	13	32.5	2	5		
Moderate adaptation.	25	62.5	3	7.5		
High adaptation.	2	5	35	87.5		

*Significant at $P \leq 0.05$.

Table (5) shows that, there was a highly statistically significant difference between the study and control group regarding their total patient's Roy Adaptation Modes in post, and follow up scores with p-value (p-value 0.00*), while there was no statistically significant difference between the study and control group regarding pretest with p-value (p-value 0.972).

Table (6): Correlation between patients' total knowledge, total quality of life and total Roy adaptation mode.

Items	Total Patients' Knowledge					
	Pre		Post		Follow Up	
	R	P	R	p	R	p
Total Quality of Life	0.279	0.01*	0.108	0.342	0.298	0.007*
Total Roy Adaptation Modes	0.338	0.00*	0.517	0.00*	0.581	0.000*

*Significant at $P \leq 0.05$.

Table (9) illustrates that; There was a significant positive correlation between total patients' knowledge and total Roy adaptation within three tests, while there was a significant positive correlation between total patients' knowledge and total quality of life within pretest and follow up test. There was a significant positive correlation between total patients' knowledge and total Roy adaptation mode within pretest, posttest and follow up test.

Discussion

In the present study, findings regarding to the patient's characteristics revealed that, the socio-demographic and medical characteristics of subjects in both study and control groups, were not significantly dissimilar; this means that the participants were selected from identical population of thalassemia patients with good random allocation obtained. The mean age of the studied patients in both study and control groups were between (18-24) years with mean age (27.8±8.83, 27.3±7.07) and more than half of them were males. From researcher point of view, the short life span is associated with iron overload, which may eventually affect the organs and lead to their failure.

This result is similar to Hossain et al., (2023) who reported that the mean age of the studied patients was (19.75 ± 8.02) their age group was between (14-24) years, in a study titled "Health-related quality of life among thalassemia patients in Bangladesh using the SF-36 questionnaire.

Regarding to social status, the study results showed more than two thirds of the studied patients in both study and control groups were single. These results may be due to most patients are fear of passing the disease to their children by heredity. These results is in the same line with Khodashenas et al., (2021) who revealed that majority of the participants were single in their study that entitled "Quality of life and related paraclinical factors in Iranian patients with transfusion-dependent thalassemia".

Concerning education level, occupation, residence and income of the studied subjects, half of studied patients in study and control groups, can't read and write and more than two thirds of patients doesn't work, live in rural areas and had insufficient income. This result may be due to that the thalassemia is a chronic disease which affect the patient's, physical, psychological and social status of patients and cause severe level of fatigue which interfere with the patient's ability to study is go through their education in addition to frequent hospitalization of patients due to the chronic nature of the disease and frequent receiving of treatment such as blood transfusion, which make patients in ability to work. Therefore, patients' income is insufficient.

This result ¹is similar to Hossain et al., (2023) who reported that the majority of the studied patients were unemployment and low income. This result also agree with Badur et al., (2021) ⁴⁰who revealed that more than half of patients were illiterate in their study that titled "Evaluation of the relationship between sociodemographic characteristics, quality of life, depression, drug compliance and biochemical parameters in patients with thalassemia major in Isparta".

In relation to past health history, chronic diseases in ⁴⁵this study showed that about more than one quarter of patients in study and control group suffer from diabetes mellitus and ³⁴hypertension and more than one third of patients had heart problem. These results may be due to pancreatitis which leads to diabetes mellitus and iron accumulation on the heart can cause cardiomyopathy. ¹These results is agree with Hamdy et al., (2021) who revealed that more than one third of patients had diabetes mellitus and cardiomyopathy in their study that titled "Assessment of quality of life among beta-thalassemia major patients attending the hematology outpatient clinics at Cairo university hospital"

In relation to family history, the current results ⁵found that the more than two thirds of the studied patients in both study and control groups had family history for thalassemia from first degree relation. ¹This result may be due to that thalassemia is genetic disease that can be transferred through heredity. ¹This result go in the same context with Abd Al-Abbass et al., (2024) who reported that two thirds of patients had family history of ¹thalassemia in a study that titled " Epidemiological characteristics and disease complications in thalassemia syndrome patients in Babylon, Iraq"

Pertaining to patients' knowledge regarding definition, causes, signs and symptoms, diagnostic test, complications, treatment, iron

therapy, diet and activity of thalassemia,³ the findings of the present study illustrated that, there was no statistical significant difference between study and control groups as regarding knowledge scores before the nursing program implementation, as the studied patients allocated randomly from the same population.

These findings are in line with Hassani et al., (2024) who conducted that no statistically significant difference between the mean scores of knowledge in the two groups before the educational intervention⁴⁶ in their study that titled "Effectiveness of educational intervention on quality of life in adults with thalassemia major".¹

The result of the present study demonstrated that, there is an improvement in patients' knowledge in the study group after the implementation of educational program and follow up, from the researcher point of view, this result may be due to training program affect patients' knowledge positively that appeared in the highest scores in the post intervention phase. The studied patients were prone to the educational program covered all the knowledge, and skills needed by the patient based on the assessment done during the pilot study and included all items related to the knowledge about the disease. **Which support research hypothesis (1).**

These findings are in line with Hassani et al., (2024) who conducted that a statistically significant difference between the mean scores of knowledge in the two groups after the educational intervention, study group demonstrated that increase knowledge and level of awareness in one week after the educational intervention and four months after follow up phase.

As regarding quality of life for patients with thalassemia, the findings of the present study revealed that there was no statistical

significant difference between study and control groups before program but found highly statistical significant between both group after educational program. From researcher point of view, Educational programs for patients with thalassemia can significantly enhance health-related quality of life by improving their understanding of the condition, promoting self-management skills, and fostering adherence to treatment plans. Studies have shown that structured educational interventions can lead to improved physical practices, psychological resilience, social integration, environmental adaptation and ultimately, quality of life.

The present study demonstrated a significant improvement in quality of life among patients in the intervention group, who received the educational program, compared to the control group that received routine care only. While the control group patients mostly reported poor quality of life, the intervention group patients showed a notable shift from poor to average quality of life after the program. This highlights the effectiveness of targeted educational interventions in enhancing patient outcomes.

This finding in line with **Dehnoalian, et al., (2017)**, who reported that comparison of the dimensions of quality of life in patients with thalassemia before and after the intervention indicated improvement in scores in all aspects of quality of life ($P > 0.05$). in their study that titled "The impact of educational counseling program on quality of life of thalassemia patients: a comprehensive short scale to assess the functional, psychosocial, and therapeutic factors of QOL among stroke survivors"

Similarly, ⁴³ this finding in the same line with **Etemad et al, (2021)**, who demonstrated that, a significant increase in the QoL score was observed in all dimensions after the implementation of the intervention in one month and three month, compared to the QoL before the intervention

in their study that titled " Quality of life and related factors in β -thalassemia patients "

At the same direction, **Imanian et al. (2024)** who conducted that, there was no significant difference between the control and intervention groups in the mean pre-test scores of quality of life, the mean post-test scores of quality of life in the intervention group were significantly increase than the control group in their study that titled "An investigation of group-based mobile learning on stress, anxiety, depression, and pain among Beta-thalassemia major patients".

As regarding Roy adaptation model, the findings of the present study revealed that there was no statistical significant difference between study and control groups before program but found highly statistical significant between both group after educational program. From researcher point of view, the educational program, based on the Roy Adaptation Model, significantly improved adaptation and quality of life in patients. By enhancing their ability to cope with challenges, the program fostered better physiological, psychological, social and environmental adaptation. The effectiveness of the educational program was evident in the improved quality of life outcomes, highlighting its potential as a valuable tool for patient care.

The present study demonstrated the effectiveness of the educational program based on the Roy Adaptation Model in improving adaptation levels in patients. The study group shows a significant shift from low to high adaptation after receiving the program, while most patients in the control group had mal adapted. This finding in line with **Hasanah, et al., (2022)**, who reported that, RAM-based interventions effectively overcome the problems experienced by patients and reduce the perceived symptoms and improve the patient's quality of life in their

study that titled " A Critical analysis of using Roy's Adaptation Model in nursing research as an empirical for clinical practice: a systematic review".

¹¹In the same context **Imani, et al., (2024)**, who ¹⁴reported that, the application of Roy model was determined in improving adaptation level in different modes and in various diseases and situations, including in hemodialysis patients, thalassemia, heart failure, breast cancer, mental disorders, burns, asthma, multiple sclerosis, diabetes, blood pressure and covid- 19 ¹⁴In their study that titled " The application of Roy adaptation model in iranian nursing care: a narrative review study ".

Similarly, this finding ¹³in the same line with **Yaghoubinia, et al., (2017)**, who demonstrated that, there was a statistically significant difference for the mean scores of physiological adaptation ²⁶of patients on pre and post-application of RAM. There was observed improvement in adaptation level among patients after application of RAM. In their study ¹⁸that titled "Effect of care plan based on Roy adaptation model on physiological adaptation in patients with thalassemia major ".

Regarding ⁶⁷the current study correlations, the findings indicated a statistically significant positive relationship between the Roy Adaptation Model (RAM), patient ¹knowledge, and quality of life (QoL) before ⁵⁵and after the implementation of the educational program.

The researcher believes that patients with greater knowledge about their condition are more likely to manage their symptoms effectively and adopt healthier lifestyle choices. This enhanced understanding helps to reduce stress and promotes better psychological and physical well-being. According to Roy's model, adaptation occurs through four modes: physiological, self-concept, role function, and interdependence. As

patients gain a deeper understanding of thalassemia, they are better equipped to adapt within each of these domains, resulting in an overall improvement in their quality of life. Therefore, increasing disease-specific awareness and education plays a critical role in supporting patient adaptation and enhancing daily living.

¹³ This finding supported with Hasanah, et al., (2022), who reported ¹⁶ that, RAM-based interventions effectively overcome the problems experienced by patients and reduce the perceived symptoms and improve the patient's quality of life. Also this result support with ⁸ Hosseini & Soltanian, (2022), who reported that application of care program based on the Roy's adaptation model increased physical and psychological adaptive (healthy) behaviors in patients with chronic diseases, which might improve the patients' psychological adaptation to the disease, proper disease control, reduction of complications, and quality of nursing care. In their study that titled ¹² "Application of Roy's adaptation model in clinical nursing ".

In the same context Hassani, et al., (2024), who demonstrated that a structured educational intervention significantly improved knowledge and quality of life among patients with thalassemia major, highlighting the importance of educational support in chronic disease management. In their study that titled ²⁷ " Effectiveness of educational intervention on quality of life in adults with thalassemia major: A quasi-experimental study based on PRECEDE model ".

² Conclusion

Based on the findings of the study, it can be concluded that, the educational program based on Roy adaptation model had a positive effect ² on the patient's outcomes among study group as regarding knowledge, quality of life and adaptation level of patient compared to control group,

as well as there was statistically positive correlation with high significance between knowledge, adaptation and quality of life.

Recommendation:

- Educational programs based on the Roy Adaptation Model should be regularly implemented to improve patients' adaptation and enhance their quality of life.
- Future studies are recommended to assess the long-term impact of adaptation-based educational program on large sample of patients with thalassemia for generalization.

Reference

- Abd Al-Abbass, H., Obaid, A., Makki, H., Abdulrasol., & Jasim, A. (2024).** Epidemiological Characteristics and Disease Complications in Thalassemia Syndrome Patients in Babylon, Iraq. Journal of Babol University of Medical Sciences, 26(1).
- Abdelrahman, B., Mohey El-Deen, H., & Sabry, S.. (2023).** Effect of Applying Roy's Adaptation Model on Quality of Life for Patients with Vitiligo. Egyptian Journal of Nursing and Health Sciences, 4(4), 215-242.
- Al_Hamadiny, S., Alqaisi, A., Al-Sarray, R., Ali, S., Trad, H. & Abood, A. 2024** Psychological Factors Associated with Major Thalassemia. Journal of Medical Genetics and Clinical Biology Volume: 1, Issue: 9, ISSN: 3032- 1085
- Badur, A., Ozbalci, D. & Alanoglu, E. (2021).** Evaluation of The Relationship between Sociodemographic Characteristics, Quality of Life, Depression, Drug Compliance and Biochemical

Parameters in Patients with Thalassemia Major in Isparta. *Annals of Medical Research*, 28(8), 1500-1507.

Chow, C., Shao, J. & Wang, H. (2007). Sample Size Calculation in Clinical Research, Second edition. Chapman and Hall/ CRC Biostatistics Series, P: 389-405.

Dehnoalian, A., Dehkordi, S., Alaviani, M., Motamedi, Z., Ahmadpour, S. & Banan-Sharifi, M. (2017). The Impact of Educational Counseling Program on Quality of Life of Thalassemia Patients. *Jundishapur Journal of Chronic Disease Care*, 6(4).

DeSanto, S. & Fawcett, J. (2009). Toward Understanding and Measuring Adaptation Level in The Context of The Roy Adaptation Model. *Nursing Science Quarterly*, 22(4), 355-359.

El Arab, R., Alzghoul, H., Abu-Mahfouz, M., Aldarwish, Z., Abbadi, M., Ghashi, E. & Seweid, M. (2025). Nurses' Perspectives on the Sleep Quality of Hospitalized Patients in Al Ahsa, Saudi Arabia. *Nursing Reports*, 15(2), 54.

Etemad, K., Mohseni, P., Aghighi, M., Bahadorimonfared, A., Hantooshzadeh, R., Taherpour, N. & Hajipour, M. (2021). Quality of Life and Related Factors in β -thalassemia Patients. *international journal for hemoglobin research*, 45(4), 245-249.

Hamdy, M., Draz, I., El Sayed, I., Ayyad, A. & Salemd, M. (2021). Assessment of Quality of Life among Beta-Thalassemia Major Patients Attending the Hematology Outpatient Clinics at Cairo University Hospital. *Open Access Macedonian Journal of Medical Sciences*, 9(E), 156-160.

Haq, N., Masood, N., Nasim, A., Riaz, S., Saood, M. & Yasmin, R. (2017). Assessment of Disease State Knowledge and Awareness Among the Guardians of Thalassemia Patients Attending Different Health Facilities in Quetta, Pakistan. In Value In Health, 20(5), 236-236.

Hasanah, I., Nursalam, N., Pandin, M., Ikhwan, D. & Rohita, T. (2022). A Critical Analysis of Using Roy's Adaptation Model in Nursing Research as an Empirical for Clinical Practice: a Systematic Review. doi:10.20944/www.preprints.org.0276.v1

Hassan, W., Soliman, H., Mohamed, M. & Fawzi, A. (2025). The Prevalence of Blood Transfusion Episodes among Thalassemia Patients at Sporting Hospital Alexandria. European Journal of Theoretical and Applied Sciences (ISSN 2786-7447) Volume 3, Number 2.

Hassani, L., Seyrafi, N., Mohammadi, S., Aghamolaei, T., Ghanbarnejad, A. & Reza, M. (2024). Effectiveness of Educational Intervention on Quality of Life in Adults with Thalassemia Major: A quasi-Experimental Study Based on PRECEDE Model. Health Science Reports, 7(9), e70075.

Hossain, M., Islam, M., Munni, U. R., Gulshan, R., Mukta, S., Miah, M. & Islam, M. (2023). Health-Related Quality of Life among Thalassemia Patients in Bangladesh Using the SF-36 Questionnaire. Scientific Reports, 13(1), 7734.

Hosseini, M., & Soltanian, M. (2022). Application of Roy's Adaptation Model in Clinical Nursing: A Systematic Review. Journal of Iranian Medical Council. 5(4).

Imani, A., Jaghour, E., Molavynejad, S. & Asadizaker, M. (2024). The Application of Roy Adaptation Model in Iranian Nursing

Care: A Narrative Review Study. *Community Health Journal*, 17(4), 27-37.

Khodashenas, M., Mardi, P., Ghahfarokhi, N., Tavakoli, B., Jamee, M. & Ghodrati, N. (2021). Quality of Life and Related Paraclinical Factors in Iranian Patients with Transfusion-Dependent Thalassemia. *Journal of Environmental and Public Health*, (2849163), 6.

Kumar, S., Jan, R., Rattani, S., & Yaqoob, A. (2022). Theory Guided Practices: An Approach to Better Nursing Care through the Roy Adaptation Model. *International Journal of Current Research and Review*, 14(14), 58-63.

Majeed, I., Sehar, S., Afzal, M., Gilani, S. A., Parveen, K., & Ahmed, R. (2020). 34. Effect of Roy's Adaptation Model Based Interventions on Quality of Life in Patients with Type II Diabetes. *Pure and Applied Biology (PAB)*, 9(1), 332-339.

Man, C. D., Maideen, S. F. K., & Rashid, A. (2019). Knowledge, Attitude and Practice towards Dietary Iron among Patients with Thalassemia and their Caregivers in Peninsular Malaysia. *Med J Malaysia*, 74(5), 365.

Olwi, D., Merdad, L. & Ramadan, E. (2018). Thalassemia: A Prevalent Disease yet Unknown Term among College Students in Saudi Arabia. *Journal of Community Genetics*, 9(3), 277-282.

Setiawan, H., Xu, W., Wang, C., Li, C., Ariyanto, H., Firdaus, F. & Hu, R. (2025). The Effect of Mobile Application Based Genetic Counseling on the Psychosocial Well-Being of Thalassemia Patients and Caregivers: A randomized Controlled Trial. *Patient Education and Counseling*, 130, 108457.

- Skevington, S., Lotfy, M. & O'Connell, K. (2004).** The World Health Organization's WHOQOL-BREF Quality of Life Assessment: Psychometric Properties and Results of the International Field Trial. *Quality of Life Research*, 13(2), 299–310.
- Su, Y., Xie, J., He, J., Shen, Y., Li, T., Huang, W. & Bian, Q. (2025).** Screening and Treatment of Thalassemia. *Clinica Chimica Acta*, 120211.
- Thein, S. & Rees, D. (2025).** Haemoglobin and the Inherited Disorders of Globin Synthesis. *Hoffbrand's Postgraduate Haematology*, 222-248.
- Wangi, K., Shaleha, R., Wijaya, E., & Birriel, B. (2025).** Psychosocial Problems in People Living with Thalassemia: A Systematic Review. *SAGE Open Nursing*, 11, 23779608251323811.
- Yaghoubinia, F., Navidian, A. & Yousefian, N. (2017).** Effect of Care Plan Based on Roy Adaptation Model on Physiological Adaptation in Patients with Thalassemia Major. *Medical-Surgical Nursing Journal*, 6(2 & 3).

The Effect of the Educational Program Based on Roy's Adaptation Model on the Quality of Life for

ORIGINALITY REPORT

51%

SIMILARITY INDEX

40%

INTERNET SOURCES

41%

PUBLICATIONS

25%

STUDENT PAPERS

PRIMARY SOURCES

1

Nora Ramadan Korany, Hanan Sayed Ali, Asmaa Abd ELRahman Abd ELRahman, Reham ELAshery Ashery. "Relationship between Knowledge of Patients with Thalassemia and their Quality of Life", Egyptian Journal of Health Care, 2022

Publication

8%

2

eurchembull.com

Internet Source

7%

3

www.eurchembull.com

Internet Source

5%

4

Yue Su, Jiahao Xie, Junjia He, Yeyu Shen, Ting Li, Weitao Huang, Xiangmin Tong, Qiong Bian. "Screening and treatment of thalassemia", Clinica Chimica Acta, 2025

Publication

3%

5

www.noveltyjournals.com

Internet Source

3%

6

Submitted to Purdue University

7

Mona Abdallah Abdel-Mordy, Wafaa Atta Mohammed, Zeinab Rabea Abd Elmordy. "Effect of Application Roy's Adaptation Model on Women's Satisfaction and Quality of Life after Mastectomy", Egyptian Journal of Health Care, 2021

Publication

1 %

8

Submitted to Herzing University

Student Paper

1 %

9

Ahmed Attia Mousa. "Th Effect of Nursing Intervention protocol on Controlling Extremities Muscle Contracture for Critically Brain Injured Patients", Helwan International Journal for Nursing Research and Practice, 2022

Publication

1 %

10

Submitted to Universiti Teknologi MARA

Student Paper

1 %

11

www.ajol.info

Internet Source

1 %

12

www.jimc.ir

Internet Source

1 %

13

nur.tanta.edu.eg

Internet Source

1 %

14	chj.rums.ac.ir Internet Source	1 %
15	onlinelibrary.wiley.com Internet Source	1 %
16	www.researchgate.net Internet Source	1 %
17	iosrjournals.org Internet Source	1 %
18	Submitted to Jersey College Student Paper	1 %
19	pmc.ncbi.nlm.nih.gov Internet Source	<1 %
20	worldwidescience.org Internet Source	<1 %
21	Submitted to University of Wales central institutions Student Paper	<1 %
22	Submitted to Wilkes University Student Paper	<1 %
23	Afaf Metwally, Amany Abdelaziz, Mohamed Ghalwash, Amany Mohamed. "Effect of Self-Care Practice Health Educational Program for Patients on Urinary Tract Infection Recurrence", Tanta Scientific Nursing Journal, 2021	<1 %

24	Zienab Hussein Ali, Ayat Mostafa Abdel Hady, Ola Abdelwahab Abdallah Srouer. "Critical Care Nurses' Performance and Obstacles towards Palliative Care for Critically Ill Patients", The Egyptian Journal of Hospital Medicine, 2022 Publication	<1 %
----	--	------

25	journals.lww.com Internet Source	<1 %
----	--	------

26	pdfs.semanticscholar.org Internet Source	<1 %
----	--	------

27	Submitted to Republic of the Maldives Student Paper	<1 %
----	--	------

28	Samah Mahmoud Mohamed Abd-EL Baky, Sabah Saad Al Sharkawi, Faten Shafik Mahmoud, Seham Mohammed Abdelaziz. "Effect of Educational Intervention on Quality of Life for Mothers' Having Children with Ventriculoperitoneal Shunt", Journal of Nursing Science Benha University, 2023 Publication	<1 %
----	---	------

29	www.ncbi.nlm.nih.gov Internet Source	<1 %
----	--	------

30	Marwa Hanafy Sayed Ahmed, Kamelia Fouad Abdalla, Jackleen Faheem Gendey, Arzak Mohamed Khalifa. "Factors Affecting	<1 %
----	--	------

Arteriovenous Fistula Survival among Hemodialysis Patients", Egyptian Journal of Health Care, 2022

Publication

31

Submitted to The University of Texas at Arlington

Student Paper

<1 %

32

Submitted to Radford University

Student Paper

<1 %

33

www.coursehero.com

Internet Source

<1 %

34

www.iosrjournals.org

Internet Source

<1 %

35

Fereshteh Hadi, Shahram Molavynejad, Nasrin Elahi, Habib Haybar, Elham Maraghi. "King's Theory of Goal Attainment: Quality of Life for People With Myocardial Infarction", Nursing Science Quarterly, 2023

Publication

<1 %

36

cuir.car.chula.ac.th

Internet Source

<1 %

37

www.researchsquare.com

Internet Source

<1 %

38

Submitted to University of Arizona

Student Paper

<1 %

39	www.isoqol.org Internet Source	<1 %
40	Submitted to Prairie View A&M University Student Paper	<1 %
41	Submitted to South University Student Paper	<1 %
42	www.slideshare.net Internet Source	<1 %
43	xianshiyoudaxuexuebao.com Internet Source	<1 %
44	"Effect of Preoperative Nursing Protocol on Pain and Daily Activities for Post Inguinal Hernia Repair Patients", Minia Scientific Nursing Journal, 2020 Publication	<1 %
45	Ahmed Mekdad. "Th Effect of Nursing Intervention on the Reduction of Pulmonary Complications after Thoracotomy", Helwan International Journal for Nursing Research and Practice, 2022 Publication	<1 %
46	SayedReza Borzou, SafuraKhan Mohammadi, GholamHossein Falahinia, Saeed Mousavi, Zahra Khalili. "Effects of roy's adaptation model in nursing practice on the quality of life	<1 %

in patients with type II diabetes", Journal of Nursing and Midwifery Sciences, 2015

Publication

47

Submitted to Southwest Minnesota State University

Student Paper

<1 %

48

Submitted to auf

Student Paper

<1 %

49

Rania Aboelnour. "Assessment of Female Students Knowledge and Self-esteem regarding Sexual Harassment at Beni-Suef University", Helwan International Journal for Nursing Research and Practice, 2022

Publication

<1 %

50

Yasmin Korany Ali, Furat Hussein Mahmoud, Sabah Nagah Hassan. "Assessment of Nurses' Performance Regarding Care of Patients with Spinal Cord Injury", The Egyptian Journal of Hospital Medicine, 2022

Publication

<1 %

51

abjournals.org

Internet Source

<1 %

52

Nahla Mohammed Hashem Elkerdawy, Neamatallah Gomaa Ahmed, Asmaa Mohamed Khorais. "Knowledge and Adherence to Oral Anticoagulant Therapy in

<1 %

Patients with Atrial Fibrillation", Egyptian Journal of Health Care, 2023

Publication

53

fnur.stafpu.bu.edu.eg

Internet Source

<1 %

54

www.dovepress.com

Internet Source

<1 %

55

www.medicopublication.com

Internet Source

<1 %

56

Mauro Porcu, Mariana Ragassi Urbano, Waldiceu A. Verri, Regina Céla Rezende Machado et al. "Comparison of the severity of depressive and anxiety symptoms, biomarkers, and childhood trauma among bipolar smokers and non-smokers, and controls", Journal of Affective Disorders Reports, 2022

Publication

<1 %

57

research.asu.edu.eg

Internet Source

<1 %

58

sciencescholar.us

Internet Source

<1 %

59

sifisheressciences.com

Internet Source

<1 %

60

Esraa Aatef Dawood Romeeh, Magda Abd El Sattar Ahmed, Wafaa Khalil Ibrahim.

<1 %

"Mothers' Awareness about Poisoning Prevention among their Children under Five Years Old", Egyptian Journal of Health Care, 2022

Publication

-
- 61 Gehan Karawan Sayed Sallam, Kameilia Fouad Abdalla, Sara Fathi Mahmoud. "Relation Between Compliance of Patients Post Coronary Artery Bypass Surgery Towards Symptoms Management Strategies and Experienced Discomforts", Egyptian Journal of Health Care, 2022

<1 %

Publication

-
- 62 Marzieh Hosseini, Mitra Soltanian. "Application of Roy's Adaptation Model in Clinical Nursing: A Systematic Review", Journal of Iranian Medical Council, 2022

<1 %

Publication

-
- 63 Shaimaa M Gallo, Om Ibrahim A Elmelegy, Wael H Mahmoud, Zeinab F Bahgat. "Effect of Massage by Using Lavender and Chamomile Oils of Intact Skin on Outcomes of Moderately Burned Patients", Egyptian Journal of Health Care, 2022

<1 %

Publication

-
- 64 Zodwa Dlamini. "Understanding Pancreatic Cancer - Global Strategies and African Perspectives", CRC Press, 2025

<1 %

-
- 65 online-journal.unja.ac.id <1 %
Internet Source
-
- 66 "Symposia", Quality of Life Research, 2005 <1 %
Publication
-
- 67 Amany Abdalla, Yasmen Mohammed, Huwida Abd elmonem. "Effect of Health Coaching Intervention on Mothers' Performance in Care of their Children with Beta Thalassemia", Assiut Scientific Nursing Journal, 2022 <1 %
Publication
-
- 68 Amira Hassan. "Effect of Educational Program on Nurses' practice Regarding Care of Adult Patients with Endotracheal Tube", Port Said Scientific Journal of Nursing, 2018 <1 %
Publication
-
- 69 Cihat Demirel, Serap Parlar KILIÇ. "The effects of education based on the Roy adaptation model on medication adherence and psychosocial adjustment in hypertensive patients", Journal of Vascular Nursing, 2024 <1 %
Publication
-
- 70 Doaa Goda, Nadia Taha, Abd El Fatah Frere, Elham Mohamed. "Effect of Health Promotion Program on Compliance of Patients with Hypertension toward Therapeutic Regimen", <1 %

71

Naglaa Saad Abd El-aty, Reda Dardier Hussein Awad, Abeer Mohamed Abdelkader, Manal Abo-elilal Abo elkiat Hamad et al. "Effect of Simulation Based-Learning on Improving Cognitive Achievement of Nursing Students during Care of Patients with Bronchial Asthma", International Egyptian Journal of Nursing Sciences and Research, 2022

Publication

72

Peter J.N. Dejonckheere, Kristof Van De Keere, Nele Mestdagh. "Training the Scientific Thinking Circle in Pre- and Primary School Children", The Journal of Educational Research, 2009

Publication

73

Wei Geng, Hongping Jia, Xianjing Zeng, Jia Li. "Effects of Nursing Interventions on Improving Patients With Heart Failure: A Systematic Review and Meta-Analysis", Research and Theory for Nursing Practice, 2025

Publication

74

Atefeh Dehnoalian, Shirin Madadkar Dehkordi, Mehri Alaviani, Zohre Motamedi, Sadaf Ahmadpour, Maryam Banan-Sharifi.

<1 %

<1 %

<1 %

<1 %

"The Impact of Educational Counseling Program on Quality of Life of Thalassemia Patients", Jundishapur Journal of Chronic Disease Care, 2017

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On