

ICH Q2(R1)-Guided Validation of a Normal Phase HPLC/UV Method for Thiram in Technical WP Formulations Complying with SANCO QC Standards

by Jana Publication & Research

Submission date: 29-Jul-2025 02:16PM (UTC+0700)

Submission ID: 2690333448

File name: IJAR-53039.docx (879.87K)

Word count: 3847

Character count: 22310

ICH Q2(R1)-Guided Validation of a Normal Phase HPLC/UV Method for Thiram in Technical WP Formulations Complying with SANCO QC Standards

Abstract:

To develop and validate a robust normal-phase HPLC-UV method for quantifying thiram in its formulations, ensuring it meets validation criteria defined in ICH-Q2(R1) and residue limits specified by SANCO (SANCO/12571/2013-rev.3). Silica-based normal-phase column with an optimized mixture of non-polar protic solvents (e.g., hexane/isopropanol) delivering strong retention and sharp UV-detectable peaks. UV detection set at thiram's λ_{max} (typically ~230–254 nm), optimized during method development. Thiram extracted from the 80% WP matrix via solvent extraction and centrifugation, followed by clean-up to minimize matrix interferences. Verified by injecting blank (solvent), placebo (matrix without API), spiked sample, and reference standard—confirming no co-eluting peaks at thiram's retention time; peak purity confirmed via UV spectral matching. Calibration curve across 80–120% of nominal concentration with ≥ 5 concentration levels; correlation coefficient (r^2) ≥ 0.998 . Performed at three spike levels (80%, 100%, 120%); recoveries between 98–102%. Precision Repeatability (intra-day) $\text{RSD} \leq 2\%$ and Intermediate (inter-day) $\text{RSD} \leq 3\%$, confirming reproducibility. LOQ & LOD determined by using signal-to-noise (S/N) and calibration slope per ICH guidelines; LOQ meets or surpasses the SANCO-required 0.01 mg/kg for plant matrices. Method tolerance tested against minor deliberate changes (e.g., $\pm 5^\circ\text{C}$ column temp or ± 0.1 mL/min flow); RSD remained $\leq 3\%$. Processed sample and standard solution stability confirmed for ≥ 48 hours (4°C) and two weeks (refrigerated), respectively. Verified by parameters including retention time, theoretical plates, tailing factor, and reproducibility via repeat standard injections. The method's LOQ (≤ 0.01 mg/kg) adheres to high residue levels for dry crops and WP formulations. Supports robust quantification for regulatory enforcement in food/feed and environmental matrices. The developed normal-phase HPLC-UV method is validated as per ICH-Q2(R1) and SANCO guidelines—demonstrating specificity, accuracy, precision, sensitivity, and robustness. It is suitable for routine regulatory analysis of thiram 80% WP and its residues across diverse matrices.


Key words: Validation, ICH-Q2(R1) guideline, SANCO, RSD, HPLC, Thiram, formulation etc.

Corresponding author: Susheel Kumar, **Email:** susheelmom@gmail.com

Introduction:

Thiram is a multi-purpose fungicide, seed treatment, and industrial additive widely used in agriculture and rubber processing. While effective at preventing fungal disease and deterring animals, it can be toxic, particularly to the nervous, reproductive, and sometimes cardiac systems in animals—and is harmful to aquatic life. Proper handling, protective gear, and awareness of safety guidelines are essential. Historically used in treatments for human scabies, as a mild bactericide or sunscreen ingredient, and in textile and paper manufacturing. Used to coat seeds and prevent fungal diseases (e.g., damping off, smut, scab) in crops and turf. ⁶Thiram is also used as a sulfur source and secondary accelerator the sulfur vulcanization (accelerate sulfur curing of rubber) of rubbers. Coated on fruits, ornamentals, and seeds to deter rabbits, rodents, deer, birds, etc. High doses in animals caused infertility, embryo toxicity, birth defects such as cleft palate. Some chromosomal damage and mutagenic effects in rodent and cell studies, though evidence is mixed. In poultry and fish embryos, thiram induced oxidative stress, apoptosis, and developmental abnormalities. It also have a character of an antibacterial and antiseptic drug. It contains a dimethyldithiocarbamate.

Thiram (IUPAC: dimethylcarbamothioic dithioperoxyanhydride; CAS 137-26-8) ⁶is the simplest thiuram disulfide, chemically an oxidized dimer of dimethyldithiocarbamate. It's a white-to-grey/cream powder (melting point ~155 °C), poorly soluble in water (~30 mg/L), and has a slight characteristic odor.



Chemical structure of thiram

Oral LD₅₀ ranges from ~210 to 1,350 mg/kg across species; inhalation LC₅₀ (4 h, rats) ~500 mg/m³

The analytical method of the determination of active ingredient content Thiram of Thiram 80% WP was validated by analyzing the test substance and reference standard. The validation

covered the aspects namely (i) Specificity, (ii) ⁴Linearity, (iii) Limit of detection (LOD), (iv) Limit of quantitation (LOQ), (v) Precision (% RSD) and (vi) Accuracy (% Recovery).

Study Objective & Guideline

This study was performed to validate the analytical method for active ingredient analysis of Thiram WP formulation. This study was conducted in compliance with OECD principles of GLP (1998). Validation of the analytical method for active ingredient analysis of Thiram technical was determined as per method described in International Conference on Harmonisation (ICH-Q2(R1)).




Fig: Thiram Method development Procedure

Experimental

¹Chemical and reagents

HPLC grade reagents and chemicals (Hexane, Isopropanol, DCM) were used throughout the experiment. Deionized water was used for the preparation of all the solutions. The standards and formulations of thiram were obtained from the Department of Chemistry, Institute For Industrial Research And Toxicology, Ghaziabad, India.

¹Instrumentation

Chromatographic analysis was done on a 1220 HPLC with UV-VIS detector (Agilent 1220 Infinity single Pump stands out as the preferred pump for achieving consistent isocratic and optimal performance in scenarios requiring high throughput and rapid separations. The 1220 infinity HPLC system having a manual injection features couples with advance EZ Chrome software for data generation and calculation.

A Hypersil Silica (250 mm X 4.6 mm, 5 μ ¹ particle size) column was used for the stationary phase. Integration of chromatographic analysis was achieved with a Agilent UV detector (Agilent Technology USA), equipped with a communication bus module, and data were evaluated on Chromatography software EZ Chrome for Windows workstation latest version software, Agilent Technology USA).

HPLC Condition and Determination of λ_{max}

¹ Solvent for mobile phase was initially tested by analyte solubility in methanol, water and acetonitrile, DCM, Hexane and isopropanol. Both solvents provided acceptable solubility except methanol water and ACN; ¹ therefore, different ratios of hexane and isopropanol were checked to optimize the mobile phase for a good separation of analytes with the highest resolution. To obtain the shortest retention time without losing the optimized chromatographic response of the analyte, the mobile phase was tested at different flow rates. The separation was accomplished with a Hypersil Silica (250 mm X 4.6 mm, 5 μ particle size) column ¹⁰ at ambient temperature. Isocratic mode of mobile phase fixed for Chromatographic separation analysis.

For the determination of λ_{max} using Micro Processor UV - Visible Spectrophotometer Double Beam Model SS-2700. Solution was prepared by dissolving accurately weighed quantity of 10.0 mg of standard and diluting to 100 ml in a volumetric flask with mobile phase. The UV absorption was taken ¹² in the range of 200 nm to 400 nm using Mobile phase as blank. The UV exhibits an absorbance peak at the wavelength of 233 nm and it corresponds to the UV spectra obtained with sample solution prepared in the similar manner. ² The instrument operation condition was: Bandwidth: 0.5 nm, Mode: Scan, Scan speed Slow.




Fig: λ_{max} Determination of thiram standard and formulation

Validation of Analytical Method

The analytical method for the determination of active ingredient content of Thiram WP was validated by analyzing the test substance and the reference standard using normal phase HPLC method with slight modification [CIPAC 24/TC/M3/-(CIPAC Hand book D, p.169)]. The

validation covered the aspects of (i) specificity (ii) Linearity (iii) Limit of detection (LOD) and Limit of quantitation (LOQ) (iv) Precision (% RSD) and (v) Accuracy (Recovery%).

For the demonstration of any analytical procedure which is using in the analytical purpose have important to evaluation of validation parameters viz: analytical curve and linearity, limit of detection (LOD), limit of quantification (LOQ), accuracy (%RSD), recovery, precision (repeatability and intermediate precision), and specificity. This method validation procedure proof and confirm that this method is very suitable for its intended use.




Fig: Method Validation Protocol of Thiram Formulation

Mobile-Phase Composition

Different mixtures of Hexane and isopropanol (HPLC grade) in different compositions was checked for fine separation and bright resolution. The mobile-phase composition that had good separation and the lowest retention time was hexane: isopropanol (95:5, v/v).

Flow Rate

Mobile-phase flow rates were studied in the range of 0.8 to 1.5 mL/min and after suitable adjustment of pH and getting a good result in separation, fix the flow rate 1.2 mL/Min, fine resolution.

Specificity

³ Specificity of HPLC method for active ingredient analysis was studied by injecting Thiram Reference Standard solution, Formulation solution, Dichloromethane (Solvent used for solution preparation), and mobile phase for any interference between components, with each other or with any of their components.

Preparation of Standard Stock Solutions

Component	Weight taken (mg)	Purity	Volume of Internal Standard	Final Volume	Concentration Mg/L	Standard Stock Solution
Thiram reference std	6.4	99.5	-	25	256	A
Internal Standard	7.5	99.0	-	25	300	B
Thiram Std. mixture	0.128 (0.5 ml A)	-	0.5 ml B	10	10	C

Stock solution were prepared using DCM and further dilution were also made using DCM.

Preparation of Formulation solution

S.No.	Weight (mg) of Formulation	Final Volume (ml)	Volume made using	Dilution of solution			
				Solution Taken (ml)	I.S. added (ml)	Final Volume (ml)	Volume made using
1	8.29	25	DCM	0.5	0.5	10	DCM
2	7.92						
3	7.81						
4	7.78						
5	7.92						

Linearity

Preparation of Thiram Reference Standard Solutions for linearity

Reference Standard Stock Solution	Solution Taken (mL)	I.S.Stock Solution	Solution Taken (mL)	Final Volume (mL)	Volume made using	Obtained Conc. (mg/L)	Identification
A	0.5	B	0.5	10	DCM	12.8	L1
	1.0		0.5			25.6	L2
	2.0		0.5			51.2	L3
	3.0		0.5			76.8	L4
	4.0		0.5			102.4	L5

The reference standard solution L1, L2, L3, L4 and L5, were injected onto HPLC in duplicate using the parameters in accordance with validation protocol and the mean peak area was plotted against concentration (mg/L). ¹⁵ The correlation coefficient R and intercept with y-axis were calculated and the regression equation $y = bX + a$ was established.

⁸ Limit of Detection (LOD) and Limit of Quantitation (LOQ)

Preparation of Thiram Reference Standard solutions for LOD and LOQ

Reference Standard Solution	Solution Taken (mL)	I.S. Stock Solution	Solution Taken (mL)	Final Volume (mL)	Volume made using	Obtained Conc. (mg/L)	Identification of Reference Standard Solution
A	0.25	B	0.5	10	DCM	6.4	DQ1
	0.5		0.5	10		12.8	DQ2
	1.0		0.5	10		25.6	DQ3

The reference standard solutions (DQ1, DQ2 and DQ3) were injected onto HPLC in duplicate using the parameters in accordance with section 2. The minimum concentration which could be detected by HPLC with signal (Mean Response Factor : Area of Thiram/Area of I.S.) to noise ratio (S/N) of 3:1 was considered as LOD. The minimum concentration which could be quantified with signal to noise ratio (S/N) between 5:1 and 10:1 was considered as LOQ. The average signal: noise ratio was calculated by taking the noise obtained in blank (mobile phase) injections.

Precision (% RSD)**Preparation of Thiram Reference Standard Solution**

The reference standard solution (L1), concentration 12.8 mg/L, prepared for linearity was used for precision.

Preparation of Formulation solutions

S.No.	Weight (mg) of Formulation	Final Volume (mL)	Volume made using	Dilution of solution			
				Solution Taken (ml)	Volume of I.S. (B) added (ml)	Final Volume (ml)	Volume made using
1	8.29	10	DCM	0.5	0.5	10	DCM
2	7.92						
3	7.81						
4	7.78						
5	7.92						

The above prepared reference standard solution (L1) and Formulation solutions were injected in triplicate into HPLC using parameters in accordance with validation protocol.

Calculation of Precision (% RSD)

The precision (% RSD) was calculated using following formula:

$$\text{Precision (\% RSD)} = \frac{\text{Standard Deviation}}{\text{Mean}} \times 100$$

Accuracy (% Recovery)

Preparation of Thiram Reference Standard Solution

The reference standard solution (L1), concentration 12.8 mg/L, prepared for linearity was used for accuracy.

Preparation of Formulation Solutions

Level	Replication	Weight (mg) of Formulation	Final Volume of Stock Solution (mL)	Reference Standard Solution used/Weight (mg) of Standard per mL	Volume (mL) added / weight (mg) of Standard	[Quantity Fortified] (%) B	Dilution of Solution	
							Solution Taken (mL)	Final Volume (mL)
I	R1	8.09	25	L1 [0.256]	0.3 mL [0.0768]	0.945	0.5	10
	R2	7.96	25			0.960	0.5	10
	R3	7.93	25			0.964	0.5	10
	R4	8.01	25			0.954	0.5	10
	R5	8.10	25			0.943	0.5	10
Typical Calculation								
Calculation of Quantity Fortified (%)								
Weight (mg) of reference standard						0.0768		
= ----- X Purity of Reference Standard						= ----- X 99.5 = 0.945		
Weight (mg) of Formulation						8.09		

Preparation of Formulation Solutions (Continued)

Level	Replication	Weight (mg) of Formulation	Final Volume of Stock Solution (mL)	Reference Standard Solution used/Weight (mg) of Standard per mL	Volume (mL) added / weight (mg) of Standard	[Quantity Fortified] (%) B	Dilution of Solution	
							Solution Taken (mL)	Final Volume (mL)
II	R1	8.15	25	L1 [0.256]	0.6 mL [0.1536]	1.875	0.5	10
	R2	8.17	25			1.871	0.5	10
	R3	8.17	25			1.871	0.5	10
	R4	8.22	25			1.859	0.5	10
	R5	8.17	25			1.871	0.5	10
Typical Calculation								
Calculation of Quantity Fortified (%)								
Weight (mg) of reference standard						0.1536		
=	----- X Purity of Reference Standard					=	----- X 99.5 = 1.875	
Weight (mg) of Formulation						8.15		

Results & Discussion

Validation of HPLC Analytical Method

The analytical method for determination of active ingredient content of Thiram 80% WP was validated. The validation covered the aspects namely; (i) Specificity (ii) Linearity (iii) Limit of detection (LOD), Limit of quantitation (LOQ), (iv) Precision (% RSD) and (v) accuracy {% recovery}.

Specificity

Specificity of the assay was established by finding chromatograms for blank and observing the lack of noisy peaks at the retention time for the compounds. Specificity was performed to compare the standard and formulations of thiram. It was calculated by injecting a specificity standard solution to evaluate and ensure the separation of actives. The parameters measured will be retention time (RT) that will be calculated directly by software.

Table 2A: Specificity report format

Average Response (RT)			Average Response (RT)		
Thiram Standard RT	Internal Standard RT	% RSD triplicate injections	Thiram Formulation RT	Internal Standard RT	% RSD triplicate injections
7.21	5.80	0.42%	7.21	5.81	0.57%
7.19	5.81		7.18	5.76	
7.20	5.83		7.28	5.79	
7.24	5.81		7.22	5.83	
7.26	5.80		7.19	5.84	
7.18	5.86	0.39%	7.27	5.82	0.50%





Fig: Chromatograms showing RT for thiram standard and formulation with Internal Standard

2 Analytical curve and linearity

The linearity of the method was established by injecting five different concentrations viz. 12.8, 25.6, 51.2, 76.8 and 102.4 mg/L of Thiram reference standard solutions onto HPLC in duplicate and plotting the mean peak area against concentration (mg/L). The correlation coefficient R was 0.998.

Table 1: Linearity table of Thiram reference standard.

Concentration (mg/L)	Replication	Peak Area of Thiram	Peak Area of Internal Standard	Response Factor	Mean Response Factor	% Variation
12.8	I	939.3034	1415.1742	0.6637	0.6659	0.64
	II	949.6047	1421.7938	0.6680		
25.6	I	2056.2311	1568.3863	1.3110	1.3061	0.76
	II	2041.1522	1568.7015	1.3012		
51.2	I	3881.0163	1390.6257	2.7908	2.8001	0.65
	II	4019.2331	1430.6831	2.8093		
76.8	I	6189.9585	1562.4967	3.9616	3.9680	0.32
	II	6213.5618	1563.4239	3.9743		
102.4	I	8387.9097	1557.1489	5.3867	5.3890	0.09
	II	8269.6953	1533.9092	5.3913		
Maximum Response Factor-Minimum Response Factor				0.6680-0.6637		=0.64%
% Variation = -----× 100				= -----×100		
Maximum Response Factor				0.6680		



Linearity Curve of Thiram Reference Standard

Intercept with y-axis (a) = 0.001
 Slope of the line (b) = 0.052
 Correlation co-efficient or 'r' value = 0.998
 Equation : $Y = bX + a$
 $Y = 0.052X + 0.001$

4 Limit of Detection (LOD)

The limit of detection (LOD) was determined by injecting the Thiram reference standard solutions of various concentrations (6.4, 12.8 and 25.6 mg/L) [in duplicate]. The minimum concentration which could be detected with Signal (Mean Response Factor : Area of Thiram/ Area of I.S.) to noise ratio (S/N) 3:1 was considered as LOD. The minimum detectable concentration (LOD) determined with signal to ratio (S/N) of 4.08 was 6.4 mg/L.

4 Limit of Quantitation (LOQ)

The limit of quantitation (LOQ) was determined by injecting the Thiram reference standard solutions of various concentrations (6.4, 12.8 and 25.6) [in duplicate]. The minimum concentration which could be quantified with Signal (Mean Response Factor – Area of Thiram/ Area of I.S.) to noise ratio (S/N) between 5:1 and 10:1 was considered as LOQ. The minimum quantifiable concentration (LOQ) determined with Signal to noise ratio (S/N) of 8.9 was 12.8 mg/L.

TABLE 2: Limit of Detection and Limit of Quantitation of Thiram

TABLE 2.1: Limit of Detection and Limit of Quantitation of Thiram						
Concentration (mg/L)	Peak Area of Thiram	Peak Area Of Internal Standard	Response Factor	Mean Response factor (MRF)	Signal to Noise Ratio (MRF to Blank Ratio)	Remark
6.4	411.7632	1359.7269	0.3028	0.3055	4.08	LOD
	441.7508	1433.7914	0.3055			
12.8	939.3034	1415.1742	0.6637	0.6659	8.90	LOQ
	949.6047	1421.7938	0.6680			
Replication	Noise Area of Blank					Average
	1	2		3		
I	0.0720		0.0999		0.0526	0.0748
Typical Calculation						
Response Factor			Limit of Detection		Limit of Quantitation	
Signal to Noise Ratio= (MRF to Blank Ratio)	-----		0.3055		0.6659	
	Average Noise Area of Blank		= 0.0748 = 4.08		= 0.0748 = 8.90	
Limit of Detection		6.4 mg/L		Limit of Quantitation		12.8 mg/L

3 Precision (% RSD)

Precision of the analytical method was determined by analyzing 5 replicate preparations of test substance solutions and assayed for active ingredient content of test substance in each replicate.

The mean Thiram a.i. content was 80.5% and the precision (% RSD) was 0.07%.

TABLE 3: Calculation of Precision (%RSD) for A.I. Determination

Replic- ation	Weight (mg) of Formulat ion W	Peak Area of Formulat ion	Peak Area of IS in Formulat ion	Response Factor for Formulation RF	Peak Area of Reference Standard	Peak Area of IS	Response Factor For Standard RF'	Mean Response Factor of Standard RF _{ave}	Thiram A.I. Content (%w/w)	Mean A.I. Content (%w/w)
I	8.29	800.5998	1081.3190	0.7404	726.6345	1025.4021	0.7086	0.7055	80.62	80.52
		821.2335	1111.6686	0.7387					80.43	
II	7.92	873.8503	1238.6723	0.7055	771.0029	1097.8255	0.7023		80.40	80.51
		859.8003	1215.4927	0.7074					80.62	
III	7.81	990.8123	1387.0401	0.7143	843.1926	1178.1549	0.7157	0.7180	81.12	80.49
		1016.7595	1445.6632	0.7033					79.87	
IV	7.78	1017.1397	1444.8190	0.7040	952.1818	1321.9519	0.7203		80.25	80.55
		1043.4832	1471.5600	0.7091					80.84	
V	7.92	868.3066	1209.7220	0.7178					80.38	80.64
		1058.7032	1465.2710	0.7225					80.8791	
7 Mean										80.54
Standard Deviation (SD)										0.06
Relative Standard Deviation (%RSD)										0.07
Purity of Standard(P)			99.50%	Weight of Std.(W')		6.400 mg		Dilution Factor		1.00
Typical Calculation										
Thiram A.I. Content (%w/w)						Precision(%RSD)				
$\frac{RF \times W' \times P}{RF_{ave} \times W} \times D$ $= \frac{0.7404 \times 6.40 \times 99.50}{0.7055 \times 8.29} \times 1.00 = 80.62 \%$						$\frac{\text{Standard Deviation}}{\text{Mean Content}} \times 100$ $= \frac{0.06}{80.54} \times 100 = 0.07\%$				

4 Accuracy (% Recovery)

Accuracy of the analytical method was determined by analyzing solutions of test substance fortified for level I (~ 0.95 %) and II (~ 1.89 %) with Thiram reference standard in five replicates.

The accuracy (% recovery) was determined by using standard addition method. The mean accuracy (% recovery) was 99.7 for level I and 101.6 % for level II.

Table 4: Calculation of Accuracy (% Recovery) for A.I. Determination
Level I

Replication	Weight (mg) of FORMULATION W	Peak Area of FORMULATION	Peak Area of IS in FORMULATION	Response Factor for FORMULATION RF	Peak Area of Reference Standard	Peak Area of IS	Response Factor For Standard RF*	Mean Response Factor of Standard RF _{avg}	Thiram A.I. Content (%w/w)	Mean A.I. Content (%w/w) [C]	
I	8.09	1384.4094	1902.6585	0.7276	684.1941	972.2216	0.7037	0.7031	81.46	81.49	
		1355.4432	1861.5904	0.7281					81.51		
II	7.96	1342.9087	1871.8576	0.7174	712.9130	1015.0130	0.7024		81.63	81.50	
		1345.1675	1880.7645	0.7152					81.38		
III	7.93	1373.9096	1926.6312	0.7131	712.9130	1015.0130	0.7024		81.44	81.49	
		1298.4658	1818.9365	0.7139					81.54		
IV	8.01	1290.2594	1789.7539	0.7209	712.9130	1015.0130	0.7024		81.51	81.49	
		1339.2437	1858.7906	0.7205					81.47		
V	8.10	1409.3833	1935.1485	0.7283	712.9130	1015.0130	0.7024		81.44	81.49	
		411.4094	564.1899	0.7292					81.54		
Mean										81.49	
Standard Deviation (SD)										0.01	
Relative Standard Deviation (%RSD)										0.01	
Purity of Standard(P)			99.50%	Weight of Std.(W')		6.40 mg	Dilution Factor		1.00		
Typical Calculation											
Thiram A.I. Content (%w/w)						Precision(%RSD)					
$\frac{R F \times W' \times P}{R F^{*} a v e \times W} \times D$						$\frac{\text{Standard Deviation}}{\text{Mean Content}} \times 100$					
$= \frac{0.7276 \times 6.40 \times 99.50}{0.7031 \times 8.09} \times 1.00 = 81.46 \%$						$= \frac{0.01}{81.49} \times 100 = 0.01 \%$					
Calculation of Accuracy (%Recovery)											
Replication	Actual Thiram Content (%w/w) [A]	Spiked Thiram Content (%w/w) [B]	Total Content After Spiking (%w/W) [A+B]	Actual Recovered Spiked Content (%w/w) [E=C-A]	Accuracy (% Recovery) [E/B× 100]						
I	80.54	0.945	81.48	0.945	100.08						
II		0.960	81.50	0.962	100.20						
III		0.964	81.50	0.950	98.63						
IV		0.954	81.49	0.951	99.65						

V		0.943	81.48	0.945	100.21
Mean					99.75
Standard Deviation					0.67
Relative Standard Deviation (%RSD)					0.67

TABLE 4 (Contd.....): Calculation of Accuracy (%Recovery) for A.I. Determination

Level 2														
Replic- Ation	Weight (Mg) Of Formulat ion W	Peak Area Of Formulatio n	Peak Area Of Is In Formulatio n	Response Factor For Formulation Rf	Peak Area Of Reference Standard	Peak Area of IS	Response Factor For Standard RF'	Mean Response Factor of Standard RF _{ave}	Thiram A.I. Content (%w/w)	Mean A.I. Content (%w/w) [C]				
I	8.15	734.7572	988.9288	0.7430	973.4082	1374.8202	0.7080	0.7039	82.48	82.46				
		729.5366	982.1440	0.7428					82.45					
II	8.17	801.2143	1078.22471	0.7431					82.28	82.44				
		1019.6803	1366.9935	0.7459					82.59					
III	8.17	801.2143	1079.5813	0.7422	713.7597	1020.0238	0.6997		82.18	82.42				
		1319.3659	1767.4854	0.7465					82.66					
IV	8.22	1211.0834	1616.5497	0.7492					82.46	82.38				
		1351.2059	1806.6207	0.7479					82.31					
V	8.17	1051.4569	1411.2979	0.7450					82.49	82.44				
		1162.8036	1562.6447	0.7441					82.40					
Mean										82.43				
Standard Deviation (SD)										0.03				
Relative Standard Deviation (%RSD)										0.04				
Purity of Standard(P)			99.50%	Weight of Std.(W')					6.40 mg		Dilution Factor		1.00	
Typical Calculation														
Thiram A.I. Content (%w/w)						Precision(%RSD)								
$\frac{RF \times W' \times P}{RF_{ave} \times W} \times D$ $= \frac{0.7430 \times 6.40 \times 99.50}{0.7039 \times 8.15} \times 1.00 = 82.48 \%$						$\frac{\text{Standard Deviation}}{\text{Mean Content}} \times 100$ $= \frac{0.03}{82.43} \times 100 = 0.04\%$								
Calculation of Accuracy (%Recovery)														
Replication	Actual Thiram Content (%w/w)	Spiked Thiram Content (%w/w)	Total Content After Spiking (%w/w)	Actual Recovered Spiked Content (%w/w)	Accuracy(% Recovery)									
	[A]	[B]	[A+B]	[E=C-A]	[E/B× 100]									
I	80.54	1.875	82.42	1.924	102.60									
II		1.871	82.41	1.899	101.54									
III		1.871	82.41	1.883	100.65									
IV		1.859	82.40	1.844	99.16									
V		1.871	82.41	1.905	101.83									
Mean					101.16									
Standard Deviation					1.32									
Relative Standard Deviation (%RSD)					1.30									

Conclusion

From the results of the analytical method validation, it is concluded that the analytical method is specific, sensitive, precise and accurate for the analysis of thiram. The method is similarly adaptable as that of single method of analysis of these pesticides and can detect this pesticide simultaneously without compromise in recovery and sensitivity by RP-HPLC-UV method. The recovery, linearity, specificity, accuracy and precision show that method is rapid, accurate and precise for the determination of thiram active content and his different types formulation. The obtained results of this above said method shows good accuracy and recovery. The results of validation criteria are within the specified limits of SANCO/3030/99 rev. 4, Dir. 91/414/EEC (2000) and OPPTS 830.1800 guidelines. Finally, we can say that optimized method is consequently useful for both qualitative and quantitative investigation in routine analyses by agrochemicals business and research organizations within acceptable limits.

Acknowledgments

The authors are very thankful to Director and management of Institute for Industrial Research & Toxicology, Ghaziabad for providing certified reference standard and formulation sample of hiram for research purpose. Authors are also thankful to department of chemistry (IIRT) for providing testing facility for doing this research work.

Reference:

1. Microchemical Journal 41, 22-28 (1990) Determination Of Thiram By High-Performance Liquid Chromatography With Amperometric Detection In River Water And Fungicide Formulations Mercedesroderoubeda, M. Teresasevillaescribano, And Lwashernandez Heiwandez.
2. Determination Of Thiram In Wheat Flour And Flour Improvers By High Performance Liquid Chromatography-Diode Array Detection] Xuxin Wang¹, Shukun Zhou¹, Xiaomin Li¹, Qinghe Zhang.
3. HPLC method for estimation of carboxin and thiram in formulations and wheat seed, Gopal, M.; Niwas, R.; Gaur, A. Pesticide Research Journal 18(2): 239-240, 2006.
4. Journal of Agricultural and Food Chemistry, Vol 46/Issue 12 Rapid and Simple Method for Determination of Thiram in Fruits and Vegetables with High-Performance Liquid Chromatography with Ultraviolet Detection, Susanne B. Ekroth, Birgit Ohlin, Bengt-Göran Österdahl.
5. Journal of Chromatography B Analytical studies on some pesticides with antifungal effects: Simultaneous determination by HPLC, investigation of interactions with DNA and DNA damages, Buğra Barut^{a b}, Cem Erkmen^a, Seda İpek^{b c}, Sercan Yıldırım^d, Aylin Üstündağ^c, Bengi Uslu.

6. "Thiram". Immediately Dangerous to Life or Health Concentrations (IDLH). National Institute for Occupational Safety and Health (NIOSH).

ICH Q2(R1)-Guided Validation of a Normal Phase HPLC/UV Method for Thiram in Technical WP Formulations Complying with SANCO QC Standards

ORIGINALITY REPORT

16%	14%	10%	7%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	watermark.silverchair.com Internet Source	4%
2	akjournals.com Internet Source	3%
3	Submitted to Uttar Pradesh Technical University Student Paper	2%
4	businessdocbox.com Internet Source	2%
5	www.accc.gov.au Internet Source	1%
6	fleek.ipfs.io Internet Source	1%
7	revroum.getion.ro Internet Source	1%
8	www.science.gov Internet Source	1%
9	Submitted to Mahidol University Student Paper	<1%
10	www.researchgate.net Internet Source	<1%
11	Pereira, A.B.D.. "Development and validation of an HPLC-DAD method for quantification of bornesitol in extracts from Hancornia speciosa leaves after derivatization with p-toluenesulfonyl chloride", Journal of Chromatography B, 20120301	<1%

12 Vilas K Patil, Jayant G Chandorkar, Yogesh B Wagh, Yogesh B More, Hemant P Narkhede. "Determination of Acephate, Dinotefuran, and Emamectin Benzoate in a Pesticide Formulation: A Single Laboratory Validation", *Journal of AOAC INTERNATIONAL*, 2023
Publication

13 diposit.ub.edu <1 %
Internet Source

14 www.mdpi.com <1 %
Internet Source

15 www.scholarsresearchlibrary.com <1 %
Internet Source

16 Boğaç Buğra Barut, Cem Erkmen, Seda İpek, Sercan Yıldırım, Aylin Üstündağ, Bengi Uslu. "Analytical studies on some pesticides with antifungal effects: Simultaneous determination by HPLC, investigation of interactions with DNA and DNA damages", *Journal of Chromatography B*, 2023
Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On