

# The comorbidity of depression and anxiety among Saudi Board Residents in different health training programs in Al-Ahsa, Saudi Arabia, 2024

*by* Jana Publication & Research

---

**Submission date:** 30-Jul-2025 12:20PM (UTC+0200)

**Submission ID:** 2722775955

**File name:** IJAR-53050.doc (1.06M)

**Word count:** 12483

**Character count:** 70034

# The comorbidity of depression and anxiety among Saudi Board Residents in different health training programs in Al-Ahsa, Saudi Arabia, 2024

## Abstract

14

### Introduction

Depression and anxiety disorders are among the most prevalent mental illnesses worldwide; they affect a person's functionality, causing long-term impairments. The prevalence of depression and anxiety in the general population is increasing, and healthcare providers are at risk of developing these mental illnesses, which impact the care provided to patients, the quality of life of the physicians, and the level of work productivity.

### Aim

to examine the prevalence and factors associated with the comorbidity of depression and anxiety among training physicians in various health training programs under the Ministry of Health (MOH) in Al-Ahsa, Saudi Arabia.

### Methods

This is a descriptive cross-sectional study conducted among training physicians in various health care training programs under the Ministry of Health in Al-Ahsa, Saudi Arabia. Data collected from different fields from a total of the study population 471 of a total population sampling. The data was collected through a validated self-administered questionnaire form. Data was analyzed using IBM SPSS Statistics for Windows, version 26 (IBM Corp., Armonk, NY, United States). Descriptive statistics were used to calculate the frequency and percentage of categorical variables.

### Results

A total of 221 professionals were included in the data collection from a total population of 471. The majority, 157 (71.0%), were between 24 and 29 years old, and 111 (50.2%), were females. Around 131 (59.3%) were married, and 111 (50.2%) reported receiving up to 5 on-call shifts per month. A total of 160 (72.4%) were not exercising, 188 (85.1%) were non-smokers, 187 (84.6%) did not complain of chronic diseases, and 195 (88.2%) did not complain of psychiatric illness. The participants showed mild 35.3%, moderate depression 33%, and moderately-severe 6.8% or severe depression 12.2%. Severe depression was highest in General surgery and orthopedics specialties (50% both). The residents, 49.8% (n=110), had no or minimal anxiety; others showed mild 22.6%, moderate 12.7%, and severe depression 14.9%. Severe anxiety was highest in Orthopedics (50%) and Psychiatry specialties (33.3%). Comorbidity of depression and anxiety

was Hughes in Orthopedics, followed by Obstetrics and Gynecology, internal medicine, and pediatrics.

#### Conclusion

The comorbidity of depression and anxiety are prevalent in residents with different level in different specialties. Improving the mental health of training residents and providing new interventions to prevent, recognize, and manage this issue in our community settings, especially in high-risk specialties, will improve overall healthcare quality

**Keywords:** Comorbidity, Depression, Anxiety, Al-Ahsa, Training Physicians in Health Training Programs, Ministry of Health (MOH)

#### 42 Introduction:

##### Depression

One of the most prevalent mental illnesses in the world, depression, can affect a person's functionality and result in long-term impairments in all facets of life. Globally, depression affects up to 5% of the population. The definition of depression is a chronically low mood that impairs many aspects of a patient's functioning and is accompanied by a loss of interest in previously enjoyed activities. Disturbances in sleep, focus, and appetite, constant weariness, guilt, and suicidal thoughts or attempts are other signs of depression (1). Suicidal ideation is a comprehensive phrase used to describe a variety of concepts, including thoughts and wants. It can be either passive or active, depending on whether the patient has plans to commit suicide or not, as well as a desire to die (2). Although the exact etiology of depression is still unknown, it is recognized to be influenced by several interrelated social, psychological, and biological factors. Research has indicated that depression is hereditary; almost one-third of all instances are thought to be inherited (1). Obesity, financial difficulties, a lack of social support, experiencing stressful life events such as parental loss or a history of abuse during childhood, having other medical conditions like multiple sclerosis, dissatisfaction with one's quality of life, excessive workload at work, and burnout at work are all risk factors linked to depression (1, 3). Gender is a significant risk factor; women are twice as likely as men to have depression. It was discovered that the prevalence of depression in relation to gender varies by region in the Gulf and Saudi Arabia. In

Saudi Arabia, up to one-third of all high school female students suffer from depression, whereas in Oman, it only affects less than one-fifth of all female adolescents. This is also the case in Egypt (3). Research indicates that there is a major risk factor for depression associated with family history; those who have first-degree relatives with a diagnosis of depression are three times more likely to be affected than those who do not (4). Research has indicated that depression is quite common throughout the world, with an estimated 300 million people affected. The percentage of people who are identified and successfully receive treatment is still low, even though there are established guidelines for diagnosing depression and effective treatment programs are available (5).

Depression screening is crucial and vital. Depression is a very common and extremely incapacitating aspect of health care. Patient health questionnaires, such as the Patient Health Questionnaire-2 (PHQ-2) or the Patient Health Questionnaire-9 (PHQ-9), are frequently used by healthcare providers to screen for depression. Both have extremely sensitive and specific levels for identifying and evaluating the severity of depression in individuals (6,7). A popular self-administered tool for determining the severity of depression is the PHQ-9. Research has demonstrated that the PHQ-9 is a viable and trustworthy tool for screening for depression and determining its severity, having been developed over many years by numerous researchers (7). Nine items on the PHQ-9 are designed to test for signs of depression and disability over the course of the previous two weeks. There are four possible answers for each question, with a score ranging from zero (not at all) to three (almost every day), for a total of twenty-seven points that examine the existence and severity of depression (6). A score of 20–27 indicates that the screening for depression is positive and that it is severe; a score of 15–19 indicates that the depression is moderately severe; a score of 10–14 indicates that the depression is moderate; a score of 5–9 indicates that the depression is mild; and a score of 4–1 indicates that the depression is minimal. When it comes to screening for depression, PHQ-9 has a greater specific level than PHQ-2, while PHQ-2 has a similar sensitivity level. It is advised that everyone over the age of eighteen be eligible for a depression test, regardless of whether they have any risk factors for the condition (5).

Two interest and mood-related multiple-choice items on the PHQ-2 have a score range of 0 to 3, with a total score of three or above being regarded as a positive screening result (6,7). The Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) criteria for Major Depressive Disorder (MDD) must be checked to confirm the diagnosis whenever the screening result is positive (6). The DSM-5 states that a patient must exhibit at least five symptoms from all the criteria for an MDD

diagnosis, with at least one of those symptoms having to be a loss of interest or a poor mood that has persisted for at least the previous two weeks. Sleep difficulty, poor attention, disturbed appetite, psychomotor agitation, low energy, guilt, and suicidal thoughts are some additional symptoms (6,8). Suicidal thoughts or attempts to ensure patient safety, together with interventions such as regular and close monitoring or even hospitalization when required, must be considered when evaluating patients with depressed symptoms (4). The DSM-5 for MDD is consistent with the diagnosis of depression and uses the total score result to determine how severe depression is (6,8).

Laboratory tests are useful in ruling out any potential medical condition that could mimic depression, even though they are not useful in diagnosing depression (4). A complete blood count (CBC), thyroid profile, analysis of iron and vitamin B12 levels for potential anemia, liver enzyme levels, and renal function tests are among the laboratory tests used to determine the patient's preferred medication based on their health (6).

There is a clear correlation between depression and substantial functional disability. All people with depression have functional impairment, according to studies, and almost two-thirds of all people with depression have a severe kind of impairment associated with the condition. It has been discovered that the functional impairment of people with depression is comparable to that of people with organic chronic diseases like diabetes or even cancer. <sup>60</sup>Patients with depression are more likely to seek medical attention and be admitted to the hospital than people without depression. <sup>65</sup>They are also more likely to attempt suicide and suffer from unemployment twice as frequently as people without depression. Because depression can worsen the condition of coexisting cardiovascular diseases, significantly increase mortality following myocardial infarction or unstable angina, and raise the risk of hypertension and stroke events, patients who suffer from <sup>66</sup>depression are at an increased risk of mortality and mobility rate. The greatest important effect of depression on death rates has been shown to be suicide; research has shown that <sup>33</sup>people with depression are eleven times more likely to attempt suicide than people without depression (9).

The true issue with depression is that the majority of those who experience it do not seek treatment; research indicates that as many as 60% of all individuals with depression do not seek treatment. The embarrassment associated with having mental illnesses or the stigma of not embracing them could be the cause of this (4). Some patients appear with a severe form of depression that requires inpatient therapy due to delusional thoughts, hallucinations, or suicidal thoughts, even though the most prevalent

types are mild to moderate and can be safely treated as an outpatient. There are pharmacological, non-pharmacological, or a combination of both types of depression treatment available. Treatment options include electroconvulsive therapy (ECT), psychotherapies, and antidepressant drugs (10). <sup>16</sup> Tricyclic antidepressants, monoamine oxidase inhibitors, selective serotonin reuptake inhibitors (SSRIs), and serotonin-norepinephrine reuptake inhibitors (SNRIs) are among the various classes of antidepressant drugs; each group has a distinct degree of safety and side effects (11). Because SSRIs are safe, effective, and have few, manageable side effects, they are regarded as the primary and first pharmacological line for treating mild to moderate depression. Venlafaxine and duloxetine are examples of SNRIs, while citalopram, sertraline, fluoxetine, and paroxetine are examples of SSRIs. As the second-line therapeutic option if SSRIs are unable to produce improvement or a response, SNRIs play a crucial role during treatment. Atypical antidepressants include, for instance, mirtazapine and bupropion. When treating difficult-to-treat depression, atypical antidepressants can be administered either alone or in combination (4). The treatment of severe forms of depression typically involves a <sup>30</sup> combination of pharmacological and non-pharmacological therapies, which can take the form of drugs and psychological therapy (11). Clinical response and improvement with drugs often take four to six weeks, which is why close follow-up is essential for continual monitoring and ensuring compliance with the treatment plan. Psychological therapy can be undertaken as a stand-alone initial <sup>30</sup> intervention for patients with mild to moderate depression who prefer not to use medication. Pregnancy, breastfeeding, or attempting to conceive are additional reasons to consider commencing treatment with psychiatric therapy. Although there are other forms of psychological therapies, <sup>59</sup> cognitive behavioral therapy (CBT) is the most helpful for depression (10,11). Helping patients recognize and alter any maladaptive beliefs or behaviors is the primary goal of cognitive behavioral therapy (4).

## Anxiety disorders

When a person encounters pressures, threatened situations, or topics where they must take action to stop the threat from harming them, they experience normal anxiety. <sup>67</sup> Anxiety is typically accompanied by physical symptoms such as tightness in the chest and palpitations, as well as bewilderment and the admission of fear. If normal anxiety continues to interfere with functioning, it may be unhealthy. According to estimates, one in ten people will have impairment due to anxiety disorders, which are

very common (12). <sup>71</sup> Anxiety disorder is the most prevalent mental health problem among all mental disorders. Anxiety disorders <sup>19</sup> can present in several forms, including generalized anxiety disorder (GAD), agoraphobia, panic disorder, and different types of phobias, such as social phobia. Certain <sup>70</sup> anxiety disorders, such as social phobia and generalized anxiety disorder, emerge during childhood. Conversely, other forms typically emerge throughout early adulthood or later, such as agoraphobia and panic disorder. Research indicates that the <sup>48</sup> prevalence of anxiety disorders is estimated to be between twenty and thirty percent among individuals in early adulthood, while it affects up to ten to fourteen percent of those in later adulthood (13). Due to the predominantly chronic and substantially debilitating nature of anxiety disorders, they rank sixth among the most burdensome health-related conditions. Anxiety disorders result in substantial and profound impairment across various facets of patients' lives, adversely affecting their quality of life, occupational performance, physical functioning, overall psychological well-being, social interactions, and leading to increased healthcare utilization (14). Concerning the risk factors associated with anxiety disorders, there are several risks that are typically multifactorial and have been linked together. The primary risk factors include gender, exposure to stressors or stressful life events, the presence of chronic health conditions such as autoimmune diseases, a history of mental illnesses, current mental health issues, low socioeconomic status, residing in impoverished areas, and experiences of trauma during childhood, such as parental separation or loss. A significant risk factor for anxiety disorders, particularly <sup>17</sup> generalized anxiety disorder (GAD), is the presence of a family history of mental health issues, especially among parents (15,16).

When comparing males and females regarding the association with anxiety disorders, female individuals face a risk of developing these disorders that is twice that of their male counterparts (17). Low self-esteem <sup>50</sup> can be influenced by a variety of factors, including a challenging family environment, <sup>32</sup> exposure to multiple traumatic events such as childhood sexual abuse, a significant family history of depression, and gender. These elements <sup>2</sup> can have a direct impact on an individual's mental health, heightening the risk for <sup>17</sup> anxiety disorders and major depressive disorder (MDD) (16). Some variables may help avoid or reduce the likelihood of acquiring anxiety disorders. These include receiving social support, learning healthy coping techniques for dealing with life's pressures, and elements relating to the good environment in which the person lives (18). Different types of anxiety illnesses have comparable symptoms. These symptoms include fear or worrying about situations or

objects that are caricatured as pathological due to their severity, being disproportionate to the threats, and causing functional impairment that requires medical attention. Patients experiencing dread of threatening circumstances or objects experience palpitations, chest tightness or pain, perspiration, abdominal discomfort, restlessness, disorientation, and impaired focus (13). Other specific symptoms of anxiety disorders include social fear or concern regarding performance, unexplained panic attacks without a known trigger, and avoidance behaviors associated with threats. The prevalence of each type of anxiety disorder that seeks medical assistance and visits primary health care clinics is six percent in the case of GAD, thirteen percent in the case of panic disorder, whether with or without agoraphobia, and approximately five percent in the case of social phobia (19). Anxiety disorders, like any other mental illness, can result in a variety of comorbidities, which have a substantial impact on the patient's quality of life. Some of these comorbidities include an increased risk of developing other mental disorders, such as depression or substance misuse, a higher risk of developing new anxiety disorders over time, and an increased risk of developing or worsening physical disorders (18). Depression is recognized as the most prevalent mental illness among various mental disorders, often occurring alongside anxiety disorders. Research indicates that around fifty percent to two-thirds of individuals experiencing anxiety disorders also contend with depression. The correlation between anxiety disorders and depression has resulted in a 3-5-fold increase in the incidence of depression among those affected by anxiety disorders (13).

Health care should identify high-risk patients for anxiety disorders and establish the diagnosis to give effective treatment to prevent disabilities and substantial life impairment. In general care, Generalized Anxiety Disorder-7 (GAD-7) should be used to screen for anxiety disorders due to its 93% sensitivity and 97% specificity. GAD-7 screens for anxiety disorders and assesses severity to allow doctors to continue treatment if positive. Anxiety disorders can be screened using GAD-2 or GAD-7. The screening methods differ in that GAD-7 is a seven-item scale and GAD-2 is a quick version of GAD-7 that contains only the first two questions, which are crucial to anxiety disorder screening. GAD-7 is a self-administered survey based on the Diagnostic and Statistical Manual of Mental Disorders-IV (DSM-IV) to identify anxiety disorders like GAD, social phobia, panic disorder, and post-traumatic stress disorder (20,21). GAD-7 contains seven questions to evaluate patients for anxiety disorder symptoms, limitations, and functional impairment over the past two weeks. The four options for each question range from "not at all" to "nearly daily" and are scored from zero to three. A total score



between 5 and 10 indicates a mild anxiety disorder screening, 10-14 indicates a moderate screening, and 14 or more indicates a severe screening. GAD-7 survey is an established and reliable diagnostic tool for primary care and all general populations. Positive GAD-2 has only the first two questions with multiple-scoring answers. If the overall score is three or higher, further evaluation is needed (21).

When screening results show anxiety disorders, DSM-5 criteria should be used to confirm the diagnosis, which includes excessive worrying about almost everything for at least six months, with difficulty controlling these worries. Three of the six other symptoms—restlessness, poor focus, weariness, irritability, muscle tension, and sleep disturbance—must also be present. Patients' occupational and social impairments are used to confirm the diagnosis. All cases of anxiety disorder must rule out other psychological or physical issues to be diagnosed (22).

Various effective treatment options exist for anxiety disorders, demonstrating the ability to alleviate overall symptoms of anxiety and enhance the quality of patients' lives. There exist both pharmacological and non-pharmacological approaches for the treatment of anxiety disorders (13). A variety of psychological therapies, including cognitive behavioral therapy (CBT) and the self-help method, are available as first-line treatments for anxiety disorders. The most well-known method for treating anxiety disorders is cognitive behavioral therapy (CBT). CBT emphasizes the need to identify and replace unhealthy beliefs with more constructive ones, as well as the need to expose patients to real or imagined frightening stimuli to break destructive patterns of behavior. When psychological therapy fails to alleviate symptoms, when anxiety disorder is severe or persistent, or when another medical condition, such as depression and anxiety disorder, is present, pharmacological treatment may be considered as a first line of defense. It is possible to use medications from different classes, such as SSRIs and SNRIs (13, 19).

Among the SSRIs is sertraline, a medicine that works well for treating anxiety disorders alone but fails to alleviate symptoms when other mental health issues, including depression, are present. Three excellent choices for treating GAD include duloxetine, escitalopram, and venlafaxine. Another anxiolytic medicine is benzodiazepine, which is useful in treating most anxiety disorders but has adverse effects, including tolerance and reliance; hence, it is only used for limited periods of time with restrictions while monitoring. When dealing with issues like treatment resistance or co-morbidities, it is common practice to employ a hybrid approach that incorporates elements of both approaches (13).

The treatment strategy should consider the patient's previous successful treatment, preference for all choices, other physical or psychological problems, age or gender, and future pregnancy planning (19). Identifying risk factors and adhering to a clear guideline for detection and treatment are essential in the prevention of anxiety disorders. Implementing effective prevention strategies is crucial, achievable through the reduction of negative risk factors associated with anxiety disorders or the enhancement of protective factors related to these disorders (23). It has been discovered that anxiety and depression are closely connected and highly comorbid (24, 25). Approximately one-fourth of patients with mental illnesses experience comorbidity of anxiety disorders and depression. Anxiety is present in over eighty percent of patients with depression, while depression is found in up to ninety percent of patients with anxiety disorders (24).

## Literature Review

We conducted a search of the PubMed database for prior studies. The determinants of prior studies were in English, published within the last decade, and related to the comorbidity of depression and anxiety among training physicians in various health training programs. Meta-analyses and systematic reviews have been conducted to examine the global prevalence of depression among adolescents in one study (26) and within general populations in two additional studies (27, 28). The studies indicated that 34% of participants exhibited depressive symptoms, with a point prevalence of Major Depressive Disorder (MDD) at 8%. The prevalence was found to be the highest in Africa, Asia, and the Middle East (26). The prevalence of depression in general populations is increasing, potentially due to changes in the calculation of depression rates, variations in help-seeking behavior, or shifts in healthcare professionals' perceptions of the condition (27). In 2017, the global incidence of depressive cases increased by 49.86 percent compared to 1990 (28). A systematic review was conducted to examine the prevalence of anxiety disorders within the general population. This study estimated 301.39 million prevalent cases globally, indicating a 50% increase in anxiety disorders since 1990 (29). A global meta-analysis and systematic review was conducted among resident physicians to

determine the prevalence of depressive symptoms and depression. The study indicated a prevalence of 28.8%, with projections suggesting an increase over time (30).

Three cross-sectional studies were conducted in South Korea among workers (31), Australian doctors (32), and in the U.S. on anesthesiology training physicians from their first year to their first year after graduation (33). The three studies focused on depression, revealing that individuals working over 69 hours per week exhibited a greater likelihood of experiencing moderate to severe depressive symptoms compared to those working 40 hours per week (31). About 60% of doctors who reported ever feeling significantly sad sought some type of professional care; privacy and confidentiality were the most often stated impediments to seeking help (32). Moreover, 32% and 12% have distress or depression, respectively, among anesthesiology residents. A higher chance of distress and depression was found to be associated with more hours worked per week (33). Findings regarding risk factors associated with depression indicated that employed women and individuals with lower income levels exhibited a more pronounced correlation between extended work hours and depressive symptoms (31), females had higher odds of seeking help, as did locally trained doctors and senior doctors (32), and the perception of workplace resource availability, and ability to maintain a work-life balance were associated with a lower overall risk of depression and distress (33). Variations in help-seeking behaviors were associated with medical specialties; psychiatrists exhibited a higher propensity to seek professional assistance, whereas surgeons, pathologists, and radiologists demonstrated a lower likelihood of seeking professional or medical help for depression (32). The prevalence of depression and anxiety among Brazilian medical residents was 46.9% and 56.6%, respectively. Risk factors such as anxiety, burnout, extended duty hours, lack of days off, prevalence of psychological abuse, and female gender were independently correlated with depression. Additionally, female gender, advanced age, prolonged work hours, work-life conflicts, unsupervised assistance, and depression were independently linked to anxiety. The findings were obtained from a cross-sectional study conducted in Brazil, focusing on medical residents in training programs (34). Two cross-sectional studies were conducted to study the prevalence of depression and anxiety among physicians in France (35) and in China (36). These studies showed similar results related to anxiety, in which 32.3% reported having an anxiety disorder, 8.1 % had an MDD, and less than one in five received psychotherapy or antidepressant medications in France (35). 25.67 % of physicians reported having anxiety symptoms, 28.13 % reported having depressed symptoms, and 19.01 % reported having both anxiety and

depression symptoms in China (36). Being a woman and drinking more coffee were directly linked to a higher rate of anxiety disorders. On the other hand, harassment, drinking alcohol, and not knowing how to deal with sickness well at first were all linked to major depression and anxiety disorders (35).<sup>85</sup> Anxiety and depressive symptoms were related to<sup>34</sup> long working hours (above 60 hours per week), frequent night shifts (twice or more per week), poor self-reported physical health, and irregular physical exercise (36). Anxiety disorders and depression were both positively correlated with each other,<sup>75</sup> according to a cross-sectional study performed among emergency physicians in China (37).<sup>7</sup> Two cross-sectional studies were conducted in Saudi Arabia focusing on anxiety disorders, depression, and anxiety among adults aged 18 and older (38), including all medical residents (39). Variations in the prevalence of depression and anxiety among the targeted populations were observed, with findings indicating that 12.4% of the nationwide population in Saudi Arabia was at risk for GAD and 12.7% for MDD. In comparison, 65.8% of medical residents reported experiencing symptoms of depression, while 58.3% reported symptoms of anxiety. <sup>78</sup>Diagnosis and treatment rates were notably low, with only 12.5% of individuals receiving treatment for anxiety and 0.5% for depression (38). Identified risk factors include female sex, lower income, and smoking, which are<sup>41</sup> associated with both Generalized Anxiety Disorder (GAD) and Major Depressive Disorder (MDD). Participation in volunteer activities, regular physical exercise, and the pursuit of daily hobbies serve as protective factors within the general population. The residency degree (R5) and gender (female) were independently associated with anxiety symptoms. Family medicine specialty and female gender were independent predictors of depressive symptoms among medical residents (39). Two cross-sectional studies were conducted involving all medical residents in Saudi Arabia. The initial study was conducted in Jeddah (40), whereas the subsequent study took place in Makkah (41). The studies<sup>2</sup> indicated that the prevalence of diagnosed depression among residents of Jeddah was 75%, categorized by severity: mild depression at<sup>44</sup> 35.6%, moderate depression at 34.2%, and severe depression at 6% (40). In Makkah, the prevalence of anxiety was recorded at 39.5%, while the prevalence of depression was noted at 20.9% (41). Residents engaged in emergency and surgical programs exhibit a heightened risk of developing moderate to severe depression relative to other specialties. Contributing factors include extended work hours, age, sleep duration, and exposure to injustice, all of which elevate the overall risk of<sup>49</sup> anxiety and depression among resident physicians (41). A recent cross-sectional study found a<sup>57</sup> 10% prevalence of moderately severe to severe depression among residents in family medicine in Riyadh, with a significant

correlation between the severity<sup>14</sup> of depression and burnout. A recent cross-sectional study conducted in Makkah examined the prevalence of anxiety disorders and depression among surgical residents and doctors. The findings revealed that 11.8% of surgeons reported a previous history of anxiety or depression, while 30.7% experienced anxiety and 27.5% experienced depression (42). Anxiety demonstrated a significant correlation with participation in the Saudi board program, working over nine hours, and experiencing work-related stress. Depression among surgeons is significantly associated with age, particularly between 50 and 59 years, in addition to factors such as career satisfaction and work-related stress (43). This study aimed to examine the prevalence and factors associated with the comorbidity of depression and anxiety among training physicians in various health training programs under the Ministry of Health (MOH) in Al-Ahsa, Saudi Arabia.

## Materials and methods

### Study design and setting

This descriptive cross-sectional study was conducted among training physicians in various health care training programs under the Ministry of Health from March to December 2024 in Al-Ahsa, in the eastern province region of the Kingdom of Saudi Arabia.

### Sample size

Considering the variables and outcome of the study, assuming the expected 50% of the study population (471), comorbidity of depression and anxiety on training physicians in various health training programs under the Ministry of Health, with an allowable margin error of 5%, at a 95% confidence interval,<sup>15</sup> a minimum sample size of 221 was calculated. After the random sampling, a total of 221 professionals from various fields were included in the data collection.

### Data collection tools<sup>51</sup>

The data was collected by a self-administered questionnaire form, sent by email, and/or WhatsApp. The questionnaire was filled out by the training physicians. The questionnaire was divided into four sections: Socio-Demographic section that included age, gender, nationality, marital status, having children, living with family, monthly income, residency year, training center, and having on-call as part of work. The second was the Lifestyle and medical history section that included exercise, smoking, current or history of chronic medical illnesses, current or history of psychiatric illnesses, ever receiving any professional psychological help, and previous

knowledge about psychological support for trainees (Daem). The third was the Depression assessment section that included <sup>82</sup> the PHQ-9 screening instrument in both <sup>1</sup> Arabic and English versions. The PHQ-9 is a valid and reliable tool for measuring symptoms of depression in all individuals, and the scores are divided into five categories: a score of <5 indicates minimal or no symptoms of depression, a score of 5 to 9 indicates mild symptoms, a score of 10 to 14 indicates moderate symptoms, a score of 15–19 indicates moderately severe symptoms, and of 20 or more indicates severe symptoms. And finally, <sup>2</sup> the fourth was the anxiety disorders assessment section that included the GAD-7 screening instrument. Arabic and English versions of the questionnaire were provided. The scores are classified into four categories: a score of <5 indicates no or minimal anxiety symptoms, a score of 5 to 9 indicates mild symptoms, a score of 10 to 14 indicates moderate symptoms, and a score of 15 or more indicates severe symptoms.

#### **Data analysis**

<sup>39</sup> Data was entered into Microsoft 365 (Microsoft® Corp., Redmond, WA, United States), and analyses were carried out using <sup>1</sup> IBM SPSS Statistics for Windows, version 26 (IBM Corp., Armonk, NY, United States). Descriptive statistics were used to summarize the data by calculating the frequency and percentage of categorical variables. Pearson's Chi-squared test revealed statistically significant differences in depression and anxiety prevalence regarding sociodemographic data, lifestyle, and history of clinical and psychiatric illness ( $p < 0.05$ ).

#### **Ethical consideration**

<sup>55</sup> Official ethical approval was obtained from <sup>2</sup> Research Ethical Committee (REC) with Reference No.: REC-? /? /???. Consent was accepted as a prerequisite for data collection. All collected data were kept confidential and were used only for research.

#### **Budget**

This study was entirely self-funded by the researchers.

## **Results**

### <sup>2</sup> Sociodemographic characteristics

Overall, a total of 242 residents responded to the online survey. Of these, 21 respondents were excluded due to missing more than one entry in the questionnaire. The majority, 157 (71.0%), were between 24 and 29 years old. Nearly half of the respondents, 111 (50.2%), were females. Nearly all the residents included in the study, 220 (99.5%), were Saudis. Around 131 (59.3%) were married, and 123 (55.7%) were without children. A high proportion of the respondents were living with their families, 192 (86.9%), with a monthly income of 15,000-20,000 SAR, 186 (84.2%), and were satisfied with their income, 115 (52.0%). The best part of the shared residents was family medicine 89 (40.3%), in Primary health care (PHC) - Al-Ahsa Academy of Family Medicine 86 (38.9%), and in the first residence year 89 (40.3%). More than half of the respondents reported receiving up to 5 on-call shifts per month, 111 (50.2%). The full sociodemographic characteristics are summarized in [Table 1].

**Table 1: Demographic data of the study respondents (n=221)**

Demographics		Frequency (%)
Age	<24 Y	1 (0.5)
	24-29 Y	157 (71.0)
	30-34 Y	57 (25.8)
	35-39 Y	6 (2.7)
Gender	Males	110 (49.8)
	Females	111 (50.2)
Nationality	Saudi	220 (99.5)
	Non-Saudi	1 (0.5)
Marital status	Single	85 (38.5)
	Married	131 (59.3)
	Divorced	3 (1.4)
	I don't prefer to tell	2 (0.9)
Do you have children?	Yes	98 (44.3)
	No	123 (55.7)
Do you live with your family?	Yes	192 (86.9)
	No	29 (13.1)
Monthly income	< 15,000 SR	5 (2.3)
	15,000-20,000 SR	186 (84.2)
	> 20,000 SR	30 (13.6)

Are you satisfied with your current income?	Satisfied	115 (52.0)
	Neutral	78 (35.3)
	Unsatisfied	28 (12.7)
Training center in Al-Ahsa, Saudi Arabia	King Fahad Hospital-Al-Hofuf (KFHH)	58 (26.2)
	Maternity & Children's Hospital in Al-Ahsa (MCH)	45 (20.4)
	Primary health care (PHC) - Al-Ahsa Academy of Family Medicine	86 (38.9)
	Psychiatric Hospital in Al-Ahsa	3 (1.4)
	Al Jabr Eye and ENT Hospital in Al-Ahsa	8 (3.6)
	Prince Sultan Cardiac Center	3 (1.4)
	PHC Al Salhea	6 (2.7)
	Others	12 (5.4)
Specialty	Family medicine	89 (40.3)
	Preventive medicine	16 (7.2)
	Internal medicine	17 (7.7)
	Pediatric	17 (7.7)
	Obe & Gynae	27 (12.2)
	Cardiac surgery	3 (1.4)
	ICU	13 (5.9)
	General Surgery	4 (1.8)
	Neurosurgery	2 (0.9)
	Orthopedic	2 (0.9)
	Radiology	9 (4.1)
	Urology	7 (3.2)
	Dermatology	3 (1.4)
	Neurology	2 (0.9)
	ENT	7 (3.2)
	Psychiatry	3 (1.4)
Residency year	R1	89 (40.3)
	R2	52 (23.5)
	R3	44 (19.9)
	R4	26 (11.8)
	R5	10 (4.5)
Number of your on-calls per month	No on-call	48 (21.7)
	5 times or less per month	111 (50.2)
	> 5 times per month	62 (28.1)

#### *Lifestyle and medical history characteristics*



A high proportion of the respondents, 160 (72.4%), were not exercising at least 150 minutes per week, and were non-smokers, 188 (85.1%). Most of the shared residents, 187 (84.6%), did not complain of any chronic diseases. The main reported diseases were G6PD and bronchial asthma, 8 (3.6%). Many of the participants, 195 (88.2%), did not complain of any psychiatric illness. The main reported disease was depression, 14 (6.3%). The bulk of them, 182 (82.4%), had not received professional psychological help. However, 128 (57.8%) have heard about SCFHS psychological support for trainees (Daem). The lifestyle and medical history are summarized in [Table 2].

**Table 2:** Lifestyle and medical history of the study respondents (n=221)


Lifestyle and medical history		Frequency (%)
Are you exercising at least 150 minutes per week?	Yes	61 (27.6)
	No	160 (72.4)
Are you a smoker?	Yes	33 (14.9)
	No	188 (85.1)
Do you have a current or past history of chronic medical illnesses?	No	187 (84.6)
	HTN	5 (2.3)
	G6PD	8 (3.6)
	PCOS	3 (1.4)
	DM	5 (2.3)
	Asthma	8 (3.6)
	Thyroid	3 (1.4)
	Anemia	2 (0.9)
Do you have a current or past history of psychiatric illnesses?	No	195 (88.2)
	Depression	14 (6.3)
	Anxiety	7 (3.2)
	ADHD	1 (0.5)
	Bipolar	1 (0.5)
	Burnout	1 (0.5)
	Overthinking	1 (0.5)
	Panic disorder	1 (0.5)

Have you ever received any professional psychological help?	No	182 (82.4)
	Yes, from the outpatient clinic	15 (6.8)
	Yes, from a psychological care app	12 (5.4)
	Yes, from psychological support for trainees (Daem)	2 (0.9)
	Yes, from a relative or friend	6 (2.7)
	Yes, from others	4 (1.8)
Have you heard about SCFHS psychological support for trainees (Daem)?	Yes	128 (57.9)
	No	93 (42.1)

### *Depression prevalence among Saudi residents*

According to the residents' response, only 12.7% (n=27) had no or minimal depression. However, many of the participants showed mild and moderate depression, 35.3% (n=78) and 33% (n=73) respectively. Moreover, some residents had moderately severe or severe depression, 6.8% (n=15) and 12.2% (n=27) respectively [Figure 1]. Depression prevalence varied across different demographic groups [Table 3]. Among residents aged 24–29 years (the highest frequency of the participants), 36.9% had mild depression, 15.2% had no or minimal depression, and 14.6% had moderate depression, while 6.4% and 10.8% exhibited moderately severe and severe depression, respectively. Regarding gender, moderately severe and severe depression were more common in females, 8.1% and 14.4% respectively. A greater number of females (35.1%) recorded mild depression, while the superiority of males (40%) showed no or minimal depression. The preponderance of Saudi residents had mild or minimal depression (35%, 33.2% respectively). The lowest percentage of severely depressed residents was the married ones (9.9%). Severely depressed residents were more likely to be divorced (33.3%) or single (14.1%). Having no children or living alone was accompanied by a higher rate of severe depression among the participants (15.4%, 20.7% respectively). More monthly income and salary satisfaction were accompanied by a lower rate of severe depression. Concerning the training center, the centers whose residents reported minimal depression were Prince Sultan Cardiac Center (66.7%), PHC Al-Salhea, and Al-Jabr Eye and ENT Hospital (50% both). The first and fourth years of residence were accompanied by the highest score of severe depression (16.9%, 15.4% respectively). More on-calls per month were accompanied by a higher rate of severe depression. Smoking and lack of regular exercise for 150 minutes per week were accompanied

by a higher rate of severe depression (15.2%, 14.4% respectively). Healthy residents with no history of chronic disease or psychiatric illness and awareness about Daem SCFHS psychological support had a higher rate of minimal depression (35.8%, 35.9% and 41.4%) respectively [Table 3].



**Table 3: Sociodemographic characteristics, lifestyle, medical history, and different levels of depression**

Sociodemographic characteristics, lifestyle, and medical history		No or minimal depression	Mild depression	Moderate depression	Moderately severe depression	Severe depression	P value
		n (%)	n (%)	n (%)	n (%)	n (%)	
Age	<24 Y	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	0.337
	24-29 Y	49 (31.2%)	58 (36.9%)	23 (14.6%)	10 (6.4%)	17 (10.8%)	
	30-34 Y	20 (35.1%)	19 (33.3%)	5 (8.8%)	5 (8.8%)	8 (14.0%)	
	35-39 Y	4 (66.7%)	1 (16.7%)	0 (0.0%)	0 (0.0%)	1 (16.7%)	
Gender	Males	44 (40.0%)	39 (35.5%)	10 (9.1%)	6 (5.5%)	6 (5.5%)	0.142
	Females	29 (26.1%)	39 (35.1%)	18 (16.2%)	9 (8.1%)	16 (14.4%)	
Nationality	Saudi	73 (33.2%)	77 (35.0%)	28 (12.7%)	15 (6.8%)	27 (12.3%)	0.765


	Non-Saudi	0 (0.0%)	1 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Marital status	Single	25 (29.4%)	30 (35.3%)	11 (12.9%)	7 (8.2%)	12 (14.1%)	0.715
	Married	48 (36.6%)	46 (35.1%)	16 (12.2%)	8 (6.1%)	13 (9.9%)	
	Divorced	0 (0.0%)	1 (33.3%)	1 (33.3%)	0 (0.0%)	1 (33.3%)	
	I don't prefer to tell	0 (0.0%)	1 (50.0%)	0 (0.0%)	0 (0.0%)	1 (50.0%)	
Do you have children?	Yes	39 (39.8%)	34 (34.7%)	12 (12.2%)	5 (5.1%)	8 (8.2%)	0.232
	No	34 (27.6%)	44 (35.8%)	16 (13.0%)	10 (8.1%)	19 (15.4%)	
Do you live with your family?	Yes	68 (35.4%)	63 (32.8%)	25 (13.0%)	15 (7.8%)	21 (10.9%)	0.051
	No	5 (17.2%)	15 (51.7%)	3 (10.3%)	0 (0.0%)	6 (20.7%)	
Monthly income	< 15,000 SR	1 (20.0%)	2 (40.0%)	1 (20.0%)	0 (0.0%)	1 (20.0%)	0.808
	15,000-20,000 SR	58 (31.2%)	68 (36.6%)	23 (12.4%)	14 (7.5%)	23 (12.4%)	
	> 20,000 SR	14 (46.7%)	8 (26.7%)	4 (13.3%)	1 (3.3%)	3 (10.0%)	
Are you satisfied with your current income?	Satisfied	48 (41.7%)	38 (33.0%)	12 (10.4%)	8 (7.0%)	9 (7.8%)	0.010*
	Neutral	18 (23.1%)	33 (42.3%)	13 (16.7%)	5 (6.4%)	9 (11.5%)	
	Unsatisfied	7 (25.0%)	7 (25.0%)	3 (10.7%)	2 (7.1%)	9 (32.1%)	
Training center in Al-Ahsa, Saudi Arabia	King Fahad Hospital-Al-Hofuf (KFHH)	16 (27.6%)	21 (36.2%)	8 (13.8%)	3 (5.2%)	10 (17.2%)	0.134
	Maternity & Children's Hospital in Al-Ahsa (MCH)	5 (11.1%)	17 (37.8%)	8 (17.8%)	7 (15.6%)	8 (17.8%)	
	Primary health care (PHC) - Al-Ahsa Academy of Family Medicine	39 (45.3%)	28 (32.6%)	8 (9.3%)	4 (4.7%)	7 (8.1%)	
	Psychiatric Hospital in Al-Ahsa	1 (33.3%)	1 (33.3%)	0 (0.0%)	0 (0.0%)	1 (33.3%)	
	Al-Jabr Eye and ENT Hospital in Al-Ahsa	4 (50.0%)	2 (25.0%)	0 (0.0%)	1 (12.5%)	1 (12.5%)	
	Prince Sultan Cardiac Center	2 (66.7%)	0 (0.0%)	1 (33.3%)	0 (0.0%)	0 (0.0%)	
	PHC Al Salhea	3 (50.0%)	2 (33.3%)	1 (16.7%)	0 (0.0%)	0 (0.0%)	
	Others	3 (25.0%)	7 (58.3%)	2 (16.7%)	0 (0.0%)	0 (0.0%)	
Residency year	R1	17 (19.1%)	35 (39.3%)	12 (13.5%)	10 (11.2%)	15 (16.9%)	0.095
	R2	21 (40.4%)	21 (40.4%)	4 (7.7%)	3 (5.8%)	3 (5.8%)	
	R3	22 (50.0%)	10 (22.7%)	7 (15.9%)	1 (2.3%)	4 (9.1%)	
	R4	9 (34.6%)	9 (34.6%)	3 (11.5%)	1 (3.8%)	4 (15.4%)	
	R5	4 (40.0%)	3 (30.0%)	2 (20.0%)	0 (0.0%)	1 (10.0%)	
Number of your on-calls per month	No on-call	24 (50.0%)	16 (33.3%)	6 (12.5%)	0 (0.0%)	2 (4.2%)	0.008*
	5 times or less per month	38 (34.2%)	39 (35.1%)	14 (12.6%)	7 (6.3%)	13 (11.7%)	
	> 5 times per month	11 (17.7%)	23 (37.1%)	8 (12.9%)	8 (12.9%)	12 (19.4%)	
Are you exercising at least 150 minutes per week?	Yes	26 (42.6%)	26 (42.6%)	3 (4.9%)	2 (3.3%)	4 (6.6%)	0.023*
	No	47 (29.4%)	52 (32.5%)	25 (15.6%)	13 (8.1%)	23 (14.4%)	
Are you a smoker?	Yes	10 (30.3%)	13 (39.4%)	3 (9.1%)	2 (6.1%)	5 (15.2%)	0.915
	No	63 (33.5%)	65 (34.6%)	25 (13.3%)	13 (6.9%)	22 (11.7%)	
Do you have a current or past history of chronic medical	No	67 (35.8%)	66 (35.3%)	21 (11.2%)	12 (6.4%)	21 (11.2%)	0.183
	HTN	1 (20.0%)	1 (20.0%)	(40.0%)	0 (0.0%)	1 (20.0%)	
	G6pD	2 (25.0%)	0 (0.0%)	1 (12.5%)	3 (37.5%)	2 (25.0%)	

illnesses?	PCOS	0 (0.0%)	2 (66.7%)	1 (33.3%)	0 (0.0%)	0 (0.0%)	
	DM	0 (0.0%)	2 (40.0%)	1 (20.0%)	0 (0.0%)	2 (40.0%)	
	Asthma	2 (25.0%)	4 (50.0%)	1 (12.5%)	0 (0.0%)	1 (12.5%)	
	Thyroid	0 (0.0%)	2 (66.7%)	1 (33.3%)	0 (0.0%)	0 (0.0%)	
	Anemia	1 (50.0%)	1 (50.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Do you have a current or past history of psychiatric illnesses?	No	70 (35.9%)	71 (36.4%)	22 (11.3%)	12 (6.2%)	20 (10.3%)	0.101
	Depression	0 (0.0%)	3 (21.4%)	5 (35.7%)	2 (14.3%)	4 (28.6%)	
	Anxiety	2 (28.6%)	2 (28.6%)	1 (14.3%)	1 (14.3%)	1 (14.3%)	
	ADHD	0 (0.0%)	1 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
	Bipolar	0 (0.0%)	1 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
	Burnout	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	
	Overthinking	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	
	Panic disorder	1 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Have you ever received any professional psychological help?	No	67 (36.8%)	63 (34.6%)	22 (12.1%)	12 (6.6%)	18 (9.9%)	0.236
	Yes, from the outpatient clinic	3 (20.0%)	5 (33.3%)	1 (6.7%)	1 (6.7%)	5 (33.3%)	
	Yes, from the psychological care app	0 (0.0%)	5 (41.7%)	3 (25.0%)	2 (16.7%)	2 (16.7%)	
	Yes, from psychological support for trainees (Daem)	0 (0.0%)	1 (50.0%)	0 (0.0%)	0 (0.0%)	1 (50.0%)	
	Yes, from a relative or friend	2 (33.3%)	2 (33.3%)	2 (33.3%)	0 (0.0%)	0 (0.0%)	
	Yes (from others)	1 (25.0%)	2 (50.0%)	0 (0.0%)	0 (0.0%)	1 (25.0%)	
Have you heard about CPHS psychological support for trainees (Daem)?	Yes	53 (41.4%)	48 (37.5%)	17 (13.3%)	4 (3.1%)	6 (4.7%)	0.001*
	No	20 (21.5%)	30 (32.3%)	11 (11.8%)	11 (11.8%)	21 (22.6%)	

\* = Significant (P-value is less than 0.05)

#### ***Specialty of the residents and different levels of depression***

General surgery and orthopedics were the specialties encountered in severe depression (50% both), followed by internal medicine, Obe & Gynae, and Pediatrics (23.5%, 18.5% & 17.6%) respectively. No or minimal depression was observed mainly in residents of cardiac surgery, followed by preventive medicine (66.7% and 56.3%), respectively with no significant variation [Fig. 2 & Table 4].



**Table 4: Specialty of the residents and different levels of depression**

Specialty		No or minimal depression	Mild depression	Moderate depression	Moderately severe depression	Severe depression	P value
		n (%)	n (%)	n (%)	n (%)	n (%)	
Specialty	Family medicine	38 (42.7%)	30 (33.7%)	10 (11.2%)	4 (4.5%)	7 (7.9%)	0.208
	Preventive medicine	9 (56.3%)	5 (31.3%)	1 (6.3%)	0 (0.0%)	1 (6.3%)	
	Internal medicine	3 (17.6%)	4 (23.5%)	4 (23.5%)	2 (11.8%)	4 (23.5%)	
	Pediatric	2 (11.8%)	2 (11.8%)	2 (11.8%)	4 (23.5%)	3 (17.6%)	
	Obe & Gynae	2 (7.4%)	11 (40.7%)	6 (22.2%)	3 (11.1%)	5 (18.5%)	
	Cardiac surgery	2 (66.7%)	0 (0.0%)	1 (33.3%)	0 (0.0%)	0 (0.0%)	
	ICU	6 (46.2%)	5 (38.5%)	0 (0.0%)	1 (7.7%)	1 (7.7%)	
	General Surgery	1 (25.0%)	1 (25.0%)	0 (0.0%)	0 (0.0%)	2 (50.0%)	
	Neurosurgery	0 (0.0%)	1 (50.0%)	1 (50.0%)	0 (0.0%)	0 (0.0%)	
	Orthopedic	0 (0.0%)	1 (50.0%)	0 (0.0%)	0 (0.0%)	1 (50.0%)	
	Radiology	3 (33.3%)	2 (22.2%)	3 (33.3%)	0 (0.0%)	1 (11.1%)	


Urology	2 (28.6%)	4 (57.1%)	0 (0.0%)	0 (0.0%)	1 (14.3%)
Dermatology	1 (33.3%)	2 (66.7%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Neurology	0 (0.0%)	2 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
ENT	3 (42.9%)	2 (28.6%)	0 (0.0%)	1 (14.3%)	1 (14.3%)
Psychiatry	1 (33.3%)	2 (66.7%)	0 (0.0%)	0 (0.0%)	0 (0.0%)

\* = Significant (P-value is less than 0.05)

### **Anxiety prevalence among Saudi residents**

According to the residents' response, nearly half, 49.8% (n=110), had no or minimal anxiety. However, the other half showed mild, moderate, and severe depression, 22.6% (n=50), 12.7% (n=28), and 14.9% (n=33) respectively [Figure 3]. Anxiety prevalence varied across different demographic groups [Table 5]. Among residents aged 24–29 years (the highest frequency of the participants), 50.3% had no or minimal anxiety, 21.7% had mild anxiety, and 12.7% had moderate anxiety, while 15.3% exhibited severe anxiety. Regarding gender, moderate and severe anxiety were more common in females, 15.3% and 19.8% respectively. A greater number of males (54.5%) recorded minimal anxiety, while the superiority of females (45%) showed no or minimal anxiety. The preponderance of Saudi residents had minimal or mild anxiety (50%, 22.3% respectively). The lowest percentage of residents with severe anxiety was the single ones (10.6%), followed by married, then divorced residents (16.8% and 33.3%), respectively. Having children or living alone was accompanied by a higher rate of severe anxiety among the participants (15.3%, 20.7% respectively). More monthly income and salary satisfaction were accompanied by a lower rate of severe anxiety. Concerning the training center, the centers whose residents reported no or minimal anxiety were Prince Sultan Cardiac Center (100%), Primary health care – Al-Ahsaa Academy of Family Medicine (57%), PHC Al-Salhea, and Al-Jabr Eye and ENT Hospital (50% both). The fourth and the years of residence were accompanied by the highest score of severe anxiety (23.1%, 19.1% respectively). More on-calls per month were accompanied by a higher rate of severe anxiety. Smoking and lack of regular exercise for 150 minutes per week were accompanied by a higher rate of severe anxiety (18.2% and 18.8% respectively). Healthy residents with no history of chronic disease or psychiatric illness and

awareness about Daem SCFHS psychological support had a higher rate of minimal anxiety (51.3%, 53.3% and 57% respectively) [Table 5].



**Table 5: Sociodemographic characteristics and different levels of anxiety**

Sociodemographic characteristics, lifestyle, and medical history		No or minimal anxiety	Mild anxiety	Moderate anxiety	Severe anxiety	P value
		n (%)	n (%)	n (%)	n (%)	
Age	<24 Y	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	0.338
	24-29 Y	79 (50.3%)	34 (21.7%)	20 (12.7%)	24 (15.3%)	
	30-34 Y	26 (45.6%)	15 (26.3%)	8 (14.0%)	8 (14.0%)	
	35-39 Y	5 (83.3%)	1 (16.7%)	0 (0.0%)	0 (0.0%)	
Gender	Males	60 (54.5%)	28 (25.5%)	11 (10.0%)	11 (10.0%)	0.087
	Females	50 (45.0%)	22 (19.8%)	17 (15.3%)	22 (19.8%)	
Nationality	Saudi	110 (50.0%)	49 (22.3%)	28 (12.7%)	33 (15.0%)	0.329
	Non- Saudi	0 (0.0%)	1 (100.0%)	0 (0.0%)	0 (0.0%)	
Marital status	Single	38 (44.7%)	23 (27.1%)	15 (17.6%)	9 (10.6%)	0.278
	Married	70 (53.4%)	27 (20.6%)	12 (9.2%)	22 (16.8%)	
	Divorced	1 (33.3%)	0 (0.0%)	1 (33.3%)	1 (33.3%)	
	I don't prefer to tell	1 (50.0%)	0 (0.0%)	0 (0.0%)	1 (50.0%)	
Do you have children?	Yes	54 (55.1%)	18 (18.4%)	11 (11.2%)	15 (15.3%)	0.437




	No	56 (45.5%)	32 (26.0%)	17 (13.8%)	18 (14.6%)	
Do you live with your family?	Yes	102 (53.1%)	36 (18.8%)	27 (14.1%)	27 (14.1%)	0.001*
	No	8 (27.6%)	14 (48.3%)	1 (3.4%)	6 (20.7%)	
Monthly income	< 15,000 SR	3 (60.0%)	0 (0.0%)	0 (0.0%)	2 (40.0%)	0.527
	15,000-20,000 SR	93 (50.0%)	42 (22.6%)	23 (12.4%)	28 (15.1%)	
	> 20,000 SR	14 (46.7%)	8 (26.7%)	5 (16.7%)	3 (10.0%)	
Are you satisfied with your current income?	Satisfied	70 (60.9%)	23 (20.0%)	8 (7.0%)	14 (12.2%)	0.012*
	Neutral	29 (37.2%)	22 (28.2%)	15 (19.2%)	12 (15.4%)	
	Unsatisfied	11 (39.3%)	5 (17.9%)	5 (17.9%)	7 (25.0%)	
Training center in Al-Ahsa, Saudi Arabia	King Fahad Hospital-Al-Hofuf (KFHH)	28 (48.3%)	17 (29.3%)	5 (8.6%)	8 (13.8%)	0.29
	Maternity & Children's Hospital in Al-Ahsa (MCH)	17 (37.8%)	9 (20.0%)	10 (22.2%)	9 (20.0%)	
	Primary health care (PHC) - Al-Ahsa Academy of Family Medicine	49 (57.0%)	15 (17.4%)	10 (11.6%)	12 (14.0%)	
	Psychiatric Hospital in Al-Ahsa	1 (33.3%)	0 (0.0%)	0 (0.0%)	2 (66.7%)	
	Al-Jabr Eye and ENT Hospital in Al-Ahsa	4 (50.0%)	2 (25.0%)	1 (12.5%)	1 (12.5%)	
	Prince Sultan Cardiac Center	3 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
	PHC Al Salhea	3 (50.0%)	3 (50.0%)	0 (0.0%)	0 (0.0%)	
	Others	5 (41.7%)	4 (33.3%)	2 (16.7%)	1 (8.3%)	
Residency year	R1	36 (40.4%)	21 (23.6%)	15 (16.9%)	17 (19.1%)	0.065
	R2	28 (53.8%)	16 (30.8%)	6 (11.5%)	2 (3.8%)	
	R3	29 (65.9%)	5 (11.4%)	3 (6.8%)	7 (15.9%)	
	R4	10 (38.5%)	6 (23.1%)	4 (15.4%)	6 (23.1%)	
	R5	7 (70.0%)	2 (20.0%)	0 (0.0%)	1 (10.0%)	
Number of your on-calls per month	No on-call	31 (64.6%)	11 (22.9%)	2 (4.2%)	4 (8.3%)	0.036*
	5 times or less per month	55 (49.5%)	26 (23.4%)	12 (10.8%)	18 (16.2%)	
	> 5 times per month	24 (38.7%)	13 (21.0%)	14 (22.6%)	11 (17.7%)	
Are you exercising at least 150 minutes per week?	Yes	33 (54.1%)	19 (31.1%)	6 (9.8%)	3 (4.9%)	0.027*
	No	77 (48.1%)	31 (19.4%)	22 (13.8%)	30 (18.8%)	
Are you a smoker?	Yes	13 (39.4%)	9 (27.3%)	5 (15.2%)	6 (18.2%)	0.643
	No	97 (51.6%)	41 (21.8%)	23 (12.2%)	27 (14.4%)	
Do you have a current or past history of chronic medical illnesses?	No	96 (51.3%)	45 (24.1%)	21 (11.2%)	25 (13.4%)	0.028*
	HTN	1 (20.0%)	1 (20.0%)	2 (40.0%)	1 (20.0%)	
	G6pD	2 (25.0%)	0 (0.0%)	2 (25.0%)	4 (50.0%)	
	PCOS	3 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
	DM	1 (20.0%)	1 (20.0%)	0 (0.0%)	3 (60.0%)	
	Asthma	3 (37.5%)	2 (25.0%)	3 (37.5%)	0 (0.0%)	
	Thyroid	2 (66.7%)	1 (33.3%)	0 (0.0%)	0 (0.0%)	
Do you have a current or past history of psychiatric illnesses?	Anemia	2 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0.052
	No	104 (53.3%)	44 (22.6%)	22 (11.3%)	25 (12.8%)	
	Depression	3 (21.4%)	3 (21.4%)	4 (28.6%)	4 (28.6%)	
	Anxiety	2 (28.6%)	2 (28.6%)	1 (14.3%)	2 (28.6%)	

	ADHD	0 (0.0%)	1 (100.0%)	0 (0.0%)	0 (0.0%)	
	Bipolar	0 (0.0%)	0 (0.0%)	1 (100.0%)	0 (0.0%)	
	Burnout	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	
	Overthinking	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (100.0%)	
	Panic disorder	1 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Have you ever received any professional psychological help?	No	100 (54.9%)	36 (19.8%)	22 (12.1%)	24 (13.2%)	0.058
	Yes, from the outpatient clinic	3 (20.0%)	5 (33.3%)	2 (13.3%)	5 (33.3%)	
	Yes, from a psychological care app	2 (16.7%)	5 (41.7%)	2 (16.7%)	3 (25.0%)	
	Yes, from psychological support for trainees (Daem)	0 (0.0%)	0 (0.0%)	1 (50.0%)	1 (50.0%)	
	Yes, from a relative or friend	3 (50.0%)	3 (50.0%)	0 (0.0%)	0 (0.0%)	
	Yes (from others)	2 (50.0%)	1 (25.0%)	1 (25.0%)	0 (0.0%)	
Have you heard about CFHS psychological support for trainees (Daem)?	Yes	73 (57.0%)	30 (23.4%)	11 (8.6%)	14 (10.9%)	0.014*
	No	37 (39.8%)	20 (21.5%)	17 (18.3%)	19 (20.4%)	

\* = Significant (*P*-value is less than 0.05)

### ***Specialty of the residents and different levels of anxiety***

Orthopedics and Psychiatry were the specialties encountered in severe anxiety (50% and 33.3% respectively), followed by pediatrics (29.4%). On the other hand, all cardiac surgery and neurology residents (100%) showed no or minimal anxiety, followed by radiology and psychiatry residents (77.8% and 66.7%), respectively, with no significant variation [Fig. 4 & Table 6].




**Table 6: Specialty of the residents and different levels of depression**

Specialty		No or minimal anxiety	Mild anxiety	Moderate anxiety	Severe anxiety	P value
		n (%)	n (%)	n (%)	n (%)	
Specialty	Family medicine	48 (53.9%)	18 (20.2%)	18 (20.2%)	5 (5.6%)	0.653
	Preventive medicine	8 (50.0%)	5 (31.3%)	1 (6.3%)	2 (12.5%)	
	Internal medicine	6 (35.3%)	5 (29.4%)	2 (11.8%)	4 (23.5%)	
	Pediatric	7 (41.2%)	2 (11.8%)	3 (17.6%)	5 (29.4%)	
	Obe & Gynae	9 (33.3%)	7 (25.9%)	7 (25.9%)	4 (14.8%)	
	Cardiac surgery	3 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	
	ICU	7 (53.8%)	6 (46.2%)	0 (0.0%)	0 (0.0%)	
	General Surgery	2 (50.0%)	1 (25.0%)	0 (0.0%)	1 (25.0%)	
	Neurosurgery	1 (50.0%)	1 (50.0%)	0 (0.0%)	0 (0.0%)	
	Orthopedic	0 (0.0%)	0 (0.0%)	1 (50.0%)	1 (50.0%)	
	Radiology	7 (77.8%)	1 (11.1%)	0 (0.0%)	1 (11.1%)	
	Urology	2 (28.6%)	2 (28.6%)	2 (28.6%)	1 (14.3%)	
	Dermatology	2 (66.7%)	1 (33.3%)	0 (0.0%)	0 (0.0%)	

Neurology	2 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
ENT	4 (57.1%)	1 (14.3%)	1 (14.3%)	1 (14.3%)
Psychiatry	2 (66.7%)	0 (0.0%)	0 (0.0%)	1 (33.3%)

#### *Specialty of the residents and comorbidity of depression and anxiety*

Orthopedics was the specialty encountered in the highest percentage of comorbidity of depression and anxiety (100%), followed by neurosurgery, Obe & Gynae, internal medicine, and pediatrics. On the other hand, cardiac surgery showed the least comorbidity of both depression and anxiety, followed by preventive medicine [Fig. 5 & Table 7].



**Table 7: Specialty of the residents and comorbidity of depression and anxiety**

	Depression	Anxiety
--	------------	---------

Family medicine	57.3%	46.1%
Preventive medicine	43.8%	50.0%
Internal medicine	82.4%	64.7%
Pediatric	88.2%	58.8%
Obe & Gynae	92.6%	66.7%
Cardiac surgery	33.3%	0.0%
ICU	53.8%	46.2%
General Surgery	75.0%	50.0%
Neurosurgery	100.0%	50.0%
Orthopedic	100.0%	100.0%
Radiology	66.7%	22.2%
Urology	71.4%	71.4%
Dermatology	66.7%	33.3%
Neurology	100.0%	0.0%
ENT	57.1%	42.9%
Psychiatry	66.7%	33.3%

## <sup>37</sup> Discussion

In the present study, the prevalence of depression, anxiety, and burnout among Saudi board residents and their possible association with predictor variables were assessed. The <sup>47</sup>PHQ-9 and <sup>5</sup>GAD-7 measures are useful for detecting anxiety and depression in this population, while not being diagnostic assessments. <sup>5</sup>This study explored anxiety and depression levels among the Saudi board residents, highlighting the profound impact of sociodemographic characteristics, medical history, and lifestyle on mental health. <sup>28</sup>A sample size of 221 participants revealed the prevalence of depression and anxiety (from mild to severe) of 67% and 50.8% <sup>1</sup>respectively, significantly higher than reported by previous Saudi studies (44,45).

### <sup>5</sup> Sociodemographic differences in mental health

Our results indicated that female Saudi residents reported significantly higher anxiety and depression rates than their male counterparts, consistent with earlier studies (46-48). Women are more likely to experience anxiety and depression during times of stress, possibly because of the various <sup>75</sup>roles they play in society, such as juggling caring and professional obligations (49,50). Furthermore, women may be more vulnerable to chronic stress due to biological processes like the impact of sex hormones on stress management (51). This contradicts the results of other

research on medical students in Jeddah, which <sup>1</sup> found no significant differences in depression rates between genders (52).

Our findings showed that unmarried people <sup>2</sup> were more likely to experience depression and anxiety than married people. This corresponds to a previous study that used nationally representative of approximately 541 million adults, <sup>6</sup> de-identified individual-level data from seven countries: the United States, the United Kingdom, Mexico, Ireland, Korea, China, and Indonesia. The study included <sup>6</sup> 106,556 cross-sectional and 20,865 longitudinal participants. The period of follow-up varied from four to eighteen years (53).

The current study found that higher income and its associated satisfaction were linked to lower rates of both depression and anxiety. <sup>23</sup> These results were in line with a previous study that revealed that low-income workers <sup>77</sup> are more vulnerable to depression and anxiety (30).

The current study uncovered a significant disparity <sup>2</sup> in the prevalence of depression and anxiety among inhabitants of various training centers. Moreover, more on-call duties were accompanied by more depression and anxiety. This was previously clarified as potentially referring to varying levels of support in the workplace. The safeguarding of laborers' health has emerged as a global concern. To enhance the mental health of employees, it is necessary to provide continuous management and global attention, as mental disorders can be prevented in the workplace (54). Moreover, workers who reported <sup>26</sup> higher levels of work and interpersonal stress were more likely to experience feelings of defeat, which in turn led to an increase in <sup>11</sup> depression and anxiety symptoms. This mediation effect was more pronounced for individuals with reduced levels of social support (55). Furthermore, Poor work ability and work stress are associated with anxiety and depression in healthcare professionals (56).

#### ***Residents' specialty and variation in mental health***

The current study showed a clear disparity <sup>2</sup> in the prevalence of depression and anxiety among inhabitants of various training specialties. The highest risk was in orthopedic, neurosurgery, and Obstetrics & Gynaecology specialties, and the lowest was in cardiac surgery and preventive medicine. <sup>23</sup> These results were in line with a previous study that found that about 30% of Saudi surgeons suffer from comorbidity <sup>76</sup> of depression and anxiety (43). The high risk of comorbidity

<sup>62</sup> of depression and anxiety among orthopedic residents was previously observed in a systematic review, which mentioned that <sup>1</sup> Depression and anxiety are highly prevalent among orthopedic residents, with significant levels of psychological distress affecting nearly one-third of this population (57). Globally, depending on the tool used, <sup>29</sup> the prevalence of depression or depressed symptoms among resident physicians ranged from 20.9% to 43.2%, with the overall estimate rising with each passing year (29). Moreover, Anxiety was assessed to be 17% and depression to be 45% during COVID-19. highest rates of depression in pediatrics, internal medicine, and anesthesiology. Pediatrics and internal medicine have the highest levels of anxiety (58).

#### ***Lifestyle and medical history differences in mental health***

The present work revealed that regular exercise, at least for 150 minutes per week, was accompanied by a lower rate of depression and anxiety among the participants. The same findings were observed by Martin-Del-Campo et al. (2023), who found that <sup>3</sup> moderate physical activity was associated with lower depression, anxiety, and stress in healthcare workers during the COVID-19 pandemic. They attributed these results to the increases in serotonin, dopamine, and norepinephrine levels (59). Also, other research found that healthcare workers' mental health can be enhanced by working out at home using a variety of fitness applications and little to no equipment. To assess interest, adoption, adherence, and the mental, physical, and financial impacts of the suite of applications, larger efficacy trials are required, in which all healthcare workers from an organization are given more opportunity to use them. Such studies could point to a low-cost, high-reward strategy that health care networks could implement, integrated with wellness initiatives, to lessen the rising mental health issue in healthcare. This <sup>2</sup> may be due to the promotion of brain-derived neurotrophic factors (BDNF), which support brain health, or due to endorphin release, which improves mood and reduces pain perception, or stress hormone regulation, which reduces the cortisol levels (60). Moreover, one of the main causes of depression is a sedentary lifestyle and inactivity. Sport and <sup>12</sup> physical activity provide a potential, evidence-based alternative for the prevention and treatment of depression because of their widespread accessibility, substantial and clinically meaningful benefits, and near-complete lack of side effects. <sup>81</sup> Therefore, in order to improve mental health, health professionals should support any increase in physical activity. Exercise is a safe and affordable way to treat depression, so interventions aimed at encouraging and maintaining its use should be given much more thought

in clinical practice. It could also be a desirable choice <sup>3</sup> for public health promotion and long-term mental health care systems around the world (61).

The current study elucidated that smoking among the participants has no role in minimizing the rate of depression or anxiety. This contradicts the results of other research, which found that the correlation between mental illness and smoking is evident, as individuals with mental health conditions exhibit both increased prevalence and greater intensity of smoking behaviors (62).

The present study concluded that healthcare workers with a pre-existing history of chronic or psychiatric disease <sup>7</sup> had more than double the risk of depression and anxiety. The same results were recorded by a prospective cohort study of nurses (2020–2022) in China reported that a history of diabetes and hypertension significantly increased anxiety and depressive symptoms (63). Previous research observed that anxiety and depression prevalence are strongly influenced by chronic illnesses. Increased prevalence of these mental health conditions is linked to cardiovascular disease, COPD, cancer, neurological disorders, including multiple sclerosis and Parkinson's, and <sup>1</sup> chronic pain conditions like fibromyalgia and arthritis. Patients with various chronic conditions need to have their mental health addressed. Furthermore, the psychological effects of chronic illnesses may be influenced by gender (64). Additionally, <sup>1</sup> the trend we observed, which indicated that females diagnosed with psychological disorders had a higher prevalence of depression than males, is consistent with previous research <sup>1</sup> that indicates that a history of psychiatric disorders substantially influences the prevalence of depression (65).

## Conclusion

Although depression and anxiety are prevalent in healthcare professionals, they are also prevalent in residents, irrespective of their specialty. These conditions are frequently referred to as "invisible diseases" due to their challenging identification and diagnosis. A significant determinant of anxiety in this study was gender, as females consistently exhibited a <sup>1</sup> higher prevalence of anxiety than their male counterparts. Numerous studies, both domestically and internationally, have also documented this trend. The <sup>1</sup> levels of anxiety and depression were substantially correlated with lifestyle factors <sup>1</sup> and health-related concerns, particularly the history of chronic diseases and psychiatric disorders. This investigation illustrated the correlation



between anxiety symptoms and depressive symptoms in residents who receive assistance at their place of employment. The safeguarding of inhabitants' health has evolved into a worldwide concern. To enhance the mental health of inhabitants, it is necessary to provide continuous management and global attention, as mental disorders can be prevented in the workplace. To address the constraints of the current investigation, additional longitudinal and clinical investigations are required.

## References

1. Otte C, Gold SM, Penninx BW, Pariante CM, Etkin A, Fava M, Mohr DC, Schatzberg AF. Major depressive disorder. *Nat Rev Dis Primers*. 2016 Sep 15;2:16065. doi: 10.1038/nrdp.2016.65. PMID: 27629598.
2. Harmer B, Lee S, Duong TVH, Saadabadi A. Suicidal Ideation. 2023 Dec 4. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. PMID: 33351435.
3. Razzak HA, Harbi A, Ahli S. Depression: Prevalence and Associated Risk Factors in the United Arab Emirates. *Oman Med J*. 2019 Jul;34(4):274-282. doi: 10.5001/omj.2019.56. PMID: 31360314; PMCID: PMC6642715.
4. Chand SP, Arif H. Depression. 2023 Jul 17. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. PMID: 28613597.
5. Trivedi MH. Major Depressive Disorder in Primary Care: Strategies for Identification. *J Clin Psychiatry*. 2020 Mar 17;81(2):UT17042BR1C. doi: 10.4088/JCP.UT17042BR1C. PMID: 32220155.
6. Maurer DM, Raymond TJ, Davis BN. Depression: Screening and Diagnosis. *Am Fam Physician*. 2018 Oct 15;98(8):508-515. PMID: 30277728.
7. Ford J, Thomas F, Byng R, McCabe R. Use of the Patient Health Questionnaire (PHQ-9) in Practice: Interactions between patients and physicians. *Qual Health Res*. 2020 Nov;30(13):2146-2159. doi: 10.1177/1049732320924625. Epub 2020 Jun 20. PMID: 32564676; PMCID: PMC7549295.

8. Tolentino JC, Schmidt SL. DSM-5 Criteria and Depression Severity: Implications for Clinical Practice. *Front Psychiatry*. 2018 Oct 2;9:450. doi: 10.3389/fpsyt.2018.00450. PMID: 30333763; PMCID: PMC6176119.
9. McLaughlin KA. The public health impact of major depression: a call for interdisciplinary prevention efforts. *Prev Sci*. 2011 Dec;12(4):361-71. doi: 10.1007/s11121-011-0231-8. PMID: 21732121; PMCID: PMC3219837.
10. Gautam S, Jain A, Gautam M, Vahia VN, Grover S. Clinical Practice Guidelines for the management of Depression. *Indian J Psychiatry*. 2017 Jan;59(Suppl 1):S34-S50. doi: 10.4103/0019-5545.196973. PMID: 28216784; PMCID: PMC5310101.
11. Kovich H, Kim W, Quaste AM. Pharmacologic Treatment of Depression. *Am Fam Physician*. 2023 Feb;107(2):173-181. PMID: 36791444.
12. Dean E. Anxiety. *Nurs Stand*. 2016 Jul 13;30(46):15. doi: 10.7748/ns.30.46.15.s17. PMID: 27406490.
13. Penninx BW, Pine DS, Holmes EA, Reif A. Anxiety disorders. *Lancet*. 2021 Mar 6;397(10277):914-927. doi: 10.1016/S0140-6736(21)00359-7. Epub 2021 Feb 11. Erratum in: *Lancet*. 2021 Mar 6;397(10277):880. PMID: 33581801; PMCID: PMC9248771.
14. Zimmermann M, Chong AK, Vechiu C, Papa A. Modifiable risk and protective factors for anxiety disorders among adults: A systematic review. *Psychiatry Res*. 2020 Mar;285:112705. doi: 10.1016/j.psychres.2019.112705. Epub 2019 Dec 4. PMID: 31839417.
15. Zhang X, Norton J, Carrière I, Ritchie K, Chaudieu I, Ancelin ML. Risk factors for late-onset generalized anxiety disorder: results from a 12-year prospective cohort (the ESPRIT study). *Transl Psychiatry*. 2015 Mar 31;5(3):e536. doi: 10.1038/tp.2015.31. PMID: 25826111; PMCID: PMC4429171.
16. Blanco C, Rubio J, Wall M, Wang S, Jiu CJ, Kendler KS. Risk factors for anxiety disorders: common and specific effects in a national sample. *Depress Anxiety*. 2014 Sep;31(9):756-64. doi: 10.1002/da.22247. Epub 2014 Feb 27. PMID: 24577934; PMCID: PMC4147018.
17. Narmandakh A, Roest AM, de Jonge P, Oldehinkel AJ. Psychosocial and biological risk factors of anxiety disorders in adolescents: a TRAILS report. *Eur Child Adolesc Psychiatry*.

- 2021 Dec;30(12):1969-1982. doi: 10.1007/s00787-020-01669-3. Epub 2020 Oct 28. PMID: 33113027; PMCID: PMC8563629.
18. Ströhle A, Gensichen J, Domschke K. The Diagnosis and Treatment of Anxiety Disorders. *Dtsch Arztebl Int.* 2018 Sep 14;155(37):611-620. doi: 10.3238/arztebl.2018.0611. PMID: 30282583; PMCID: PMC6206399.
  19. Szuhany KL, Simon NM. Anxiety Disorders: A Review. *JAMA.* 2022 Dec 27;328(24):2431-2445. doi: 10.1001/jama.2022.22744. PMID: 36573969.
  20. Sapra A, Bhandari P, Sharma S, Chanpura T, Lopp L. Using Generalized Anxiety Disorder-2 (GAD-2) and GAD-7 in a Primary Care Setting. *Cureus.* 2020 May 21;12(5):e8224. doi: 10.7759/cureus.8224. PMID: 32582485; PMCID: PMC7306644.
  21. Munir S, Takov V. Generalized Anxiety Disorder. 2022 Oct 17. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. PMID: 28722900.
  22. Cabral MD, Patel DR. Risk Factors and Prevention Strategies for Anxiety Disorders in Childhood and Adolescence. *Adv Exp Med Biol.* 2020;1191:543-559. doi: 10.1007/978-981-32-9705-0\_27. PMID:32002945.
  23. Tiller JW. Depression and anxiety. *Med J Aust.* 2013 Sep 16;199(S6):S28-31. doi: 10.5694/mja12.10628. PMID: 25370281.
  24. Kalin NH. The Critical Relationship Between Anxiety and Depression. *Am J Psychiatry.* 2020 May 1;177(5):365-367. Doi10.1176/appi.ajp.2020.20030305. PMID: 32354270.
  25. Shorey S, Ng ED, Wong CHJ. Global prevalence of depression and elevated depressive symptoms among adolescents: A systematic review and meta-analysis. *Br J Clin Psychol.* 2022 Jun;61(2):287-305. doi: 10.1111/bjc.12333. Epub 2021 Sep 26. PMID: 34569066.
  26. Moreno-Agostino D, Wu YT, Daskalopoulou C, Hasan MT, Huisman M, Prina M. Global trends in the prevalence and incidence of depression: a systematic review and meta-analysis. *J Affect Disord.* 2021 Feb 15;281:235-243. doi: 10.1016/j.jad.2020.12.035. Epub 2020 Dec 9. PMID: 33338841.
  27. Liu Q, He H, Yang J, Feng X, Zhao F, Lyu J. Changes in the global burden of depression from 1990 to 2017: Findings from the Global Burden of Disease study. *J Psychiatr Res.* 2020 Jul;126:134-140. doi: 10.1016/j.jpsychires.2019.08.002. Epub 2019 Aug 10. PMID: 31439359.

28. Yang X, Fang Y, Chen H, Zhang T, Yin X, Man J, Yang L, Lu M. Global, regional and national burden of anxiety disorders from 1990 to 2019: results from the Global Burden of Disease Study 2019. *Epidemiol Psychiatr Sci.* 2021 May 6;30:e36. doi: 10.1017/S2045796021000275. PMID: 33955350; PMCID: PMC8157816.
29. Mata DA, Ramos MA, Bansal N, Khan R, Guille C, Di Angelantonio E, Sen S. Prevalence of Depression and Depressive Symptoms Among Resident Physicians: A Systematic Review and Meta-analysis. *JAMA.* 2015 Dec 8;314(22):2373-83. doi: 10.1001/jama.2015.15845. PMID: 26647259; PMCID: PMC4866499.
30. Choi E, Choi KW, Jeong HG, Lee MS, Ko YH, Han C, Ham BJ, Chang J, Han KM. Long working hours and depressive symptoms: moderation by gender, income, and job status. *J Affect Disord.* 2021 May 1;286:99-107. doi: 10.1016/j.jad.2021.03.001. Epub 2021 Mar 4. PMID: 33714177.
31. Muhamad Ramzi NSA, Deady M, Petrie K, Crawford J, Harvey SB. Help-seeking for depression among Australian doctors. *Intern Med J.* 2021 Dec;51(12):2069-2077. doi: 10.1111/imj.15035. PMID: 32833296.
32. Sun H, Warner DO, Macario A, Zhou Y, Culley DJ, Keegan MT. Repeated Cross-sectional Surveys of Burnout, Distress, and Depression among Anesthesiology Residents and First-year Graduates. *Anesthesiology.* 2019 Sep;131(3):668-677. doi: 10.1097/ALN.0000000000002777. PMID: 31166235.
33. de Melo Silva Júnior ML, Valença MM, Rocha-Filho PAS. Individual and residency program factors related to depression, anxiety, and burnout in physician residents - a Brazilian survey. *BMC Psychiatry.* 2022 Apr 19;22(1):272. doi: 10.1186/s12888-022-03916-0. PMID: 35436910; PMCID: PMC9016975.
34. Fond G, Boulangeat C, Messiaen M, Duba A, Boucekine M, Auquier P, Lançon C, Boyer L. Anxiety and depression in young physicians: Prevalence and associated factors. The MESSIAEN national study. *Encephale.* 2022 Feb;48(1):26-30. doi: 10.1016/j.encep.2021.02.005. Epub 2021 Apr 20. PMID: 33892920.
36. Gong Y, Han T, Chen W, Dib HH, Yang G, Zhuang R, Chen Y, Tong X, Yin X, Lu Z. Prevalence of anxiety and depressive symptoms and related risk factors among physicians

- in China: a cross-sectional study. *PLoS One*. 2014 Jul 22;9(7):e103242. doi: 10.1371/journal.pone.0103242. PMID: 25050618; PMCID: PMC4106870.
37. Xu H, Peng L, Wang Z, Liu X. Effects of psychological capital and social support availability on anxiety and depression among Chinese emergency physicians: Testing moderated mediation model. *Front Psychol*. 2022 Dec 7; 13:991239. doi: 10.3389/fpsyg.2022.991239. PMID: 36571060; PMCID: PMC9768176.
  38. Alhabeeb AA, Al-Duraihem RA, Alasmay S, Alkhamaali Z, Althumiri NA, BinDhim NF. National screening for anxiety and depression in Saudi Arabia 2022. *Front Public Health*. 2023 Jun 27;11:1213851. doi: 10.3389/fpubh.2023.1213851. PMID: 37441650; PMCID: PMC10333514.
  39. Alawad HS, Amin HS, Alfariis EA, Ahmed AM, Alosaimi FD, BaHammam AS. Anxiety and depression symptoms among medical residents in KSA during the COVID-19 pandemic. *J Taibah Univ Med Sci*. 2022 Apr;17(2):192-202. doi: 10.1016/j.jtumed.2022.01.005. Epub 2022 Feb 5. PMID: 35153646; PMCID: PMC8816958.
  40. Alshardi A, Farahat F. Prevalence and Predictors of Depression Among Medical Residents in Western Saudi Arabia. *J Clin Psychol Med Settings*. 2020 Dec;27(4):746-752. doi: 10.1007/s10880-019-09667-7. PMID: 31595403.
  41. Bondagji D, Fakeerh M, Alwafi H, Khan AA. The Effects of Long Working Hours on Mental Health Among Resident Physicians in Saudi Arabia. *Psychol Res Behav Manag*. 2022 Jun 20;15:1545-1557. doi: 10.2147/PRBM.S370642. PMID: 35757011; PMCID: PMC9231535.
  42. AlNahedh A, BinRusayis A, Al-Tannir M, AlFayyad I. Prevalence of Depression and Burnout among Family Medicine Residents in Riyadh City, Saudi Arabia. *Health Psychol Res*. 2023 Dec 28;11:90620. doi: 10.52965/001c.90620. PMID: 38162541; PMCID: PMC10756857.
  43. Hariri N, Bawahab N, Banoon E, Abo Alshamat R, Almadani N, AlQashqri H. Prevalence of Depression and Anxiety Disorders Among Surgical Doctors in Public Hospitals in Makkah City, Saudi Arabia: An Analytical Cross-Sectional Study. *Cureus*. 2023 Jan 1;15(1):e33225. doi: 10.7759/cureus.33225. PMID: 36733556; PMCID: PMC9888731.

44. Nour MO, Alharbi KK, Hafiz TA, Alshehri AM, Alyamani LS, Alharbi TH, Alzahrani RS, Almalki EF, Althagafi AA, Kattan ET, Tamim HM. Prevalence of Depression and Associated Factors among Adults in Saudi Arabia: Systematic Review and Meta-Analysis (2000-2022). *Depress Anxiety*. 2023 Sep 14; 2023:8854120. doi: 10.1155/2023/8854120. PMID: 40224601; PMCID: PMC11921833.
45. Alhalafi AH. Prevalence of Anxiety and Depression during the Coronavirus Disease 2019 Pandemic in Riyadh, Saudi Arabia: A Web-Based Cross-Sectional Survey. *J. Pharm. Res. Int.* [Internet]. 2020 Nov. 7 [cited 2025 Jun. 25];32(27):65-73. Available from: <https://journalipri.com/index.php/JPRI/article/view/1694>
46. Islam, M., George, P., Sankaran, S., Su Hui, J. L. & Kit, T. Impact of COVID-19 on the mental health of healthcare workers in different regions of the world. *BJPsych Open*. 2021, 7 (S1), S258–S259. <https://doi.org/10.1192/bjo.2021.690>.
47. Kisely, S. et al. Occurrence, prevention, and management of the psychological effects of emerging virus outbreaks on healthcare workers: rapid review and meta-analysis. *BMJ*. 5, m1642. 2020. <https://doi.org/10.1136/bmj.m1642>.
48. Al Hourri, H. N. et al. Stress, depression, anxiety, and quality of life among the healthcare workers during COVID-19 pandemic in Syria: a multi-center study. *Ann. Gen. Psychiatry*, 2023, 22 (1), 41. <https://doi.org/10.1186/s12991-023-00470-1>.
49. Prowse, R. et al. Coping with the COVID-19 pandemic: examining gender differences in stress and mental health among university students. *Front. Psychiatry*, 2021, 12, 650759. <https://doi.org/10.3389/fpsyt.2021.650759>.
50. Graves, B. S., Hall, M. E., Dias-Karch, C., Haischer, M. H. & Apter, C. Gender differences in perceived stress and coping among college students. Dalby AR, ed. *PLoS One*, 2021 16(8), e0255634. <https://doi.org/10.1371/journal.pone.0255634>.
51. Matud, M., Díaz, A., Bethencourt, J. & Ibáñez, I. Stress and psychological distress in emerging adulthood: A gender analysis. *JCM*, 2020, 9 (9), 2859. <https://doi.org/10.3390/jcm9092859>.
52. Malebari Azizah M., Alamoudi Saeed O., AL-Alawi Talal I., Alkhateeb Anas A., Albuqayli Adnan S., and Alothmany Hamzah N. Prevalence of depression and anxiety among university students in Jeddah, Saudi Arabia: exploring sociodemographic and associated

- factors. *Frontiers in Public Health*, 2024, 12 <https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2024.1441695>. DOI=10.3389/fpubh.2024.1441695
53. Zhai X, Tong HHY, Lam CK, Xing A, Sha Y, Luo G, Meng W, Li J, Zhou M, Huang Y, Wong LS, Wang C, Li K. Association and causal mediation between marital status and depression in seven countries. *Nat Hum Behav*. 2024 Dec;8(12):2392-2405. doi: 10.1038/s41562-024-02033-0. Epub 2024 Nov 4. PMID: 39496771.
  54. Oh J, Lee S, Sim J, Kim S, Cho A, Yun B, Yoon JH. Association between Self-Perceived Social Support in the Workplace and the Presence of Depressive/Anxiety Symptoms. *Int J Environ Res Public Health*. 2021 Sep 30;18(19):10330. doi: 10.3390/ijerph181910330. PMID: 34639639; PMCID: PMC8508368.
  55. Wang R, Liu Y, Liu S, Chen H, Yu X, Xu C, Chen Y, Xia D, Ge X, Wang Z, Chang R, Hu F, Shen T, Wang Y, Wang Z, Cao B, Zhang K, Zou H, Qin J, Li S, Cai Y. Relationships of work stress and interpersonal needs with industrial workers' mental health: a moderated mediation model. *BMC Public Health*. 2023 Jul 12;23(1):1341. doi: 10.1186/s12889-023-16002-1. PMID: 37438732; PMCID: PMC10339573.
  56. Magnavita N, Meraglia I, Riccò M. Anxiety and depression in healthcare workers are associated with work stress and poor work ability. *AIMS Public Health*. 2024 Dec 13;11(4):1223-1246. doi: 10.3934/public health, 2024063. PMID: 39802561; PMCID: PMC11717537.
  57. Amini, M.J., Shafiee, A., Mirhoseini, M.S. *et al.* Prevalence of depression and anxiety in orthopedic residents: a systematic review and meta-analysis. *BMC Med Educ* **24**, 1261 (2024). <https://doi.org/10.1186/s12909-024-06213-4>
  58. Martínez-García JA, Aguirre-Barbosa M, Mancilla-Hernández E, -Hernández-Morales MDR, Guerrero-Cabrera MB, Schiaffini-Salgado LG. Prevalencia de depresión, ansiedad y factores asociados en médicos residentes de centros hospitalarios durante la pandemia de COVID-19 [Prevalence of depression, anxiety, and associated factors in residents from hospital centers during COVID-19 pandemic]. *Rev Alerg Mex*. 2022 May 29;69(1):1-6. Spanish. doi: 10.29262/ram.v69i1.903. PMID: 36927745.
  59. Martín-Del-Campo F, Ramírez-Pineda JL, Ávila-Cornejo RM, Gutiérrez-Casillas SB, Sánchez-Soriano A, Cueto Manzano AM. Moderate physical activity is associated with

- lower depression, anxiety, and stress in healthcare workers during the COVID-19 pandemic. *J Psychosom Res.* 2023 Nov; 174:111488. doi: 10.1016/j.jpsychores.2023.111488. Epub 2023 Sep 8. PMID: 37690331.
60. Boucher VG, Haight BL, Hives BA, Zumbo BD, Merali-Dewji A, Hutton S, Liu Y, Nguyen S, Beauchamp MR, Black AT, Puterman E. Effects of 12 Weeks of At-Home, Application-Based Exercise on Health Care Workers' Depressive Symptoms, Burnout, and Absenteeism: A Randomized Clinical Trial. *JAMA Psychiatry.* 2023 Nov 1;80(11):1101-1109. doi: 10.1001/jama.psychiatry.2023.2706. PMID: 37556150; PMCID: PMC10413218.
  61. Lange KW, Nakamura Y, Lange KM. Sport and exercise as medicine in the prevention and treatment of depression. *Front Sports Act Living.* 2023 Mar 9; 5:1136314. doi: 10.3389/fspor.2023.1136314. PMID: 36969961; PMCID: PMC10033769.
  62. Lawrence D, Mitrou F, Zubrick SR. Smoking and mental illness: results from population surveys in Australia and the United States. *BMC Public Health.* 2009 Aug 7; 9:285. doi: 10.1186/1471-2458-9-285. PMID: 19664203; PMCID: PMC2734850.
  63. Hu X, Mu W, Zhou J, Zhou H, Yan X, Yue K, Liu T, Huang W, Ren L, Zou F, Zhang A, Sun X, Zeng H. Risk factors for anxiety, depression, stress, and their comorbidities among nurses: a prospective cohort from 2020 to 2022. *BMC Nurs.* 2024 Dec 18;23(1):916. doi: 10.1186/s12912-024-02577-6. PMID: 39696389; PMCID: PMC11658259.
  64. Clarke DM, Currie KC. Depression, anxiety and their relationship with chronic diseases: a review of the epidemiology, risk and treatment evidence. *Med J Aust.* (2009) 190: S54–60. doi: 10.5694/j.1326-5377.2009.tb02471.x
  65. Mirza AA, Milaat WA, Ramadan IK, Baig M, Elmorsy SA, Beyari GM, et al. Depression, anxiety and stress among medical and non-medical students in Saudi Arabia: an epidemiological comparative cross-sectional study. *Neurosciences.* (2021) 26:141–51. doi: 10.17712/nsj.2021.2.20200127



# The comorbidity of depression and anxiety among Saudi Board Residents in different health training programs in Al-Ahsa, Saudi Arabia, 2024

## ORIGINALITY REPORT

16%

SIMILARITY INDEX

13%

INTERNET SOURCES

14%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

1

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

3%

2

[www.frontiersin.org](http://www.frontiersin.org)

Internet Source

2%

3

[pmc.ncbi.nlm.nih.gov](http://pmc.ncbi.nlm.nih.gov)

Internet Source

1%

4

Ana Chico, Rosa Corcoy. "Intensive Insulin Therapy (Basal-Bolus)", American Journal of Therapeutics, 2022

Publication

<1%

5

[www.nature.com](http://www.nature.com)

Internet Source

<1%

6

[pubmed.ncbi.nlm.nih.gov](http://pubmed.ncbi.nlm.nih.gov)

Internet Source

<1%

7

[www.researchgate.net](http://www.researchgate.net)

Internet Source

<1%

8

Abu Bakkar Siddik, Morioum Sarkar Munmun, Nafiul Hasan, Khalid Syfullah et al. "Physio-psychosocial risk of depression among college-going adolescents: A cross-sectional study in Bangladesh", Journal of Affective Disorders Reports, 2024

Publication

<1%

9

[www.ophrdc.org](http://www.ophrdc.org)

Internet Source

<1%

10	<a href="http://www.jfmpc.com">www.jfmpc.com</a> Internet Source	<1 %
11	<a href="http://bmcpublichealth.biomedcentral.com">bmcpublichealth.biomedcentral.com</a> Internet Source	<1 %
12	Submitted to Tung Wah College Student Paper	<1 %
13	Khawlah A. Alzaben, Bayan S. Alshuhayb, Sulaiman M. Alsulaiman, Maram Alenazi, Mashaal A. Alkhayyal. "Coexistence of Congenital Lacrimal Gland Agenesis and Congenital Aniridia: Case Report", American Journal of Ophthalmology Case Reports, 2025 Publication	<1 %
14	<a href="http://assets.cureus.com">assets.cureus.com</a> Internet Source	<1 %
15	<a href="http://www.dovepress.com">www.dovepress.com</a> Internet Source	<1 %
16	<a href="http://www.science.gov">www.science.gov</a> Internet Source	<1 %
17	Gia Merlo, Christopher P. Fagundes. "Lifestyle Psychiatry - Through the Lens of Behavioral Medicine", CRC Press, 2023 Publication	<1 %
18	Submitted to Rutgers University, New Brunswick Student Paper	<1 %
19	"Care of Adults with Chronic Childhood Conditions", Springer Science and Business Media LLC, 2024 Publication	<1 %
20	Submitted to 2U University of Southern California- MSW Student Paper	<1 %

21	Internet Source	<1 %
22	docs.google.com Internet Source	<1 %
23	migrationletters.com Internet Source	<1 %
24	aquila.usm.edu Internet Source	<1 %
25	www.thelancet.com Internet Source	<1 %
26	iris.unipa.it Internet Source	<1 %
27	journals.lww.com Internet Source	<1 %
28	www.researchsquare.com Internet Source	<1 %
29	www.the-rheumatologist.org Internet Source	<1 %
30	oda.oslomet.no Internet Source	<1 %
31	www.medrxiv.org Internet Source	<1 %
32	Andrea Cook, Jennifer Champion. "Nutritional Psychology - Understanding the Relationship Between Food and Mental Health", CRC Press, 2025 Publication	<1 %
33	Submitted to Tufts University Student Paper	<1 %
34	journals.plos.org Internet Source	<1 %
35	Submitted to CSU, Fullerton Student Paper	

<1 %

36

Submitted to University of Alabama at  
Birmingham

Student Paper

<1 %

37

[www.scielo.br](http://www.scielo.br)

Internet Source

<1 %

38

[ijbmc.org](http://ijbmc.org)

Internet Source

<1 %

39

Submitted to West Coast University

Student Paper

<1 %

40

[www.cell.com](http://www.cell.com)

Internet Source

<1 %

41

[www.scienceopen.com](http://www.scienceopen.com)

Internet Source

<1 %

42

Daisy Follett, Caitlin Hitchcock, Tim Dalgleish,  
Jason Stretton. "Reduced Social Risk-Taking in  
Depression", PsyArXiv, 2021

Publication

<1 %

43

[gpsych.bmj.com](http://gpsych.bmj.com)

Internet Source

<1 %

44

[link.springer.com](http://link.springer.com)

Internet Source

<1 %

45

[pppkam.org.my](http://pppkam.org.my)

Internet Source

<1 %

46

Yan Zhang, Li Zhang, Jianyu Que, Miao Jia, Xi  
Nan, Juanniu Zhang, Haifei Gao, Lixia Chen.  
"Prevalence and Correlates of Mental Health  
Problems in Resident Physicians in Inner  
Mongolia Autonomous Region: A Cross-  
Sectional Study", Risk Management and  
Healthcare Policy, 2025

Publication

<1 %

47 Zhen Tian, Yike Huang, Tao Yue, Jiaqing Zhou, Lu Tao, Ling Han, Kexiang Yan, Qiong Huang, Zhenghua Zhang, Chunhong Shao. "A Chinese cross-sectional study on depression and anxiety symptoms in patients with psoriasis vulgaris", Psychology, Health & Medicine, 2018  
Publication

---

48 [iacc.hhs.gov](http://iacc.hhs.gov)  
Internet Source

---

49 [jamanetwork.com](http://jamanetwork.com)  
Internet Source

---

50 [opendentistryjournal.com](http://opendentistryjournal.com)  
Internet Source

---

51 [valleyinternational.net](http://valleyinternational.net)  
Internet Source

---

52 [worldwidescience.org](http://worldwidescience.org)  
Internet Source

---

53 [www.netce.com](http://www.netce.com)  
Internet Source

---

54 [www.verywellhealth.com](http://www.verywellhealth.com)  
Internet Source

---

55 Ahmad A. Mirza, Waleed A. Milaat, Iman K. Ramadan, Mukhtiar Baig et al. "Depression, anxiety and stress among medical and non-medical students in Saudi Arabia: An epidemiological comparative cross-sectional study", Neurosciences, 2021  
Publication

---

56 Andrew J. Green, Debbra D. Bazata, Kathleen M. Fox, Susan Grandy. "Quality of Life, Depression, and Healthcare Resource Utilization among Adults with Type 2 Diabetes Mellitus and Concomitant Hypertension and

## Obesity: A Prospective Survey", Cardiology Research and Practice, 2012

Publication

- 57 Deemah AlAteeq, Shouq Mohammed AlNuja'im, Abdullah H. AlGharbi. "Prevalence of depression and anxiety among working women in Saudi Arabia: psychosocial and perinatal correlates", BMC Women's Health, 2025

Publication

- 58 Lei Shi, Guoqiang Li, Jiatong Hao, Weidong Wang et al. "Psychological depletion in physicians and nurses exposed to workplace violence: A cross-sectional study using propensity score analysis", International Journal of Nursing Studies, 2020

Publication

- 59 Mossler, Ron . "Child and Adolescent Development, Third Edition", UAGC, 2024

Publication

- 60 Robbert Sanderman, Karen Morgan. "The Routledge International Handbook of Health Psychology - Global and Contemporary Issues", Routledge, 2025

Publication

- 61 Zheng Zhang, Wenhua Wang, Xiaoxiao Yuan, Xue Wang, Yi Luo, Linfei Dou, Lei Zhang, Mingyang Wu. "Adverse childhood experiences and subsequent physical and mental health among young adults: Results from six universities in China", Psychiatry Research, 2024

Publication

- 62 [bmcmmededuc.biomedcentral.com](https://bmcmmededuc.biomedcentral.com)

Internet Source

63	Internet Source	<1 %
64	<a href="https://brieflands.com">brieflands.com</a> Internet Source	<1 %
65	<a href="https://childmind.org">childmind.org</a> Internet Source	<1 %
66	<a href="https://jkms.org">jkms.org</a> Internet Source	<1 %
67	<a href="https://mentalhealthgateway.org">mentalhealthgateway.org</a> Internet Source	<1 %
68	<a href="https://ojs.zu.edu.pk">ojs.zu.edu.pk</a> Internet Source	<1 %
69	<a href="https://scholarworks.gsu.edu">scholarworks.gsu.edu</a> Internet Source	<1 %
70	<a href="https://www.medicinaantroposofica.it">www.medicinaantroposofica.it</a> Internet Source	<1 %
71	Abay Woday Tadesse, Getinet Ayano, Berihun Assefa Dachew, Kim Betts, Rosa Alati. "Maternal perinatal cannabis use disorder and the risk of anxiety disorders in offspring: Insights from a longitudinal data-linkage cohort study", Journal of Affective Disorders, 2025 Publication	<1 %
72	Abdulelah M. Aldhahir. "Prevalence and sociodemographic factors of depression, anxiety and stress in Saudi Arabia: a survey of respiratory therapists", Frontiers in Psychiatry, 2024 Publication	<1 %
73	Dimitrios Tsatiris. "Physician Burnout - How to Rise Above a Broken Healthcare System as a Practicing Clinician", Routledge, 2025 Publication	<1 %

74 Heather Katherine Scott-Marshall. "Safe limits on work hours for the nursing profession: a rapid evidence review", *Frontiers in Global Women's Health*, 2024 <1 %

---

Publication

75 Lin Zhang, Jingli Wen, Ling Yuan, Youde Yan, Zhenjiang Zhang, Kai Li, Zuoling Tang. "Anxiety and Depression in Healthcare Workers Two Years After COVID-19 Infection and Scale Validation", *Springer Science and Business Media LLC*, 2024 <1 %

---

Publication

76 M Subramaniam, JA Vaingankar, B Tan, E Abdin et al. "Examining Psychological Distress Among Youth in Singapore: Insights from the National Youth Mental Health Study", *Asian Journal of Psychiatry*, 2025 <1 %

---

Publication

77 Rui Tao, Wenzheng Li, Kaiyuan Min, Daming Mo et al. "Prevalence and associated factors of depression, anxiety and stress among clinical therapists in China in the context of early COVID-19 pandemic", *Frontiers in Psychiatry*, 2024 <1 %

---

Publication

78 Amani Busili, Azizah Makrami, Amnah Keni, Alaa Khardily, Dalyah Alzahrani, Idris Busaily, Fatimah Busayli. "COVID-19 exposure and depression-anxiety levels among Saudi adults in the Jazan region: a study with a predominantly female and undergraduate sample", *Frontiers in Psychiatry*, 2023 <1 %

---

Publication

79 Huiyoung Shin, Chaerim Park. "Perceived stress shapes symptom and social network dynamics: a network analysis of depression,



anxiety, and relationship-specific support and strain", BMC Psychiatry, 2025

Publication

80

Kathleen A. Kendall-Tackett. "Depression in New Mothers, Volume 2 - Screening, Assessment, and Treatment Alternatives", Routledge, 2024

Publication

<1 %

81

Klaus W. Lange, Yukiko Nakamura, Katharina M. Lange. "Sport and exercise as medicine in the prevention and treatment of depression", Frontiers in Sports and Active Living, 2023

Publication

<1 %

82

Lamers, F.. "Summed score of the Patient Health Questionnaire-9 was a reliable and valid method for depression screening in chronically ill elderly patients", Journal of Clinical Epidemiology, 200807

Publication

<1 %

83

Mohammad Javad Amini, Arman Shafiee, Mohammad Sajjad Mirhoseini, Mahdi Mohammaditabar et al. "Prevalence of depression and anxiety in orthopedic residents: a systematic review and meta-analysis", BMC Medical Education, 2024

Publication

<1 %

84

Seong-Hi Park, Sun Kyung Park. "An updated systematic review and meta-analysis of the predictive validity of the general anxiety disorder (GAD)-7 and GAD-2 in screening for anxiety disorders", Journal of Affective Disorders, 2025

Publication

<1 %

85

Yanhong Gong, Tieguang Han, Wei Chen, Hassan H. Dib, Guoan Yang, Runsen Zhuang, Yuqi Chen, Xinyue Tong, Xiaoxv Yin, Zuxun Lu.

<1 %

"Prevalence of Anxiety and Depressive  
Symptoms and Related Risk Factors among  
Physicians in China: A Cross-Sectional Study",  
PLoS ONE, 2014

Publication

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On