

Quality of life of cancer patients followed in the medical oncology department of a National Hospital Center of Senegal and associated factors

by Jana Publication & Research

Submission date: 01-Aug-2025 12:12PM (UTC+0200)

Submission ID: 2723686167

File name: IJAR-53083.docx (491.47K)

Word count: 8246

Character count: 41301

53 Quality of life of cancer patients followed in the medical oncology department of a National Hospital Center of Senegal and associated factors

Summary

Introduction: Faced with the highly aversive symptoms and incapacitating side effects of anticancer therapy, the assessment of the Quality of Life (QoL) of patients is essential to better inform clinical decisions. Using a mixed design, the QoL of cancer patients followed in the medical oncology department of the Dalal University Hospital Center Jam m of Senegal was assessed, as well as the factors associated with it. **Method:** The data collected using the self-questionnaire QLQ-C30 version 3.0 of EORTC, were analyzed according to the guidelines of the "QLQ-C30 Scoring Manual" by calculating the average of the raw scores of each of the 30 items completed, then the Standardized Scores (SN) for the multi-item dimensions ranging from 0 to 100. To determine the overall QoL level of each respondent based on their SN, an assessment scale consisting of four different levels corresponding to four class intervals with an amplitude equal to 25, was proposed: SN € [0-25[= Qdv heavily degraded; SN € [25-50[= Qdv degraded; SN € [50-75[= Qdv moderately degraded; SN € [75-100] = Qdv slightly degraded. Finally, the categorical variables were summarized in numbers and percentages and by calculating the deviations from independence. Bivariate statistical analyses (Chi2 test or Fisher's Exact test) were carried out to verify the existence of a statistical dependence between the socio-professional, demographic, clinical and therapeutic characteristics and the QoL of the patients (P-value < 0.05 indicates a significant dependence). **Results:** 60 cancer patients (mean age = 54.42 years) were included in the study, 47% of whom were at SBR stage II of the cancerous disease. QoL was slightly degraded in 52% of the respondents, moderately degraded in 22%, degraded in 16% and significantly degraded in 10%. Disease stage (p=0.003), presence of metastases (p=0.004), location of metastases (p=0.004), curative chemotherapy (p=0.003), palliative chemotherapy (p=0.004), curative radiotherapy (p=0.003) and marital status (p=0.02) significantly influence their QoL. **Discussions:** Taking these factors into account could improve the QoL of cancer patients.

Keywords: Quality of life; Cancer; Dalal National Hospital Center Jam m ; Senegal

Abstract

Introduction: Faced with the highly aversive symptoms and incapacitating side effects of anticancer therapy, the assessment of patients' Quality of Life (QoL) is essential to better inform clinical decisions. Using a mixed design, the QoL of cancer patients followed in the medical oncology department of the Dalal Jamm University Hospital in Senegal was assessed, as well as the factors associated with it. **Method:** Data collected using the EORTC QLQ-C30 version 3.0 self-administered questionnaire were analyzed according to the guidelines of the "QLQ-C30 Scoring Manual" by calculating the average of the raw scores for each of the 30 items completed, then the Standardized Scores (SN) for the multi-item dimensions ranging from 0 to 100. To determine the overall QoL level of each respondent based on their SN, an assessment scale consisting of four different levels corresponding to four class intervals with an amplitude equal to 25 was proposed: SN € [0-25[= QoL severely degraded; SN € [25-50[= QoL degraded; SN € [50-75[= QoL moderately degraded; SN € [75-100] = QoL slightly degraded. Finally, categorical

variables were summarized in numbers and percentages and by calculating deviations from independence. Bivariate statistical analyzes (Chi-square test or Fisher's Exact test) were performed to verify the existence of a statistical dependence between socio-professional, demographic, clinical and therapeutic characteristics and the QoL of patients (P-value < 0.05 indicates a significant dependence). **Results:** 60 cancer patients (mean age = 54.42 years) were included in the study, 47% of whom were at SBR stage II of the cancerous disease. QoL was slightly degraded in 52% of the respondents, moderately degraded in 22%, degraded in 16% and significantly degraded in 10%. Disease stage ($p=0.003$), presence of metastases ($p=0.004$), location of metastases ($p=0.004$), curative chemotherapy ($p=0.003$), palliative chemotherapy ($p=0.004$), curative radiotherapy ($p=0.003$) and marital status ($p=0.02$) significantly influence their QoL. **Discussions:** Taking these factors into account could improve the QoL of cancer patients.

Keywords: Quality of life; Cancer ; Dalal Jamm National Hospital Center; Senegal

2 Introduction

Cancer is and remains a major public health problem due to its seriousness and the complexity of its therapeutic management. Its occurrence affects all aspects of the cancer patient's life, as well as those around them (AL Septans , 2014). The World Health Organization [WHO], reported by P. Aubery and BA Gauzere (2020), reminded us that cancer is responsible for one in six deaths worldwide, but that up to 50% of all its forms can be prevented. Despite the progress made in recent years in diagnosis, therapeutic management and prevention, cancer remains a serious disease that wreaks havoc (P. Amsalhem *et al.* , 2007) and the most common symptoms resulting from the disease itself and its treatments include: distress (PA Ganz *et al.* , 2002), pain (P.J. Christo and D. Mazloomdoost , 2008), depression (M. Sharpe and J. Walker, 2009), fatigue (M. Andrea *et al.* , 2008) and insomnia (J. Savar *et al.* , 2005). Cancer therefore remains a challenge for humanity, in the face of which it is still powerless.

Therapeutic management of cancer is multidisciplinary, interdisciplinary and transdisciplinary. Depending on the type of cancer and its stage of development, and given that each cancer patient is unique, it includes different types of treatment which can be offered alone or in combination, namely: oncological surgery ; chemotherapy ; targeted therapies ; radiotherapy ; hormone therapy ; immunotherapy. These treatments generally affect the patient's Quality of Life (QoL) and can be responsible for side effects or late sequelae , while it is crucial that cancer patients maintain the best possible QoL through adjuvant care that goes beyond the discipline of oncology, including rehabilitation, supportive care and palliative care.

Indeed, according to the WHO (1993), QoL is an individual's perception of their place in life, in the context of the culture and value system in which they live, in relation to their goals, expectations, standards and concerns; QoL is a very broad concept that can be influenced in a complex way by the subject's physical health, psychological state and level of independence, social relationships and relationship to the essential elements of their environment. As for QoL related to health, it is defined as a state of well-being which is based on two components: 1- the ability to carry out daily activities which reflect the well-being physical, psychological And social and; 2- the satisfaction of patient with her level of functioning, control of his disease and related symptoms has her treatment (R. Feld , 1995; CC Gotay *et al.* , 1992). For Anon (2018), the concept of health-related quality of life refers to the subject's judgment of their own health. It includes physical, emotional, psychological and social dimensions as well as symptoms related to

the disease and treatments and is based on objective (living conditions, functional health) and subjective (satisfaction, happiness, well-being) items integrating the person as a whole and their environment.

Improving the QoL of cancer patients is one of the two criteria used by the Food and Drug Administration (FDA) for the marketing authorization of new cancer drugs (JR Johnson and R. Temple, 1985). Better still, among the judgment criteria used in cancerology, after overall survival, is the evaluation of QoL which reports a direct clinical benefit as perceived and quantified by the patient (A. Bryand *et al.* , 2015). The question of the evaluation of QoL therefore seems to be a major issue today and its study is currently part of most therapeutic trials validating new molecules or therapeutic protocols (KA Reiss *et al.*, 2015). Also, according to DF Cella (2007), the measurement of quality of life in cancer patients is done for three main reasons: 1- the evaluation of rehabilitation needs; 2- as a parameter in the evaluation of treatment results and; 3- as a predictive factor of response to future treatment . The evaluation of the QoL of cancer patients is therefore essential to better inform clinical decisions, hence the interest of this study whose objectives are to: 1- evaluate the impact of cancer and its treatments on the different dimensions of QoL and; 2- determine the factors associated with the quality of life of patients with cancer and followed in the medical oncology department of the Dalal National Hospital Center Jam m in the Republic of Senegal.

Materials and Methods

To achieve the objectives of the study, the philosophical orientation chosen (research design) and which seems to us to be the most suitable is that of a mixed study (qualitative and quantitative). The qualitative approach of the study is inspired by hermeneutic or interpretative phenomenology (HG Gadamer , 1976 and P. Ricœur, 1976) in which descriptive and interpretative perspectives coexist. As a qualitative research method in Nursing Sciences and Social Sciences, phenomenology aims at understanding and describing the human experience as lived by the participants (here, the participants in the study are cancer patients followed in the medical oncology department of the Dalal National Hospital Center Jam m in the Republic of Senegal) to a study (PL Munhall , 2012) and it seeks to discover the essence of phenomena, their nature and the meaning that human beings attribute to them (M. Van Manen , 1990) .

Indeed, the phenomenon under study, within the framework of this research, concerns the quality of life of cancer patients. In other words, what are the experiences of cancer patients of the impact of the cancerous disease and its treatments on the different dimensions [1- symptomatological; 2- physical; 3- psychological (mental) and emotional; 4- social and environmental; 5- economic; 6- sexual and body image and; 7- spiritual and psychological] of their quality of life?

Through documentary analysis (analysis of patient records) and on the basis of simple random probability sampling, patients of both sexes diagnosed with cancer and receiving therapeutic care for at least six months before the survey period (July 1st to August 31, 2021), followed in the medical oncology department of the Dalal National Hospital Center Jamm from Senegal, who gave their free and informed consent to participate in the survey and who are physically and mentally able to complete the data collection tool (self-questionnaire) themselves in order to observe the greatest neutrality, were included in the study.

The self-administered questionnaire " Quality of Life of Cancer Patients " from the European Organization for Research and Treatment of Cancer (European Organization for Research and Treatment of Cancer) [EORTC QLQ-

C30], version 3.0 (EORTC, 1995), was adapted and used as the data collection tool for the study. Developed in 1996 by the EORTC (1995), this tool was chosen for the study because it is currently the most widely used in French and European studies on QoL (NK Aaronson et al., 1993). Validated in 2009 (S. Gentile et al., 2008); D. Beauger et al., 2013), the EORTC QLQ-C30 version 3.0 (EORTC, 1995) is a standard data collection tool for all types of cancer and consists of a main questionnaire that allows exploring several dimensions of QoL (NW Scott et al., 2008). Also, it is a general questionnaire usable for all cancer locations, a useful, reliable, interesting measurement tool in current clinical practice and suitable for routine use in measuring QoL in cancer patients and having been the subject of a long development which has approved its validity and reproducibility (A. Georgakopoulos, 2013). The European Organization for Research and Treatment of Cancer (EORTC) has also demonstrated that the quality of life measurement system based on the QLQ-C30 questionnaire is a practical, reliable, valid and intercultural method (NK Aaronson et al., 1993). It is a tool widely recognized for its high reliability and validity (BA Ayana et al., 2016; MN Azmawati et al., 2014; B. McCarthy, 2011; Michels et al., 2014).

The EORTC QLQ-C30 questionnaire version 3.0 (EORTC, 1995) consists of thirty (30) items whose responses allow the establishment of different scales, each of them exploring a different dimension of quality of life. It is structured as follows :

- **Five (5)** functional scales to explore the dimensions: physical (items 1 to 5); executive (items 6 and 7); social (items 26 and 27); cognitive (items 20 and 25) and; emotional (items 21 to 24) of the quality of life of the person interviewed;
- **Nine (9)** symptomatic scales to explore: fatigue (items 10, 12 and 18); nausea symptoms (items 14 and 15); pain (items 9 and 19); dyspnea (item 8); insomnia (item 11); anorexia (item 13); diarrhea (item 17); constipation (item 16) and; financial difficulties (item 28) felt by the patient and;
- **One (1)** scale measuring the overall quality of life of each patient (items 29 and 30).

Each dimension explored includes between one and five different items. For the majority of questions, there are four possible response types: 1 - Not at all; 2 - A little; 3 - Quite a bit; 4 - A lot. The response methods, however, are different for the two items constituting the measurement of the patient's overall health status, the latter having to self-assess their overall health status and quality of life on a scale ranging from 1 (very poor) to 7 (excellent).

In addition to the 30 different specific variables (items) constituting the EORTC QLQ-C30 version 3.0 questionnaire (EORTC, 1995) for the assessment of the patient's QoL, we adapted the questionnaire to the study by supplementing sociodemographic and clinical variables of each patient included in the study from the documentary analysis of their care file in order to look for possible significant associations between them (sociodemographic and clinical variables) and the quality of life of the cancer patient (Factors associated with the quality of life of the cancer patient). These include the following variables: Age; Sex; Place of residence; Marital status; Professional status; Level of education; Cancer locations; Types of treatment received; Types of analgesics received; Stages of development (SBR [Scarff Bloom Richardson] grade of the cancerous disease and; Surgical and medical history of the patient (See the adapted questionnaire in appendix 1).

raw scores for each of the 30 items completed, then the Standardized Scores (SN) ranging from 0 to 100, were calculated according to the procedure described in the EORTC QLQ-C30 Scoring Manual (P. Fayers et al., 2001)

(Figure 1)

Figure 1: Method of calculating raw scores and standardized scores

Technical Summary

In practical terms, if items I_1, I_2, \dots, I_n are included in a scale, the procedure is as follows:

Raw score
Calculate the raw score
 $RawScore = RS = (I_1 + I_2 + \dots + I_n) / n$

Linear transformation
Apply the linear transformation to 0-100 to obtain the score S ,
Functional scales: $S = \left\{ 1 - \frac{(RS - 1)}{range} \right\} \times 100$
Symptom scales / items: $S = \left\{ \frac{(RS - 1)}{range} \right\} \times 100$
Global health status / QoL: $S = \left\{ \frac{(RS - 1)}{range} \right\} \times 100$
 $Range$ is the difference between the maximum possible value of RS and the minimum possible value. The QLQ-C30 has been designed so that all items in any scale take the same range of values. Therefore, the range of RS equals the range of the item values. Most items are scored 1 to 4, giving $range = 3$. The exceptions are the items contributing to the global health status / QoL, which are 7-point questions with $range = 6$, and the initial yes/no items on the earlier versions of the QLQ-C30 which have $range = 1$.

To determine ¹⁰the overall quality of life level of each patient surveyed from their Normalized Score obtained, an assessment scale consisting of four different levels corresponding to four class intervals with an amplitude equal to 25, was proposed (table 1).

Table 1: Ranking scale of each cancer patient surveyed according ⁷to their overall quality of life level based on the standardized score obtained ⁶²

Total Normalized Score obtained for the patient for the multi-item dimensions between	The overall quality of life of the cancer patient is:
[0-25[Heavily degraded
[25-50[Degraded
[50-75[Moderately degraded
[75-100]	Little degraded

⁶³It should be noted that ⁶⁵the standardized scores range from 0 to 100: a standardized score of zero corresponds to the worst quality of life and a standardized score of 100 reflects the best quality of life for the multi-item dimensions (NK Aaronson et al., 1993). In other words, a ⁸high global health score reflects good health and

quality of life and a high score for a functional scale reflects optimal function of the measured variables. On the other hand, a high score for a symptomatic scale reflects, conversely, a high level of symptoms.

Regarding the quantitative approach of the study, the categorical variables were summarized in numbers (absolute frequencies) and percentages (relative frequencies) and by calculations of arithmetic mean and deviations from independence (P. Cibois , 2003). Bivariate statistical analyses (Chi2 test or Fisher 's Exact test) were carried out to verify the existence of a statistical dependence between the socio-professional, demographic, clinical and therapeutic characteristics and the quality of life of the patients (P-value < 0.05 indicates a significant dependence).

Results

60 cancer patients, 51% of whom were male, were included in the study. Their average age was 54.42 years, with extremes of 13 and 84 years.

Epidemiological profile of patients and therapeutic management methods

Table 2: Distribution of respondents according to the location of the cancer, the types of treatment and the types of analgesic received (n=60)

No.	Cancerous locations	Frequency	Percentage
01	Cervical cancer	19	31%
02	Prostate cancer	16	27%

03	Cancer of the ENT sphere	08	13%
04	Cavum cancer	04	06%
05	Breast cancer	05	09%
06	Rectal cancer	04	07%
07	Other cancerous locations	04	07%
Total		60	100%
No.	Types of treatment received	Frequency	Percentage
01	Radio-chemotherapy	33	55%
02	Chemotherapy alone	11	18%
03	Exclusive radiotherapy for curative purposes	10	17%
04	Hormone therapy	03	05%
05	Adjuvant chemotherapy	03	05%
Total		60	100%
No.	Types of painkillers received	Frequency	Percentage
01	1st ^{level} analgesics	11	19%
02	2nd ^{level} analgesics	08	13%
03	3rd ^{level} analgesics	02	03%
04	No pain therapy received	39	65%
Total		60	100%

According to the data in Table 2, the most common cancers were cervical (31%) and prostate (27%). Radiochemotherapy was the most dominant type of treatment (55%), and the majority of patients had not received any pain relief therapy.

Furthermore, still from the documentary analysis of patient files, 11% of cancers were SBR grade I or low grade (least aggressive tumors for total histoprognostic scores of 3, 4 or 5); 47% were SBR grade II (for histoprognostic scores of 6 or 7) and; 42% were SBR grade III or high grade (most aggressive tumors for histoprognostic scores of 8 or 9) . Also, 42% of patients had a secondary cancer location (metastasis), 54% had a history of surgery, 43% had various medical histories and 3% had neither a history of surgery nor a medical history.

Overall quality of life of cancer patients surveyed (n = 60)

Table 3: Raw mean scores and standardized scores ¹¹calculated according to the procedure described in the EORTC QLQ-C30 Scoring Manual (P. Fayers et al, 2001)

Items/scales	Raw Average Scores (n = 60)	Normalized scores (n = 60)	ddl	Test p	alpha
	M(SD)	M(SD)			
Global Health Scale / QoL	59.2	61.7	19	t = -.424	.676
Functional scales					
Physical functioning	72.3	68.3		z = -0.803	.422
Daily activity	65.8	56.7		z = -1.31	.190
Emotional functioning	53.8	67.5		z = -2.033	.042
Cognitive functioning	57.5	72.5		z = -1.807	.071

Social functioning	65.0	65.8	z = -0.045	.964
Symptom scales/items				
Fatigue	42.8	46.7	t = .694	.496
Nausea and vomiting	13.3	12.5	z = -0.503	.615
Pains	30.0	28.3	z = .000	1.000
Dyspnea	16.7	28.3	z = -1.725	.084
Insomnia	50.0	50.0	z = -0.064	.949
Loss of appetite	13.3	21.7	z = -0.85	.396
Constipation	21.7	28.3	z = -0.791	.429
Diarrhea	11.7	16.7	z = -1.134	.257
Financial difficulties	18.3	20.0	z = -0.214	.831

Notes. M: mean; SD: standard deviation; df: degrees of freedom; Student's t for paired samples; Wilcoxon's z

Table 4: Distribution of cancer patients according to their level of overall quality of life based on the standardized score obtained for the multi-item dimensions (n=60)

Normalized score between	Assessment scale of the level of overall quality of life	Frequency	Percentage
[0-25[Quality of life significantly deteriorated	06	10%
[25-50[Degraded quality of life	10	16%
[50-75[Moderately poor quality of life	13	22%
[75-100]	Little deterioration in quality of life	31	52%
Total		60	100%

According to the data in Tables 3 and 4, more than half (52%) of the cancer patients surveyed have a slightly degraded overall quality of life. On the other hand, 10% of them have a significantly degraded overall quality of life and 16% have a degraded overall quality of life.

Correlations between socio-professional, demographic, clinical and therapeutic variables and the quality of life of cancer patients surveyed

Table 5: Bivariate analysis between socio-professional, demographic, clinical and therapeutic characteristics and the overall quality of life score

No.	Sociodemographic, clinical and therapeutic variables	p-value 5%	No.	Sociodemographic, clinical and therapeutic variables	p-value 5%
01	Age	0.837	09	Presence of metastases	0.004
02	Sex	0.326	10	Location of metastases	0.004
03	Marital status	0.217	11	Surgery	-
04	Place of residence	0.525	12	Curative chemotherapy	0.003
05	Occupation	0.898	13	Curative radiotherapy	0.003
06	Educational level	0.426	14	Hormone therapy	0.152
07	Location of the cancerous disease	0.769	15	Palliative chemotherapy	0.004
08	Stage of cancer disease	0.003	16	Painkillers	0.153

From the data presented in Table 5, it appears that: 1 - the stage of the cancerous disease ($p = 0.003$); 2 - the presence of metastases ($p = 0.004$); 3 - the location of metastases ($p = 0.004$); 4 - curative chemotherapy ($p = 0.003$); 5 - palliative chemotherapy ($p = 0.004$) and; 6 - curative radiotherapy ($p = 0.003$), are factors that significantly influence the QoL of cancer patients because their P-Values are all < 0.05

Table 6: Bivariate analysis between socio-professional and demographic characteristics and the different levels of quality of life of cancer patients (n=60)

Features socio- professional and demographic	Effective Overall (N = 60)	Quality of life				P-Value
		Strongly degraded (n = 6)	Degraded (n = 10)	Moderately degraded (n = 13)	Little degraded (n = 31)	
Sex						0.12
Women	29 (49)	5 (8)	7 (12)	6 (10)	12 (20)	
Man	31 (51)	1 (2)	3 (5)	7 (12)	19 (32)	
Place of residence						0.3
Urban	29 (49)	4 (7)	7 (12)	5 (8)	13 (22)	
Rural	31 (51)	2 (3)	3 (5)	8 (13)	18 (30)	
Marital status						0.02
Singles	15 (25)	2 (3)	6 (10)	4 (7)	3 (5)	
Divorced	11 (8)	1 (2)	2 (3)	2 (3)	6 (10)	
Married	27 (45)	1 (2)	1 (2)	6 (10)	19 (32)	
Widowers	7 (12)	2 (3)	1 (2)	1 (2)	3 (5)	
Professional status						0.59
Active	22 (37)	1 (2)	2 (3)	6 (10)	13 (22)	
Unemployed	16 (27)	3 (5)	4 (7)	4 (7)	5 (8)	
Retirees	12 (20)	1 (2)	3 (5)	1 (2)	7 (12)	
Housewives	10 (17)	1 (2)	1 (2)	2 (3)	6 (10)	
Level instruction						0.36
Illiterate	9 (15)	2 (3)	4 (7)	1 (2)	2 (3)	
Preschool	11 (18)	1 (2)	3 (5)	3 (5)	4 (7)	
Primary	9 (15)	1 (2)	1 (2)	1 (2)	6 (10)	
Secondary	13 (22)	1 (2)	1 (2)	3 (5)	8 (13)	
University	18 (30)	1 (2)	1 (2)	5 (8)	11 (18)	

According to the data in Table 6, the predominant marital status was married (45%), followed by single (25%). 51% of the respondents stated that they resided in rural areas. Only 3.7 % of them were still professionally active, followed by the unemployed (27%). University and secondary school graduates represented more than 50% of the cancer patients surveyed.

Among the socio-professional and demographic variables, only **marital status was the most significant factor related to the quality of life of cancer patients** ($P\text{-value} < 0.05$) while gender, residential area, professional status and educational level did not show significant differences. Furthermore, the majority of married people (19/27); professionally active people (13/22); people living in rural areas (18/31); men (19/31); university-educated people (11/18) and secondary-educated people (8/13), seem to have a little degraded quality of life.

The analysis of contingency tables 7, 8 and 9 below shows that almost **the deviations from independence calculated** for the different modalities of the socio-professional and demographic variables: 1 - Sex; 2- Place of residence; 3- Marital status; 4- Professional status and; 5- Level of education, **are either greater than or less than zero**, which confirms an association relationship between these variables and the different levels of the overall

quality of life of the cancer patients surveyed with the exception of the "Housewives" modality of the "Professional status" variable whose deviation from independence is zero (Table 7) in patients with a highly degraded Quality of life (standardized score between [0-25[). In other words, it seems that there is no association relationship between being a cancer woman at home and the strong degradation of the overall quality of life.

Table 7: Correlation between the level of quality of life of the cancer patients surveyed and the variables sex and place of residence

Variables socio demographics	Quality of life level										Total	
	Significantly degraded quality of life (standardized score between [0-25])			Degraded quality of life (normalized score between [25-50])			Moderately degraded quality of life (normalized score between [50-75])			Little deterioration in quality of life (normalized score between [75-100])		
	Effective	%	Effective	%	Effective	%	Effective	%	Effective	%		
Sex												
Women	05	8%	07	11%	06	10%	12	20%	29	49%		
Man	01	2%	03	5%	07	12%	19	32%	31	51%		
%	06	10%	10	16%	13	22%	31	52%	60	10%		
Theoretical frequencies (f _t) = Observed margin products ÷ Observed total												
Women	(06÷60) × 29 = 2.9			(10÷60) × 29 = 04.83			(13÷60) × 29 = 06.28			(31÷60) × 29 = 14.98		
Man	(06÷60) × 31 = 3.1			(10÷60) × 31 = 05.17			(13÷60) × 31 = 06.72			(31÷60) × 31 = 16.02		
Total	06			10			13			31		
Deviation from independence = observed frequency (f _o) - theoretical frequency (f _t)												
Women	f _o	f _t	f _o - f _t	f _o	f _t	f _o - f _t	f _o	f _t	f _o - f _t	f _o	f _t	
Man	05	2.9	+2.1	07	04.83	+2.17	06	06.28	-0.28	12	14.98	
%	01	3.1	-2.1	03	05.17	-2.17	07	06.72	+0.28	19	16.02	
Place of residence												
Urban	04	7%	07	12%	05	8%	13	22%	29	49%		
Rural	02	3%	03	4%	08	14%	18	30%	31	51%		
%	06	10%	10	16%	13	22%	31	52%	60	10%		
Theoretical frequencies (f _t) = Observed margin products ÷ Observed total												
Urban	(06÷60) × 29 = 2.9			(10÷60) × 29 = 04.83			(13÷60) × 29 = 06.28			(31÷60) × 29 = 14.98		
Rural	(06÷60) × 31 = 3.1			(10÷60) × 31 = 05.17			(13÷60) × 31 = 06.72			(31÷60) × 31 = 16.02		
Total	06			10			13			31		
Deviation from independence = observed frequency (f _o) - theoretical frequency (f _t)												
Urban	f _o	f _t	f _o - f _t	f _o	f _t	f _o - f _t	f _o	f _t	f _o - f _t	f _o	f _t	
Rural	04	2.9	+1.1	07	04.83	+2.17	05	06.28	-1.28	13	14.98	
%	02	3.1	-1.1	03	05.17	-2.17	08	06.72	+1.28	18	16.02	
										+1.98		

Table 8: Correlation between the level of quality of life of the cancer patients surveyed and the variables "Marital status" and "Professional status"

Variables socio demographics	Quality of life level										Total
	Significantly degraded quality of life (standardized score between [0-25])		Degraded quality of life (normalized score between [25- 50])		Moderately degraded quality of life (normalized score between [50-75])		Quality of life slightly degraded (normalized score between [75-100])				
	Effective	%	Effective	%	Effective	%	Effective	%			
	Marital status										
Singles	02	4%	06	9%	04	7%	03	5%	15	25%	
Divorced	01	2%	02	3%	02	3%	06	10%	11	18%	
Married	01	1%	01	2%	06	10%	19	31%	27	44%	
Widowers	02	3%	01	2%	01	2%	03	6%	07	13%	
%	06	10%	10	16%	13	22%	31	52%	60	100%	
Theoretical frequencies (f _i) = Observed margin products ÷ Observed total											
Singles	(06+60) x 15 = 1,5		(10+60) x 15 = 02,50		(13+60) x 15 = 03,25		(31+60) x 15 = 07,75		15		
Divorced	(06+60) x 11 = 1,1		(10+60) x 11 = 01,83		(13+60) x 11 = 02,38		(31+60) x 11 = 05,68		11		
Married	(06+60) x 27 = 2,7		(10+60) x 27 = 04,50		(13+60) x 27 = 05,85		(31+60) x 27 = 13,95		27		
Widowers	(06+60) x 07 = 0,7		(10+60) x 07 = 01,17		(13+60) x 07 = 01,52		(31+60) x 07 = 03,62		07		
Total	06	10	13	31	31						60
Deviation from independence = observed frequency (f _o) - theoretical frequency (f _i)											
Singles	f _o	f _i	f _o - f _i	f _o	f _i	f _o - f _i	f _o	f _i	f _o - f _i	f _i	
Divorced	02	1,5	+0,5	06	02,50	+3,5	04	03,25	03	07,75	-4,75
Married	01	1,1	-0,1	02	01,83	+0,17	02	02,38	06	05,68	+0,32
Widowers	01	2,7	-1,7	01	04,50	-3,50	06	05,85	19	13,95	+05,05
	02	0,7	+1,3	01	01,17	-0,17	01	01,52	03	03,62	-0,62
Professional status											
Active	01	1%	02	3%	06	10%	13	21%	22	37%	
Unemployed	03	5%	04	7%	04	7%	05	9%	16	27%	
Retirees	01	2%	03	4%	01	2%	07	12%	12	20%	
Housewives	01	2%	01	2%	02	3%	06	10%	10	16%	
Total	06	10%	10	16%	13	22%	31	52%	60	100%	
Theoretical frequencies (f _i) = Observed margin products ÷ Observed total											
Active	(06+60) x 22 = 2,2		(10+60) x 22 = 03,86		(13+60) x 22 = 04,76		(31+60) x 22 = 11,36		22		
Unemployed	(06+60) x 16 = 1,6		(10+60) x 16 = 02,67		(13+60) x 16 = 03,47		(31+60) x 16 = 08,27		16		
Retirees	(06+60) x 12 = 1,2		(10+60) x 12 = 02		(13+60) x 12 = 02,60		(31+60) x 12 = 06,20		12		
Housewives	(06+60) x 10 = 0,1		(10+60) x 10 = 01,67		(13+60) x 10 = 02,17		(31+60) x 10 = 05,17		10		
Total	06	10	13	31	31						60
Deviation from independence = observed frequency (f _o) - theoretical frequency (f _i)											
Active	f _o	f _i	f _o - f _i	f _o	f _i	f _o - f _i	f _o	f _i	f _o - f _i	f _i	
Unemployed	01	2,2	-1,2	02	03,86	-1,66	06	04,76	13	11,36	+1,64
Retirees	03	1,6	+1,4	04	02,67	+1,33	04	03,47	05	08,27	-3,27
Housewives	01	1,2	-0,2	03	02	+0,1	01	02,60	07	06,20	+0,80
	01	0,1	00	01	01,67	-0,67	02	02,17	06	05,17	+0,83

Table 9: Correlation between the level of quality of life of the cancer patients surveyed and the variable "Level of education"

Variables socio demographics	Quality of life level										Total		
	Significantly degraded quality of life (standardized score between [0-25])		Degraded quality of life (normalized score between [25-50])		Moderately degraded quality of life (normalized score between [50-75])		Little deterioration in quality of life (normalized score between [75-100])						
	Effective	%	Effective	%	Effective	%	Effective	%					
	ve	ve	ve	ve	ve	ve	ve	ve					
Educational level													
Illiterate	02	5%	04	7%	01	2%	02	3%	09	17%			
Preschool	01	1%	03	4%	03	4%	04	7%	11	16%			
Primary	01	1%	01	2%	01	3%	06	10%	09	16%			
Secondary	01	2%	01	1%	03	4%	08	13%	13	20%			
University	01	1%	01	2%	05	9%	11	19%	18	31%			
%	06	10%	10	16%	13	22%	31	52%	60	10%			
Theoretical frequencies (f _t) = Observed margin products ÷ Observed total													
Illiterate	(06+60) x 09 = 0.9		(10+60) x 09 = 01.50		(13+60) x 09 = 01.95		(31+60) x 09 = 04.65		09				
Preschool	(06+60) x 11 = 1.1		(10+60) x 11 = 01.83		(13+60) x 11 = 02.38		(31+60) x 11 = 05.68		11				
Primary	(06+60) x 09 = 0.9		(10+60) x 09 = 01.50		(13+60) x 09 = 01.95		(31+60) x 09 = 04.65		09				
Secondary	(06+60) x 13 = 1.3		(10+60) x 13 = 02.17		(13+60) x 13 = 02.82		(31+60) x 13 = 06.72		13				
University	(06+60) x 18 = 1.8		(10+60) x 18 = 03		(13+60) x 18 = 03.90		(31+60) x 18 = 09.30		18				
Total	06		10		13		31		60				
Deviation from independence = observed frequency (f _o) - theoretical frequency (f _t)													
Illiterate	f _o	f _t	f _o -f _t	f _o	f _t	f _o -f _t	f _o	f _t	f _o -f _t	09			
Preschool	02	0.9	+1.1	04	01.50	+02.50	01	01.95	-0.95	02	04.65	-2.65	11
Primary	01	1.1	-0.1	03	01.83	+1.17	03	02.38	+0.62	04	05.68	-1.68	09
Secondary	01	0.9	+0.1	01	01.50	-0.50	01	01.95	-0.95	06	04.65	+1.36	13
University	01	1.3	-0.3	01	02.17	-01.17	03	02.82	+0.18	08	06.72	+1.28	18
University	01	1.8	-0.8	01	03	-2	05	03.90	+1.1	11	9.30	+1.17	60

Discussion and conclusion

The quality of life of a person in general and that relating specifically to his health, is today a very important parameter to take into account in clinical and therapeutic decisions, especially in the process of caring for patients with a chronic illness such as cancer in palliative care. If the definition of quality of life of the WHO (1993) as used in this study as a theoretical framework of reference seems to have an international consensus, it is not easy to identify the concept of quality of life and to develop a clear and precise definition that would receive the approval of all scientific circles (C. Mercier & J. Filion , 1987).

The results of the study showed that quality of life (QoL) is affected to varying degrees (Table 4) by cancer, which corroborates the results of a study conducted by the Directorate of Research, Studies, Evaluation and Statistics (DREES) of the French Ministry of Health (DREES, 2005) which reported that the QoL of people with cancer is logically worse than that of the general population. According to DREES (2005), while physical quality of life is more or less impaired depending on the location of cancer, mental quality of life is not influenced by any clinical variable, which seems to be a new result that no other study had highlighted.

The study showed that socio-professional, demographic, therapeutic and clinical variables could, in general, influence either negatively or positively the quality of life of a cancer patient. C. Mercier & J. Filion (1987) are of the same opinion and affirm that it is undeniable that socio-demographic and economic variables play an important role in the evaluation of the quality of life. In our study, among the respondents who have a significantly degraded quality of life (Table 7), women were dominant (5/6), which corroborates the results of a Tunisian survey (R. Khalladi et al., 2020) which reported to us that women with cancer [...] had a poor quality of life.

Almost half of our respondents (27/60 or 45%) were married. This life as a couple would seem to be a factor of psychological support for cancer patients according to the DREES study (2005) which reported that the occurrence of cancer in one of the spouses preserves, or even strengthens the couple. The majority (22/60 or 37%) of our respondents were professionally active and more than half (31/60 or 51.67%) had a secondary or university level of education. All this further corroborates the results of the DREES study (2005) which informs us that having [...] a high level of education, high income [...], promotes a fighting spirit and reduces the feeling of distress in cancer patients. Being married, therefore living as a couple (45% of respondents), still being professionally active (37% of respondents) or being a housewife (17% of respondents), are all social resources that could protect cancer patients from the deterioration of their quality of life. F. Cousson-Gélie (2014) also believe that social resources seem to be a powerful predictor of the quality of life of patients with the disease. Better still, the social support perceived by cancer patients and its protective role with regard to emotional distress and quality of life has been well established by several studies (SM Alferi et al, 2001; C. Dunkel-Schetter , 1984; CN Hoskins et al, 1996; A. Moyer and P. Salovey , 1999; SJ Neuling and HR Winefield , 1988; PA Parker, 2003).

More than half (55%) of our surveyed patients received radiochemotherapy or chemotherapy alone (18%) as treatment types. According to a publication of the Canadian Cancer Society (2024) , chemotherapy agents , while destroying cancer cells, can also damage normal cells causing several side effects such as : fatigue; nausea and vomiting; loss of appetite; hair loss; diarrhea; constipation; sore mouth and throat; pain in muscles, joints and

digestive tract; mucus; changes in ¹² taste and smell; skin problems; eye and vision problems; fertility problems; problems at the injection site; hearing problems; organ damage and nerve damage; memory and attention problems and other cognitive disorders; sexual disorders; negative reactions to medications and; secondary cancers. The treating clinician must therefore constantly monitor the occurrence of these side effects in patients, which are also ³ factors that significantly influence their quality of life. This is all the more true since in our study, curative chemotherapy ($p=0.003$), palliative chemotherapy ($p=0.004$) and curative radiotherapy ($p=0.003$) were identified as ²³ factors that significantly influence the QoL of the cancer patients surveyed.

In our study, the stage of the cancer disease ($p=0.003$) was also identified as a factor that significantly influences ¹⁷ the quality of life of the cancer patients surveyed, which corroborates the results of the study by O. Popa-Velea et al (2017) who inform us that the overall deterioration of ⁴⁵ the quality of life was positively correlated with the stage and evolution of the cancer disease.

Limitations of the study

³³ Like other quality of life assessment questionnaires for cancer patients, the EORTC QLQ-C30 version 3.0 questionnaire (EORTC, 1995) used in this study has methodological limitations, but this does not reduce the quality of the results obtained. Although the value of ²⁰ quality of life assessment questionnaires in current practice and clinical trials is no longer in doubt, there are unknowns and methodological weaknesses that hinder their use and/or interpretation, in particular the subjectivity of the data collected given the self-assessment nature of these questionnaires for assessing the different dimensions of quality of life, which casts doubt on their reliability.

Indeed, the complexity of quality of life definitions and the heterogeneity in the practical ²⁵ use of quality of life questionnaires is a major limitation to the exploitation of data in clinical trials (EA Hahn et al, 2007; IB Wilson and PD Cleary (1995). Questionnaires for ³⁵ assessing quality of life in cancer patients are often standardized and therefore cannot ensure a personalized assessment of patients. They thus hinder the performance of meta-analyses that would allow greater statistical power (EK Donovan, 2019). Missing data often result from the non-existence of a specific questionnaire validated for a location.

A new approach therefore consists of the use of direct patient perceptions ⁵ (Patient - reported outcomes [PROs]) based on the quality of life questionnaires previously described ³⁵ (Patient reported outcome measures [PROMs]) in order to better assess the consequences of treatments on the functioning and well-being of cancer patients; the patient will thus be better placed at the heart of their care and therapeutic decision-making. Indeed, PROs are carried out with the aim of taking into account patients' perceptions of treatments and they ²² play a role in the management of symptoms during the administration of different therapies, but also in improving communication with different health professionals and even in patient survival (A. Caissie et al, 2018). As for PROMs, they are questionnaires completed by patients and which allow obtaining a score that can guide the clinician towards the importance of certain symptoms or conditions related to the patient's cancer (CF Snyder et al, 2013).

List of references

1. Aaronson Neil, Ahmedzai Sam, Bergman Bengt, Bullinger Monika, Cull Ann, Duez Nicole et al, (1993), *The European Organization for Research and Treatment of Cancer QLQ-C30: A quality-of-life instrument for use in international clinical trials in oncology*. Journal of the National Cancer Institute 1993; 85:365-376.
2. Azmawati Mohammed Nawi , Najibah Endut , Ahmad Zailani Hatta Mohd Dali and Norfazilah Ahmad (2014), *Quality of life by stage of cervical cancer among Malaysian patients* . Asian Pacific Journal of Cancer Prevention, 15(13), 5283-5286.
3. Aubry Pierre and Gaüzère Bernard-Alex (2023), *Cancers in developing countries* . René Labusquière Center , Institute of Tropical Medicine, University of Bordeaux, 33076 Bordeaux (France)
4. Andrea Barsevick , Tracey Newhall and Susan Brown (2008), *Management of Cancer-Related Fatigue*. Clin J Oncol Nurs . 2008 October ; 12(5 Supp): 21–25. doi:10.1188/08.CJON.S2.21-25.
5. Alferi Susan, Carver Charles, Antoni Michael, Weiss, S. and Durèn , RE (2001), *An exploratory study of social support, distress, and life disruption among low-income Hispanic women under treatment for early stage breast cancer* , Health Psychology, 20 (1), 41-46.
6. Amsalhem , P., Etessami , R., and Morère , JF (2007). Epidemiology, risk factors, screening. In *Breast cancer* (pp. 11-22). Springer, Paris.
7. Anon. Health-Related Quality of Life (HRQOL). Atlanta, GA: Centers for Disease Control and Prevention (CDC); 2018. URL: <https://www.cdc.gov/hrqol/index.htm> [accessed May 9, 2021]
8. Bryand , A., Hamidou, Z., Paget-Bailly, S., Bonnetain, F., Mathelin, C., Baldauf , JJ and Akladios , C. (2015). *Study of quality of life in ovarian neoplasia: tools and challenges* . Gynecology Obstetrics & Fertility, 43(2), 151-157
9. Birhanu Abera Ayana, Shiferaw Negash, Lukman Yusuf, Wondemagegnehu Tigeneh, Wondemagegnehu Haile (2016) *Reliability and Validity of Amharic Version of EORTC QLQ-C 30 Questionnaire among Gynecological Cancer Patients in Ethiopia*. PLoS One. 15;11(6); DOI: 10.1371/journal.pone.0157359
10. Beauger Davy, Gentile Stéphanie, Jouve Elisabeth, Dussol Bertrand, Jacquelinet Christian and Briançon Serge (2013), *Analysis , evaluation and adaptation of the ReTransQoL : a specific quality of life questionnaire for renal transplant recipients* . Health Qual Life Outcomes 11:148.
11. Christo Paul and Mazloomdoost Danesh (2008), *Interventional pain treatments for cancer pain* . Annals of the New York Academy of Sciences, 1138(1), 299-328.
12. Cousson-Gélie Florence (2014). *Psychosocial determinants of quality of life in patients with serious and/or chronic illness*. In F. Bacro (ed.), *Quality of life* (1 -). Rennes University Press. <https://doi.org/10.4000/books.pur.61277> , pp. 103-115
13. Cashier Amanda, Brown Erika, Olson Rob, Barbera Lisa, Davis Carol-Anne, Brundage Michael et al (2018), *Improving patient outcomes and radiotherapy systems: A pan-Canadian approach to patient-reported outcome use*. Med Phys;45(10):e841 -4.

14. Donovan Elysia, Sienna Julianna, Mitera Gunita , Kumar-Tyagi Nidhi, Parpia Sameer, Swaminathan Anand (2019), *Single versus multifraction radiotherapy for spinal cord compression: A systematic review and meta-analysis*. *Radiother Oncol* 2019;134:55 -66.
15. David Cella (2007), *The concept of quality of life: palliative care and quality of life* , *Nursing Research* 2007/1 No. 88 | pages 25 to 31 ISSN 0297-2964 DOI 10.3917/rsi.088.0025
16. Directorate of Research, Studies, Evaluation and Statistics [DREES] of the French Ministry of Health (2005). *Life two years after cancer diagnosis*.
17. Dunkel - Schetter Christine (1984), *Social support and cancer: Findings based on patient interviews and their implications* . *Journal of Social Issues*, 40 (4), 77-98.10.1136/jech.2006.056028 _
18. European Organization for Research and Treatment of Cancer (1995). *Quality of Life of Cancer Patients [EORTC QLQ-C30], version 3.0* . EORTC Study Group on Quality of Life.
19. Feld Ronalde (1995). *Endpoints in cancer clinical trials: is there has need for measuring quality of life?* *Supportive Care in Cancer* ; Volume 3 (1), pages 23–27, (1995)
20. Fayers Peter, Aaronson NK, Bjordal K, Groenvold M, Curran D, Bottomley A, on behalf of the EORTC Quality of Life Group. *The EORTC QLQ-C30 Scoring Manual (3rd Edition)*. Published by: European Organization for Research and Treatment of Cancer, Brussels 2001
21. Ganz Patricia, Desmond Katherine , Leedham Beth, Rowland Julia, Meyerowitz Beth & Belin Thomas (2002). *Quality of life in long-term, disease-free survivors of breast cancer: a follow-up study* . *Journal of the National Cancer institute* , 94(1), 39-49
22. Georgakopoulos Alexandros , Kontodimopoulos Nick, Chatzioannou Sofia & Niakas Dimitris (2013). *EORTC QLQ-C30 and FACT- Lym for the assessment of health-related quality of life of newly diagnosed lymphoma patients undergoing chemotherapy* . *European Journal of Oncology Nursing*, 17(6), 849-855
23. Gentile Stéphanie, Jouve Elisabeth, Dussol Bertrand, Moal Valerie , Berland Yvon, Sambuc Roland (2008). *Development and validation of a French patient-based health-related quality of life instrument in kidney transplant : the ReTransQoL* . *Health Quality Life Outcomes* 6:78
24. Gotay , CC, Korn, EL, McCabe, MS, Moore, TD, & Cheson , BD (1992). *Quality of life assessment in cancer treatment protocols: Research issues in protocol development*. *JNCI Journal of the National Cancer Institute*, 84(8), 575-579
25. Gadamer Hans-Georg (1976) , *Philosophical hermeneutics* . Berkeley, CA. University of California Press
26. Hahn Elizabeth, Cella David, Chassany Olivier, Fairclough Diane, Wong Gilbert, Hays Ron, et al. (2007), *Precision of health-related quality-of-life data compared with other clinical measures* . *Mayo Clin Proc*; 82(10):1244 -54
27. Hoskins, CN, Baker, S., Sherman, D. and Bohlander , J., (1996), "Social support and patterns of adjustment to breast cancer", *Scholarly Inquiry for Nursing Practice*, 10 (2), 99-123.
28. Johnson, J.R., & Temple, R. (1985). Food and Drug Administration requirements for approval of new anticancer drugs. *Cancer treatment reports* , 69 (10), 1155-1159.

29. **Khalladi Rania, Imen Gargouri , Chekib Zedini and Helmi Ben Saad** (2020), What factors influence the quality of life of Tunisian patients with lung cancer? J Fac Med Or 4(2):597-608. DOI:10.5281/zenodo.4282512
30. **Munhall Patricia** (2012), *Nursing research: A qualitative perspective (5th edition)* . Sudbury, MA: Jones & Bartlett.
31. **Michels Fernanda Alessandra Silva, Latorre Maria do Rosario Dias de Oliveira and Maciel Maria do Socorro** (2013). *Validity, reliability and understanding of the EORTC-C30 and EORTC-BR23, quality of life questionnaires specific for breast cancer* . Rev Bras Epidemiol .(2):352-63.
32. **McCarthy Bridie** (2011). *Family members of patients with cancer: what they know, how they know and what they want to know* . Europe an Journal Of Oncology Nursing . 2011 Dec;15(5):428-41.
33. **Mercier Céline and Filion Jocelyne** (1987). *Quality of life: theoretical and empirical perspectives* . Mental Health in Quebec, 12(1), 135–143. <https://doi.org/10.7202/030380ar>
34. **Moyer Anne and Salovey Peter** (1999), *Predictors of social support and psychological distress in women with breast cancer* , Journal of Health Psychology, 4 (2), 177-191.10.1016/0277-9536(88)90273-0:
35. **Neuling Sandra and Vignoble Hélène** (1988). *Social support and recovery after surgery for breast cancer: Frequency and correlates of supportive behaviors by family, friends and surgeon*. Social Science and Medicine , 27 (4), 385-392.
10.1002/pon.635
36. **Neil Scott, Peter Fayers , Neil Aaronson, Andrew Bottomley, Alexander de Graeff , Mogens Groenvold , Michael Koller , Morten Petersen and Mirjam Sprangers** (2008), *The relationship between overall quality of life and its subdimensions was influenced by culture: analysis of an international database*. Journal of Clinical Epidemiology Volume 61, Issue 8 , August 2008, Pages 788-795.
<https://doi.org/10.1016/j.jclinepi.2007.08.015>
37. **World Health Organization [WHO]** (1993), *Study protocol for the World Health Organization project to develop a Quality of Life assessment instrument (WHOQOL)* . Quality of life Research , 2, 153-159.
38. **Philippe Cibois** (2003), *Deviations from Independence. Simple Techniques for Analyzing Survey Data* . Human Sciences. 102 pages.
39. **Parker Patricia, Baile Walter, Carl de De Moor and Cohen Lorenzo** (2003), *Psychosocial and demographic predictors of quality of life in a large sampler of cancer patients* , Psycho-Oncology, 12 (2), 183-193.
40. **Popa- Velea Ovidiu , Diaconescu Liliana , Jidveian Popescu Mara and Carmen Truțescu** (2017), *Resilience and active coping style: Effects on the self-reported quality of life in cancer patients*. Int. J. Psychiatry Med. , flight. 52(2), 124–136.
41. **Reiss Kim, Joseph Herman , Marianna Zahurak , Anthony Brade, Laura Dawson , Angela Scardina et al.,** (2015). *A Phase I Study of Veliparib (ABT-888) in Combination with Low-Dose Fractionated Whole Abdominal Radiation Therapy in Patients with Advanced Solid Malignancies and Peritoneal Carcinomatosis* , Clinical Cancer Research., vol. 21, no. 21, p. 22–25

42. **Ricœur Paul** (1976), *Interpretive theory. Discourse and the surplus of the meaning* . Fort Worth, TX: Texas Christian University Press ; 1976.
43. **Sharpe Michael and Walker Jane** (2009), *Symptoms: A new approach*. Psychiatry , 8, 146-148.
<http://dx.doi.org/10.1016/j.mpps.2009.03.016>
44. **Savard, J., Simard, S., Ivers , H., and Morin, CM** (2005). *Randomized study of the effectiveness of cognitive behavioral therapy for insomnia secondary to breast cancer, part I: sleep and psychological effects*. *Journal of Clinical Oncology* , 23 (25), 6083–6096 .
45. **Septans Anne-Lyse** (2014). *Methodology for the assessment of quality of life adapted to the end of life of cancer patients* (Doctoral dissertation, University of Franche-Comté).
46. **Canadian Cancer Society** (2024). *Side effects of chemotherapy* . Available :
<https://cancer.ca/fr/treatments/treatment-types/chemotherapy/side-effects-of-chemotherapy>
47. **Snyder C laire, Blackford Amanda, Okuyama Toru , Akechi Tatsuo , Yamashita Hiroko , Toyama Tatsuya et al.** (2013), *Using the EORTC-QLQ-C30 in clinical practice for patient management: identifying scores requiring a clinician's attention* . *Qual Life Res* 2013;22(10):2685 -91.
48. **Van Manen Max** (1990), *Researching lived experience* . Human science for an action sensitive pedagogy. Buffalo, NY: State of University of New York Press .
49. **Wilson IB, Cleary PD.** Linking clinical variables with health-related quality of life. A conceptual model of patient outcomes . *JAMA* 1995;273(1): -5965.

Quality of life of cancer patients followed in the medical oncology department of a National Hospital Center of Senegal and associated factors

ORIGINALITY REPORT

20%

SIMILARITY INDEX

18%

INTERNET SOURCES

16%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1

flylib.com

Internet Source

2%

2

cegh.net

Internet Source

1%

3

worldwidescience.org

Internet Source

1%

4

www.dovepress.com

Internet Source

1%

5

"Abstract Supplement 2018 ACR/ARHP Annual Meeting", Arthritis & Rheumatology, 2018

Publication

1%

6

N. Balakrishnan. "Advances on Theoretical and Methodological Aspects of Probability and Statistics", CRC Press, 2019

Publication

1%

7

theses.gla.ac.uk

Internet Source

1%

8

www.hrpub.org

Internet Source

1%

9

Lamiaie Hallab, Asma Azzouzi, Bassima Chami. "Quality of life after extraction of mandibular wisdom teeth: A systematic review", Annals of Medicine and Surgery, 2022

Publication

1%

10

C. R. B. Joyce, Hannah M. McGee, Ciaran A. O'Boyle. "Individual Quality of Life - Approaches to Conceptualisation and Assessment", Routledge, 2013

Publication

1%

11

raybaser.r-universe.dev

Internet Source

<1%

12	Submitted to Purdue University Student Paper	<1 %
13	Submitted to University of Sydney Student Paper	<1 %
14	ebpj.e-iph.co.uk Internet Source	<1 %
15	www.mdpi.com Internet Source	<1 %
16	Cyrus Mowdawalla, Nina Kogekar, Joshua Wynne, S. Sharareh Dehghani et al. "S1819 Liver Decompensation and Mortality Outcomes Using Baveno VII Cutoffs for Transient Elastography Among Patients With Compensated Advanced Chronic Liver Disease in a Diverse, Urban US Cohort With Obesity and MASLD", American Journal of Gastroenterology, 2024 Publication	<1 %
17	d.docksci.com Internet Source	<1 %
18	assets-eu.researchsquare.com Internet Source	<1 %
19	bmcwomenshealth.biomedcentral.com Internet Source	<1 %
20	ebooks.cambridge.org Internet Source	<1 %
21	www.ncbi.nlm.nih.gov Internet Source	<1 %
22	Dagmara Kuliś, María Arnott, Elfriede R Greimel, Andrew Bottomley, Michael Koller. "Trends in translation requests and arising issues regarding cultural adaptation", Expert Review of Pharmacoeconomics & Outcomes Research, 2014 Publication	<1 %
23	dspace.ubvu.vu.nl Internet Source	<1 %
24	estudogeral.sib.uc.pt	

<1 %

- 25 Carol Ott. "Osteoporosis risk and interest in strength training in men receiving androgen ablation therapy for locally advanced prostate cancer", *Journal of the American Academy of Nurse Practitioners*, 3/2005

Publication

<1 %

- 26 ecancer.org

Internet Source

<1 %

- 27 umsha.ac.ir

Internet Source

<1 %

- 28 www.insee.fr

Internet Source

<1 %

- 29 etheses.whiterose.ac.uk

Internet Source

<1 %

- 30 jbd.or.kr

Internet Source

<1 %

- 31 kipdf.com

Internet Source

<1 %

- 32 etd.aau.edu.et

Internet Source

<1 %

- 33 hdl.handle.net

Internet Source

<1 %

- 34 tel.archives-ouvertes.fr

Internet Source

<1 %

- 35 "INTERNATIONAL COLUMN: Supportive care needs in predicting the quality of life among gynecological cancer patients", 'Pappin Communications'

Internet Source

<1 %

- 36 Diego A.S. Yamazaki, Andrew M.F. Rozada, Paula Baréa, Elaine C. Reis et al. "Novel arylcarbamate-N-acylhydrazones derivatives as promising BuChE inhibitors: Design, synthesis, molecular modeling and biological evaluation", *Bioorganic & Medicinal Chemistry*, 2021

Publication

<1 %

37	Submitted to Brunel University Student Paper	<1 %
38	epublications.uef.fi Internet Source	<1 %
39	link.springer.com Internet Source	<1 %
40	www.yumpu.com Internet Source	<1 %
41	5dok.net Internet Source	<1 %
42	Lena Rosenlund, Eskil Degsell, Asgeir Store Jakola. " Moving from clinician-defined to patient-reported outcome measures for survivors of high-grade glioma ", Patient Related Outcome Measures, 2019 Publication	<1 %
43	ir.library.louisville.edu Internet Source	<1 %
44	mafiadoc.com Internet Source	<1 %
45	pmc.ncbi.nlm.nih.gov Internet Source	<1 %
46	Journal of Services Marketing, Volume 27, Issue 6 (2013-09-14) Publication	<1 %
47	Kidane Dinku, Dereje Chala, Firaol Regea, Matiyos Lema. "Health-related quality of life and associated factors among individuals with heart failure attending public hospitals in Nekemte town, Western Oromia, Ethiopia: A cross-sectional study", Springer Science and Business Media LLC, 2024 Publication	<1 %
48	brieflands.com Internet Source	<1 %

49	Internet Source	<1 %
50	Farwa Chaudhary, Saima Dawood. "Personality traits, coping strategies and quality of life in patients of hepatitis C", European Review of Applied Psychology, 2023 Publication	<1 %
51	Gus A. Baker, Ann Jacoby. "Quality of Life in Epilepsy - Beyond seizure counts in assessment and treatment", Psychology Press, 2013 Publication	<1 %
52	Zodwa Dlamini. "Understanding Pancreatic Cancer - Global Strategies and African Perspectives", CRC Press, 2025 Publication	<1 %
53	dokumen.pub Internet Source	<1 %
54	journals.plos.org Internet Source	<1 %
55	studentsrepo.um.edu.my Internet Source	<1 %
56	synapse.koreamed.org Internet Source	<1 %
57	www.esmo.org Internet Source	<1 %
58	www.hirsla.lsh.is Internet Source	<1 %
59	www.researchsquare.com Internet Source	<1 %
60	Moustafa El-Melook, Amany Shebl, Raefa Alam, Nazem Ali. "Coping Behaviors and Health Related Quality of Life of Geriatric Patients with Cancer", Alexandria Scientific Nursing Journal, 2015 Publication	<1 %
61	Richard P. McQuellon. "Quality-of-Life Measurement in Women with Breast Cancer", The Breast Journal, 7/1997	<1 %

62	canadianoncologynursingjournal.com Internet Source	<1 %
63	dergipark.org.tr Internet Source	<1 %
64	dl.cm-uj.krakow.pl:8080 Internet Source	<1 %
65	ijpsat.org Internet Source	<1 %
66	ouci.dntb.gov.ua Internet Source	<1 %
67	repository-api.brieflands.com Internet Source	<1 %
68	ro.uow.edu.au Internet Source	<1 %
69	talenta.usu.ac.id Internet Source	<1 %
70	www.theses.fr Internet Source	<1 %
71	Berkay Ertin, Seda Kurt. "Evaluation of symptom control and functional quality of life in Turkish patients with cancer", European Journal of Oncology Nursing, 2022 Publication	<1 %
72	Yannic Volz, Lennert Eismann, Paulo Pfitzinger, Thilo Westhofen et al. "Long-term health-related quality of life (HRQOL) after radical cystectomy and urinary diversion - A propensity score-matched analysis", Clinical Genitourinary Cancer, 2022 Publication	<1 %
73	Carey, M.S.. "The prognostic effects of performance status and quality of life scores on progression-free survival and overall survival in advanced ovarian cancer", Gynecologic Oncology, 200801 Publication	<1 %

74 Dirk Vordermark, Jörn Wulf, Klaus Markert, Kurt Baier et al. "3-D conformal treatment of prostate cancer to 74 Gy vs. high-dose-rate brachytherapy boost: A cross-sectional quality-of-life survey", Acta Oncologica, 2009
Publication

<1 %

75 Won Beom Jung, Chang Sik Yu, Seok Byung Lim, In Ja Park, Yong Sik Yoon, Jin Cheon Kim. "Anastomotic Recurrence After Curative Resection for Colorectal Cancer", World Journal of Surgery, 2016
Publication

<1 %

76 e-revista.unioeste.br
Internet Source

<1 %

77 koreascience.or.kr
Internet Source

<1 %

Exclude quotes On
Exclude bibliography On

Exclude matches Off