

**QUATERNARY TERRACES OF DIFFERENT DOMAINS OF ALAKNANDA & ITS
TRIBUTARIES, BEHAVIOUR OF STATISTICAL PARAMETERS DOWN THE CURRENT
IN BADRINATH - RESHIKESH SECTION, GARHWAL HIMALAYA, UTTARAKHAND STATE
INDIA**

ABSTRACT

The sedimentological study in Alaknanda and Bhagirathi valley in upper Ganga basin has been attempted in parts of Uttarkashi, Chamoli, Pauri and Tehri districts in parts of QA sheet 53J and 53N on 1:50000 scale of Garhwal Himalaya U.P; presently known as Uttarakhand State of Union of India. The area of Upper Ganga basin consisting of Alaknanda, Bhagirathi, Bhilangna, Nandakini, Mandakini, Pindar, Dhauliganga, Balganga, Madhmeshwar Ganga and Berhi Ganga. The Alaknanda is characterised by six terraces followed by Bhagirathi with five terraces, Bhilangna Nandakini four terraces Mandakini /Pindar/Dhauliganga /Balganga three terraces, Madhmshwar Ganga two terraces and Berhi Ganga one terrace, amidst these Alaknanda is trunk stream and others are tributaries.

The Alaknanda is the trunk stream of Ganga system, it drains the eastern part of the area of study. The rocks of Alaknanda valley and adjoining area consist of three units viz Central Crystalline, Garhwal Group and Dudatoli Groups which from north to south are separated by thrust or fault. The Central Crystalline Group in this area consists of northerly dipping sequence of Kyanite schist, Garnet mica schist, quartzites and para amphibolites of Tugth formation, it is intruded by granite at Ragsi. The main Central Thrust separates it from Garhwal Group of rocks. The Dudatoli Group is represented Pauri Phyllite and Kirsu Quartzite which forms the northern limb of Dudatoli syncline. The north Almora Thrust makes its boundary with Garhwal group. The latter is divisible into Rudraprayag, Lamri, Chamoli and Gawangarh and Patrali Formation which occurs in normal stratigraphic order. It is intruded by biotite granite by biotite granite at Nainidevi and Mohankal, with tourmaline granite around Chirpatikhal and also by basic intrusive. The Rudraprayag, Lamri and Chamoli formations are equivalent to Uttarkashi Shyalnan and Nagnithank Formation respectively in Bhagirathi valley.

The Garhwal Group has been subjected to three phases of tectonic deformation. The south east to southerly plunging folds such as Marithanasa and Pingapani synclines, Karanprayag anticline were developed during the second phase of movements. The Alaknanda fault which cuts off set of the formation and earlier structures between Sunala is the strike slip fault in western part, appears to be the youngest elements. The impact of this fault is manifested in alignment of river terraces and landscape profile in Alaknanda valley. Geologically, the Bhagirathi valley and adjoining areas comprises of four distinct units namely from north to south the Central Crystalline Group, the Deoban Group, the Simla Group, and Krol belt rock separating from one another by thrust or faults. The main Central Thrust passing through Sainj upstream Uttarkashi in northern part brings the northerly dipping crystalline rocks in sharp contact with underlying Deoban Group (Garhwal Group) sedimentaries which comprises a lower Deoban Formation of Phyllite, slate, Meta basics, minor quartzite and lime stone, the middle Deoban formation of lime stone and upper Deoban formation of Quartzite and basics. The southern contact of Deoban Group is faulted one with comprising mainly siltstone, greywacks and slates dipping south.

This fault called Sringar Nalupani fault is of fundamental nature. In the southern and eastern part of the area this fault marks the contact between Deoban and Chandpur formation. In the western part south heading Ton Thrust separates the underlying Chandpur formation from the underlying Simla slates which shows abundant development of slump balls, rods etc. indicating syndepositional disturbances in the basin of sedimentation; while in Chandpur formation, is mainly argillaceous, becoming arenaceous towards the top. The Tons thrust passes through Laluli in Nagun Gad and is probably truncated by Tehri Nalupani fault at Chandpur in Bhagirathi valley.

55 The study of geology, geomorphology, Quaternary terraces and landscape profile section revealed
56 that there is sharp curvilinear break in morphogenetic expression of the area, North of Wazri in Jamuna
57 valley, North of Uttarkashi in Bhagirathi valley, around Tugnath and Chamoli and South of Joshimuth
58 in Alaknanda valley, which is significant element, appears to be due to horizontal movement of a sub-
59 tectonic plates towards south., It may be causative factor in dislocation in tectonic ecology of the
60 area, related to recent micro shocks in Joshimuth in Niti and Hellong area, mass failure of landscape
61 profile, landslide and mass wasting activities and other natural hazards; it is matter of serious
62 concerned and needs further attention.

63 The area genetically comprised of terraces of three domains, viz. Glacial,
64 Fluvio-glacial and Fluvial which represent distinct environment of sedimentation of Pleistocene, late
65 Pleistocene and Holocene time during Quaternary period. The Glacial terraces are identified at an
66 average elevation of 1150 m above MSL, the fluvio-glacial terraces at an average elevation of 975m
67 above MSL and fluvial terraces at an average elevation between 650 to 900 m above m.s.l. amidst
68 these sedimentological studies of Alaknanda and Bhagirathi terraces in type area is attempted in to
69 understand the nature of erosional and depositional processes, sedimentary pattern, behavior of
70 transporting agencies, load characteristics, current capacity, energy condition to decipher over all
71 history of Quaternary sedimentation in these valleys in increasing antiquity.

72 The average mean size of sediments of terraces of glacial domain is 0.09 ϕ (Coarse sand). It varies
73 from -2.81 ϕ to 2.8 ϕ i.e. the sediment consist of very coarse sand to fine sand. The maximum value of
74 (MZ) is -2.81 ϕ near the origin of the river and minimum 2.8 ϕ near the outer limit of these terraces
75 around Chamoli. The (MZ) shows sharp decrease in size in Badrinath and Vishnuprayag, section
76 corresponding to the steep slope of river. Down the stream Vishnuprayag although it decrease in its
77 value but display strong variation in size, which is attributed to the mixing of sediments brought by the
78 sub-glacier joining the main Alaknanda at various points. It is a measure of sorting which reflects the
79 consistency in the energy level of depositing medium. In respect of glacial terraces the average
80 standard deviation is 3.34 ϕ (very poorly sorted). It varies from 2.00 ϕ to 4.30 ϕ i.e. the sediments are
81 poorly sorted to extremely poorly sorted. The sediments of these terraces are extremely assorted and
82 are heterogenous in nature and multi source of their derivation.. The average (SKI) for glacial
83 sediments is 0.064 ϕ i.e. the sediments are negative skewed. It ranges from -0.450 ϕ to + 0.52 ϕ i.e.
84 the sediment are negative skewed to positive skewed, which indicate the tendency of gradual decrease
85 in value of (SKI) in upstream direction as result of retreat of glacier and decrease in the transport
86 capacity. The 56% of the sample shows the negative value and 44% positive value. The sediments are
87 very positive skewed to very negative skewed which seems to be due to mixing of sediments brought
88 by various glaciers. The sediments down the stream of Badrinath have the strong tendency to be
89 positive skewed.. The value of (KG) in the sediments of glacier terraces is highly variable. The average
90 value is 0.716 ϕ (platykurtic); whereas it varies from 0.49 ϕ to 1.10 ϕ (very platykurtic to
91 leptokurtic).. In Alaknanda valley along the stretch of 110 km between Chamoli and Badrinath there is
92 general uniformity in (KG) value except in the confluence area.

93

94 The average mean size of sediments of fluvio-glacial domain is 2.15 ϕ (medium sand). It varies from
95 -2.53 ϕ to 3.12 ϕ i.e. the sediments consist of very coarse to very fine sand. The size distribution of
96 these deposits in the study area is extremely irregular and erratic. The sediments near the outer edge of
97 glacial deposit downstream of Chamoli ranges 0.75 ϕ to 0.50 ϕ i.e. coarse to very coarse sand which
98 constantly show decrease in (MZ) along the stretch of 30 km up to Karanprayag. The sediments
99 between Chamoli and Karanprayag along 45 km show range of order of 1.25 ϕ to 2.75 ϕ with local
100 variation. The sudden rise in (MZ) is noticed around Nandaprayag and downstream and downstream of
101 confluence of Alaknanda and Nandakini (MZ) values are of order of 0.50 ϕ to 0.15 ϕ , which indicates
102 the intensive mixing of sediments brought from the flash stream resulting from the retreating glacier at
103 different point in the valley..The standard deviation of fluvio-glacial sediments varies from 0.95 to 2.50
104 ϕ i.e. the sediments are poorly sorted to very poorly sorted. The average standard deviation is 1.563 ϕ
105 (extremely poorly sorted). Out of 50 samples 8% are moderately sorted, 10% poorly sorted and 32%
106 are very poorly sorted. The sediments near the source area conspicuously exhibit poor sorting and show
107 significant improvement down the stream with local variation. As a whole the sediments are poorly
108 sorted to very poorly sorted and heterogeneous in nature.

109 The fluvio glacial sediments show skewness ranging from -0.48 ϕ to + 0.97 ϕ i.e. the sediments are
110 skewed very negative to skewed very positive. The average of (SKI) is 0.078 ϕ i.e., the sediments are
111 fine skewed. Out of the total samples of these terraces 56% are skewed positively, 16, skewed positive
112 and 22, are skewed very negative. The assemblage of variable value of (SKI) suggests the
113 heterogeneous association of the sediments ranging from fine sand to gravel size. The (SKI) value in
114 general increase downstream with occasional variation. It is perhaps due to repeated reworking of the
115 sediments towards downstream side by flash stream resulting from the glacier. The average (KG) is
116 1.316 (leptokurtic). It ranges from 0.76 ϕ to 1.52 ϕ (platykurtic to very leptokurtic) among these 75%
117 of the sample fall in very platykurtic class 47.50 (mesokurtic) and 45, (leptokurtic). The assemblage of
118 these different classes of kurtosis suggests the dominance of coarse sediments (Folk & Ward, 1957).
119 Most of the samples between Chamoli and Karanprayag section along the stretch of about 45 km show
120 the Kurtosis value ranging between 0.90- 1.20 ϕ except in the area around Nandaprayag Nagrasu,
121 where the sedimentation is perhaps affected by lateral mixing of sediments brought by the sub-glaciers.
122 It seems that sediments were transported and deposited in the oscillating kinetic condition. The average
123 mean size for the sediments of terraces of fluvial of Alaknanda is 2.458 ϕ . The maximum value of
124 (MZ) is -0.491 ϕ is noticed near Karanprayag while minimum 4.545 ϕ at Deoprayag, near the
125 confluence of Alaknanda and Bhagirathi River. The (MZ) shows the significant consistency in its
126 value in the first stretch of about 35 km between the Karanprayag and Nagrasu, corresponding to the
127 flatter and gentle slope of the river bed. Down the stream of Nagrasu for about 75 km up to
128 Deoprayag (MZ) sharply decreases perhaps due to sharp change in bed slope of Alaknanda. In this
129 section (MZ) strongly fluctuates around Kaliyasour, Srinagar and Kirtinagar for about 25 km which
130 correlated to the sudden convexity in the river bed due to Neotectonic activity in the vicinity of
131 Srinagar fault/ North Almora thrust, which traverses across the Alaknanda around Kaliyasour. It seems
132 that the mean size of fluvial sediments sharply follow the bed slope, pointing to exponential
133 longitudinal profile, thus decrease in (MZ) in the downstream of the Alaknanda is result of decrease in
134 both transporting capacity and velocity of the river towards the later phases of sedimentation in the
135 valley.

136 The average standard deviation of sediment is 0.691 ϕ (moderately sorted) and it ranges from 0.15 ϕ to
137 1.52 ϕ i.e. the sediments are very well sorted to poorly sorted. In the upper Alaknanda, it shows
138 consistency in value down the current except around, Karanprayag and Rudraprayag, where Nandakini
139 Pindar and Mandakini joined Alaknanda respectively. The variation in and around these places appears
140 to be due to mixing of sediments brought by these tributaries. The sharp improvement in sorting is
141 noticed downstream of Rudraprayag upto Srinagar which seems to be related with the repeated
142 reworking of sediments and slope element. The sudden decline in sorting co-efficient in the stretch of
143 about 15 km between Srinagar and Kirtinagar appears to be due to either the non-transport of larger
144 grain down current or due to loss of bed slope of Alaknanda in this segment of valley. The significant
145 increase in sorting in down current of Kirtinagar indicates cyclic reworking of sediments appears due to
146 re-activation of channel in this part of valley. The average (SKI) of fluvial sediments is 0.00281 ϕ It
147 ranges from -0.99 ϕ to 0.99 ϕ i.e. the sediments are coarse to fine skewed. The (SKI) exhibits tendency
148 of gradual increase in value downstream with local variation. This suggests relative increase of fine
149 grains, down the stream. The sediment upstream of Karanprayag is negative skewed perhaps due to
150 mixing of sediments of fluvio glacial origin. The kurtosis of fluvial sediments of Alaknanda is highly
151 variable; it ranges from 0.72 ϕ to 1.72 ϕ (leptokurtic) and an average value 1.264 ϕ (platykurtic to
152 very (leptokurtic). The average values suggest the fluctuation in the energy condition of the channel
153 system. The mean value of kurtosis revealed the more intensive sorting of central part of size
154 distribution curve than the tails. Along the course of Alaknanda in Nadaprayag and Deoprayag section
155 for a distance of about 75 km except local variation around Karanprayag, Rudraprayag, there is strong
156 tendency in increase of kurtosis value downstream.

157 The longitudinal profile of Alaknanda River and its terraces of various domain is overall concave,
158 smooth and gentle, except in the area between Karanprayag, Dharkot, Rudraprayag and Kaliyasour,
159 where it is slightly upward convex. The gradient of the river between Chamoli and Karanprayag is
160 1:6.6, between Karanprayag to Rudraprayag 1:2.25, and between Rudraprayag and Srinagar is 1:1. The
161 average gradient of terraces AT1, to AT6, between Karanprayag and Dharkot and Dharkot and
162 Kaliyasour is 1:2.29, 1:1.66, 1:1.87, 1:2.20 and 1:1.25, respectively. The upward convexity in the area as
163 mentioned above indicates some differential up warping of some of the terrace blocks possibly due to
164 some movements along the Srinagar-Tehri Fault/Alaknanda Fault (Sinha & Khan .1975, Khan
165 1981). The profile of terraces of glacial domain is restricted up Chamoli and is of hanging in nature ,
166 whereas of fluvio-glacial terraces mostly confined to the upstream of Karanprayag, it pinches out

167 downstream against the terraces of fluvial domain, thereby indicating an intensive down cutting of the
168 valley floor by Alaknanda through cyclic rejuvenation consequent to recession of in post Pleistocene
169 time. The profile of glacial terraces demonstrate intensive dissection of terraces and isolated pockets
170 and lenses of occurrence at higher level in the valley and profile is of hanging in nature, whereas
171 of fluvio- glacial is of suspended in nature and suddenly abuts up stream against the profile of glacial
172 terraces and downstream against the profile of fluvial terrace. Down the current profile of fluvial
173 terraces display consistency and smoothness down the current. The profile of fluvio-glacial terraces
174 represent transitional phase of sedimentation and major in Quaternary time the valley.

175 The longitudinal profile of Bhagirathi is over all concave except in the upper reaches between Seansu
176 and Bhatwari, where the upward convexity in the alternate segment is very conspicuous. The upward
177 convexity and development of nick points, corresponds to major tectonic elements, Main Central
178 Thrust (Khan et.al. 1982 1988). Srinagar Nalupani fault and Tons thrust in the area. The gradient of the
179 channel bed in upper Bhagirathi in Bhatwari Uttarkashi section is 1:1.98, in Uttarkashi Dharasu section
180 1:1.75, in Dharasu-Tehri 1:1.55 and Tehri and Devprayag 1:1.20. The profile of domain of glacial
181 terraces extends up to Gagnani and Fluvio-glacial terraces up to Nakuri and down the current profile
182 Fluvial terraces is gentle and smooth and in conformity of river bed. The profile of glacial terraces is of
183 hanging in nature, whereas of fluvio- glacial is of suspended in nature and fluvial display consistency
184 and regularity down the current. The profile of fluvio-glacial terraces represent transitional phase of
185 sedimentation in the valley.

186 The study of Statistical parameters and their correlation with various Thrust, fault, lineament and
187 longitudinal profile of Alaknanda and its tributaries, revealed that there is strong impact and influence of
188 tectonic and Neotectonic activity on Quaternary sedimentation in the area

189 **1.0.0 INTRODUCTION**

190 The Geological and sedimentological study in Alaknanda and its
191 tributaries has been attempted first time in parts of Uttarkashi, Chamoli, Pauri and Tehri districts, an
192 area of 10000 Sq.kms in parts of QA sheet 53J and 53 N on 1:50000 scale has been covered in
193 Garhwal Himalaya U.P; presently known as Uttarakhand State of Union of India.

194 The area of study is approachable via Dehradun and Rishikesh, which is nearest rail heads of Northern
195 Railway. These heads are connected by good motorable roads leading to famous pilgrimage centre
196 Badrinath, Kedarnath, Gangotri and Jannnonthi. The state highway No 54 which is connecting
197 Rishikesh and Badrinath bifurcates at Rudraprayag along Mandakini river and terminates at Kedarnath
198 via Sonprayag.. The Tehri is about 85 kilometers from Rishikesh on state high way No 53 connecting
199 Rishikesh Tehri Uttarkashi Gangotri. This road runs along Bhagirathi River between Tehri and
200 Gangotri, it bifurcates at Dharasu and connects Bhagirathi valley to Yamuna valley crossing the water
201 divide at Ravi Pass. In addition to these there are all weather roads which connect Kathgodam to
202 Karanprayag via Ranikhet, Dwarhat and Adi-Badri from east and Mussoori via Dhanaulti to Tehri from
203 west to Alaknanda and Bhagirathi valleys respectively. (Plate No. 1 & 2)

204 **1.1.0 Previous work**

205 The area of Upper Ganga basin consisting of Alaknanda, Bhagirathi, Bhilangna, Nandakini, Mandakini,
206 Pindar, Dhauliganga, Balganga, Madhmeshwar Ganga and Berhi Ganga. Amidst these Alaknanda
207 is trunk stream and other are tributaries. These streams emerge from different glaciers in Himalaya
208 descend in sinuous to meandering channel pattern, in their courses they traverse through entrenched
209 valleys, and deep gorges leaving glacial fluvio-glacial and fluvial terraces in decreasing antiquity; due
210 to uplift and climatic changes in the area; representing different phases of sedimentation in Quaternary
211 period.
212

213

214 Padhi and Sharan (1972) Dubey (1972), Dubey (1974a), Shukla Khan & Dubey (1974) Khan (1972-
215 73) Khan (1974), Khan et.al (1974-75) Sinha & Khan (1975), Sinha & Khan (1975-76), Dubey
216 (1974) Sinha & Khan (1976) Khan (1981), Khan (1987) (Khan, 1974, 2022, 2023). have carried
217 out geological and geomorphological and sedimentological studies in parts of
218 Alaknanda, Bhagirathi, Bhilangna, Nandakini, Mandakini, Pindar, Dhauliganga, Balganga,

219 Madhmeshwar Ganga and Berhi Ganga. The present work is part of comprehensive study carried out
220 about two decade as part of official assignment (1972 to 1981) of geological survey of India.

221

222 **1.2.0 Present work**

223 The present paper is an attempt to trace integrated evolution of fluvial terrace their stratigraphy, their
224 correlation and sedimentological aspects in type area of Alaknanda and Bhagirathi in Upper Ganga
225 Basin during Quaternary period.

226 The area under study has witnessed the intensive erosional and depositional activity subsequent to
227 recession to glaciers which has entirely modified the pre-existing, topography and given rise to present
228 shape to the area., Khan, (1975) and Khan et al, 1981). The fluvial terraces of Alaknanda and its
229 tributaries are developed and evolved in response to tectonic changes and cyclic uplift of watershed
230 region of upper Ganga during Quaternary times, (Khan 1987) The glacial, inter glacial and post glacial
231 climatic conditions, have also played the vital role in morphogenetic shaping of present day complex.
232 (Khan 1981). On the merits of evolution of fluvial terraces the sequential order of valley development
233 in upper Ganga basin is established (Khan 1987). (Table No_1 &_2)

234

235 **2.0.0 QUATERNARY GEOLOGY**

236

237 Quaternary deposits of Alaknanda and its tributaries, sequence of terraces and valley development and
238 stratigraphyt in upper Ganga Basin (Khan 1981) is described below:

239

240 **2.1.0 Glacial Terraces:**

241

242 These are the high level terraces and their occurrences are restricted above an average elevation of
243 1150 m above the m.s.l. upstream of Alaknanda, Karanpryag and upstream of Uttarkashi in Bhagirathi,
244 upstream of Kund-Chatti in Mandakini and upstream of Thirpak in Nandakini valley. These terraces
245 constitute the oldest sequence of Quaternary sediments in the area and are seen in stage of high
246 denudation as isolated pockets and lenses along the higher parts of valley flank.

247

248 In Alaknanda valley the occurrences of these terraces are noticed around Chamoli, Pipalkoti, Marwari,
249 Pandukeshar, Hanuman Chatti, Joshimuth and Badrinath. In the vicinity of Badrinath, four levels of the
250 glacial terraces (lateral Moraine) have been identified, besides the terminal moraine and cirque
251 moraine. These terraces on left flank of Vishnu Ganga constitute stepped sequence representing the
252 former levels of valley floor. Each segment of this terrace is separated by ill-preserved and highly
253 dissected scarp, which have mostly subdued sharpness and convex slopes due to prolonged erosion and
254 debris slides subsequent to their deposition in the valley.

255

256 The cirque moraine comprises both coalescing and isolated talus cones descending in the valley from
257 the sides. These cones characteristically taper off upward and terminate against the eroded thresholds
258 of abandoned cirque. At places, in the valley, such as around Badrinath and Hanuman Chatti, Mana
259 Hellong and Gobindghat, the apex of these cones partly touches the outlets of cirque depression which
260 is suggestive of activeness of these glaciers in recent time.

261

262 The type development of this cirque moraine as talus is seen around Joshimuth Badrinath section in
263 Vishnu Ganga, Joshimuth and Tapoban in Dhauli Ganga. These were mostly formed during the
264 descend of numerous cirque glacier in the valley in the pleistocene time.

265

266 The glacial terraces consist of heterogeneous assemblage of sub angular to angular, unsorted,
267 unstratified rock fragments ranging from big boulders to small pebbles in size predominantly of
268 gneiss, granite, quartzite and highly weathered biotite muscovite and chlorite schist in the matrix of
269 very coarse to fine sand, silt and clay. These rock clastics are largely angular, very poorly sorted;
270 display isotropic imbrications pattern and are devoid of bedding.

271

272 The fine sediments comprise of light smoky coarse to fine sand with appreciable amount of mica
273 flakes; light to dark maroon silt and clay, light brown coarse sand with sub-ordinate amount of silt and
274 dark red and yellow sand with silt. The sand matrix contain fairly good amount of quartz and feldspar
275 grains and loosely composed mica flakes. These sediments around Chamoli, Joshimuth and Hanuman

Chatti in Alaknanda valley display lamination and cross lamination indicative of trough like sedimentation which might have been formed by chocking of the valley during the glacier advance in Pleistocene time.

2.2 Fluvio-glacial Terraces:

The fluvio-glacial terraces are noticed at an average elevation of 975 m above the m.s.l. and their occurrences are restricted to a small stretch between Nandaprayag and Chamoli in Alaknanda. These terraces are sandwiched between the glacial and fluvial terraces as they abut against the glacial terraces in the upstream and fluvial terraces in the downstream and as such these represent the transitional phase of sedimentation.

These are characterized by sub angular to sub rounded boulder, cobble, pebble of quartzite, gneiss granite, slate and decomposed highly weathered biotitic, muscovite and chlorite schist in the matrix and coarse to fine sand, silt and clay. Although the sediments of these terraces are similar in texture and composition to glacial terraces, these exhibit entirely different order of sedimentary pattern and sediment character. In contrast to the glacial terraces, these sediments display moderate degree of sphericity roundness and sorting and show preferred orientation pattern. The ill-preserved sedimentary features and long interval cyclic development of bedding is also conspicuous. These associated diagnostic sediment characters, sedimentary features and disposition of these deposit indicate an intermittent and rapid reworking of the sediments from the glacial front subsequent to the melting of glacier during the late Pleistocene times.

2.3.0 Fluvial Terraces:

The fluvial terraces or alluvial topographic benches of Alaknanda and its tributaries form the prominent Quaternary landscape in the valley breaking the monotony of vast rugged hilly tract. These terraces are formed by a combined intermittent process of aggravation and degradation in the valley associated with different phase of sedimentation of the fluvial domain. As such these are the abandoned flood plains of the river representing the former levels of valley floors and formed due to tectonic eustatic and climatic changes during the Holocene time, (Khan 1975, Khan 1981).

These terraces are comprised of sub rounded to well rounded boulder, cobble, pebble of predominantly quartzite, gneiss, granite and schist basic, slate, phyllite, limestone, in the matrix of sand and silt with subordinate amount of clay.

In Alaknanda valley, these terraces are characterized by the association of fossiliferous boulder, cobble, and pebble of quartzite with the luxuriant assemblage of products, spirifer and other species of Brachiopods and Mullosc group typically of permo-carboniferous age. Although the other rock constituents display variation in in these terraces, but the association of fossiliferous cobble pebble is conspicuous in terraces AT₃, AT₅, around Devprayag, Srinagar, Nagrosu, Gauchar, Karanprayag and Nandprayag, which suggest the presence of some fossiliferous horizon of quartzite in the headwords' ends of Alaknanda. It appears that these fossiliferous horizons were actively subjected to erosion subsequent to recedes of glacier in post Pleistocene time. It is a marker horizon of Quaternary terraces and used a tool in correlation of terraces in valley. The type development of these terraces are seen Nagrosu, Karanprayag Gauchar, Nagrasu Gulabrai, and Nandprayag Rudrapryag and Srinagar.

The sediments of these terraces are mostly look similar in composition to the terraces of glacial and fluvio-glacial origin, but are characterized by higher order of sphericity, roundness, sorting, imbrications pattern and sedimentary features, viz. graded bedding, cross bedding, both planner and tough type, lamination, minor ripples, cut and fill features and cyclic sedimentation typically of fluvial domain. The higher order of sphericity, roundness, sorting of these sediments indicate their derivation from the distant and mixed provenances during the different phases of sedimentation in the Quaternary times.

.

Table 1:- Stratigraphy Of Quaternary Deposit In Upper Ganga Basin, Garhwal Himalaya U.P.

Age	Quaternary	Environment	of	Geomorphic	land	Composition
-----	------------	-------------	----	------------	------	-------------

	Formation	sedimentation	forms	
	Younger Alluvium	Channel and Flood Plain	Flood Plain Point Bar, Channel Bar Sand Bar	Well rounded boulder, cobble, pebble of quartzite, gneiss, schist, granite, slate, limestone, phyllite and basics in the matrix of coarse to fine micaceous sand.
Holo cone Older Alluvium	Fluvial deposit	Channel and flood plain	River terraces of Alaknanda and its tributaries	Sub-rounded to well rounded boulder, cobble, pebble of quartzite granite, gneiss, schist, phyllite, slate, limestone and basic in the matrix of coarse to fine sand silt with subordinate amount of clay
Late pleis-tocene	Fluvioglacial glacial deposit	Fluvio glacial	Fluvio glacial terraces of Alaknanda & its tributaries	Sub- Angular boulders, cobbles, pebbles of quartzite gneiss, granite biotite schist, muscovite, chlorite schist, slate and basic, in the matrix of coarse to fine sand, silt and clay.
Early pleis-tocene	Glacial deposit	Glacial	Glacial terraces, terminal moraines medial moraines, cirque moraines of Alaknanda & its tributaries.	Sub-Angular to angular boulder, cobble, pebble of gneiss, granite, quartzite, granite, quartzite, biotite, muscovite, chlorite schist, in the matrix of very coarse to very fine sand, silt and clay

The successive development of Quaternary terraces of Alaknanda is as follows:

Table 2:- Stratigraphy of Quaternary terraces and stages of valley development in Upper Ganga basin, Garhwal Himalaya U.P. India.

Age	Stages of valley development	Terraces	Environment of sedimentation
Recent to Holo-cene		Flood Plain and adjoining low land area of present day course of channel and associated geomorphic features, point bar, sand bar, channel braids etc.	Channel and Flood plain environment
	VI	Berhi Ganga terraces	BRT ₁
	V	Madhmeshwar Ganga terraces	MDT ₁ to MDT ₂
	IV	Bal Ganga terraces	MDT ₁ to BGT ₂
		Mandakini terraces	MT ₁ to MT ₃
		Pindar terraces	PT ₁ to PT ₃
		Dhaulti Ganga terraces	DGT ₁ to DGT ₃
	III	Bhilanga terraces	BHT ₁ to BHT ₄
	III	Nandakini terraces	NT ₁ to NT ₄
	II	Bhagirathi terraces	BGT ₁ to BGT ₅

	I	Alaknanda terraces	AT ₁ to AT ₃
--	---	--------------------	------------------------------------

342
343 Late – Pleistocene Fluvio-glacial terraces Fluvio-glacial of Berhi Ganga, Madhmeshwar Ganga,
344 Balganga, Mandakini, Pindar, Dhaul Ganga, Bhilangna, Nandakini, Bhagirathi and Alaknanda

345
346 Pleistocene - Glacial terraces of Berhi Ganga, Glacial Madhmeshwar Ganga, Bal Ganga,
347 Mandakini, Pindar, Dhaul Ganga, Bhilangna, Nandakini, Bhagirathi and Alaknanda.
348

349 **2.4.0 The Fluvial terraces of Alaknanda Valley**

350 The Alaknanda originates at a height of 3641 meters below Bala Kun peak 16 km. upstream of
351 Badrinath from the two glaciers of Bhagirath Kharak and Satopanth. The two glaciers rise from the
352 eastern slopes of Chaukhamba (7140 Meters) peak, Badrinath and its satellite peaks.

353 These peaks separate the Gangotri group of glaciers in the west. The major portion of the Alaknanda
354 basin falls in Chamoli district from its source upto Hellang (58 Km), the valley is treated as upper
355 Alaknanda valley. The remaining part of the area is known as lower Alaknanda valley. While moving
356 from its source, the river flows in a narrow deep gorge between the mountain slopes of Alkapuri, from
357 which it derives its name. All along its course, it drains with its tributaries. Saraswati joins the
358 Alaknanda 9 Km downstream from Mana, .Khilrawan Ganga join it below the Badrinath shrine and
359 Bhuynder Ganga below Hanuman Chatt. It is Alaknanda, the trunk stream of Ganga System forms at
360 Vishnuprayag by two tributaries, viz. Vishnu Ganga and Dhaul Ganga rising from snowy peaks north
361 of Badrinath and Niti at average elevation of 3,897 m and 5,330 m respectively in Central Himalayas.
362 The river descends in straight/sinuuous to meandering channel pattern through the deep gorges across
363 the Himalayan ranges, with sinuosity index ranging from 1.95 to 2.6 for meandering segment. It is
364 joined by Bhagirathi at Devprayag. It is further joined by numerous other tributaries in its traverse in
365 Himalaya till it finally debouches in the intermountain valley at Rishikesh. The important tributaries
366 joining Alaknanda between Vishnuprayag and Rishikesh are, Berhi Ganga at Chinka, Nandakini at
367 Nandaprayag, Pindar at Karanprayag, Mandakini at Rudraprayag and Bhagirathi at Deoprayag,
368 downstream of Deoprayag it is known as Ganga.

369 The Alaknanda has formed six prominent regional terraces AT₁-AT₆ in the valley. The AT₁, being the
370 youngest and AT₆, being the oldest terrace in the area. Each of these terraces is separated by the scarp
371 both of linear and curvilinear in nature facing towards river. These terraces are both erosional and
372 depositional in nature and display divergence and convergence in their relative disposition. The
373 depositional terraces are widely developed and have occupied the larger area in the valley around
374 Srinagar Kaliyasaur, Kirtinagar Phrases, Dungri, Gulab Rai, Nagresu, Gauchar and Langasu, Sunala
375 and characteristically found to be restricted within the meander of Alaknanda. The complete sequence
376 of terraces in the valley is seen at very few places viz. around Srinagar, Koteswar, Rudraprayag,
377 Nagrasu and Gauchar which give almost the complete account of tectonic and climatic changes in the
378 area. Whereas at other places one two or three terraces are preserved and many other have been eroded
379 away. This therefore, indicates incisive as well as rapid migrating nature of the stream.

380 The average total thickness of fluvial terraces in Alaknanda is 118 m. The highest terrace i.e. AT₆ is
381 noticed at an average elevation of 795 m from m.s.l. and 150 m from the present day course of the
382 river. The average thickness of individual terrace is 16.25 m., 22.25 m., and 27.22 m., 25 m., 21.25 m.,
383 15 m., respectively. In the upstream areas, the thickness of these terraces is constantly reduced and
384 ultimately the profile of these terraces pinches out against the glacial terraces.

385 The longitudinal profile of Alaknanda River and its terraces of different domain is based on the
386 leveling carried out in the different section in the valley. The shape of the profile is overall concave,
387 smooth and gentle, except in the area between Karanprayag, Dharkot, Rudraprayag and Kaliasaur,
388 where it is upward convexity. The gradient of the river between Chamoli and Karanprayag is 1:6.6,
389 between Karanprayag and Rudraprayag it is 1:2.25, and between Rudraprayag and Srinagar is 1:1.
390 The average gradient of terraces AT₁, to AT₆, between Karanprayag and Dharkot and Dharkot and
391 Kaliasaur is 1:2.29, 1:1.66, 1:1.87, 1:2.20 1:35 and 1:1.25, respectively. The upward convexity in the
392 river bed as mentioned indicates some differential up warping of rock blocks due major thrust/ fault in
393 the area. It indicates some recent movements along the fault and thrust / traversing the area (Srinagari
394 Fault /Alaknanda Fault and others. (Sinha et.al.1975, Khan 1981).

395 The profile of glacial and fluvio-glacial terraces mostly confined to the upstream of Karanprayag. It
396 pinches out downstream against the terraces of fluvio-glacial domain and fluvial domain down the
397 stream, thereby indicating an intensive down cutting of the valley floor by Alaknanda through cyclic
398 rejuvenation in order to achieve the base level, consequent upon to recedes of glacier, due to major
399 climatic changes in post Pleistocene time. (Plate No _3).

400 **2.5.0 The Fluvial terraces of Bhagirathi Valley**

401 The Bhagirathi is the major tributary of Alaknanda. It rises from Gangotri glacier north of Uttarkashi
402 around Gaumukh in the Central Himalaya at an elevation of about 3665 m and joins Alaknanda at
403 Deoprayag the river descends in sinuous to meandering with an average sinuosity index of order of
404 1.30. The river all along the length of 160 km. It has formed the terraces of three domains Viz. glacial
405 fluvio-glacial and fluvial.

406 The fluvial domain comprised of five major terraces which are time equivalent to the five terraces of
407 Alaknanda, the trunk stream of Ganga system. These terraces are designated BT1, to BT5. The BT1, is
408 youngest terrace and BT5, being the oldest in the valley. These terraces are mostly fill and cut type and
409 are both erosional and depositional in nature. The older terraces BT3 to BT5 are elongated, rectangular
410 in shape and have paired equivalents on both the flanks of valleys, whereas the younger terraces BT1
411 and BT2 are semi circular and crescent in shape, non-cyclic in nature and are restricted within the
412 meander of Bhagirathi.

413 The type development of these terraces is seen at very few places in the valley Viz. Uttarkashi, Dunda,
414 Chinyalisaur Chamb, Tehri, whereas other places such as Sarot Seansu and Nagor one or two levels of
415 terraces were seen, which appears to be due to exposure of the area to extensive post depositional
416 activities. The total average thickness of these terraces in the valley is 36 m. The average relative
417 thickness of these terraces in Uttarkashi and Tehri section is 6.5, and 5 m respectively. (Khan, 1974,
418 2022, 2023).

419 The longitudinal profile of Bhagirathi is over all concave except in the upper reaches between Seansu
420 and Bhatwari, where the upward convexity in the alternate segment is very conspicuous. The upward
421 convexity and development of nick points, corresponds to major tectonic elements, Main Central
422 Thrust (Khan et.al. 1982). Srinagar Nalupani fault and Tons thrust in the area. The gradient of the
423 channel bed in upper Bhagirathi in Bhatwari Uttarkashi section is 1:1.98, in Uttarkashi Dharasu section
424 1:1.75, in Dharasu-Tehri 1:1.55 and Tehri and Devprayag 1:1.20. The profile of terraces also follows
425 the channel profile.

426 The type development of terraces BT₁, to BT₅, is seen in Uttarkashi and Tehri section, in the upstream
427 of Uttarkashi. The thickness of terraces squeezed out and about two kilometers upstream of Maneri
428 where the fluvial terraces are not traceable. In the upstream Delsaur and Gagnani the development of
429 fluvio-glacial terraces are seen. The profile of fluvio-glacial terraces is of suspended in nature, It
430 pinches out upstream against the glacial terraces and downstream against the fluvial terraces and
431 represent the transitional phase of sedimentation in the valley (Sinha& Khan et.al.1975).

432 In lower Bhagirathi in Tehri and Devprayag section Bhagirathi descend down through very tight valley
433 and deep gorges, as such the development of terraces is very scanty, except in the few meandering
434 loops. There are few breaks and scar and relict rock cut terraces in the valley flanks. These breaks
435 represent the former levels of valley floors corresponding to major terraces of Bhagirathi.

436

437

438

439 **2.6.0 The Fluvial terraces of Bhilangna Valley**

440 The Bhilangna is a tributary of Bhagirathi. It originates from ice clad peaks of Central Himalaya at an
441 elevation of about 3200 m and joins Bhagirathi at Tehri. It display sinuous to meandering channel
442 pattern all along its course of 75 km and embraces the terraces of three regimes, viz. glacial, fluvio-
443 glacial and fluvial, deposited in three distinct environments during the Quaternary times.

444

445 The Fluvial terraces are designated as BHT₁, to BHT₄, the BT₁, are being the youngest terrace and
446 BHT₄, is the oldest in the valley. These terraces are mostly deposition in nature and exhibit divergence
447 and convergence in their relative disposition, the former is more conspicuous in older terraces BHT₃
448 and BHT₄, whereas the later in the younger terraces. The type development of these terraces in the
449 valley is seen around Tehri, Dewal, Asena and Ghansali. Dubey, (1972) Khan (1981). ((Khan, 1974,
450 1975 2022, 2023). The total average thickness of these terraces is about 55 m whereas the relative
451 thickness is 16, 15, 18, 16 m respectively.

452 The longitudinal profile of Bhilangna is over all concave and gentle. The gradient of terraces and river
453 is 1:120. The terraces (BHT₁, to BHT₄), are well developed between Ghansali and Tehri, the type
454 development seen at Pilichi, Asena, Dewal and Tebri. These terraces have a little lesser gradient than
455 the channel which indicates that presently the river is slowly undercutting its bed. There is significant
456 reduction in the gradient of terraces between Dewal and Asena, the gradient being slightly more
457 towards both upstream and downstream. This appears to be related with Neotectonic movement along
458 the Srinagar Nalupani fault (Sinha & Khan et.al.1975), which runs oblique to the river through Dewal
459 or along the minor fault between Dewal and Asena running parallel to the river.

460 **2.7.0 The Fluvial terraces of Nandakini Valley**

461 The Nandakini, rises from Semudra Glaciers drainage the western slopes of Trishul mountains (3660
462 m) in the Central Himalaya. It descends down in sinuous to meandering pattern, with sinuosity index
463 ranging from 1.20 to 1.25. It also passes through the straight segment of the valley and tight meanders
464 and joins Alaknanda at Nandaprayag. It comprises three distinct groups of terraces deposited entirely in
465 different environment, viz. glacial, fluvio-glacial and fluvial. In the fluvial domain four prominent
466 regional terraces in Nandakini have been identified which are time equivalent of the four younger
467 terraces of Alaknanda the trunk stream and the Bhagirathi, Bhilangna, and other major tributaries of the
468 Ganga system.

469 These Fluvial terraces are designated as NT₁, NT₂, NT₃, and NT₄, NT₁ is the youngest and NT₄ being
470 the oldest in the valley, NT₀ is the low surface of the present day flood plain of the stream. These
471 terraces are both erosional and depositional in nature. The terraces NT₄, and NT₃, and have generally
472 the divergent in mutual relation, whereas NT₂ and NT₁, have convergent relation. The NT₄ and NT₃
473 are mostly cyclic in nature and NT₁ is non-cyclic and characteristically restricted within the meander
474 of channel. The full sequence of terraces is very rarely preserved in the valley such as around
475 Nandprayag, Rajwaki, whereas at other places one or two level of terraces is seen. It is possibly due to
476 frequent lateral shift of the channel, extensive lateral cutting and subsequently repeated reworking of
477 the terrace sediments in the environments of increasing discharge and steep gradient towards the later
478 phases of river sedimentation. The full sequence of terraces NT₁- NT₄, are observed between
479 elevation of 1080 to 1100 m above m.s.l The total thickness of fluvial terraces in the valley is 60 m
480 whereas the average relative thickness of individual terraces 10, 20.25, 23.21 and 22.40 respectively.
481 ((Khan, 1974, 1975 2022, 2023). .

482 The longitudinal profile of Nandakini is quite simple as compared to Alaknanda. The slope of profile is
483 slightly concave, smooth and has gradual slope. The profile of fluvial terraces in general follows the
484 profile of present day channel. It appears that Nandakini has adjusted its course along some weaker
485 planes during upraise of head ward ends during the Holocene times (Khan 1981).

486 The gradient of river bed between Nanala and Nandprayag is 1:0.55 and 1:2.77 respectively. Between
487 Nanala and Ghat and around Chamtali there is a sudden fall in the bed slope indicating some up
488 warping neotectonic activity in the area in recent past.

489 The profile of fluvio-glacial terraces is suspended in nature; it pinches out upstream against the terraces
490 of glacial and downstream against the terraces of fluvial domain and represents the transitional phase of
491 sedimentation in the valley.

492 The profile of glacial terraces is restricted upstream of Nanala, it is mostly dissected and discontinuous
493 in nature and thereby indicating extensive erosion of these terraces by renewed depositional activities
494 subsequent to the recede of glacier in post-Pleistocene times

495

496

497 **2.8.0 The Fluvial terraces of Mandakini Valley**

498 The Mandakini rises from the Gangotri group of glaciers in the north of Sonprayag around Kedarnath
499 at an elevation of 3562 m and mingle with Alaknanda at Rudraprayag. All along its length it has
500 formed the stepped sequence of terraces of three distinct domains viz. glacial, fluvio-glacial and fluvial.
501 In fluvial domain three prominent regional terraces have been identified which are designated as MT1,
502 to MT3 and are time equivalent to the three terraces of Alaknanda and other tributaries (Khan1981).
503 These are polycyclic depositional terraces and their wide development is seen around Barhi Bhatwari,
504 Sauri, Agustmuni, Rampur and Tilwara. These terraces portrays divergent and convergent relation
505 amidst each other and are generally semicircular, elongated, semi-circular to circular in shape and are
506 both cyclic and non-cyclic in nature. These terraces conspicuous embraces sedimentary features such
507 as graded bedding, cross bedding both planer and trough type, lamination, graded lamination, cut and
508 fill features, around Agustmuni, Tilwara, Behri, Saurgarh, Sauri and Bhatwari. The total average
509 thickness of fluvial terraces in the valley between Kund Chatti and Rudraprayag is about 40 m. The
510 highest terrace is observed at an average elevation of 810 m above m.s. (Khan, 2022, 2023)

511 The longitudinal profile of Mandakini terraces in general is concave with mild convexity in the area
512 around Rampur and Tilwara. The profile of river bed is steep to gentle, between Kund Ghatti and
513 Augustmuni, it has gradient 1:10, Agustmuni and Rampur, 1:5, whereas between Rampur and
514 Rudraprayag, the gradient is considerably reduced and profile has become nearly flat. The profile of
515 both river terraces and river bed is slightly convex upward between Agustmuni and Tilwara. It possibly
516 indicates some differential up warping in the area (Khan 1988), caused by recent movement and
517 neotectonic along some lineaments traversing across in valley.

518 The fluvial terraces generally have uniform thickness and gradual gradient in Agasmuni and
519 Rudraprayag section, whereas upstream of Agasmuni it is considerably reduced. The average gradient
520 of these terraces between Agasmuni and Rudraprayag is 1:4.4, 1:4.4 and 1:5.56, respectively.

521 The profile of fluvio-glacial terraces is restricted in the middle part of the valley. It is truncated
522 upstream against the profile of glacial and downstream against the fluvial terraces, which suggests an
523 intermediate stage of sedimentation between the domains of these two terraces in the valley during the
524 quaternary times. The average gradient of these terraces in the valley is 1:52. The profile of glacial
525 terraces restricted upstream of Agustmuni, in between Kund Chatti and Bhatwari it has gradients of
526 1:9.33, and in between Bhatwari and Agustmuni is 1:92.

527 **2.9.0 The Fluvial terraces of Pindar Valley**

528 The Pindar rises from Milamand Pindar glacier from the Nandadevi group in Central Himalaya, at an
529 elevation of 3621 m. It traverses across the Himalayan ranges and descends down through steep gorges
530 forming straight sinuous to meandering channel pattern and joins its trunk stream Alaknanda at
531 Karanprayag.

532 The Pindar all along its major part of traverse maintains straight course, between Kheta and Dewal, it
533 almost drains in WNW - ESE direction. At Dewal it takes an acute turn changing its course to NNE -
534 SSW, which further down stream of Nandikesri again swings in WNW- ESE direction and maintains it
535 upto Karanprayag, where it mingles with Alaknanda. In between Dewal and Nandikesri the course of

Pindar appears to have been controlled by NNE - SSW trending fault, which up stream of Dewal also control the course of Kali Ganga a small tributary of the Pindar. All along its length it shows swelling and pinching in width, which varies from 500 to 680 m between Kheta and Dewal, 600 to 700 m between Dewal and Theralli, 500 to 650 between Theralli to Narayanbag and between Narayanbag to Nalgaon the stream passes through the straight segment of deep gorge, having steep sides of valley flanks and smallest average width is 260 m. The Pindar downstream of Nalgaon the width of valley broadens out to the range from 670 to 850 m. (Khan, 1974, 1975 2022, 2023).

In Pindar valley between Theralli - Simli and Karanprayag three prominent regional terraces have been identified. These are designated as PT₁, to PT₃. The PT₁, being the youngest and PT₃ being the oldest terrace in the valley (Khan, 1975). These terraces are correlated with the three younger terraces of Alaknanda and other tributaries. These terraces were formed by combined and intermittent processes of aggradations and degradation associated with different phases of sedimentation of fluvial regime. The process is repeated thrice in the valley during the Holocene times. The highest terrace is observed at an elevation of 830 m above m.s.l. and 60 m from the present course of channel. The total thickness of these terraces in the valley is 55.5 m whereas the average relative thickness of individual terrace is 15 , 20 and 20 m (PT₁, to PT₃) respectively.

The general shape of profile of Pindar is concave and gentle with isolated convexity and steepness at places. The concavity and gentle shape of profile indicates the graded nature and balanced cutting of the river bed, while the convexity indicates some up lift along lineaments traversing the area. At places it is also assumed to be due to good competence of bed rock constituting the river bed which is still under active cutting. The convexity in the channel profile in lower Pindar deposition of coarser material in the river bed by tributaries joining Pindar the trunk stream in the lower segment of valley. The association of innumerable channel braids under loading of Pindar is perhaps due to loss of bed slope and low energy condition. The convexity in a river profile in upper segment in catchment area related to recent movements along the lineaments traversing the area which have affected Quaternary land form and various terrace block and tilt in older terraces in Pindar valley (Khan 1974 and 1975)

2.10.0 The Fluvial terraces of Dhauli Ganga Valley

The river Dhauli Ganga rises from the Nitti Pass at about 5070 meters. It lies between the Kamet groups of peaks in the west and Nandadevi group in the east. The Dhauli Ganga takes a northern course at Malari. Between Malari and Tapoban, it is almost a narrow gorge with perpendicular cliffs on either side. The Dhauli Ganga is fed by Girthi Ganga at Kurkuti and Rishi Ganga 500 m. below Reni. It joins Vishnu Ganga, near Joshimukh and down the Vishnuprayag it is known as the Alaknanda. The river has, conspicuous straight sinuous to meandering course and descends down through the tight gorges with the steeply rising valley flanks. The river has formed three prominent fluvial terraces besides several channel and land form elements. These terraces are genetically both erosional and depositional in nature and have been designated as DHT₁, to DHT₃. The DHT₁ is being the youngest and DHT₃ is oldest terrace. The total average thickness of these terraces is 22 m and relative average thickness of these terraces is 10, 8 and 6 m respectively ((Khan, 1974, 1975 2022, 2023).

2.11.0 The Fluvial terraces of Bal Ganga Valley

The Bal-Ganga is a small tributary of Bhilagna. It rises from the ice clad peak (10746) in the Central Himalaya southwest of Kedarnath. This stream has formed three prominent fluvial terraces, beside the glacial and fluvio-glacial terraces in the upper reaches which occur in the form of isolated and dissected caps. These fluvial terraces are mainly depositional in nature and are designated as BLT₁, to BLT₃, the BLT₁, it is being the lowest and youngest terrace, whereas BLT₃, the oldest terrace. The relative average thickness of these terraces is 5, 10 and 7 m respectively. ((Khan, 1974, 1975, 2023).

2.12.0 The Fluvial terraces of Madhmeshwar Ganga Valley

The Madhmeshwar Ganga is a tributary of the Mandakini. The stream originates from glacier southwest of Kedarnath and joins Mandakini near Okhimut. It is controlled by NNE-SSW trending lineament. The stream has formed two prominent river terraces, each separated by scarp. These are depositional in nature and are fill and cut type. The shapes of these terraces are semi-circular, which is mostly restricted in the

channel meander and non-cyclic in nature. These terraces are designated as MDT1, to MDT2, from younger to older terraces respectively. The relative thickness of these terraces is 8 and 10 m respectively (Khan 1981).

These terraces constitute the complete sequence of sediments of fluvial domain measuring about 12 m deposited in Madhmeshwar Ganga valley during Quaternary period. ((Khan, 1974, 1975 2022, 2023).

2.13.0 Berhi Ganga Valley.

The Behri Ganga is a tributary of Alaknanda, it originates from glaciated top in the Central Himalaya in upstream of south east of Tapoban and joins Alaknanda between Joshimath and Chamoli it is known as the Alaknanda. The river has conspicuous straight ENE-course and descends down through the tight gorges with the steeply rising valley flanks. The river has formed one prominent fluvial terrace besides several channel and land form elements. These terraces are genetically both erosional and depositional in nature and have been designated as BRT1 is being the youngest is oldest terrace. The average thickness of these terraces is 5 m and relative average height is 8 from river bed. ((Khan, 1974, 1975 2022, 2023)

3.1.0 The Statistical Computations

The statistical analysis of sediment sample of the Alaknanda valley and particle size distribution curves were expressed on a Φ scale. Folk and Ward's (1957) graphical method was adopted to calculate mean size (M_z), sorting (σ_1), Skewness (SK_1) and Kurtosis (K_G). This method involves the measurement of several percentiles from cumulative curves (Φ_5 , Φ_{16} , Φ_{25} , Φ_{50} , Φ_{75} , Φ_{84} and Φ_{95}). The formulae are as follows:

$$\begin{aligned} \Phi &= -\log_2 G \\ \text{where } G &= \text{the grain size (mm)} \\ &(\text{i.e. sieve mesh opening}) \\ \text{Mean size} \quad M_z &= \frac{\Phi_{16} + \Phi_{50} + \Phi_{84}}{3} \\ \text{Sorting} \quad \sigma_1 &= \frac{\Phi_{84} - \Phi_{16}}{4} + \frac{\Phi_{95} - \Phi_5}{6.6} \\ \text{Skewness} \quad SK_1 &= \frac{\Phi_{16} + \Phi_{84} - 2\Phi_{50}}{2(\Phi_{84} - \Phi_{16})} + \frac{\Phi_5 + \Phi_{95} - 2\Phi_{50}}{2(\Phi_{95} - \Phi_5)} \\ \text{Kurtosis} \quad K_G &= \frac{\Phi_{95} - \Phi_5}{2.44(\Phi_{75} - \Phi_{25})} \end{aligned}$$

The computed textural parameters of sediments and their binary relation applied as tool in differentiating the various environments of Quaternary sedimentation in Moila R.J. et.al. (1968) the same key is used as tool to analyze and differentiate sediments of various domains in Narmada valley.

615

616 **3.2.0 Statistical Parameters of Glacial, Fluvio-glacial and Fluvial terraces of Alaknanda valley**
617 **:**

618 In Alaknanda valley 150 sand samples were collected, 50 each from glacial, fluvio-glacial and fluvial
619 terraces from the stretch of 225 km between Badrinath and Deoprayag. The results & findings are
620 discussed below (Plate No_1,2,3 & 4).

621 **3.2.1 Glacial Terraces:**

622 **Mean Size (MZ)**

623 The average mean size of Glacial terraces is 0.09 ϕ (Coarse sand). It varies from -2.81 ϕ to 2.8 ϕ i.e.
624 the sediment consist of very coarse sand to fine sand. The maximum value of (MZ) is -2.81 ϕ near the
625 origin of the river and minimum 2.8 ϕ near the outer limit of these terraces around Chamoli. The (MZ)
626 shows sharp decrease in size in first 25 km between Badrinath and Vishnuprayag, corresponding to the
627 steep slope of river. Down the stream Vishnuprayag although it show decrease in its value but display
628 strong variation in size, which is attributed to the mixing of sediments brought by the sub-glacier
629 joining the main Alaknanda at various points. It is seen that between Badrinath and Vishnuprayag the
630 (MZ) constantly decreases; whereas downstream, the (MZ) between Joshimuth and Chinka show sharp
631 fall, where as close to the Chamoli, it again increases. The variation in the (MZ) in glacial terrace
632 indicates the extensive mixing of sediment brought by sub-glacier meeting Alaknanda at different
633 points.

634 **Inclusive Graphic Standard Deviation (δ)**

635 It is a measure of sorting which reflects the consistency in the energy level of depositing medium. In
636 respect of glacial terraces the average standard deviation is 3.34 ϕ (very poorly sorted). It varies from
637 2.00 ϕ to 4.30 ϕ i.e. the sediments are poorly sorted to extremely poorly sorted. The
638 relative variation and average distribution indicate that 20% of samples are poorly sorted, 58% very
639 poorly sorted and 22% are extremely poorly sorted. The sediments however, show slight improvement
640 in sorting downstream with fluctuation. The sediments of glacial terraces in the vicinity of confluence
641 of major stream with Alaknanda show highly variable sorting with insignificant improvement. As a
642 whole the sediments of these terraces are extremely assorted and are heterogenous in nature and multi
643 source of their derivation.

644 **Inclusive Graphic Skewness (SKI)**

645 It denotes the symmetry of grain size frequency distribution. The symmetry curves possess zero value,
646 these with excess fine material show positive value with these excessive coarse material have negative
647 value. The average (SKI) for glacial sediments is 0.064 ϕ i.e. the sediments are negative skewed. It
648 ranges from -0.450 ϕ to + 0.52 ϕ i.e. the sediment are negative skewed to positive skewed, which
649 indicate the tendency of gradual decrease in value of (SKI) in upstream direction as result of retreat of
650 glacier and decrease in the transport capacity. The 56% of the sample shows the negative value and
651 44% positive value. Around Vishnuprayag, Chinka the sediments are very positive skewed to very
652 negative skewed which seems to be due to mixing of sediments brought by various glaciers. The
653 sediments down the stream of Badrinath have the strong tendency to be positive skewed.

654 **Graphic Kurtosis (KG)**

655 It indicates the peakedness of curve lower value of (KG) that sediment are (Platykurtic) points towards
656 broad peak, while value of (KG) (Leptokurtic) denotes pronounced peak in the centre. The value of
657 (KG) in the sediments of glacier terraces is highly variable. The average value is 0.716 ϕ (platykurtic);
658 whereas it varies from 0.49 ϕ to 1.10 ϕ (very platykurtic to leptokurtic).

659 The average value suggests the fluctuation in the energy condition of the glacier and most intense
660 sorting in the sediments prevailed during the deposition. In Alaknanda valley along the stretch of 110
661 km between Chamoli and Badrinath there is general uniformity in (KG) value except in the confluence
662 area where sudden variation are common.

663 In spite of strong variation in (KG) in the vicinity around Chinka Vishnuprayag there is tendency in
664 decrease in (KG) value towards upstream.

665 3.2.2: Fluvio-glacial terraces:

666 MEAN SIZE (MZ)

667 The average mean size of fluvio-glacial sediments is (medium sand). It varies from -2.53ϕ to 3.12ϕ
668 i.e. the sediments consist of very coarse to very fine sand. The size distribution of these deposits in the
669 study area is extremely irregular and erratic. Out of 50 samples 10% of sample show range of (MZ) of
670 order of $0.75 \phi - 0.50 \phi$, 21% 0.25 to 0.75ϕ , 25% 0.75 to 1.75ϕ , 35% 1.75ϕ to 2.50ϕ and 19%
671 beyond 2.50ϕ . The sediments near the outer edge of glacial deposit downstream of Chamoli ranges
672 0.75ϕ to 0.50ϕ i.e. coarse to very coarse sand which constantly show decrease in (MZ) along the
673 stretch of 30 km upto Karanprayag. The sediments between Chamoli and Karanprayag along a stretch
674 of 45 km show range of order of 1.25ϕ to 2.75ϕ with local variation. The sudden rise in (MZ) is
675 noticed around Nandaprayag and downstream and downstream of confluence of Alaknanda and
676 Nandakini (MZ) values are of order of 0.50ϕ to 0.15ϕ , which indicates the intensive mixing of
677 sediments brought from the flash stream resulting from the retreating glacier at different point in the
678 valley. The decrease in size downstream up to Karanprayag along a distance of about 30 km is without
679 any anomaly. The variation in (MZ) down the confluence of Alaknanda and Pindar is very
680 conspicuous, which is assumed to be the adding of a large bulk of sediments perhaps brought by sub-
681 glacier along the Pindar valley, from the close proximity of provenances.

682 Inclusive Graphic Standard Deviation (δ)

683 The standard deviation of fluvio-glacial sediments varies from 0.95 to 2.50ϕ i.e. the sediments are
684 poorly sorted to very poorly sorted. The average standard deviation is 1.563ϕ (extremely poorly
685 sorted). Out of 50 samples 8% are moderately sorted, 10% poorly sorted and 32% are very poorly
686 sorted. The sediments near the source area conspicuously exhibit poor sorting and show significant
687 improvement down the stream with local variation. As a whole the sediments are poorly sorted to very
688 poorly sorted and heterogeneous in nature.

689 Inclusive Graphic Skewness (SKI)

690 The fluvio glacial sediments show skewness ranging from -0.48ϕ to $+0.97 \phi$ i.e. the sediments are
691 skewed very negative to skewed very positive. The average of (SKI) is 0.078ϕ i.e., the sediments are
692 fine skewed. Out of the total samples of these terraces 56% are skewed positively, 16, skewed positive
693 and 22, are skewed very negative. The assemblage of variable value of (SKI) suggests the
694 heterogeneous association of the sediments ranging from fine sand to gravel size. The (SKI) value in
695 general increase downstream with occasional variation. It is perhaps due to repeated reworking of the
696 sediments towards downstream side by flash stream resulting from the glacier. The 75% of the sample
697 show skewness between the ranges -0.40ϕ to 0.30ϕ which indicates conspicuous heterogeneous
698 assemblage of sediments in terraces of fluvio-glacial domain of Alaknanda.

699 Graphic Kurtosis (KG)

700 The average (KG) is 1.316 (leptokurtic). It ranges from 0.76ϕ to 1.52ϕ (platykurtic to very
701 leptokurtic) among these 75% of the sample fall in very platykurtic class 47.50 (mesokurtic) and 45,
702 (leptokurtic). The assemblage of these different classes of kurtosis suggests the dominance of coarse
703 sediments (Folk & Ward, 1957). Most of the samples between Chamoli and Karanprayag section along
704 the stretch of about 45 km show the Kurtosis value ranging between $0.90 - 1.20 \phi$ except in the area
705 around Nandaprayag Nagrasu, where the sedimentation is perhaps affected by lateral mixing of
706 sediments brought by the sub-glaciers. It seems that sediments were transported and deposited in the
707 oscillating kinetic condition.

708

709

710

711 **3.2.3: Fluvial Terraces:**

712 **MAN SIZE (MZ)**

713 The average mean size for the sediments of fluvial terraces of Alaknanda is 2.458 ϕ . The maximum
714 value of (MZ) is -0.491 ϕ is noticed near Karanprayag while minimum 4.545 ϕ at Deoprayag, near the
715 confluence of Alaknanda and Bhagirathi River. The (MZ) shows the significant consistency in its
716 value in the first stretch of about 35 km between the Karanprayag and Nagrasu, corresponding to the
717 flattered and gentle slope of the river bed. Down the stream of Nagrasu for about 75 km upto
718 Deoprayag (MZ) sharply decreases perhaps due to sharp change in bed slope of Alaknanda. In this
719 section (MZ) strongly fluctuates around Kaliyasour, Srinagar and Kirtinagar for about 25 km which
720 correlated to the sudden convexity in the river bed due to Neotectonic activity in the vicinity of
721 Srinagar fault/ North Almora thrust, which traverses across the Alaknanda around Kaliyasour.

722 It seems that the mean size of fluvial sediments sharply follow the bed slope, pointing to exponential
723 longitudinal profile, thus decrease in (MZ) in the downstream of the Alaknanda is result of decrease in
724 both transporting capacity and velocity of the river towards the later phases of sedimentation in the
725 valley.

726 **Inclusive Graphic Standard Deviation (δ)**

727 The average standard deviation is 0.691 ϕ (moderately sorted) and it ranges from 0.15 ϕ to 1.52 ϕ i.e.
728 the sediments are very well sorted to poorly sorted. In the upper Alaknanda it shows consistency in
729 value along the greater length, except around, Karanprayag and Rudraprayag, where Nandakini Pindar
730 and Mandakini joined Alaknanda respectively. The variation in and around these places appears to be
731 due to mixing of sediments brought by these tributaries. The sharp improvement in sorting is noticed
732 downstream of Rudraprayag upto Srinagar which seems to be related with the repeated reworking of
733 sediments and slope element. The sudden decline in sorting co-efficient in the stretch of about 15 km
734 between Srinagar and Kirtinagar appears to be due to either the non-transport of larger grain down
735 current or due to loss of bed slope of Alaknanda in this Segment of valley. The significant increase in
736 sorting in down current of Kirtinagar indicates cyclic reworking of sediments appears due to re-
737 activation of channel in this part of valley.

738 **Inclusive Graphic Kewness (SKI)**

739 The average (SKI) of fluvial sediments is 0.00281. It ranges from -0.99 ϕ to 0.99 ϕ i.e. the sediments
740 are coarse to fine skewed. The (SKI) exhibits tendency of gradual increase in value downstream with
741 local variation. This suggests relative increase of fine grains, down the stream. The sediment upstream
742 of Karanprayag is negative skewed perhaps due to mixing of sediments of fluvio glacial origin.

743 **Inclusive Graphic Kurtosis (KG)**

744 The kurtosis of fluvial sediments of Alaknanda is highly variable, it ranges from 0.72 ϕ to 1.72 ϕ
745 (leptokurtic) and an average value 1.264 ϕ (platykurtic to very (leptokurtic). The average values
746 suggest the fluctuation in the energy condition of the channel system. The mean value of kurtosis
747 revealed the more intensive sorting of central part of size distribution curve than the tails. Along the
748 course of Alaknanda in Nadaprayag and Deoprayag section for a distance of about 75 km except local
749 variation around Karanprayag, Rudraprayag, there is strong tendency in increase of kurtosis value
750 downstream.

751 **4.1.0 Statistical Parameters of Glacial, Fluvio-glacial and Fluvial terraces of Bhagirathi valley :**

752 In Bhagirathi valley 150 sand samples were collected, 50 each from glacial, fluvio-glacial and fluvial
753 terraces from from the stretch of 175 km between Gangotri and Deoprayag. (Plate No_ 1, 2 4).

754 The results are discussed here under.

755

756 **4.2.0 GLACIAL TERRACES:**

757 **Mean Size (MZ)**

758 The average mean size of glacial terraces is 2.55 ϕ (Fine sand). It ranges from -0.87 ϕ to 3.42 ϕ i.e.
759 the sediments consist of coarse to fine sand with silt and clay. The (MZ) in the area around Gaumukh
760 and Gagnani section in the upper Bhagirathi significantly decreases downstream, except with very little
761 variation around Sukhi, where as a sudden rise in (MZ) is conspicuous features around Malla . The
762 behavior of mean size in this segment represents the steeper slope of Bhagirathi and anomaly in (MZ)
763 around Sukhi assumed to the lateral mixing of sediments brought by sub-glacier from close proximity.
764 In Gagnani and Bhatwari section along the length of 45 km the (MZ) shows consistency with no
765 significant variation.

766 **Inclusive Graphic standard deviation (δ)**

767 The average standard deviation is 2.35 ϕ (very poorly sorted). It varies from 1.420 ϕ to 3.885 ϕ i.e. the
768 sediments are poorly sorted to extremely poorly sorted. The majority of the sample shows the range of
769 standard deviation of order ranging from 2.00 ϕ to 3.255 ϕ . The 60% of sample show poor sorting,
770 25% very poor sorting and 35% extremely poor sorting. The sediments show significant improvement
771 in sorting downstream of Gaumukh in upper Bhagirathi, with anomalous variation around Sukhi,
772 Gagnani and Uttarkashi. The variation in sorting appears to have been related with mean size, which is
773 greatly affected by lateral mixing of sediments brought by sub-glacier joining main valley at various
774 points.

775 **Inclusive Graphic Skewness (SKI)**

776 The average skewness is 0.258 ϕ (very negative skewed). It ranges from -0.425 ϕ to + 0.215 ϕ , which
777 ascribe that the sediments are very negative to positive skewed. The 15% samples are very negative
778 skewed, 30% negative skewed, 15% nearly symmetrical and 15% positive skewed and 19% very
779 positive skewed. The 54 samples along the length of 85 km in upper Bhagirathi valley are coarse
780 skewed and 46% fine skewed. The sediments as a whole show strong departure towards coarseness as
781 well as with fineness. The departure from symmetry appears to be related with deviation of mean size.
782 In general the asymmetry passes from upstream to downstream as coarse skewed to fine skewed, which
783 indicates constant decrease in energy condition of the system during sedimentation.

784 **Graphic Kurtosis (KG)**

785 The average value of kurtosis is 0.98 (Masokurtic). It ranges from 0.65 ϕ to 1.84 % (very platykurtic to
786 very leptokurtic). The 6% samples are very leptokurtic, 50% mesokurtic, 21% leptokurtic, 23% are
787 platykurtic. The average value suggests fluctuation in the energy condition of the system. In spite of
788 variation (KG) increases downstream, which is perhaps related with sorting deviation.

789 **4.3.0 Fluvio-glacial Terraces :**

790 **Mean Size (MZ)**

791 The average mean size is 1.225 ϕ . It ranges from -2.580 ϕ to 2.255 ϕ i.e. the sediments consist of very
792 coarse sand to fine silt and clay. The maximum value - 2.580 ϕ is noticed around Bhatwari and the
793 minimum 2.255 ϕ at Nakuri. The (MZ) shows significant decrease in size from Gagnani to Uttarkashi
794 for the distance of about 35 km which corresponds to the steep slope of river bed of Bhagirathi. The
795 downstream of Uttarkashi show further significant decline in the size but with strong fluctuation, which
796 appears to be due to strong making of sediments brought by the streams resulted consequent upon the
797 meeting of glacier at various points along the length of valley.

798 **Inclusive Graphic Standard Deviation (δ)**

799 The average standard deviation is 1.752 ϕ (poorly sorted). It ranges from 1.002 ϕ to 3.421 ϕ . The 54%
800 of samples are poorly sorted and rest 44% very poorly sorted. The 94% of the samples show the sorting

801 beyond 1.302 ϕ i.e. the majority of sediments are poorly sorted to very poorly sorted. The sediments
802 show strong fluctuation in the sorting but decline to exhibit any significant improvement downstream.

803

804

805 **Inclusive Graphic Skewness (SKI)**

806 The average skewness value is -0.215 ϕ (very negative skewed). The skewness of fluvio-glacial
807 sediments varies from -0.415 ϕ to 0.325 ϕ i.e. sediments are strongly coarse to fine skewed. The 24%
808 of the samples are negative to very negative, skewed, 30% negative skewed, 30% nearly symmetrical,
809 15% positive skewed and 1% is very positive skewed. The skewness value of these sediments indicates
810 the diverse and heterogeneous association of the sediments ranging in size from fine sand to gravel.
811 The (SKI) shows sharp increase in its value downstream with local variation.

812 **Graphic Kurtosis (KG)**

813 The average kurtosis is 1.221 ϕ (very leptokurtic). It varies from 0.872 ϕ to 2.112 ϕ . The 3 samples
814 are platykurtic, 18% mesokurtic, 24% are leptokurtic, and 55% are very leptokurtic. The sediments
815 were poorly sorted in the central part of size distribution curve than the tails. In the stretch of 45 km
816 despite the local variation the (KG) does not show steady decrease in its value downstream. The local
817 variation seems to be due to local mixing of sediments brought by the net flash streams resulted due to
818 melting of glacier at different places and joining the main stream Bhagirathi at various points in the
819 valley.

820 **4.4.0 Fluvial Terraces:**

821 **Mean Size (MZ)**

822 The average mean size is 1.499 ϕ (very coarse sand). It varies from 1.252 ϕ to 2.58 ϕ i.e. the
823 sediments consist of very coarse to very fine sand, silt and clay. The (MZ) inspite of variation sharply
824 decreases downstream. In Nakuri and Tehri section for the distance of about 60 km, there is not much
825 variation of (MZ) but downstream of Tehri upto Deoprayag, there is constant decrease in the (MZ), it is
826 correlated with the former, appears to be due to gentle to flattened slope of river bed, where as the later
827 to the steep fall in the gradient of the Bhagirathi in the Tehri Deoprayag section. The local variation
828 around Nakuri, Dharasu, Tehri indicates the mixing of the sediments brought by subsequent streams
829 towards the later phase of sedimentation.

830 **Inclusive Graphic Standard Deviation (δ)**

831 The average standard deviation is 0.462 ϕ (moderately sorted). It varies from 0.220 ϕ to 1.3340 ϕ i.e.
832 the sediments are moderately sorted to well sorted. The sorting of the sediments all along the section
833 between Uttarkashi to Deoprayag to the distance of about 125 km show mark improvement
834 downstream, but the strong local fluctuations around Dharasu, Uttasu and Tehri are very conspicuous.
835 It seems to be due to lateral mixing of sediments brought by tributaries viz. Jalkhur, Bhilangna joining
836 Bhagirathi at different points. In the upper Bhagirathi in Uttarkashi Tehri section, sorting improves
837 with distance; whereas in the lower Bhagirathi in Tehri and Deoprayag section it becomes poorer. The
838 down current improvement of sorting in the upper Bhagirathi is probably related to decrease in size due
839 to non transport of the larger grains downstream. Decrease in the sorting of the lower Bhagirathi is
840 related to the increase in the fine grain sediments.

841 **Inclusive Graphic Skewness (SKI)**

842 The average skewness is + 0.265 ϕ (positive skewed). It varies from -0.423 ϕ to + 0.632 ϕ i.e. the
843 sediments are negative skewed to very positive skewed. The 70% samples are negative to very negative
844 skewed, 60% are near symmetrical, 12% is positive skewed and 14% are very positive skewed. The
845 skewness shows sharp fall in its value downstream. It suggests the in downstream increase at fine
846 fraction in the sediment load due to low carrying capacity of channel system. In the upper Bhagirathi
847 the samples show the negative value and the sediments show strong departure from symmetry towards

848 coarseness and fines around Dharasu, Seansu and Tehri. In Tehri, Deoprayag section the sediments are
849 sharply positively skewed.

850 **Graphic Kurtosis (KG)**

851 The average value of kurtosis is 1.212 ϕ (leptokurtic). It varies from 0.752 ϕ to 1.423 ϕ i.e. the
852 sediments are platykurtic to leptokurtic in nature. The 30% of the samples are platykurtic; 52% are
853 mesokurtic and 18% are leptokurtic. The average value 1.212 ϕ suggest frequential change in the
854 energy condition of channel system of Bhagirathi and more active and intense sorting of central part of
855 size distribution curve than the tail. Along the entire length of about 180 km of Bhagirathi, from its
856 source at Gaumukh to Deoprayag, where it joins Ganga, except local variation there is strong tendency
857 of decrease in (KG) downstream.

858 **5.0.0 Statistical Parameters of Fluvial terraces of Bhilangna valley :**

859 In Bhilangna valley 50 sand samples were collected from the different terraces of fluvial regime in
860 Bhilangna valley from a stretch of 70 km between Ghansali and Tehri.(Plate No.1, 2_ & 3& 5)

861 The results are are dicussed below.

862 **Mean Size (MZ)**

863 The average mean size of the sediments fluvial terraces of Bhilangna is 1.752 ϕ , whereas it ranges
864 from 0.525 ϕ to 3.22 ϕ i.e. the sediments consist of coarse to fine sand. The maximum value of (MZ)
865 is 6.525 ϕ near Ghanshali, while minimum is 3.22 ϕ near the confluence with Bhagirathi at Tehri.
866 Inspite of local variation (MZ) it decreases downstream, which corresponds to the steep slope of
867 Bhilangna. The variation in (MZ) indicates the mixing of sediments brought by subsequent stream
868 joining Bhagirathi at various point.

869 Nevertheless, the (MZ) of fluvial terraces of Bhilangna broadly follows the bed slope pointing to
870 exponential longitudinal profile. The decrease in (MZ) size indicates both decrease in transporting
871 capacity and velocity of the channel system.

872 **Inclusive Graphic Standard Deviation (δ)**

873 The average standard deviation (δ) of sediments is 1.552 ϕ . It ranges from 0.581 ϕ to 2.121 ϕ i.e. the
874 sediments are poorly sorted to well sorted. Except with little local variation, the sorting of sediments
875 increases downstream. The variation suggests the mixing of local sediments brought by contributories
876 streams from close proximity. The 75% of the samples are represented by the range of sorting of order
877 of 1.225 ϕ to 1.852 ϕ , the 20% 1.852 ϕ to 2.122 ϕ and 5% 0.580 ϕ to 1.852 ϕ .down thestream
878 respectively.

879 **Inclusive Graphic Skewness (SKI)**

880 The average skewness is +0.355 ϕ . It varies from -0.525 ϕ to +0.555 ϕ i.e. the sediments are negative
881 to positive skewed. The 35% samples are nearly symmetrical, 45% are positive skewed. 12% very
882 positive skewed 3% negative skewed and 3% very negative skewed. The skewness value sharply
883 increased downstream, which suggest the constant increase of fines fraction of sediments and low load
884 capacity of the channel system.

885 **Inclusive Graphic Kurtosis (KG)**

886 The average value of kurtosis is 1.285 ϕ . It ranges from 0.552 ϕ to 1.560 ϕ i.e. the sediments are very
887 platykurtic to leptokurtic. The average value of (KG) indicates fluctuation in the energy of the channel.
888 The 66% samples are leptokurtic, 12% are mesokurtic 12% are platykurtic, 10% are very platykurtic.
889 Inspite of strong fluctuation the (KG) shows steady increase in its value downstream.

890

891 **6.0.0 Statistical parameters of Fluvial terraces of Nandakini valley:**

892 In Nandakini valley 50 sediment samples collected for statistical analysis from from the different
893 terraces of fluvial regime from the stretch of 70 km between Ghat and Nandaprayag for
894 sedimentological study. (Plate No.1, 2_ & 3& 6)

895 **Mean Size (MZ)**

896 The average mean size of the sediments of fluvial terraces of Nandakini is 1.765 ϕ , whereas it ranges
897 from -0.330 ϕ to 3.255 ϕ i.e. the sediments consist of very coarse sand to very fine sand. The
898 maximum size of sediment -0.330 ϕ is noticed around Ghat, whereas the minimum size 3.255 ϕ is
899 around Nanda- prayag. The (MZ) except little variation it display steady decreases downstream,
900 which suggest repeated reworking of sediments from the source and also decline in the steady load
901 carrying capacity of the channel system towards downstream.

902 **Inclusive Graphic Standard Deviation (δ)**

903 The average standard deviation (δ) is 1.655 ϕ of sediment it varies from 0.625 ϕ to 2.850 ϕ i.e. the
904 sediment are moderately sorted to poorly sorted. The sediments show improvement in sorting
905 downstream. The 20% samples are moderately sorted 35% are poorly sorted and 45% are very poorly
906 sorted. The 80% of samples in the upper Nandakini are poorly sorted to very poorly sorted, which show
907 sharp improvement downstream. The poor sorting in the upper Nandakini valley is due to close
908 proximity of sediments source and improvement in sorting-downstream by repeated reworking of
909 sediments and steady increase of finer fractions.

910

911 **Inclusive Graphic Skewness (SKI)**

912 The average value of skewness is +0.285 ϕ it varies from -0.525 ϕ to + 0.550 ϕ i.e. the sediments are
913 negative skewed to very positive skewed. The 42% samples are negative skewed, 30% samples are
914 nearly symmetrical 18% are positive skewed and 10% samples are very positive skewed. The (SKI)
915 value increases downstream indicates calm and stable energy condition towards late history of
916 sedimentation in valley .

917 **Graphic Kurtosis (KG)**

918 The average value of kurtosis is 0.745 ϕ (platykurtic) and it varies from 0.525 ϕ to 1.385 ϕ i.e. the
919 sediments are very platykurtic to leptokurtic. Out of the samples 10% are very platykurtic 18% samples
920 are platykurtic 26% samples mesokurtic, 20% are leptokurtic and 26% are very leptokurtic, except
921 local variation (KG) has got strong tendency to decrease downstream

922

923 **7.0.0. Statistical Parameters of Fluvial terraces Mandarini valley:** .(Plate No.1, 2_ & 3& 7)

924 In Mandakini valley 70 sand samples were collected from the different terraces of fluvial regime from a
925 stretch of 70 km between Guptkashi and Rudraprayag (Plate No.1, 2_ & 3& 7)

926 The results are discussed below

927 **Mean Size (MZ)**

928 The average mean size is 2.688 ϕ . It varies from 0.821 ϕ to 4.225 ϕ i.e. the sediments consist of very
929 coarse to very fine sand. The (MZ) except a little variation it progressively decreases downstream. In
930 the upper stretch between Guptkashi and Kund-Chatti, it shows steady value, whereas downstream it
931 constantly decreases upto Rudraprayag. The sharp fall in (MZ) is noticed between Agastmuni, and
932 Rudraprayag which appears to be related with steep gradient of the Mandakini River.

933

Inclusive Graphic Standard Deviation (δ)

934 The average standard deviation is 0.985 ϕ (moderately sorted). It varies from 0.452 ϕ to 1.955 ϕ
935 (poorly sorted to well sorted). The standard deviation of sediment except some local variation, show
936 steady improvement downstream. The fluctuation in sorting around Kund-Chatti and Agastmuni are of
937 very strong nature, these appear to have been related with the mixing of sediments, brought by net
938 work of streams, joining Mandakini in this segment of valley. In the upper Mandakini, sorting
939 significantly increases with the distance, whereas in the middle segment of Mandakini it strongly
940 fluctuates and in the lower part of valley downstream of Agastmuni, it shows sharp improvement. The
941 overall improvement down the current is related to the increase in fine grained fraction of sediments
942 and repeated reworking of sediments.

943

Inclusive Graphite Skewness (SKI)

944 The average value of skewness is 0.438 ϕ . This varies from 200 ϕ to +0.82 ϕ i.e. the sediments are
945 negative skewed to very positive skewed. The 55% of samples show, skewness ranging between -0.10 ϕ
946 to 0, 10 ϕ , the 6% beyond +0.410 ϕ , 35% 0.10 ϕ to +0.40 ϕ and 5% between -0.20 ϕ to 0.30 ϕ . The
947 strongly skewed positive to strong skewed negative tendency of sediments from upper to lower
948 Mandakini revealed constant increase in finer sediments downstream. The little variation in skewness
949 around Agastmuni and Rampur, appears to be due to local mixing of sediments brought by small
950 stream from close proximity.

951

Graphite Kurtosis (KG)

952 The average value of kurtosis is 1,135 ϕ (leptokurtie). It ranges from 0,60 ϕ to 1.345 ϕ . It ranges from
953 The 10% samples are (platykurtie), 30% (platykurtio, to mesokurtie), 37% are leptokurtie, and 23%
954 are very leptokurtic. The kurtosis value, except local variation around Kund-Chatti, and Rampur
955 constantly decreases downstream along the length of 70 km from Guptkashi to Rudraprayag. (Khan
956 1985).

957

8.0.0 Statistical Parameters of Fluvial terraces of Pindar valley:

958 In Pindar valley 30 sand samples were collected from the different terraces of fluvial regime from a
959 stretch of 65 km between Thanala Karanpryag for sedimentological study. (Plate No.1, 2_ & 3& 9)

960 The results are discussed below

961

962

Mean Size (MZ)

963 The average mean size of sediments of fluvial terraces is 1.552 ϕ . It ranges from -0.752 ϕ to 3.255 ϕ
964 i.e. the sediments consist of coarse to very fine sand. Except local variation (MZ) it decrease
965 downstream. The variation of (MZ) in the valley appears to have been related with the local addition of
966 the sediments to the main sediment regime of the channel by small tributaries. The 27% of the samples
967 show the (MZ) of order ranging between 0.752 ϕ to 1.255 ϕ , whereas the 63% between 1,255 ϕ , to
968 3.552 ϕ . The sediment samples in the lower Pindar valley show average value of (MZ) around 1.550
969 ϕ , which indicate repeated reworking of sediments downstream in considerably stabilized energy
970 condition of the channel perhaps due to less variation in the channel gradient.

971

Inclusive Graphic Standard Deviation (δ)

972 The average standard deviation is 1.855 ϕ . It varies from 0.625 ϕ to 2.820 ϕ i.e. the sediments are
973 poorly sorted to well sorted. The sorting of sediments in general except local variation increases
974 downstream. The 30% samples are very poor sorted, 25% moderately sorted and 45% are well sorted.
975 As a whole the sediments show improvement downstream in valley.

976

Inclusive Graphic Skewness (SKI)

977 The average skewness is +0.258 ϕ . It varies from -0.352 ϕ to 0.425 ϕ i.e. the sediments are negative
978 skewed to very positive skewed. The 40% samples are negative skewed, 30% nearly symmetrical and
979 18% positive skewed and 12% are very positive skewed. The skewness values except local variation
980 increases downstream. The strong tendency of the sediment from very negative skewed to very positive
981 skewed indicate the steady increase of finer sediments towards down current in the valley.

982 **Inclusive Graphic Kurtosis (KG)**

983 The average kurtosis value is 1.165 ϕ . It varies from 0.752 ϕ to 1.255 ϕ i.e. the sediments are
984 platykurtic to very leptokurtic. The 30% samples are platykurtic, 50% are mesokurtic, 20% are
985 leptokurtic and very leptokurtic. In spite of local variation in value of (KG) there is a strong tendency
986 for decrease in its value downstream.

987 **9.0.0 Statistical Parameters of the sediment of Fluvial terraces of Dhauli -Ganga valley:**

988 In Dhauli Ganga valley 19 sediment samples were collected from the different terraces of fluvial
989 regime from a stretch of 65 km between Niti pass and Joshimuth for sedimentological study.
990 (Plate No.1, 2_ & 3). The results are discussed below

991

992 **Mean size (1)**

993 The average mean size for sediments of fluvial terraces of Dhauli-Ganga is 2.251 ϕ (fine sand). It
994 varies from 1.370 ϕ to 3.158 ϕ i.e. the sediments consist of pre-dominantly medium to fine sand. The
995 mean size in general decrease down the stream, except local variation, in the middle part of valley
996 between Malari and Lata and thereafter the mean size decrease downstream. The downstream of Lata
997 upto Joshimuth the sediment exhibit steep fall in mean size, which seems appears to be related with
998 the steep slope of valley. The mean size show inverse relation with standard deviation all along the
999 length of 110 km in the Dhauli-Ganga between Niti pass and Joshimuth as the mean size decreases
1000 down the current sediment show improvement in sorting.

1001 **Inclusive Graphic Standard Deviation (δ)**

1002 The average standard deviation for the sediments of fluvial terraces of Dhauli-Ganga is 0.269 ϕ
1003 (moderately sorted) and it varies from 0.55 ϕ to 4.92 ϕ i.e. the sediments are extremely poorly sorted
1004 to moderately sorted. In the upper part of Dhauli-Ganga between Niti pass and Malari the sediments
1005 show little variation and majority of samples display extremely poor sorting. Although sediments show
1006 little variation in values, but exhibit significant improvement in sorting down the current. In Lata and
1007 Joshimuth in the lower part of the valley, the sediments show fluctuation, which appears to be related
1008 with decrease in (MZ) and increase in coarser fraction of sediment in the valley. In general sorting of
1009 sediment except local variation show improvement down the stream.

1010 **Inclusive Graphic skewness (SKI)**

1011 The average value of skewness is -0.425 ϕ (very positive skewed). It varies from +0.541 ϕ to -0.562 ϕ
1012 i.e. the sediment are negative skewed to very positive skewed. In the upper reaches in Niti pass and
1013 Malari section the sediments are strongly negative skewed i.e. the sediment predominantly consist of
1014 coarse sediments. In the middle part of valley in Malari and Lata section it shows strong variation
1015 i.e. the sediments are both strongly negative skewed and positive skewed. In Lata and Joshimuth section
1016 the sediment display steep rise in the skewness values i.e. the sediment become, strongly positive
1017 skewed. In the upper reaches strongly negative skewed nature of sediment reveal the close proximity of
1018 provenance. The variation in skewness in Malari and Lata section indicates strong lateral mixing of
1019 sediments by subsequent streams. In Lata and Joshimuth the strongly positive skewed nature of
1020 sediments reveal relatively increase in fine grained fraction in the down the stream which is correlated
1021 with decline in transporting capacity due to low energy condition of sedimentation.

1022 **Graphic Kurtosis KG)**

1023 The average value of kurtosis is 0.825 ϕ (platykurtic). It varies from 0.432 ϕ to 1.312 ϕ very
1024 platykurtic to leptokurtic. In upper reaches between Niti pass and Malari, except local variation the
1025 value of kurtosis in general decreases. In Malari and Lata section it show strong fluctuation, whereas in
1026 Lata and Joshimuth section local variation of kurtosis sharply decrease i.e. the sediments have strong
1027 tendency to become from leptokurtic to very platykurtic down the stream.

1028 **10.0.0 Statistical Parameters of the sediments of Fluvial terraces of Bal-Ganga valley:**

1029

1030 In Bal Ganga valley 16 sediment samples collected from fluvial terraces for statistical analysis from
1031 the stretch of 40 km between between Danwali and Ghansyali . . (Plate No.1, 2_ & 3& 8)

1032 The results are discussed below

1033 **Mean Size (MZ)**

1034 The average mean size for sediments of fluvial terraces of Bal-Ganga is 2.392 ϕ (fine sand). It varies
1035 from -2.98 ϕ to 3.10 ϕ i.e. the sediment mainly comprises of very small pebbles to very fine sand. The
1036 (MZ) in upper reaches upstream of Thathikathur, except little variation in the middle part of valley it
1037 progressively decrease down the stream. In between Thathikathur and Chamyal a it shows strong
1038 variation, whereas down the stream of Chamyal a a significant steep fall in mean size is noticed. The
1039 variation in the Thathikathur and Chamyal a seem to be related with lateral mixing of sediments. The
1040 conspicuous decrease in (MZ) down the stream of Chamyal a appears to have related both with the
1041 steep change in the valley gradient and repeated reworking of sediments in the valley. The (MZ) has
1042 inverse relation with sorting in the valley, as the mean size decreases down the stream the sediments
1043 show significant in sorting.

1044 **Inclusive Graphic Standard Deviation (δ)**

1045 The average standard deviation for sediment of terraces of fluvial domain is 0.388 ϕ (well sorted),
1046 where as it varies from 0.327 ϕ to 0.51 ϕ i.e. the sediment of fluvial terraces are moderately sorted to
1047 very well sorted. In upper reaches upstream of Thathikathur except little variation the sorting values
1048 decrease down the stream which indicates improvement in sorting. In the Thathikathur and Chamyal a
1049 section it show strong fluctuation which is perhaps due to lateral mixing of sediments, whereas down
1050 the stream of Chamyal a the sediment show distinct and sharp improvement in sorting which is
1051 inversely related with the (MZ). The steep fall in the valley gradient down the stream of Thathikathur
1052 also has important bearing on these two parameters in the valley.

1053 **Inclusive Graphic Skewness (SKI)**

1054 The average value of skewness is +0.385 ϕ that the sediment are very positive skewed. It varies from -
1055 0.553 ϕ to + 0.551 ϕ i.e. the sediments are negative skewed to positive skewed. In the upper stream of
1056 Thathikathur the sediment generally show little variation in skewness value and are negative skewed.
1057 In between the Thathikathur and Chamyal a skewness show strong variation in values i.e. sediments are
1058 nearly symmetrical and are both negative and positive skewed, whereas down the stream of Chamyal a,
1059 there is steep increase in values of skewness and the sediment have got very strong tendency to become
1060 positive skewed. In general the skewness values increases down the current indicating progressive
1061 increase of finer sediments in the lower part of valley, it seems to be due to with repeated reworking
1062 and rapid transport of sediments from the source area during sedimentation.

1063 **Graphic Kurtosis (KG)**

1064 The average value of kurtosis is 0.898 ϕ (Platykurtic). It varies from 0.525 ϕ to 1.521 ϕ i.e. sediments
1065 are leptokurtic to very platykurtic in nature. In upstream of Thathikathur, except little variation the
1066 overall value of kurtosis decreases down the stream i.e. the sediments become from leptokurtic to
1067 mesokurtic in nature. Thathikathur and Chamyal a section inspite of fluctuation the value of kurtosis
1068 increases i.e. sediments show tendency to become leptokurtic, whereas down the stream of Chamyal a
1069 the values of kurtosis sharply decreases and the sediment show strong tendency to become platykurtic
1070 in nature. In general the kurtosis except local variation between Thathikathur and Chamyal a section it

1071 displays steady decrease in values i.e. the sediments show the tendency to become from leptokurtic to
1072 platykurtic in nature down the current in the valley.

1073 **11.0.0 Statistical Parameters of the sediments of Fluvial Terraces of Madhmeshwar Ganga**
1074 **valley:**

1075 In Madhmeshwar Ganga valley 16 sediment samples were collected from fluvial terraces for statistical
1076 analysis from the stretch of 45 km between Bantoli and Okhimuth. (Plate No.1, 2_ & 3&10)

1077 The results are discussed below:

1078 **Mean size (MZ)**

1079 The average mean size for sediments of fluvial terraces of Madhmeshwar Ganga is 1.199 ϕ (medium
1080 sand). It varies from 1.522 ϕ to 1.989 ϕ i.e. the sediments pre-dominantly medium to very coarse
1081 sand. In the upstream of Ransi the (MZ) constantly decreases down the stream, perhaps due to steeper
1082 slope of the valley. In between Ransi and Rawa it shows strong variation, whereas down the stream of
1083 Ransi a significant and progressive decrease in (MZ) is noticed. The variation, in (MZ) between Ransi
1084 and Rawa appear to related with the lateral mixing of sediments brought by net work of subsequent
1085 stream joining the central part of valley at various places, whereas the decrease of (MZ) downstream of
1086 Rawa is seem to be with due to steep bed slope in the lower valley.

1087 **Inclusive Graphic Standard Deviation (S)**

1088 The average standard deviation of the sediments of fluvial terraces of Madhmeshwar Ganga is 1.359
1089 (very poorly sorted). It varies from 0.521 ϕ to 3.489 ϕ i.e. the sediments are very poorly sorted to
1090 moderately sorted. In the upstream of Ransi the values of sorting decrease down the stream, which
1091 indicate improvement in sorting of sediments. In between Ransi and Rawa it shows strong fluctuation,
1092 which seem to be due to lateral mixing of sediments in central part of valley. The downstream of Ransi
1093 values of sorting progressively decreased depicting the sharp improvement in sorting of the sediment
1094 down the current of valley. This shows inverse relation of (δ) with (MZ) down the current i.e. as the
1095 sorting of sediments increases (MZ) decreases.

1096 **Inclusive Graphic skewness (SKI)**

1097 The average value of skewness is + 0.496 ϕ (very positive skewed). It varies from +0.22 ϕ to + 0.725
1098 ϕ i.e. the sediments are positive skewed to very positive skewed. In the upstream between Rawa and
1099 Banloli the values of skewness show variation i.e. the sediments are both coarse skewed and fine
1100 skewed. The downstream of Rawa the sediment show constant increase in values of skewness which
1101 reveal the strong tendency of sediments to become fine skewed as the result decrease in load carrying
1102 capacity of channel system during sedimentation perhaps due to loss in bed slope in the lower part of
1103 valley.

1104 **Graphic kurtosis (KG)**

1105 The average value of kurtosis is 0.799 ϕ (platykurtic). It varies from 0.320 ϕ to 1.210 ϕ i.e. the
1106 sediments are leptokurtic to very platykurtic in nature. In upstream of Ransa the values of kurtosis
1107 show strong fluctuation, whereas the down the stream of Ransa there is marked decrease in the value of
1108 kurtosis i.e. the sediments show strong tendency to become platykurtic in nature down the current in
1109 the Madhmeshwar Ganga valley.

1110 **12.0.0 Statistical Parameters of the sediments of Fluvial terraces of Berhi-Ganga valley:**

1111 In Dhauli Ganga valley 10 sediment samples were collected from fluvial terraces for statistical
1112 analysis from the stretch of 25 km. (Plate No.1, 2_ & 3)

1113 The results are discussed below:

1114

1115 **Mean size (MZ)**

1116 The average mean size for the sediments of fluvial terraces of Berhi Ganga is 0.2695 ϕ (very coarse
1117 sand). It varies from -0.329 ϕ to +0.325 ϕ i.e. the sediments consist of very coarse sand to very fine
1118 sand. In the Berhi Ganga except local variation around Irni the (MZ) of sediment decrease down the
1119 current which appears to be related with the steep slope of valley.

1120 **Inclusive Graphic Standard Deviation (δ)**

1121 The average standard deviation of sediments of fluvial terraces in Berhi Ganga is 2.430 ϕ (very poorly
1122 sorted). It varies from 0.521 ϕ to 3.272 ϕ i.e. sediments in general are poorly sorted to moderately
1123 sorted. Except little variation in sorting around Irni the sorting of sediments show progressive and
1124 sharp improvement down the current assumed to be due to repeated reworking of sediment and steep
1125 slope of valley segment

1126 **Inclusive Graphic skewness (SK)**

1127 The average value of skewness is + 0.188 ϕ (positive skewed). It varies from -0.285 ϕ to 0.248 ϕ i.e.
1128 the sediments are negative skewed to positive skewed. In the upstream of Irni the skewness of
1129 sediments show strong fluctuation indicating that the sediments are both strongly fine skewed and
1130 coarse skewed. Down the stream of Irni significant increase in skewness value reveal the tendency of
1131 sediments to become fine skewed indicating decrease in transporting capacity of channel down the
1132 current in the valley.

1133 **Graphic kurtosis (KG)**

1134 The average value of kurtosis is 1.256 ϕ (very leptokartie). It varies from 0.546 ϕ to 1.589 ϕ (very
1135 leptokurtic to very platykurtic). The kurtosis value except variation upstream of Irni constantly
1136 decreases which indicate strong tendency of sediments to become platykurtic in nature down the
1137 current in the Berhi Ganga.

1138 **13.0.0 Statistical Parameters of the sediments of Fluvial terraces of Ganga River :**

1139 In Ganga valley 30 sediment samples collected from fluvial terraces for statistical analysis from the
1140 stretch of 85 km down the stream of Deoprayag to Reshikesh .(Plate No.1, 2_ & 3)

1141 The results are discussed below:

1142

1143 **Mean size (MZ)**

1144 The average mean size of sediments is 2,288 ϕ (fine sand) it varies from 0.480 ϕ to 3.720 ϕ i.e. the
1145 sediments consist of coarse sand to fine silt and clay. The 55 samples show mean size between 2.50 ϕ
1146 to 3.50 ϕ , 21% between 1.00 to 2.00 ϕ and 24% below 1.00 ϕ and 14%. -2.98 ϕ to 3.10 ϕ i.e. the
1147 sediment mainly comprises of very small pebbles to very fine sand. The mean size except local
1148 variation constantly decreases down the stream.

1149 **Inclusive Graphic Standard Deviation (δ)**

1150 The averages standard deviation of sediment is 0.881 ϕ (moderately sorted), whereas it varies from -
1151 0.310 ϕ to 0.140 ϕ i.e. the sediments are moderately sorted to very well sorted. The 55% of sample
1152 show sorting between 0.40 ϕ to 1.20 ϕ the 38% between 1.20 ϕ to 1.80 ϕ and 7% between 2.00 ϕ to
1153 2.32 ϕ . The sediments show sharp improvement in sorting down the stream of Ganga valley.

1154 **Inclusive Graphic Skewness (SKI)**

1155 The average skewness value is -0.185 ϕ (negative skewed). It varies from -0.312 ϕ to 0.145 ϕ i.e. the
1156 sediments are negative skewed to very positive skewed. The 20% samples show skewness between

1157 0.040 Ø to 0.152 Ø, 35% - 0.045 Ø to 0.098 Ø and 45% samples between 0.450 Ø to 0.210 Ø except
1158 little variation the skewness value increase downstream.

1159 **Inclusive Graphic Kurtosis (KG)**

1160 The average kurtosis value is 1.716 Ø (very leptokurtic). It ranges from 0.550 to 1.990 Ø i.e. the
1161 sediments are very platykurtic to very leptokurtic. The 65% samples show kurtosis value of order
1162 between 0.860 Ø to 1.120 Ø, 25% between 1.262 Ø to 1.700 Ø and 10% between 0.620 Ø to 0.650 Ø
1163 except local variation the kurtosis value decrease downstream.

1164 **14.0.0 CONCLUSION**

1165 The sedimentological study in Alaknanda and its tributaries in upper Ganga basin
1166 has been attempted in parts of de Uttarkashi, Chamoli, Pauri and Tehri districts in parts of QA sheet 53J
1167 and 53 N on 1:50000 scale of Garhwal Himalaya U.P; presently known as Uttarakhand State of Union
1168 of India. The area of Upper Ganga basin consisting of Alaknanda, Bhagirathi, Bhilangna, Nandakini,
1169 Mandakini, Pindar, Dhauliganga, Balganga, Madhmeshwar Ganga and Berhi Ganga. The
1170 Alaknanda is characterised by six terraces followed by Bhagirathi with five terraces, Bhilangna
1171 Nandakini four terraces Mandakini /Pindar/Dhauliganga/Balganga three terraces, Madhmeshwar
1172 Ganga two terraces and Berhi Ganga one terrace, amidst these Alaknanda is trunk stream and others
1173 are tributaries. The Alaknanda is trunk stream and other are tributaries.

1174 The Alaknanda is the trunk stream of Ganga system, it drains the eastern part of the area of study. The
1175 rocks of Alaknanda valley and adjoining area consist of three units viz Central Crystalline, Garhwal
1176 Group and Dudatoli Groups which from north to south are separated by thrust or fault. The Central
1177 Crystalline Group in this area consist of northerly dipping sequence of Kyanite schist, Garnet mica
1178 schist quartzites and para amphibolites of Tugnath formation, it is intruded by granite at Ragsi. The
1179 main Central Thrust separates it from Garhwal Group of rocks. The Dudatoli Group is represented Pauri
1180 Phyllite and Kirsu Quartzite which forms the northern limb of Dudatoli syncline. The north Almora
1181 Thrust makes its boundary with Garhwal group. The latter is divisible into Rudrapur, Lamri,
1182 Chamoli and Gawangarh and Patrali Formation which occurs in normal stratigraphic order. It is
1183 intruded by biotite granite by biotite granite at Nainidevi and Mohankal, with tourmaline granite
1184 around Chirpatikhal and also by basic intrusive. The Rudrapur, Lamri and Chamoli formations are
1185 equivalent to Uttarkashi Shyaln and Nagnithank Formation respectively in Bhagirathi valley.
1186

1187 The Garhwal Group has been subjected to three phases of tectonic deformation. The south east to
1188 southerly plunging folds such as Marithanasa and Pingapani synclines Karanprag anticline were
1189 developed during the second phase of movements. The Alaknada fault which cuts off set of the
1190 formation and earlier structures between Sunala is the strike slip fault in western part, appears to be the
1191 youngest elements. The impact of this fault is manifested in alignment of river terraces and land scape
1192 profile in Alaknanda valley. Geologically, the Bhagirathi valley and adjoining areas comprises of four
1193 distinct units namely from north to south the Central Crystalline Group, the Deoban Group, the Simla
1194 Group, and Krol belt rock separating from one another by thrust or faults. The main Central Thrust
1195 passing through Sainj upstream Uttarkashi in northern part brings the northerly dipping crystalline
1196 rocks in sharp contact with underlying Deoban Group (Garhwal Group) sedimentary which comprises
1197 a lower Deoban Formation of Phyllite, slate, Meta basics, minor quartzite and lime stone, the middle
1198 Deoban formation of lime stone and upper Deoban formation of Quartzite and basics. The southern
1199 contact of Deoban Group is faulted one with comprising mainly siltstone, greywacks and slates dipping
1200 south
1201
1202

1203 This fault called Sringer Nalupani fault is of fundamental nature. In the southern and eastern part of the
1204 area this fault marks the contact between Deoban and Chandpur formation. In the western part south
1205 heading Ton Thrust separates the underlying Chandpur formation from the underlying Simla slates
1206 which shows abundant development of slump balls, rod etc. indicating syndepositional disturbances in
1207 the basin of sedimentation; while in Chandpur formation, is mainly argillaceous, becoming arenaceous
1208 towards the top. The Tons thrust passes through Laluli in Nagun Gad and is probably truncated by
1209 Tehri Nalupani fault at Chandpur in Bhagirathi valley.
1210

1211 The study revealed that there is sharp curvilinear break in morphogenetic expression of the area In
1212 North of Wazri in Jamuna valley, North of Uttarkashi in Bhagirathi valley, around Tugnath and
1213 Chamoli and South of Joshimuth in Alaknanda valley, which appears to be due to horizontal movement
1214 of a sub - tectonic plates towards south, which is causative factor in dislocation in tectonic ecology of
1215 the area, it is matter of serious concerned and needs further attention.

1216 The area genetically comprised of terraces of three domains, viz. Glacial,
1217 Fluvio-glacial and Fluvial which represent distinct environment of sedimentation of Pleistocene, late
1218 Pleistocene and Holocene time during Quaternary period.

1219 The Glacial terraces are identified at an average elevation of 1150 m above MSL, the fluvio-glacial
1220 terraces at an average elevation of 975m above MSL and fluvial terraces at an average elevation
1221 between 650 to 900 m above m.s.l, amidst these sedimentological studies of Alaknanda and Bhagirathi
1222 terraces in type area is attempted in to understand the nature of erosional and depositional processes,
1223 sedimentary pattern, behavior of transporting agencies, load characteristics, current capacity, energy
1224 condition to decipher over all history of Quaternary sedimentation in these valleys in increasing
1225 antiquity.

1226 The statistical parameters of sediment glacial terraces, Fluvio-glacial terraces and Fluvial terraces are
1227 studied down the current in Badrinath and Reshikesh section along the entire length of Alaknanda and
1228 Ganga in Garhwal Himalaya.

1229 The average mean size of sediments of terraces of glacial domain is 0.09 ϕ (Coarse sand). It varies
1230 from -2.81 ϕ to 2.8 ϕ i.e. the sediment consist of very coarse sand to fine sand. The maximum value of
1231 (MZ) is -2.81 ϕ near the origin of the river and minimum 2.8 ϕ near the outer limit of these terraces
1232 around Chamoli. The (MZ) shows sharp decrease in size in Badrinath and Vishnuprayag, section
1233 corresponding to the steep slope of river. Down the stream Vishnuprayag although it decrease in its
1234 value but display strong variation in size, which is attributed to the mixing of sediments brought by the
1235 sub-glacier joining the main Alaknanda at various points. It is a measure of sorting which reflects the
1236 consistency in the energy level of depositing medium. In respect of glacial terraces the average
1237 standard deviation is 3.34 ϕ (very poorly sorted). It varies from 2.00 ϕ to 4.30 ϕ i.e. the sediments are
1238 poorly sorted to extremely poorly sorted. The sediments of these terraces are extremely assorted and
1239 are heterogenous in nature and multi source of their derivation.. The average (SKI) for glacial
1240 sediments is 0.064 ϕ i.e. the sediments are negative skewed. It ranges from -0.450 ϕ to + 0.52 ϕ i.e.
1241 the sediment are negative skewed to positive skewed, which indicate the tendency of gradual decrease
1242 in value of (SKI) in upstream direction as result of retreat of glacier and decrease in the transport
1243 capacity. The 56% of the sample shows the negative value and 44% positive value. The sediments are
1244 very positive skewed to very negative skewed which seems to be due to mixing of sediments brought
1245 by various glaciers. The sediments down the stream of Badrinath have the strong tendency to be
1246 positive skewed.. The value of (KG) in the sediments of glacier terraces is highly variable. The average
1247 value is 0.716 ϕ (platykurtic); whereas it varies from 0.49 ϕ to 1.10 ϕ (very platykurtic to leptokurtic).

1248 The average mean size of sediments of fluvio-glacial domain is 2.15 ϕ (medium sand). It varies from -
1249 2.53 ϕ to 3.12 ϕ i.e. the sediments consist of very coarse to very fine sand. The size distribution of
1250 these deposits in the study area is extremely irregular and erratic. The sediments near the outer edge of
1251 glacial deposit downstream of Chamoli ranges 0.75 ϕ to 0.50 ϕ i.e. coarse to very coarse sand which
1252 constantly show decrease in (MZ) along the stretch of 30 km up to Karanprayag. The sediments
1253 between Chamoli and Karanprayag along 45 km show range of order of 1.25 ϕ to 2.75 ϕ with local
1254 variation. The sudden rise in (MZ) is noticed around Nandaprayag and downstream and downstream of
1255 confluence of Alaknanda and Nandakini (MZ) values are of order of 0.50 ϕ to 0.15 ϕ , which indicates
1256 the intensive mixing of sediments brought from the flash stream resulting from the retreating glacier at
1257 different point in the valley..The standard deviation of fluvio-glacial sediments varies from 0.95 to 2.50
1258 ϕ i.e. the sediments are poorly sorted to very poorly sorted. The average standard deviation is 1.563 ϕ
1259 (extremely poorly sorted). Out of 50 samples 8% are moderately sorted, 10% poorly sorted and 32%
1260 are very poorly sorted. The sediments near the source area conspicuously exhibit poor sorting and show
1261 significant improvement down the stream with local variation. As a whole the sediments are poorly
1262 sorted to very poorly sorted and heterogeneous in nature.

1263 The fluvio glacial sediments show skewness ranging from -0.48 ϕ to + 0.97 ϕ i.e. the sediments are
1264 skewed very negative to skewed very positive. The average of (SKI) is 0.078 ϕ i.e., the sediments are
1265 fine skewed. Out of the total samples of these terraces 56% are skewed positively, 16, skewed positive

1266 and 22, are skewed very negative. The assemblage of variable value of (SKI) suggests the
1267 heterogeneous association of the sediments ranging from fine sand to gravel size. The (SKI) value in
1268 general increase downstream with occasional variation. It is perhaps due to repeated reworking of the
1269 sediments towards downstream side by flash stream resulting from the glacier. The average (KG) is
1270 1.316 (leptokurtic). It ranges from 0.76 ϕ to 1.52 ϕ (platykurtic to very leptokurtic) among these 75%
1271 of the sample fall in very platykurtic class 47.50 (mesokurtic) and 45, (leptokurtic). The assemblage of
1272 these different classes of kurtosis suggests the dominance of coarse sediments (Folk & Ward, 1957).
1273 Most of the samples between Chamoli and Karanprayag section along the stretch of about 45 km show
1274 the Kurtosis value ranging between 0.90- 1.20 ϕ except in the area around Nandaprayag Nagrasu,
1275 where the sedimentation is perhaps affected by lateral mixing of sediments brought by the sub-glaciers.
1276 It seems that sediments were transported and deposited in the oscillating kinetic condition. The average
1277 mean size for the sediments of terraces of fluvial of Alaknanda is 2.458 ϕ . The maximum value of
1278 (MZ) is -0.491 ϕ is noticed near Karanprayag while minimum 4.545 ϕ at Deoprayag, near the
1279 confluence of Alaknanda and Bhagirathi River. The (MZ) shows the significant consistency in its
1280 value in the first stretch of about 35 km between the Karanprayag and Nagrasu, corresponding to the
1281 flattered and gentle slope of the river bed. Down the stream of Nagrasu for about 75 km up to
1282 Deoprayag (MZ) sharply decreases perhaps due to sharp change in bed slope of Alaknanda. In this
1283 section (MZ) strongly fluctuates around Kaliyasour, Srinagar and Kirtinagar for about 25 km which
1284 correlated to the sudden convexity in the river bed due to Neotectonic activity in the vicinity of
1285 Srinagar fault/ North Almora thrust, which traverses across the Alaknanda around Kaliyasour. It seems
1286 that the mean size of fluvial sediments sharply follow the bed slope, pointing to exponential
1287 longitudinal profile, thus decrease in (MZ) in the downstream of the Alaknanda is result of decrease in
1288 both transporting capacity and velocity of the river towards the later phases of sedimentation in the
1289 valley.

1290 The average standard deviation of sediment is 0.691 ϕ (moderately sorted) and it ranges from 0.15 ϕ to
1291 1.52 ϕ i.e. the sediments are very well sorted to poorly sorted. In the upper Alaknanda, it shows
1292 consistency in value down the current except around, Karanprayag and Rudraprayag, where Nandakini
1293 Pindar and Mandakini joined Alaknanda respectively. The variation in and around these places appears
1294 to be due to mixing of sediments brought by these tributaries. The sharp improvement in sorting is
1295 noticed downstream of Rudraprayag upto Srinagar which seems to be related with the repeated
1296 reworking of sediments and slope element. The sudden decline in sorting co-efficient in the stretch of
1297 about 15 km between Srinagar and Kirtinagar appears to be due to either the non-transport of larger
1298 grain down current or due to loss of bed slope of Alaknanda in this segment of valley. The significant
1299 increase in sorting in down current of Kirtinagar indicates cyclic reworking of sediments appears due to
1300 re-activation of channel in this part of valley. The average (SKI) of fluvial sediments is 0.00281 ϕ It
1301 ranges from -0.99 ϕ to 0.99 ϕ i.e. the sediments are coarse to fine skewed. The (SKI) exhibits tendency
1302 of gradual increase in value downstream with local variation. This suggests relative increase of fine
1303 grains, down the stream. The sediment upstream of Karanprayag is negative skewed perhaps due to
1304 mixing of sediments of fluvio glacial origin. The kurtosis of fluvial sediments of Alaknanda is highly
1305 variable; it ranges from 0.72 ϕ to 1.72 ϕ (leptokurtic) and an average value 1.264 ϕ (platykurtic to
1306 very (leptokurtic). The average values suggest the fluctuation in the energy condition of the channel
1307 system. The mean value of kurtosis revealed the more intensive sorting of central part of size
1308 distribution curve than the tails. Along the course of Alaknanda in Nadaprayag and Deoprayag section
1309 for a distance of about 75 km except local variation around Karanprayag, Rudraprayag, there is strong
1310 tendency in increase of kurtosis value downstream.

1311 The longitudinal profile of Alaknanda River and its terraces of various domain is overall concave,
1312 smooth and gentle, except in the area between Karanprayag, Dharkot, Rudraprayag and Kaliasour,
1313 where it is slightly upward convex. The gradient of the river between Chamoli and Karanprayag is
1314 1:6.6, between Karanprayag to Rudraprayag 1:2.25, and between Rudraprayag and Srinagar is 1:1. The
1315 average gradient of terraces AT₁, to AT₆, between Karanprayag and Dharkot and Dharkot and
1316 Kaliasour is 1:2.29, 1:1.66, 1:1.87, 1:2.20 and 1:1.25, respectively. The upward convexity in the area as
1317 mentioned above indicates some differential up warping of some of the terrace blocks possibly due to
1318 some movements along the Srinagar-Tehri Fault/Alaknanda Fault (Sinha & Khan .1975, Khan
1319 1981). The profile of terraces of glacial domain is restricted up Chamoli and is of hanging in nature ,
1320 whereas of fluvio-glacial terraces mostly confined to the upstream of Karanprayag, it pinches out
1321 downstream against the terraces of fluvial domain, thereby indicating an intensive down cutting of the
1322 valley floor by Alaknanda through cyclic rejuvenation consequent to recession of in post Pleistocene
1323 time. The profile of glacial terraces demonstrate intensive dissection of terraces and isolated pockets

1324 and lenses of occurrence at higher level in the valley and profile is of hanging in nature, whereas
 1325 of fluvio- glacial is of suspended in nature and suddenly abuts up stream against the profile of glacial
 1326 terraces and downstream against the profile of fluvial terrace. Down the current profile of fluvial
 1327 terraces display consistency and smoothness down the current. The profile of fluvio-glacial terraces
 1328 represent transitional phase of sedimentation and major in Quaternary time the valley.

1329

1330 The longitudinal profile of Bhagirathi is over all concave except in the upper reaches between Seansu
 1331 and Bhatwari, where the upward convexity in the alternate segment is very conspicuous. The upward
 1332 convexity and development of nick points, corresponds to major tectonic elements, Main Central
 1333 Thrust (Khan et.al. 1982 1988). Srinagar Nalupani fault and Tons thrust in the area. The gradient of the
 1334 channel bed in upper Bhagirathi in Bhatwari Uttarkashi section is 1:1.98, in Uttarkashi Dharasu section
 1335 1:1.75, in Dharasu-Tehri 1:1.55 and Tehri and Devprayag 1:1.20. The profile of domain of glacial
 1336 terraces extends up to Gagnani and Fluvio-glacial terraces up to Nakuri and down the current profile
 1337 Fluvial terraces is gentle and smooth and in conformity of river bed. The profile of glacial terraces is of
 1338 hanging in nature, whereas of fluvio- glacial is of suspended in nature and fluvial display consistency
 1339 and regularity down the current. The profile of fluvio-glacial terraces represent transitional phase of
 1340 sedimentation in the valley.

1341 The study of geology, geomorphology, Quaternary terraces and landscape profile section revealed
 1342 that there is sharp curvilinear break in morphogenetic expression of the area, North of Wazri in Jamuna
 1343 valley, North of Uttarkashi in Bhagirathi valley, around Tugnath and Chamoli and South of Joshimuth
 1344 in Alaknanda valley, which is significant element, appears to be due to horizontal movement of a sub -
 1345 tectonic plates towards south., It may be causative factor in dislocation in tectonic ecology of the
 1346 area, related to recent micro shocks in Joshimuth in Niti and Hellong area, mass failure of landscape
 1347 profile, landslide and mass wasting activities and other natural hazards; it is matter of serious
 1348 concerned and needs further attention

1349 The study of Statistical parameters and their correlation with various Thrust, fault, lineament and
 1350 longitudinal profile of Alaknanda and its tributaries, revealed that there is strong impact and influence
 1351 of tectonic and Neotectonic activity on Quaternary sedimentation in the area..


1352 **References:-**

- 1353
- 1354
- 1355 1. Dubey, U.S. (1972) Progress Report on Geomorphological study of river terraces of Alaknanda ,
- 1356 Bhagirathi, and Ganga valley, Garhwal Himalaya U.P.(Unpublished G.S.I Report 1972)
- 1357 2.Dubey, U.S.(1974) Progress Report on Geomorphological studies of river terraces of Alaknanda and
- 1358 Bhagirath valley system Garhwal Himalaya U.P. .(Unpublished G.S.I Report 1972)
- 1359 3. Dubey, U.S (1974a) Report on Geomorphological and sedimentological studies of the river terraces of
- 1360 Alaknanda , Bhagirathi, Ganga valley , Garhwal Himalaya U.P.(Unpublished G.S.I Report)
- 1361
- 1362 4.Padhi, R.N and Sharan, R.B (1972), A study of River Terraces of Alaknanda Valley Rec.Geol.surv
- 1363 India V 104pt 1 pp 39
- 1364
- 1365 5.Shukla , H.N Khan, A.A. and Dubey, U.S (1973) Report on Geomorphological studies of the River
- 1366 Terraces of , Bhagirathi and other Pleistocene landform .(Unpublished G.S.I Report).
- 1367
- 1368 6.Sinha, K.K. and Khan, A.A.(1976) A Report on Geomorphological studies of parts of Garhwal
- 1369 Himalaya , U.P. (based on Photo Interpretation with selected field Checks) 11 Geomorphology Course
- 1370 I.P.I. Dehra Dun).
- 1371
- 1372 7. Khan, A.A (1981) Geology and Geomorphology of Eastern part of Ganga Complex in parts of Tehri,
- 1373 Chamoli, and Pauri District Uttar Pradesh . (Unpublished G.S.I Report FS 1976-77)
- 1374
- 1375 8. Khan, A.A.(1974) Report on Geomorphological studies of river terraces and other Pleistocene
- 1376 landform of Alaknanda and Pindar Valley . (Unpublished G.S.I Report)


- 1377
1378 9. Khan , A.A.(1974)Report on Geomorphological studies of river terraces and other Pleistocene
1379 landform of Alaknanda and Pindar Valley . .(Unpublished G.S.I Report)
- 1380 10. Khan, A.A and Balchandran, V (1974-75) Record Vol 109 of Geological Survey of India, Part I PP
1381 59
- 1382 11. Khan, A.A. (19 75) Report on Geomorphological studies of river terraces and other Pleistocene
1383 land forms in Alaknanda, Pindar and Attagad valley . (Unpublished G.S.I Report of FS 1973-74)
- 1384 12. Khan, A.A (1981) Geology and Geomorphology of Eastern part of Ganga Complex in parts of
1385 Tehri,Chamoli, and pauri District Uttar Pradesh . (Unpublished G.S.I Report FS 1976-77)
- 1386 13. Khan, A.A and Dubey, U.S (1981) Evaluation of Quaternary Landscape of intermontane Jamuna
1387 valley in Dheradun U.P;) and Sirmur (H. P). Rec.Geol.Surv.Indi; Vol112, pt 3 pp 125- 135.
- 1388 14. Khan, A.A. and Nawani , P.C (1981.) Quaternary geological and geomorphological Evaluation of
1389 Eastern part of Ganga Complex, Garhwal Himalaya U.P India (Abstract) IV Regional Conference
1390 Geol. SE Asia Manila Philippines
- 1391 15. Khan, A.A and Dubey, U.S (1985) An attempt to study the sedimentary structures and paleo current
1392 of Quaternary terraces of Alaknanda valley of Garhwal Himalaya U.P (Accepted for publication in
1393 GSI Record of 1985).
- 1394 16. Khan.A.A (.1987) Geomorphic Evaluation and Quaternary Sedimentation in upper Ganga Basin
1395 Garhwal Himalaya (U.P.). Ph.D. Unpublished thesis submitted to Vikram University Ujjain M.P. India
1396
- 1397 17 . Khan.A.A (.2018) Quaternary deposits of Narmada valley in relation to Geomorphology
1398 and Neotectonic of the area with special reference to Hominid occurrence D.sc. Unpublished
1399 thesis submitted to Barkatullaha University Bhopal, M.P. India.
1400

KEY MAP OF ALAKNANDA AND ITS TRIBUTARIES. GARHWAL HIMALAYA U. P.

Plate No. 1



- 1401
- 1402
- 1403
- 1404
- 1405
- 1406



1407

1408

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418


1419

1420

1421

1422

1423
1424
1425
1426
1427



1428







PLATE NO. 8

1441



1442

1443

1444

1445

1446




DIAGRAM SHOWING LONGITUDINAL RIVER PROFILE AND VARIATION IN STATISTICAL PARAMETERS IN
QUATERNARY TERRACES OF FLUVIAL DOMAIN DOWN THE STREAM - BAL-GANGA VALLEY,
GARHWAL HIMALAYA, (U.P.)



1453


1454

1455

1456

1457

1458



1459

1460

1461

1462

1463

1464

