

Beyond field success: insights into functional and numerical response of the predatory mite *Typhlodromalus aripo* and the role of cassava exudate in predatory performance

by Jana Publication & Research

Submission date: 23-Aug-2025 12:00PM (UTC+0700)

Submission ID: 2690326558

File name: IJAR-53486.docx (413.98K)

Word count: 5974

Character count: 33151

Beyond field success: insights into ¹¹functional and numerical response of the
predatory mite *Typhlodromalus aripo* and the role of cassava exudate in
predatory performance

Abstract

The predatory mite *Typhlodromalus aripo* is a key natural enemy of cassava green mite (*Mononychellus tanajoa*) on cassava in Sub-Saharan Africa, yet its functional and numerical responses remain poorly understood. We evaluated, under controlled laboratory conditions (25 °C, 70–90% RH, 12L:12D photoperiod), the functional and numerical responses of *T. aripo* to four prey stages (eggs, larvae, protonymphs, deutonymphs) at four densities (5–60), with and without cassava exudate. *T. aripo* exhibited a Type II functional response across all prey stages, with highest attack rates on larvae and protonymphs, and shortest handling times on eggs and larvae. Predation rates were consistently higher on early prey stages, but cassava exudate reduced attack rates and increased handling times, except on larvae where both parameters increased. Oviposition and efficiency of conversion of ingested food (ECI) rose with prey stage, peaking on deutonymphs, and exudate enhanced reproduction and ECI particularly on early stages. These results demonstrate the effectiveness of *T. aripo* as biological control agent against cassava green mite, capable of exploiting a range of prey stages. Cassava exudate, while influencing predatory behavior, can enhance reproductive performance, thereby demonstrating the predator's long-term persistence in cassava fields. The functional and numerical response data presented here provide valuable reference information for selecting and developing new predatory mite species for use as biological control agents in similar agroecological contexts.

Keys words: Predation, oviposition, efficiency of conversion, prey stage, biological control, cassava

Introduction

The functional and numerical responses are among the most studied attributes of the predator-prey relationships in the context of biological control of arthropod pests (i.e., Dalir *et al.*, 2020; Fathipour *et al.*, 2017). Functional response describes how the rate of prey consumption by a predator changes with prey density, while numerical response refers to changes in predator reproduction based on prey availability. These metrics are fundamental for predicting the impact of a predator on pest populations and are increasingly used as trait-based tools for selecting and comparing candidate biological control agents. Holling (1959, 1961) described three types of functional responses that characterize how a predator's rate of prey consumption changes with prey density. A type I describes a linear increase in prey consumption with increasing prey density, until a maximum is reached (due to predator satiation), a type II describes a decelerating curve – prey consumption increases with prey density but at a decreasing rate due to handling time, and a Type III describing a sigmoidal curve – slow increase at low prey densities, rapid increase at intermediate densities, and saturation at high densities.

Predatory mites of the family Phytoseiidae have been by far the most extensively studied around the world, including Africa, because of their potential as biological control agents of important pests such as spider mites, thrips, and whiteflies in agricultural crops (Gerson *et al.* 2003; McMurtry *et al.* 2013). They have been used in biological control programs with two success stories; the classical biological control of the citrus and avocado mites by *Euseius stipulatus* (Athias-Henriot) in California and the classical biological control of the cassava green mite by *Typhlodromalus aripo* (Deleon) in Africa (Yaninek and Hanna, 2003; McMurty *et al.*, 2015). In the latter case – biological control of the cassava green mite *Monomychellus tanajoa* (Bondar) (Acari: Tetranychidae) in Africa – several species of predatory mites were introduced from Brazil, origin of the pest, for release in Africa (Yaninek and Hanna, 2003). The predator *T. aripo* was the most successfully established predator (along with followed by *T. manihoti*) due to its high dispersal rate and its ability to persist in cassava fields, hence effectively controlling the pest *M. Tanajoa* (Yaninek and Hanna, 2003). A series of ecological studies were conducted prior and post-release of *T. aripo* in cassava fields for a better understanding of the host-predator and predator-prey interactions, as well as predator-predator interactions (i.e., Gnanvossou *et al.*, 2003; Onzo *et al.*, 2003; Hanna and Onzo, 2009; Onzo *et al.*, 2012; Onzo *et al.*, 2013; Onzo *et al.*, 2014;). However, despite the several ecological studies, a good understand of the functional response of *T. aripo* is still missing. Therefore, it is essential to provide data on the functional response of this effective biocontrol agent, which can be important when developing a new species for use as a biological control agent in a similar context.

Moreover, functional response is not static, but rather influenced by biotic factors, including the predator's age (Fathipour *et al.*, 2017; Dalir *et al.*, 2020), the prey's developmental stage (Dalir *et al.*, 2020), and the

availability of alternative non-prey food sources (Fathipour et al. 2020). The availability of alternative plant-based foods – such as pollen, plant exudates – can modulate predator predation and enhance the biological control of target pest (McMurtry and Scriven, 1966; Nomikou et al, 2010; Konstantinos Samaras, 2021). In the context of cassava agroecosystems, cassava plants produce large quantities of exudates, which have been identified as important alternative food sources for predatory mites, allowing their survival during periods of prey scarcity (Klein, 1990; Bakker and Klein, 1990; Loko et al., 1994; Gnanvossou et al., 2005). These findings suggest that cassava exudates may play a critical role in modulating the predator-prey dynamics between *T. aripo* and *M. tanajoa*. Therefore, evaluating the influence of prey stage and cassava exudate on the functional and numerical responses of *T. aripo* provides a better insight into its effectiveness as a robust and adaptable biological control agent in sub-Saharan Africa cassava agroecosystems.

This study was conducted to provide a comprehensive assessment of the functional and numerical responses of *T. aripo* to its prey, *M. tanajoa*, under controlled laboratory conditions, and evaluate the influence of additional exudates on predatory performance.

Material and methods

Mite sources and rearing

The predators (*T. aripo*) used in this study were obtained from a laboratory colony originally established from individuals collected in July 1998 from cassava fields near Sè, Cotonou, Benin (Gnanvossou et al. (2003). The colony has been maintained under controlled environmental conditions: $25 \pm 1^\circ\text{C}$, 70–90% relative humidity, and a 12:12 h light–dark photoperiod. The rearing method followed the procedure described by McMurtry and Scriven (1965) and Friese et al. (1987), with minor adaptations. Predators were maintained on black PVC tiles (4 x 4 cm) placed atop a 1 cm-thick foam pad of the same dimensions inside a 14.5 cm diameter Petri dish filled with water. To prevent mite escape and contamination, absorbent paper bands were arranged along the edges of the tile, with one edge immersed in water to serve as a continuous moisture source. An oviposition site was prepared using a square of hydrophobic cotton wool, lightly stretched to form an open central platform, and covered with a transparent plastic sheet of matching shape. Fresh cassava leaves infested with all developmental stages of *M. tanajoa* were provided as a food source. These leaves were collected from greenhouse-grown cassava plants reared in pots under fine-mesh cages. For the experiments, specific developmental stages of *M. tanajoa* were obtained from colonies maintained in a screenhouse on 2- to 3-week-old potted cassava plants. These colonies were initiated with cohorts of eggs laid by *M. tanajoa* females collected from cassava fields in southwestern Benin.

Functional response bioassay

To initiate the bioassay, a cohort of 1-day-old *T. aripo* eggs was obtained by placing 100 mated females from the mother colony onto 2.5 cm diameter black PVC disks and allowing them to oviposit for 24 hours. The eggs laid were then carefully transferred to new rearing units and reared to adulthood, following the procedure described by Mégevand et al. (1993), with *Tetranychus urticae* provided as food during development. Upon emergence, adult mites were sexed, and females were isolated individually for experimentation. Each experimental unit consisted of a 2.5 cm diameter cassava leaf disc, placed on a layer of solidified agar-agar (1% solution) at the bottom of a Petri dish of similar dimensions. To allow adequate ventilation, two 2 mm-diameter holes were made on opposite sides of each dish and covered with fine nylon mesh. All experiments were conducted under controlled conditions: $25 \pm 1^\circ\text{C}$, 70–90% relative humidity, and a 12:12 h light:dark photoperiod. To assess the functional and numerical responses of *T. aripo* to different prey stages, newly emerged adult females were transferred singly to the prepared experimental units and offered one of the four immature stages (egg, larvae, protonymph, and deutonymph) of *M. tanajoa* at densities of 5, 15, 30, and 60 individuals per unit. Only immature stages of *M. tanajoa* were used, as these are typically the stages most targeted by phytoseiid mites (Reis et al., 2000; Forero et al., 2008; Marafeli et al., 2011). Each prey stage and density combination was tested in both the presence and absence of cassava exudate, which was provided *ad libitum* within each unit. A total of 20 replicates (females) were tested per prey stage and density combination. Prey consumption and oviposition was recorded daily over a 7-day period, with both prey and cassava exudate replenished every 24 hours.

Data analysis

Data on functional response were analyzed in two steps (Juliano, 2001). First, a logistic regression Model of the proportion of prey consumed (N_e/N_0) as a function of prey density (N_0) was used to determine the type of the functional response of the predator to individual immature stage (equation 1):

$$N_e / N_0 = \frac{\exp(P_0 + P_1 N_0 + P_2 N_0^2 + P_3 N_0^3)}{1 + \exp(P_0 + P_1 N_0 + P_2 N_0^2 + P_3 N_0^3)} \quad (\text{Equation 1})$$

where N_0 is the initial density of prey offered, N_e is the number eaten, P_0 , P_1 , P_2 , and P_3 are the intercept, linear, quadratic, and cubic coefficients, respectively, estimated using the maximum likelihood method. The type of the functional response curve is determined based on the sign of P_1 and P_2 from equation 1 (Juliano, 2001): if $P_1 < 0$, it implies Type II functional response. If $P_1 > 0$ and $P_2 < 0$, it is Type III functional response. After the shape of functional response was determined (Type II), the next step was to

estimated the handling time (T_h) and the attack rate (a) of parameters using the random predator equation (equation 2; Roger, 1972):

$$N_e = N_0 \{1 - \exp[a(T_h N_e - T)]\} \quad (\text{Equation 2})$$

where T is the total time available for the predator for search during the experiment (24 h), a is the attack rate (searching efficiency), T_h is the handling time of prey by the predator, and N_e and N_0 as described above. A nonlinear least-squares regression (PROC NLIN, SAS Institute, 2004) was used to fit the random predator equation to data and to estimate the parameters describing the type II response.

Numerical response

To assess the numerical response of *T. aripo*, the number of eggs laid by the predator was recorded concurrently with the number of prey individuals consumed, using the same experimental units over a 7-day observation period. The predator's daily oviposition rate was then regressed against the initial prey densities, following the approach of Fathipour and Maleknia (2016). This regression analysis quantified the relationship between prey density and predator fecundity, offering insights into the predator's reproductive potential in response to increasing prey availability.

The efficiency of conversion of ingested food (ECI) into egg biomass was calculated using the equation of Omkar and Pervez (2004) (Equation 3):

$$ECI = \frac{\text{Number of eggs laid}}{\text{Number of prey consumed}} \times 100 \quad (\text{Equation 3})$$

To evaluate the effects of prey density, prey developmental stage, and the presence of cassava exudate on predation, oviposition, and the efficiency of conversion of ingested food (ECI), data were analyzed using analysis of variance (ANOVA), and Tukey's Honestly Significant Difference (HSD) test was used for multiple comparisons at the 5% significance level ($\alpha = 0.05$).

Results

Functional response

Analysis of variance revealed that prey developmental stage ($F_{3,1213} = 515.5$, $P < 0.0001$), prey density ($F_{3,5665} = 2406.9$, $P < 0.0001$), and the presence of additional cassava exudate ($F_{1,820.9} = 348.7$, $P < 0.0001$) had significant effects on the predation activity of *T. aripo* (Table 1, Table 2). Across all developmental stages, predation rates increased with prey density. However, *T. aripo* consistently showed higher predation on earlier stages (eggs and larvae) compared to later ones (protonymphs and

deutonymphs). Mean prey consumption ranged from 4.07 to 24.41 on eggs, 5.00 to 26.29 on larvae, 4.95 to 15.12 on protonymphs, and 4.57 to 14.15 on deutonymphs.

In general, the presence of cassava exudate significantly reduced predation across all prey stages and densities, although some exceptions were observed. The largest reduction (59%) was recorded on egg predation at a density of 15, while the smallest reduction (1.06%) occurred on deutonymphs at a density of 60 (Table 1).

The negative and significant linear coefficient from Equation 1 ($P < 0.05$) confirms that *T. aripo* exhibited a Type II functional response across all prey developmental stages (Table 3; Figure 1). The estimated attack rate (a) and handling time (Th) are summarized in Table 4. The lowest attack rates were recorded on eggs and deutonymphs (0.0910 and 0.0894, respectively), while the highest values were observed on larvae and protonymphs (0.9301 and 0.7951, respectively). In contrast, handling time was about 2-fold longer for predators preying on protonymphs or deutonymphs than for those feeding on eggs or larvae (Table 1).

The presence of exudate markedly influenced *T. aripo*'s predatory performance. On eggs, the attack rate decreased by six-fold and the handling time increased four-fold. On deutonymphs, the attack rate was reduced by 1.4-fold, accompanied by an approximately one-fold increase in handling time. For protonymphs, exudate reduced the attack rate by 1.4-fold and increased the handling time by 1.1-fold. In contrast, when feeding on larvae, both parameters increased in the presence of exudate, with attack rate rising by 1.6-fold and handling time by 1.3-fold.

Numerical response and food conversion efficiency

Analysis revealed that prey developmental stage, prey density, and the presence of cassava exudate all had significant effects on the oviposition of *T. aripo*, as did most of their interactions, except for the interaction between exudate and prey density, which was not significant ($P < 0.05$; Tables 1 & 2).

Oviposition increased progressively with prey developmental stage, with the lowest number of eggs laid on the egg stage (ranging from 0 eggs/female at density 5 to 0.69 eggs/female at density 60), and the highest on deutonymphs (from 0.62 eggs/female at density 5 to 1.71 eggs/female at density 30). With respect to prey density, oviposition exhibited a curvilinear pattern on early stages (eggs and larvae), increasing with density (Figure 2). On later stages (protonymphs and deutonymphs), oviposition peaked at intermediate density (30) and declined at the highest density (60).

Regression analysis showed significant and moderate correlations between prey density and oviposition ($P < 0.05$, $0.5 < R^2 < 0.75$), except for the deutonymph stage, where a weaker but significant relationship was found ($R^2 = 0.413$).

³⁷ The efficiency of conversion of ingested food (ECI) by *T. aripo* increased with the prey's developmental stage across all densities. Overall, ECI was lowest for females fed on eggs and highest for those fed on deutonymphs. For early stages (eggs, larvae, and protonymphs), ECI tended to increase with prey density, whereas for deutonymphs, it generally declined as density increased. (Table 1).

The presence of cassava exudate resulted in a statistically significant but modest increase in oviposition, and a substantial enhancement of ECI, particularly when *T. aripo* ⁶³ fed on eggs, larvae, or protonymphs. The largest increase in oviposition (71%) was observed on eggs at density 60, while the greatest ECI enhancement (63%) occurred on larvae. In contrast, no significant effect of exudate was detected on either oviposition or ECI when predators fed on deutonymphs (Table 1).

Discussion

This study addresses a critical knowledge gap regarding ³⁹ the functional and numerical responses of the predatory mite *Typhlodromalus aripo*, a key ³⁶ biological control agent against the cassava green mite in sub-Saharan Africa. Beyond quantifying predation and reproductive performance, our findings underscore the influence of prey developmental stage and plant-associated foods such as cassava exudate, in shaping both the predator's functional and numeric responses. These insights contribute ⁶⁵ to a deeper understanding of predator-prey interactions and offer valuable implications for predictive evaluation of any predatory mite ¹⁶ for use as a biological control agent in sub-Saharan Africa (Yaninek and Hanna, 2003). Overall, our findings underscore the complex interplay between prey stage, prey density, and plant-derived foods in shaping both ¹⁰ the functional and numerical responses of *T. aripo*.

Our results demonstrate that the predatory mite *T. aripo* exhibits ¹³ a Type II functional response when feeding on all immature stages of its prey, consistent with patterns reported for many phytoseiid mites (Piyani ⁴¹ *et al.*, 2021; Fathipur *et al.*, 2017; Fathipur *et al.*, 2020; Carrillo and Pena, 2011; Park *et al.*, 2021). This functional response type, characterized by a decelerating prey consumption rate at higher prey densities, suggests that handling time imposes a limit on predation, as confirmed by our parameter estimates. The longer handling times recorded for protonymphs and deutonymphs compared with eggs and larvae likely reflect increased prey size, mobility and defensive behaviors in later developmental stages, which can slow down prey capture and processing (Sabelis, 1981). Moreover, Sabelis (1990) demonstrated that, rather than being constrained solely by time-budget rules, functional response parameters are largely determined by predator satiation, which depends on prey consumption and gut clearance.

Predatory mites are categorized into four lifestyle types, with most generalist predators classified as type III. These predators ¹⁴ feed and reproduce on a wide variety of prey, but they ¹ can also thrive on alternative food sources such as pollen, plant exudates, and honeydew – either as survival food in the absence of prey

or as a complementary food source that boosts reproduction when prey is present (McMurtry *et al.*, 2013). Previous studies have shown that type III generalists may prefer *T. urticae* larvae over eggs, or exhibit no clear preference between the two (Blackwood *et al.*, 2001). For instance, *Amblyseius eharai*, a type III generalist, was reported to favor *T. urticae* larvae over other stages (Park *et al.*, 2021). Similarly, *Neoseiulus californicus* consumed more larvae, whereas *N. cucumeris* consumed more eggs (Zhang *et al.*, 2017). Similarly, in our study, *T. aripo*, also classified a type III generalist predator (McMurtry *et al.*, 2013), displayed higher predation rates on early developmental stages (eggs and larvae) than on later stages (protonymphs and deutonymphs), with larvae being the most consumed stage. Strikingly, all predator females (100%) consumed all available larvae at prey densities of 5 and 15, and 82% at higher densities, demonstrating a strong preference for the larval stage of its prey. This pattern is likely due to differences in prey size, mobility, and defensive capacity. Eggs and larvae are immobile and have softer integuments, facilitating capture and consumption, whereas protonymphs and deutonymphs are more mobile, better defended, and require longer handling times. The approximately two-fold longer handling times recorded on protonymphs and deutonymphs compared to eggs and larvae support this interpretation, and are consistent with earlier observations for other phytoseiids preying on spider mite nymphs or adults (Sabelis, 1986).

Numerical response patterns further revealed that *T. aripo*'s reproductive output is strongly influenced by prey stage. The highest oviposition rates on deutonymphs, despite their lower consumption rates compared with early stages, suggest that this prey stage provides the greatest nutritional benefit per individual consumed. This is supported by the observed trends in efficiency of conversion of ingested food (ECI), already peaking on deutonymphs at the lowest density. Similar stage-specific effects on predator fecundity have been reported for other phytoseiids, where later prey stages supply more biomass and higher-quality nutrients, promoting egg production (Fathipour and Maleknia, 2016; Fathipour *et al.*, 2022).

The presence of cassava exudate had stage-specific effects on the predatory performance of *T. aripo*. On eggs, for instance, it reduced the attack rate six-fold and increased handling time four-fold. Similar patterns have been reported for *Amblyseius swirskii* feeding on spider mite eggs, where the addition of pollen reduced the attack rate 13-fold and increased handling time 1.4-fold (Fathipour *et al.*, 2017). Moreover, while exudate generally reduced predation rates, it modestly increased oviposition and substantially enhanced ECI for early prey stages, consistent with earlier observations for other phytoseiids (Nomikou *et al.*, 2010; Fathipour *et al.*, 2020). This result might indicate that in the presence of exudate, *T. aripo* invests more efficiently in reproduction per prey consumed, by allocating energy toward egg production rather than increased foraging activity. For deutonymphs, however, exudate had no significant effect on either oviposition or ECI, suggesting that the large size and nutritional content of this later

developmental stage buffer the predator's reproductive performance against foraging interference from exudate.

Conclusion

The predatory mite *T. aripo* exhibits a type II functional response against all immature stages of its prey *M. tanajoa*, demonstrating strong potential as an effective biological control agent against cassava green mite, given its high attack rates, preference for vulnerable prey stages, and capacity to convert prey consumption into reproductive output. While cassava exudate can reduce predation rate in certain contexts, it also enhances oviposition and food conversion efficiency on early prey stages, suggesting it plays an important role in sustaining predator populations in period of prey scarcity. This dual effect highlights the importance of integrating plant traits, such as exudate production, into biological control strategies to optimize predator performance and ensure long-term suppression of pest populations in cassava systems.

Acknowledgments. We thank R. Houndafoche, P. Sovimi, for their assistance in maintaining the predator colonies and conducting experiments. This research was supported by the International Institute of Tropical Agriculture (IITA).

References

1. Bakker, F., Klein, M. (1990). The significance of cassava exudate for predacious mites. *Symp. Biol. Hung.* 39: 437-439
2. Blackwood, J.S., Schausberger, P., Croft, B.A. (2001). Prey-stage preference in generalist and specialist phytoseiid mites (Acari: Phytoseiidae) when offered *Tetranychus urticae* (Acari: Tetranychidae) eggs and larvae. *Environmental Entomology* 30: 1103-1111.
3. Carrillo, D., Pena, J.A. (2011). Prey-stage preferences and functional and numerical responses of *Amblyseius largoensis* (Acari: Phytoseiidae) to *Raoiella indica* (Acari: Tenuipalpidae). *Experimental and Applied Acarology* 55(3-4): 361-372, DOI 10.1007/s10493-011-9488-7
4. Cohen, J. (1988). Statistical power analysis for the behavioral sciences (2nd Ed.). New York: Routledge.
5. Dalir, S., Hajiqaibar, H., Fathipour, Y., Khanamani, M. (2020). Age-Dependent Functional and Numerical Responses of *Neoseiulus cucumeris* (Acari: Phytoseiidae) on Two-Spotted Spider Mite (Acari: Tetranychidae). *Journal of Economic Entomology*, 114(1), 50-61. <https://doi.org/10.1093/jee/toaa266>
6. Ding-Xu, L., Juan, T., Zuo-Rui, S. (2007). Functional response of the predator *Scolothrips takahashii* to hawthorn spider mite, *Tetranychus viennensis*: effect of age and temperature. *BioControl* 52: 41-61.
7. Fan, Y., Pettit, F. L. (1994). Functional response of *Neoseiulus barkeri* Hughes on two-spotted spider mite (Acari: Tetranychidae). *Experimental and Applied Acarology* 18: 613-621.
8. Farazmand, A., Fathipour, Y., Kamali, K. (2012). Functional response and mutual interference of *Neoseiulus californicus* and *Typhlodromus bagdasarjani* (Acari: Phytoseiidae) on *Tetranychus urticae* (Acari: Tetranychidae). *Int. J. Acarol.* 38: 369-376.
9. Fathipour, Y., Maleknia, B. (2016). Mite predators, pp. 329-366. In Omkar (ed.), *Ecofriendly pest management for food security*. Elsevier, San Diego, CL.
11. Fathipour, Y., B. Maleknia, M. Soufbaf, Reddy, G.V.P. (2020). Functional and numerical responses, mutual interference and resource switching of *Amblyseius swirskii* on two-spotted spider mite. *Biological Control* 146: 104266.

12. Fathipour, Y., Karimi, M., Farazmand, A., Talebi, A.A. (2017). Age-specific functional response and predation rate of *Amblyseius swirskii* (Phytoseiidae) on two-spotted spider mite. *Systematic and Applied Acarology*, 22(2), 159–169. <https://doi.org/10.11158/saa.22.2.1>
13. Fathipour, Y., M. Karimi, A. Farazmand, Talebi, A.A. (2017). Age-specific functional response and predation rate of *Amblyseius swirskii* (Phytoseiidae) on two-spotted spider mite. *Systematic and Applied Acarology* 22: 159–169.
14. Fathipour, Y., Maleknia, B. (2016). Mite Predators. In: Omkar (Ed.), *Ecofriendly Pest Management for Food Security*. Elsevier, San Diego, USA, pp. 329-366
15. Forero, G., Rodríguez, M., Cantor, F., Rodríguez, D., Cure, J.R. (2008). Criterios para el manejo de *Tetranychus urticae* Koch (Acari: Tetranychidae) con el ácaro de predador *Amblyseius* (*Neoseiulus*) sp. (Acari: Phytoseiidae) en cultivos de rosas. *Agronomía Colombiana* 26:78-86.
16. Friesse, D.D., Mégevand, B., Yaninek, J.S. (1987). Culture maintenance and mass production of exotic phytoseiids. *Insect Science and its Application* 8: 875-878
17. Gerson, U., Smiley, R.L., Ochoa, R. (2003) Mites (Acari) for pest control. Blackwell Science Ltd., UK, 539 pp.
18. Gnanvossou, D., Hanna, R., Toko, M. (2005). Comparative life history traits of three neotropical phytoseiid mites maintained on plant-based diets. *Biological control* 35(1): 32-39
19. Gnanvossou, D., Yaninek, S., Hanna, R., Dicke, M. (2003). Effects of prey mite species on life history of the phytoseiid predators *Typhlodromalus manihoti* and *Typhlodromalus aripo*. *Experimental and Applied Acarology* 30: 265-278
19. Hanna, R., Onzo, A. (2009). Within-plant migration of the predatory mite *Typhlodromalus aripo* from the apex to the leaves of cassava: response to day–night cycle, prey location and prey density. *Journal of Insect behavior* 22: 186-195.
20. Holling, C. S. (1961). Principles of insect predation. *Annual Review of Entomology* 6, 163-182
21. Holling, C. S. 1959. Some characteristics of simple types of predation and parasitism. *Can. Entomol.* 91: 385–398.
22. Jafari, S., Fathipour, Y., Faraji, F. (2012). The influence of temperature on the functional response and prey consumption of *Neoseiulus barkeri* (Phytoseiidae) on two-spotted spider mite. *J. Entomol. Soc. Iran* 31: 39–52
23. Juliano, S. A. 2001. Nonlinear curve fitting: predation and functional response curves, pp. 178–196. In S. M. Scheiner and J. Gurevitch (eds.), *Design and analysis of ecological experiments*. Oxford University Press, New York
23. Klein, M. (1990). Is cassava exudate a factor in the persistence of *Typhlodromalus limonicus* (Garman & McGregor) in the field? M.Sc. Thesis, University of Amsterdam, Amsterdam, The Netherlands, 34 pp. Cited by Gnanvossou, D., Hanna, R., Yaninek, J.S., Toko, M., 2005. Comparative life history traits of three neotropical phytoseiid mites maintained on plant-based diets. *Biological Control* 35, 32–39.
24. Madadi, H., Enkegaard, A., Brodsgaard, H.F., Kharrazi-Pakdel, A., Mohaghegh, J. Ashouri, A. (2007). Host plant effects on the functional response of *Neoseiulus cucumeris* on onion thrips larvae. *Journal of Applied Entomology* 131: 728–733.
25. Marafeli, P.P., Reis, P.R., Silveira, E.C., Toledo, M.A., Souza-Pimentel, G.C. (2011). *Neoseiulus californicus* (McGregor, 1954) preying in different Life stages of *Tetranychus urticae* Koch, 1836 (Acari: Phytoseiidae, Tetranychidae). *Acarologia* 51: 499-506.
26. McMurtry, J.A., Scriven, G.T. (1966). Effects of pollen and prey density on the number of prey consumed by *Amblyseius hibisci* (Acarina: Phytoseiidae). *Annals of Entomological Society of America* 59: 147-149.
27. McMurtry J.A., de Moraes, G.J., Famah Sourassou, N. (2013). Revision of the lifestyles of phytoseiid mites (Acari: Phytoseiidae) and implications for biological control strategies. *Systematic & Applied Acarology* 18(4): 297–320. <http://dx.doi.org/10.11158/saa.18.4.1>
28. Megevand, B., Klay, A., Gnanvossou, D., Paraiso, G. (1993). Maintenance and mass rearing of phytoseiid predators of the cassava green mite. *Experimental and Applied Acarology* 17: 115-128.
29. Nikbin, R., Sahragard, A., Hosseini, M. (2014). Age-specific functional response of *Trichogramma brassicae* (Hymenoptera: Trichogrammatidae) parasitizing different egg densities of *Ephesia kuehniella* (Lepidoptera: Pyralidae). *Journal of Agriculture Science and Technology* 16: 1205–1216.
30. Nomikou, M., Sabelis, M.W., Arne Janssen, A. (2010). Pollen subsidies promote whitefly control through the numerical response of predatory mites. *BioControl* 55:253–260
31. Omkar, O., Pervez, A. (2004). Functional and numerical responses of *Propylea dissecta* (Col., Coccinellidae). *Journal of Applied Entomology* 128: 140–146.

32. Onzo A, Sabelis M.W., Hanna R. (2003). Interactions in an acarine predator guild: impact on *Typhlodromalus aripo* abundance and biological control of cassava green mite in Benin, West Africa. *Experimental and Applied Acarology* 31(3-4): 225-231
33. Onzo A, Sabelis M.W., Hanna R. (2014). Single versus multiple enemies and the impact on biological control of spider mites in cassava fields in West-Africa. *Experimental and Applied Acarology* 62: 293-311
34. Onzo, A., Hanna, R., Sabelis, M.W. (2012) The predatory mite *Typhlodromalus aripo* prefers green-mite induced plant odours from pubescent cassava varieties. *Experimental and Applied Acarology* 58:359-370
34. Onzo, A., Hanna, R., Toko, M. (2013). Mix-planting pubescent and glabrous cassava affects abundance of *Typhlodromalus aripo* and its prey mite *Mononychellus tanajoa*. *Journal of Applied Entomology* 138: 297-306.
35. Park, Y.-G., Lee, J.-H., Lim, U.T. (2011) Functional response of *Amblyseius eharai* Acari: Phytoseiidae) on *Tetranychus urticae* (Acari: Tetranychidae). *PLoS ONE* 16(12): e0260861.
36. Patel, K., Zhang, Z.Q. (2017). Functional and numerical responses of *Amblydromalus limonicus* and *Neoseiulus cucumeris* to eggs and first instar nymph of tomato/potato psyllid (*Bactericera cockerelli*). *Systematic and Applied Acarology* 22: 1476-1488.
37. Reis, P.R., Teodoro, A.V., Pedro Neto, M. (2000). Predatory cctivity of phytoseiid mites on developmental stages of coffee ring spot mite (Acari: Phytoseiidae:Tenuipalpidae). *Neotropical Entomology*29: 547-553.
38. Rogers. D. (1972). Random search and insect population models. *Journal of Animal Ecology* 41(2): 369-383.
39. Sabelis, M.W. (1986). The functional response of predatory mites to the density of two-spotted spider mites. In: Dynamics of Physiologically Structured Populations Metz JAJ, Diekmann O (eds) Lecture Notes in Biomathematics 68, Springer, Berlin Heidelberg New York pp 298-321
40. Sabelis, M.W. (1990). How to analyze prey preference when prey densities varies? A new method to discriminate between effects of gut fullness and prey type composition. *Oecologia* 82: 289-298.
41. Samaras, K., Pappas, M.L., Pekas, A. (2022). Benefits of a balanced diet? Mixing prey with pollen is advantageous for the phytoseiid predator *Amblydromalus limonicus*. *Biological control* 155, 104531
42. Toko, M., O'Neil, R.J., Yaninek, J. S. (1994). Effects of cassava exudate and prey densities on the survival and the reproduction of *Typhlodromalus limonicus* (Acari: Phytoseiidae), a predator of cassava green mite *Mononychellus tanajoa* (Acari: Tetranychidae). *Experimental and Applied Acarology* 18: 221-231
43. Yaninek S., Hanna, R. (2003). Cassava green mite in Africa - a unique example of successful classical
44. Zheng, Y., De Clercq, P., Song, Z.-W., Li, D., Zhang, B.-X. (2017). Functional response of two *Neoseiulus* species preying on *Tetranychus urticae* Koch. *Systematic & Applied Acarology* 22(7): 1059-1068

Figure legend

Figure 1: Functional response of *Typhlodromalus aripo* to eggs, larvae, protonymphs, and deutonymphs of its prey *Mononychellus tanajoa*, in the absence or presence of cassava foliar exudate

Figure 2: Numerical response of *Typhlodromalus aripo* to eggs, larvae, protonymphs, and deutonymphs of its prey *Mononychellus tanajoa*, in the absence or presence of cassava foliar exudate

Table 1: Mean ($\pm$ SE) daily consumption, oviposition, and efficiency of conversion onto egg biomass of the predatory mite *Typhlodromalus aripo* at different densities of *Mononychellus tanajoa* immature stages in the presence or absence of additional exudate

Density	Prey Stage	Mean consumption / day		Mean oviposition / day		Mean ECI (%)	
		Exudate absent	Exudate present	Exudate absent	Exudate present	Exudate absent	Exudate present
5	Egg	4.07 $\pm$ 0.17 cA	3.67 $\pm$ 0.16 cA	0 dA	0.08 $\pm$ 0.02 bB	0 cB	2.34 $\pm$ 0.60 bA
	Larva	5.00 $\pm$ 0.00 aA	4.81 $\pm$ 0.05 aA	0.05 $\pm$ 0.01 cB	0.63 $\pm$ 0.05 bA	1.05 $\pm$ 0.39 cB	13.10 $\pm$ 1.01 aA
	Protonymph	4.95 $\pm$ 0.02 aA	4.80 $\pm$ 0.03 aB	0.30 $\pm$ 0.03 bB	0.48 $\pm$ 0.05 bA	6.20 $\pm$ 0.62 bB	10.10 $\pm$ 0.93 aA
	Deutonymph	4.57 $\pm$ 0.11 bA	4.12 $\pm$ 0.09 bB	0.62 $\pm$ 0.05 aA	0.62 $\pm$ 0.08 aA	13.82 $\pm$ 1.32 aA	14.86 $\pm$ 2.30 aA
15	Egg	9.46 $\pm$ 0.59 bA	6.87 $\pm$ 0.41 cB	0.02 $\pm$ 0.01 dA	0.16 $\pm$ 0.04 cB	0.14 $\pm$ 0.09 dB	2.03 $\pm$ 0.57 dA
	Larva	15.00 $\pm$ 0.00 aA	13.64 $\pm$ 0.15 aB	0.39 $\pm$ 0.04 cB	0.83 $\pm$ 0.04 bA	2.59 $\pm$ 0.30 cB	6.07 $\pm$ 0.31 cA
	Protonymph	14.19 $\pm$ 0.12 aA	12.30 $\pm$ 0.16 bB	0.82 $\pm$ 0.06 bB	1.09 $\pm$ 0.04 aA	5.78 $\pm$ 0.44 bB	8.91 $\pm$ 0.36 bA
	Deutonymph	8.50 $\pm$ 0.25 bA	7.46 $\pm$ 0.19 cB	1.18 $\pm$ 0.06 aA	1.11 $\pm$ 0.08 aA	14.08 $\pm$ 0.65 aA	14.78 $\pm$ 1.04 aA
30	Egg	20.91 $\pm$ 0.83 bA	8.80 $\pm$ 0.37 dB	0.23 $\pm$ 0.03 dA	0.39 $\pm$ 0.06 cA	1.12 $\pm$ 0.17 dB	4.25 $\pm$ 0.74 dA
	Larva	24.66 $\pm$ 0.51 aA	22.10 $\pm$ 0.30 aB	1.13 $\pm$ 0.07 cB	1.61 $\pm$ 0.05 bA	4.61 $\pm$ 0.26 cB	7.36 $\pm$ 0.24 cA
	Protonymph	14.70 $\pm$ 0.43 cA	12.50 $\pm$ 0.36 bB	1.39 $\pm$ 0.07 bB	1.63 $\pm$ 0.04 bA	9.48 $\pm$ 0.39 bB	13.27 $\pm$ 0.67 bA
	Deutonymph	13.35 $\pm$ 0.50 cA	10.75 $\pm$ 0.24 cB	1.76 $\pm$ 0.05 aA	1.86 $\pm$ 0.06 aA	13.39 $\pm$ 0.57 aB	17.52 $\pm$ 0.60 aA
60	Egg	24.41 $\pm$ 0.98 bA	21.14 $\pm$ 0.35 aB	0.69 $\pm$ 0.09 cB	1.18 $\pm$ 0.08 bA	2.79 $\pm$ 0.42 dB	5.62 $\pm$ 0.37 cA
	Larva	26.29 $\pm$ 0.42 aA	20.08 $\pm$ 0.50 aB	1.71 $\pm$ 0.46 bB	1.55 $\pm$ 0.04 aA	4.60 $\pm$ 0.25 cB	7.81 $\pm$ 0.27 bA
	Protonymph	15.12 $\pm$ 0.43 cA	13.85 $\pm$ 0.41 bA	1.36 $\pm$ 0.08 aB	1.71 $\pm$ 0.05 aA	8.96 $\pm$ 0.47 bB	12.55 $\pm$ 0.50 aA
	Deutonymph	14.15 $\pm$ 0.33 cA	14.30 $\pm$ 0.34 bA	1.56 $\pm$ 0.05 aA	1.61 $\pm$ 0.06 aA	11.15 $\pm$ 0.47 aA	11.36 $\pm$ 0.47 aA

Means followed by different letters are statistically different (Turkey HSD test, $P < 0.05$), small letters and capital letters signify differences among prey stages within each density and between exudate treatments for each prey stage, respectively.

Table 2: Analysis of variance of the effects of prey stage, prey density and additional exudate on the predation, oviposition and efficiency of conversion of *Typhlodromalus aripo* in functional response experiment under laboratory conditions

Source	df	Predation			Oviposition rate			Efficiency of conversion		
		MS	F value	P value	MS	F value	P value	MS	F value	P value
Exudate	1	479	203.03	<0.0001	7.35	46.14	<0.0001	0.1387	137.52	<0.0001
Prey stage	3	1178	499.41	<0.0001	23.16	145.32	<0.0001	34.78	344.93	<0.0001
Prey density	3	5615	2380.35	<0.0001	37.03	232.35	<0.0001	0.0122	12.11	<0.0001
Exudate x Prey density	3	117	49.69	<0.0001	0.03	0.184	0.9070	0.0066	6.52	0.0002
Exudate x Prey stage	3	70	29.59	<0.0001	0.68	4.26	0.0054	0.076	7.49	0.0052
Prey density x Prey stage	9	356	151.11	<0.0001	1.51	9.46	<0.0001	0.0099	9.84	<0.0001
Exudate x Prey stage x Prey density	9	61	25.72	<0.0001	0.47	2.92	0.0021	0.0061	6.07	0.0022
Residuals	554	2	-	-	0.16	-	-	0.0010	-	-

Table 3: Maximum-likelihood estimates from logistic regression of the proportion of *Mononychellus tanajoa* stages consumed by *Typhlodromalus aripo* as a function of initial prey density in the absence or presence of additional exudate

Prey stage	Parameter	Exudate absent			Exudate present			
		Estimate	SE	Z _{value}	Estimate	SE	Z _{value}	P _{value}
Egg	P ₀	2.5420	0.5739	4.430	1.4222	0.2432	5.847	<0.0001
	P ₁	-0.2588	0.0768	-3.370	-0.1219	0.0159	-7.661	<0.0001
	P ₂	0.0095	0.0028	3.510	0.0014	0.0002	7.135	<0.0001
	P ₃	-0.0001	0.00002	-3.748	-	-	-	-
Larva	P ₀	7.6932	2.6443	2.909	3.8913	0.4504	8.639	<0.0001
	P ₁	-0.2773	0.1045	-2.652	-0.1145	0.0250	-4.577	<0.0001
	P ₂	-	-	-	0.0006	0.0002	2.132	0.0330
	P ₃	-	-	-	-	-	-	-
Protonymph	P ₀	7.1092	0.6913	10.288	4.2831	0.3702	11.567	<0.0001
	P ₁	-0.3395	0.0364	-9.314	-0.2163	0.0211	-10.231	<0.0001
	P ₂	0.0033	0.00004	8.030	0.0021	0.0002	8.099	<0.0001
	P ₃	-	-	-	-	-	-	-
Deutonymph	P ₀	4.2350	0.7564	5.599	2.8650	0.5688	5.036	<0.0001
	P ₁	-0.4378	0.0951	-4.600	-0.3035	0.0755	-4.016	<0.0001
	P ₂	0.0134	0.0033	3.999	0.0086	0.0027	3.114	0.0018
	P ₃	-0.0001	0.00003	-3.842	-0.00007	0.00002	-2.80	0.0051

Table 4: Estimated from logistic regression of the proportion of *Mononychellus tanajoa* stages consumed by *Typhlodromalus aripo* as a function of initial prey density in the absence or presence of cassava foliar exudate

Prey stage	Parameter	Exudate absent		Exudate present	
		Estimate±SE	T/Th	Estimate ± SE	T/Th
Egg	a (h ⁻¹)	0.0910 ± 0.0142 (0.0626–0.1194)		0.0150 ± 0.0017 (0.0154–0.0225)	
	T_h (h)	0.7129 ± 0.0584 (0.5961–0.8296)	33.6	2.7881 ± 0.1188 (2.5512–3.0250)	8.6
Larva	a (h ⁻¹)	0.7951 ± 0.2417 (0.3128–1.2773)		1.2762 ± 0.4920 (0.2966–2.2556)	
	T_h (h)	0.8852 ± 0.0174 (0.8505–0.9199)	27.1	1.1179 ± 0.0212 (1.0758–1.1601)	24.4
Protonymph	a (h ⁻¹)	0.9301 ± 0.3361 (0.2596–1.6005)		0.6353 ± 0.2323 (0.1727–1.0979)	
	T_h (h)	1.5579 ± 0.0320 (1.4940–1.6218)	15.4	1.7627 ± 0.0400 (1.6832–1.8423)	13.6
Deutonymph	a (h ⁻¹)	0.0894 ± 0.0105 (0.0685–0.1103)		0.0521 ± 0.0040 (0.0440–0.0601)	
	T_h (h)	1.4463 ± 0.0530 (1.3405–1.5521)	16.5	1.3457 ± 0.0504 (1.2451–1.4463)	17.8




Figure 1




Figure 2

Beyond field success: insights into functional and numerical response of the predatory mite *Typhlodromalus aripo* and the role of cassava exudate in predatory performance

ORIGINALITY REPORT

33%

SIMILARITY INDEX

27%

INTERNET SOURCES

30%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1

[bioone.org](https://www.bioone.org)

Internet Source

3%

2

academic.oup.com

Internet Source

3%

3

www.ncbi.nlm.nih.gov

Internet Source

2%

4

escholarship.org

Internet Source

1%

5

www1.montpellier.inra.fr

Internet Source

1%

6

journals.plos.org

Internet Source

1%

7

www.frontiersin.org

Internet Source

1%

8

"Prospects for Biological Control of Plant Feeding Mites and Other Harmful Organisms", Springer Nature, 2015

Publication

1%

9

Submitted to Elmira College

Student Paper

1%

10

Niloufar Doostalizadeh, Ali Asghar Talebi, Yaghoub Fathipour, Ary Hoffmann, Mohammad Mehrabadi. "Lack of impact of Wolbachia on foraging behavior and morphological characteristics of the

1%

parasitoid wasp, *Habrobracon hebetor*
(Braconidae)", Biological Control, 2024

Publication

-
- | | | |
|----|--|-----|
| 11 | hdl.handle.net
Internet Source | 1 % |
|----|--|-----|
-
- | | | |
|----|---|-----|
| 12 | Bonaventure Vidjannagni Agboton, Rachid Hanna, Alexis Onzo, Stefan Vidal, Andreas von Tiedemann. "Interactions between the predatory mite <i>Typhlodromalus aripo</i> and the entomopathogenic fungus <i>Neozygites tanajoae</i> and consequences for the suppression of their shared prey/host <i>Mononychellus tanajoa</i> ", <i>Experimental and Applied Acarology</i> , 2012
Publication | 1 % |
|----|---|-----|
-
- | | | |
|----|---|-----|
| 13 | Reisig, D. D.. "Insecticidal management and movement of the brown stink bug, <i>Euschistus servus</i> , in corn", <i>Journal of Insect Science</i> , 2011.
Publication | 1 % |
|----|---|-----|
-
- | | | |
|----|--|-----|
| 14 | www.researchgate.net
Internet Source | 1 % |
|----|--|-----|
-
- | | | |
|----|---|------|
| 15 | Augustine Bosomtwe, George Opit, Carla Goad, Kristopher Giles, Brad Kard. "Numerical Responses of <i>Xylocoris flavipes</i> (Reuter) (Hemiptera: Anthocoridae) on a Diet of <i>Liposcelis decolor</i> (Pearman) (Psocodea: Liposcelididae)", <i>Insects</i> , 2025
Publication | <1 % |
|----|---|------|
-
- | | | |
|----|--|------|
| 16 | erepository.uonbi.ac.ke:8080
Internet Source | <1 % |
|----|--|------|
-
- | | | |
|----|--|------|
| 17 | Submitted to University of Exeter
Student Paper | <1 % |
|----|--|------|
-
- | | | |
|----|--|------|
| 18 | repositorio.ufc.br
Internet Source | <1 % |
|----|--|------|
-

19	link.springer.com Internet Source	<1 %
20	www.sparkl.me Internet Source	<1 %
21	Ashley Leach, Marc Fuchs, Riley Harding, Brian A Nault. "Iris Yellow Spot Virus Prolongs the Adult Lifespan of Its Primary Vector, Onion Thrips (Thrips tabaci) (Thysanoptera: Thripidae)", Journal of Insect Science, 2019 Publication	<1 %
22	C.E.R.E.N. DOĞAN, N.A.B.İ.A.L.P.E.R. KUMRAL. "Comparative Age-Stage, Two-Sex Life Table Analysis of Native and Abamectin and Bifenazate-Resistant Populations of Amblyseius swirskii Athias-Henriot (Acari: Phytoseiidae) under Laboratory Conditions", Crop Protection, 2025 Publication	<1 %
23	Yaghoub Fathipour, Bahador Maleknia, Abdoolnabi Bagheri, Mahmoud Soufbaf, Gadi V.P. Reddy. "Functional and numerical responses, mutual interference, and resource switching of Amblyseius swirskii on two-spotted spider mite", Biological Control, 2020 Publication	<1 %
24	Azadeh Jarrahi, Seyed Ali Safavi. " Effects of pupal treatment with Proteus® and sensu lato on functional response of parasitising in an enclosed experiment system ", Biocontrol Science and Technology, 2015 Publication	<1 %
25	Submitted to Mansoura University Student Paper	<1 %
26	Submitted to National Chung Hsing University Student Paper	<1 %

27 Yaghoub Fathipour, Fariba Mehrkhou, Mohammad Ali Mirhosseini, Mehran Rezaei, Ahmad Pervez. "Comprehensive foraging behavior of acarophagous ladybird, *Stethorus gilvifrons* (Coleoptera: Coccinellidae) on *Tetranychus urticae* (Trombidiformes: Tetranychidae): implications for biological control", *International Journal of Tropical Insect Science*, 2024

<1 %

Publication

28 M. Zaher, F. M. Momen, A. H. Rasmy, M. S. Nawar, G. Abou-Elella. " Some factors affecting reproduction and sex-ratio of the predacious mite (Muma and Denmark) (Acari: Phytoseiidae) ", *Archives Of Phytopathology And Plant Protection*, 2007

<1 %

Publication

29 Young-Gyun Park, Joon-Ho Lee, Un Taek Lim. "Functional response of *Amblyseius eharai* (Acari: Phytoseiidae) on *Tetranychus urticae* (Acari: Tetranychidae)", *PLOS ONE*, 2021

<1 %

Publication

30 edoc.unibas.ch
Internet Source

<1 %

31 Şengül BİLGİN, Levent İZCİ, Ali GÜNLÜ, Gürken DİKEN, İsmail Yüksel GENÇ. "Effects of gutting process on the shelf life of cultured meagre (*Argyrosomus regius* ASSO, 1801) stored at 4 ± 1 °C", *Food Science and Technology*, 2016

<1 %

Publication

32 Alexis Onzo, Rachid Hanna, Koffi Negloh, Muaka Toko, Maurice W. Sabelis. "Biological control of cassava green mite with exotic and indigenous phytoseiid predators—Effects of

<1 %

-
- 33 Fereshteh Bazgir, Jahanshir Shakarami, Shahriar Jafari. " Prey-stage preferences, functional and numerical responses, and mutual interference of (Acari: Phytoseiidae) on (Tetranychidae) ", International Journal of Acarology, 2020

Publication

-
- 34 M A de Toledo, P R Reis, E C da Silveira, P de P Marafeli, G C de Souza-Pimentel. "Predatory Potential of Euseius alatus (Phytoseiidae) on Different Life Stages of Oligonychus ilicis (Tetranychidae) on Coffee Leaves Under Laboratory Conditions", Neotropical Entomology, 2013

Publication

-
- 35 Shima Yazdanpanah, Yaghoub Fathipour, Elham Riahi, Myron P. Zalucki. " Generation-dependent functional and numerical responses of (Acari: Phytoseiidae) long-term reared on almond pollen ", Biocontrol Science and Technology, 2022

Publication

-
- 36 pubmed.ncbi.nlm.nih.gov

Internet Source

-
- 37 www.scienceopen.com

Internet Source

-
- 38 Karin F. S. Collier. "Neoseiulus idaeus (Acari: Phytoseiidae) as a potential biocontrol agent of the two-spotted spider mite, Tetranychus urticae (Acari: Tetranychidae) in papaya: performance on different prey stage – host plant combinations", Experimental and Applied Acarology, 03/14/2007

39	hal.science Internet Source	<1 %
40	www.biorxiv.org Internet Source	<1 %
41	www.icesse.ase.ro Internet Source	<1 %
42	www.scielo.br Internet Source	<1 %
43	www.scielo.org.co Internet Source	<1 %
44	Alexis Onzo, Rachid Hanna, Maurice W. Sabelis. "Interactions in an acarine predator guild: impact on Typhlodromalus aripo abundance and biological control of cassava green mite in Benin, West Africa", Experimental & Applied Acarology, 2003 Publication	<1 %
45	David Wari, Jun Yamashita, Hidenari Kishimoto, Shoji Sonoda. "Utilization of plant food resources by phytoseiid mite species with different feeding habits", Applied Entomology and Zoology, 2016 Publication	<1 %
46	Isabelle Vantornhout. "Influence of diet on life table parameters of Iphiseius degenerans (Acari: Phytoseiidae)", Experimental and Applied Acarology, 03/2005 Publication	<1 %
47	Konstantinos Samaras, Maria L Pappas, Apostolos Pekas, Felix Wäckers, George D Broufas. "Benefits of a balanced diet? Mixing prey with pollen is advantageous for the	<1 %

-
- 48 Mohammed M. E. Elmoghazy. " density on variety of plant leaves influencing predatory mite functional response ", International Journal of Acarology, 2022 $<1\%$
- Publication
-

- 49 Reham Abo-Shnaf, Enrico de Lillo. "Phytoseiid mites (Acari: Mesostigmata) from Apulia (Southern Italy), with a key to the species reported from Italy", International Journal of Acarology, 2021 $<1\%$
- Publication
-

- 50 Sajjad Dalir, Hamidreza Hajiqanbar, Yaghoub Fathipour, Mostafa Khanamani. " A comprehensive picture of foraging strategies of and on western flower thrips ", Pest Management Science, 2021 $<1\%$
- Publication
-

- 51 Syed Usman Mahmood, Xiaoyi Huang, Runqian Mao, Huihua Hao, Xiaoduan Fang. "Integrated Biological Control Strategies for Citrus Rust Mites: Distribution, Impact on Mandarin Quality, and the Efficacy of *Amblyseius largoensis*", Insects, 2024 $<1\%$
- Publication
-

- 52 XiaoTong Fu, YuanZhi Cao, XinTong Dong, Jing Chang, Zhijia Huo, RuiXia Meng. "Functional responses of two species of predatory mites (Acari: Phytoseiidae) to eggs and first-instar nymphs of *Bactericera Gobica Logniova* (Homoptera: Psyllidae)", Experimental and Applied Acarology, 2024 $<1\%$
- Publication
-

<1 %

54

ejbpc.springeropen.com

Internet Source

<1 %

55

entomotropica.org

Internet Source

<1 %

56

sherekashmir.informaticspublishing.com

Internet Source

<1 %

57

www.researchsquare.com

Internet Source

<1 %

58

www.tandfonline.com

Internet Source

<1 %

59

zidapps.boku.ac.at

Internet Source

<1 %

60

İsmail Döker, Hatice Yavaş, Zeinab Shirvani, Mahmut Mete Karaca, Kamil Karut, Dejan Marčić, Cengiz Kazak. "Toxicity and risk assessment of pesticides on the predatory mite, *Amblyseius swirskii* Athias-Henriot (Acari: Phytoseiidae) under laboratory conditions", *Phytoparasitica*, 2024

Publication

<1 %

61

Bruna L. Merlin, Lucia P. Ferreira, Wesley A.C. Godoy, Gilberto J. Moraes, Fernando L. Cônsoli. "Functional response of *Neoseiulus californicus* preying on *Tetranychus urticae* is affected by prey quality and host-plant acclimation", *Biological Control*, 2022

Publication

<1 %

62

Eduardo Pereira de Sousa Neto, Rosenya Michely Cintra Filgueiras, Jairo de Almeida Mendes, Neville Vieira Monteiro et al. "A drought-tolerant *Neoseiulus idaeus* (Acari: Phytoseiidae) strain as a potential control

<1 %

agent of two-spotted spider mite,
Tetranychus urticae (Acari: Tetranychidae)",
Biological Control, 2021

Publication

63

Farazmand, Azadeh, Yaghoub Fathipour, and Karim Kamali. "Intraguild predation among *Scolothrips longicornis* (Thysanoptera: Thripidae), *Neoseiulus californicus* and *Typhlodromus bagdasarjani* (Acari: Phytoseiidae) under laboratory conditions", *Insect Science*, 2013.

<1 %

Publication

64

Fereshteh Bazgir, Jahanshir Shakarami, Shahriar Jafari. " Life table and predation rate of (Acari: Phytoseiidae) fed on (Tetranychidae) on apple leaves ", *International Journal of Acarology*, 2019

<1 %

Publication

65

Harold Andrew Patrick, Ravichandran Krishnamoorthy. "Applied Research for Growth, Innovation and Sustainable Impact", *Routledge*, 2025

<1 %

Publication

66

Koffi Negloh, Rachid Hanna, Peter Schausberger. "Comparative demography and diet breadth of Brazilian and African populations of the predatory mite *Neoseiulus baraki*, a candidate for biological control of coconut mite", *Biological Control*, 2008

<1 %

Publication

67

pure.uva.nl
Internet Source

<1 %

68

Abd-Elrady K. Nasr, Faten M. Momen, Abd-Elstar M. Metwally, Mohamed Gesraha, Awad A. Abdallah, Khaled M. Saleh. "Eignung der Eier von *Corcyr a cephalonica*

<1 %

(Lepidoptera: Pyralidae) für die Entwicklung, Vermehrung und das Überleben von vier Raubmilbenarten der Familie Phytoseiidae (Acari: Phytoseiidae)", Gesunde Pflanzen, 2015

Publication

69

Elias Soares de Figueiredo, Marcela Massaro, Stephanie do Carmo, Gilberto José de Moraes. "Rearing system for the predatory phytoseiid *Euseius concordis* (Acari: Phytoseiidae)", *Experimental and Applied Acarology*, 2018

Publication

<1 %

70

Jonathan G. Lundgren. "Relationships of Natural Enemies and Non-Prey Foods", Springer Science and Business Media LLC, 2009

Publication

<1 %

71

Navid Sehat-Niaki, Azadeh Zahedi Golpayegani, Ehssan Torabi, Alireza Saboori, Behnam Amiri-Besheli, Yaghoub Fathipour. "Sublethal effects of chlorfenapyr and acequinocyl on the functional and numerical responses of the predatory mites *Phytoseiulus persimilis* and *Neoseiulus californicus* (Acari: Phytoseiidae) feeding on *Tetranychus urticae* (Acari: Tetranychidae)", *Experimental and Applied Acarology*, 2024

Publication

<1 %

72

Nikos E. Papanikolaou, George D. Broufas, Theodore Kypraios, Triantafyllia Liaka, Maria L. Pappas. "Intraspecific interactions at high predator densities affect the predation efficiency of (Acari: Phytoseiidae) when prey density is low ", *Ecological Entomology*, 2022

Publication

<1 %

73 Pinto-Zevallos, Delia M., Martín Pareja, and Bianca G. Ambrogi. "Current knowledge and future research perspectives on cassava (*Manihot esculenta* Crantz) chemical defenses: An agroecological view", *Phytochemistry*, 2016.

Publication

<1%

74 SU-QIN SHANG, LI WEI-ZHEN, YAO-NIAN CHEN, ZHU TONG. "The influence of Ahy1 strain of *Acremonium hansfordii* on functional response of *Neoseiulus barkeri* to *Tetranychus urticae*", Research Square Platform LLC, 2022

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On