

Spatio-temporal variability of the onset and end dates of the rainy seasons in the Bounkani region (North-East of Côte d'Ivoire).

by Jana Publication & Research

Submission date: 29-Aug-2025 01:17PM (UTC+0700)

Submission ID: 2690351138

File name: IJAR-53584.docx (2.94M)

Word count: 6809

Character count: 35121

Spatio-temporal variability of the onset and end dates of the rainy seasons in the Bounkani region (North-East of Côte d'Ivoire).

Abstract

In Côte d'Ivoire, 80% of rural activities are based on rain-fed agriculture, although the country has been experiencing a disruption of its rainfall regime for several decades. This study was conducted in this context in order to deepen the previous knowledge on the rainfall regime of the Bounkani region through the spatio-temporal distribution of the dates of the beginning and end of the rainy season. The rainfall data used are of the CHIRPS type for the 1981-2020 period. The application of rainfall indices allowed us to observe surplus and deficit years. These variations indicate a downward trend. The average rainfall has decreased from 1050 mm during the decade 1985-1995 to 950 mm in 2011-2020, a loss of 100 mm. This deficit was reflected in the onset and end dates as well as the number of rainy days of the rainy seasons. The study of the number of rainy days of the rainy season reveals that they have a very low downward trend. Indeed, the starting dates of the rainy season are increasingly late. On average, the start of the season, which was between April 6 and 10 in the years 1981-1995, has shifted from May 1 to 5 in the decade 2011-2020 in the North of the region. While, the South records an average date of season start from April 1 to 5 in the years 1981-1995 and from May 6 to 10 in the decade 2011-2020, a shift of more than 30 days. The start of the rainy season is progressive along the South-North gradient. Conversely, the end of the rainy season occurs first in the North and then extends to the South.

Keywords: GIS and remote sensing, early and late season, rainfall variability, CHIRPS, IDW, Côte d'Ivoire.

Introduction

Since the 1970s, West Africa has experienced one of the greatest rainfall variations observed on a global scale. These variations are reflected in rainfall deficits that have affected the countries of the Gulf of Guinea (Brou and Chaléard, 2007 ; Paturel *et al.*, 1995 ; Servat *et al.*, 1997 ; Balliet *et al.*, 2016). Côte d'Ivoire is not immune to this reality, particularly in the north of the country. The new climatic context has deteriorated considerably and the situation is more alarming in the North-East sector (Brou, 2010). This is the most deficient area in terms of cumulative annual rainfall in Côte d'Ivoire (Brou, 2009). The implications of these climatic variations on water resources are particularly high and affect many sectors of activity such as

livestock and agriculture (Ardoin-Bardin, 2004). Agriculture, which occupies a predominant place in the national economies, remains one of the activities most affected by these climate variations.

Nearly 80% of Ivorian agricultural production is rain-fed (Noufè, 2012). However, changes in the descriptive elements of the rainy season led to disruptions in crop production. This situation is therefore worrying, especially for rural communities whose agricultural activities represent the bulk of their income. According to Ahoussi et al.(2013)The disruption of existing cropping systems, such as crop calendars, is already known and observed by farmers in several localities in the country. The dating of new calendars is therefore necessary for better crop planning. In addition, the determination of the onset and end dates of the rainy seasons are crucial issues in tropical Africa, where farming practices are closely dependent on the rains, as they also determine the agricultural calendar and, to a large extent, the quality of the crops. There are several definitions in the literature relating to the determination of the onset and end dates of the rainy season. In particular, those found in studies on rainfall variability and the determination of the onset and end dates of the rainy season in West Africa (Sivakumar, 1988, Joseph et al. 1994, Traore et al. 2000, Camberlin et al. 2003, Ndong, 2003, Marteau et al. 2010) and in Côte d'Ivoire (Balme et al. 2005, Kouassi, 2007, Goula et al. 2010, Noufè, 2011, Dekoula et al. 2018, etc.). Indeed, there is no universally accepted criterion for defining the onset and end of rainy seasons. Kouassi (2007) defines according to Ozer and Erpicum (1995) defines the beginning of the rainy season as the time (from March 1^{er} to October 31) when the probability of having a rainy day during a given pentad is greater than that of having a dry day belonging to a dry episode of more than 7 days. While Dekoula et al. (2018) and Kouamé et al. (2018) in their studies in the north of Côte d'Ivoire on the variability of the rainfall regime define the beginning of the rainy season in a bimodal regime as the first day from 1^{er} February when at least 20 mm of rainfall is recorded over two successive days for some and over three successive days for others. And this, without however that it does not have a dry episode of more than 7 consecutive days during the 30 days that follow. It is worth noting that most of the methods used in these studies define a single empirical threshold and set a time interval within which the count must be made. However, none of them have studied the spatial and temporal distribution of these dates in the Bounkani region or even at the national level.

Also, the rainfall data used in these studies come from the 'Société d'Exploitation et de Développement Aéroportuaire Aéronautique et Météorologique '(SODEXAM). Ivorian researchers agree that the SODEXAM station data are not regularly distributed and have gaps over a decade or so due to the socio-political crisis of 2002. In Côte d'Ivoire, the station grid

indicates an average density of one station per 3000 km² (Lienou, 2007, Noufè, 2011, Dekoula, et al, 2018etc.).

For an in-depth study of the rainfall regime, daily data are indispensable, although they are difficult to access, particularly in Côte d'Ivoire, where the network of stations is not very dense.

In this study, the method used to determine the onset and end dates of the rainy season is suitable for all regions (universal). It is very accurate. Indeed, this method allows the calculation of a threshold for each locality according to its rainfall. In addition, the rainfall data used in this work are of satellite type. They cover the entire study area, thus allowing a good spatial analysis. This dataset (CHIRPS) is reliable, accurate with high resolution and freely available. They have been proven in several studies (Seregina et al. 2018 ; Didi et al. 2020 ; Sie et al. 2021).

In light of the above, it is essential to know and understand the onset and end dates of the rainy seasons in order to control and define a new agricultural calendar. This study contributes to the knowledge of the rainfall regime in the Bounkani region through the determination and spatial and temporal distribution of the onset and end dates of the rainy season.

Study area

The Bounkani region is located in the northeast of Côte d'Ivoire between 2.60° and 4.30° west longitude, and 8.20° and 9.95° north latitude (Fig. 1). It is located 550 km from Abidjan, the economic capital, with an area of 22,091 km² or 6.9% of the national territory. However, half of its area is occupied by the Comoé National Park, which covers 11,090 km². The region covers the departments of Nassian, Doropo, Tehini, and Bouna. This area is bordered to the north by Burkina Faso, to the east by Ghana, to the west by the Tchologo region and to the south by the Gontougo region with which it forms the Zanzan district. It is the largest region of Côte d'Ivoire with a population of 427037 inhabitants in 2021 (INS, 202121) or 38.8hbs/km². The main activity of the population is mainly agricultural. Its climate is Sudanian with a rainy season and a dry season. The average annual rainfall is between 900 and 1,200 mm (Sie et al, 2021).




Figure 1: Location of the Bounkani region

Data and methods

Study data

The data used are the Climate Group Hazards InfraRed Precipitation (CHIRPS) satellite products. They are available for over 40 years (1981 to present) with monthly, decadal, pentadal, and daily rainfall (Funk et al., 2015). In this study, annual, monthly, pentadal and daily products from 1981 to 2020 were used for studies of rainfall variability, rainy season onset and end dates.

There are 12 SODEXAM rainfall or hydrometric stations in the region. These stations are irregularly distributed and do not cover the entire area, especially in the park. Also, the data from these stations are incomplete. To overcome these problems, the coordinates of existing and fictitious stations were used to extract rainfall data from the CHIRPS products. This allowed for a homogeneous distribution of stations with reliable and accurate data (5 km resolution) for a better analysis of rainfall variability.

The rainfall data from these 33 stations, homogeneously distributed in the region, were used for the spatial distribution (Figure 2). These data are freely available online (<https://www.chc.ucsb.edu/data/chirps>).




Figure 2: Spatial distribution of rainfall stations

Methods

Interannual and seasonal variation in precipitation

The study of rainfall variability was carried out using the analysis of annual and monthly rainfall data from the CHIRPS products. In order to assess the evolution of rainfall during the different years of the study period, the Nicholson rainfall index and the Hanning low-pass filter of order 2 were applied. These methods have the advantage of highlighting the periods of surplus and deficit in rainfall. The rainfall evolution or rainfall trend was observed using the regression line.

Nicholson Rainfall Index (I_p)

It allows the observation of interannual rainfall variations over a long series of observations (Nicholson et al., 1988). The calculation of this index was carried out on the rainfall data of the four departments of the study area for a period of 40 years of observation. The Nicholson index is expressed by equation (1):

$$I_p = \frac{(X_i - \bar{X})}{\sigma} \text{ (Eq. 1)}$$

X_i the precipitation of year i ,

\bar{X} the average annual precipitation over the observation period and

σ the standard deviation over the same period.

2nd order Hanning low pass filter

According to Soro *et al* (2011) the elimination of seasonal variations allows a better observation of interannual fluctuations. In this case, the annual rainfall totals are weighted using the following equations recommended by Assani (1999):

$$\bar{X}_t = 0.06x_{(t-2)} + 0.25x_{(t-1)} + 0.38x_{(t)} + 0.25x_{(t+1)} + 0.06x_{(t+2)} \text{ for } 3 \leq t \leq (n-2) \quad \text{(Eq. 2)}$$

where $x_{(t)}$, the weighted rainfall total of term t , $x_{(t-2)}$ and $x_{(t-1)}$ are the rainfall totals of the two terms immediately preceding term t , and $x_{(t+2)}$ and $x_{(t+1)}$ are the rainfall totals of the two terms immediately following term t .

The weighted rainfall totals of the first two terms [$x_{(1)}$, $x_{(2)}$] and the last two terms [$x_{(n-1)}$, $x_{(n)}$] of the series are calculated using the following expressions (n being the size of the series):

$$\bar{X}_{(1)} = 0.54x_{(1)} + 0.46x_{(2)} \quad \text{(Eq. 3)}$$

$$\bar{X}_{(2)} = 0.25x_{(1)} + 0.50x_{(2)} + 0.25x_{(3)} \quad \text{(Eq. 4)}$$

$$\bar{X}_{(n-1)} = 0.25x_{(n-2)} + 0.50x_{(n-1)} + 0.25x_{(n)} \quad \text{(Eq. 5)}$$

$$\bar{X}_{(n)} = 0.54x_{(n)} + 0.46x_{(n-1)} \quad \text{(Eq. 6)}$$

To better observe the months of rainfall deficit and surplus in the four departments (Bouna, Doropo, Tehini and Nassian), the moving averages were centered and reduced using the formula (Eq. 1). This method appears to be more efficient because it allows for a perceptible division of the wet and dry seasons.

Variability in the onset and end dates of the rainy seasons

To identify the dates of the beginning and end of the rainy season for the different stations in our study area, the determination criteria as used by Marengo *et al* (2001) and Giráldez *et al* (2020) were applied. This method is done in two steps:

- First, there was a question of determining specific precipitation thresholds for each station above or below which the season will start or end. These thresholds were calculated by an analysis of the frequency and intensity of smoothed pentadal precipitation. The precipitation threshold is obtained in two steps: i) the 3rd order moving average was applied to smooth out irregularities in the time series. ii) The 50th percentile of precipitation between January and June was used as the threshold for identifying the start of the rainy season. To identify the end of the rainy season, the threshold was the 50th percentile of rainfall between July and December (Giráldez et

al., 2020). The onset and end dates at each station are defined by a particular rainfall threshold (Table 2).

- Second, the rainy season onset date was defined when the first pentad of a set of 6-8 consecutive pentads exceeds the precipitation threshold previously identified in Table 2. Similarly, the end date of the rainy season was determined as the date before the first pentad of a set of 6 to 8 successive pentads that do not exceed the previously identified rainfall threshold (Table 2).

Table 2: Onset and end date thresholds for each station

N°	Stations	Long.	Lat.	Early season threshold (mm)	End of season threshold (mm)
1	Awahikro	-3.923	8.310	16.03	17.00
2	Bouna	-2.983	9.267	11.09	19.72
3	Doropo	-3.317	9.817	9.58	23.10
4	Gawi	-3.811	8.935	13.46	21.87
5	Kafolo	-4.415	9.583	10.78	24.81
6	Kakpin	-3.783	8.667	14.96	19.02
7	Kokpingue	-3.146	8.981	13.33	21.04
8	Kolon	-4.076	9.067	13.40	22.21
9	Koutouba	-3.200	8.700	15.99	18.37
10	Laoudi Ba	-3.027	8.372	18.28	16.66
11	Nassian	-3.480	8.430	16.12	17.84
12	Ouango Fitini	-4.078	9.617	10.40	24.29
13	Saye	-3.350	9.450	10.93	23.29
14	Tehini	-3.683	9.617	10.36	22.82
15	Varale	-3.320	9.670	10.24	22.88
16	STA0	-3.489	9.782	9.62	23.23
17	STA1	-3.637	9.894	9.10	23.43
18	STA2	-3.856	9.803	9.40	23.02
19	STA3	-3.864	9.477	10.87	22.63
20	STA4	-4.091	9.370	11.53	23.53
21	STA5	-3.613	9.358	11.66	22.82
22	STA6	-3.072	9.556	10.43	24.16
23	STA7	-3.295	9.205	12.18	23.32
24	STA8	-2.841	8.916	13.55	19.16
25	STA9	-2.920	8.681	15.55	18.54
26	STA10	-2.709	9.403	10.83	23.68
27	STA11	-3.716	9.155	12.58	22.63
28	STA12	-3.501	9.007	13.35	21.37
29	STA13	-3.456	8.718	15.70	17.98
30	STA14	-3.716	8.479	15.65	18.37
31	STA15	-3.543	8.194	15.93	15.57
32	STA16	-2.561	8.731	13.92	18.72

Analysis of rainy days

The determination of the number of rainy days per year and per season was made on grounds of the daily rainfall data. A day is considered rainy if the amount of precipitation received that day is greater than or equal to 1 mm. The number of rainy days is evaluated on a seasonal basis.

Spatial and temporal variation in the onset and end dates of the rainy season

In this study, we used the Inverse Distance Weighting (IDW) interpolation method for the spatial distribution of data from the thirty-three (33) stations in the area of interest. This method is effective and robust for data (rain, temperature, etc.) that are continuous and evenly distributed over the area as is the case in this work.

Inverse Distance Weighting (IDW) is based on the assumption that features that are close together are more similar than those that are farther apart. To predict the value of an unmeasured sample point, IDW will use the values of the known samples surrounding that prediction point. The known values of the samples closest to the prediction point will have more influence on the predicted value than those further away. The advantage of the IDW method is that it is intuitive and efficient. This interpolation method works best with uniformly distributed points that have no outliers (Azpurua and Dos Ramos, 2010).

Thus, it weights nearby points more than those that are farther away, hence the name inverse distance weighting. The value of the prediction point is given by the following general equation:

$$\hat{Z}(s_o) = \sum_{i=1}^n \lambda_i Z(s_i) \quad (\text{Eq. 17})$$

where $\hat{Z}(s_o)$ is the value, we are trying to predict for the location s_o ; n is the number of measured sample points surrounding the prediction location that will be used in the prediction;

λ_i is the weight assigned to each measured point we will use and these weights will decrease with distance;

$Z(s_i)$ is the value observed at the location s_i .

The weights at the location s_i can be determined by:

$$\lambda_i = d_{i0}^{-p} / \sum_{i=1}^N d_{i0}^{-p} \sum \lambda_i = 1 \quad (\text{Eq. 18})$$

As the distance increases, the weight is reduced by a factor p . d_{i0} the distance between the prediction point s_o and each of the measured sample points s_i . The power parameter p influences the weighting of the measured point value on the prediction point value; that is, as the distance between the measured sample points and the prediction point increases, the

weight (or influence) that the measured point will have on the prediction will decrease exponentially. The weights of the measured point that will be used for the prediction are scaled so that their sum is equal to 1 (Hodam *et al.*, 2017).

Results and discussion

Results

Interannual variability of rainfall

Figure 3 illustrates perfectly that the amount of rainfall is decreasing in the Bounkani region. The observed reduced centered indices (Fig. 3a) show a decrease in rainfall over the entire chronicle with a small non-significant slope ($a = -0.007$) and Figure 3b also presents the average cumulative rainfall amounts per quinquennium. This figure shows that the 1986-1990 five-year period received a large amount of rain (about 1090 mm) in the region, unlike the three following five-year periods (1991-1995, 1996-2000 and 2001-2005), which recorded a gradual decline in rainfall to 985 mm, or a loss of 100 mm. A recovery of rainfall from 2006 to 2010 with an average rainfall of 1057 mm, is followed by a considerable decline from 2011 to 2020 corresponding to a drop of about 80 mm of rainfall.

For more details on the evolution of rainfall in the region, we have determined the deficit and surplus months of the different quinquennia. For the study of these months, we used monthly data.




Figure 3: Interannual variability of rainfall from 1981 to 2020

Seasonal variation in rainfall

The second-order Hanning low-pass filter and the reduced centered index applied to the five-year average of the rainfall series makes it possible to observe the surplus and deficit months in the study area (Figure 4). It can be seen that in the Bounkani region the rainy season comprises six (06) months that generally begin in May and end in October, except in the Nassian department, where the rainy season lasts seven months (Fig. 4d). It extends from April to October with a wet transition generally observed in August. This period precedes the main dry season, which lasts 5 months (November-March). There is a clear variation in the

rainy season in Bouna department (Fig. 4a). From 1981 to 1985 and from 1991 to 1995, the rainy season began in April, but from 1986 to 1990 and from 1996 to 2020, it began in May. For a better precision of the seasonal variation, the dates of onset and end of the rainy season of the different departments and the 33 stations of the study area were determined.




Figure 4: Seasonal variability of rainfall from 1981 to 2020

Variations in the onset and end of rainy seasons

The variation in the dates of the onset and end of the season is illustrated in Figures 5 and 6 respectively. Three (3) departments in the region, namely Bouna, Doropo and Tehini have an increasing trend (Fig. 5). This reveals that the rainy season is starting later and later in these areas over time. The start date varies between pentad 14 (6-10 March) and pentad 30 (26-31 May) in Bouna department. Doropo is located between the 18^{ème} (26-31 March) and 26^{ème} (6-10 May) pentad with two (02) years (1998 and 2003) of late onset (29^{ème} pentad), i.e., more than two weeks later. Like Doropo, in Tehini department, the start date oscillates between the 18^{ème} (26-31 March) and the 26^{ème} (6-10 May) pentad with the year 2013 having an early start (6-10 March) of rain and the year 2003 having a late start (21-25 May).

Unlike the first three departments, Nassian has a declining trend in season onset date. This declining trend shows that in the southern part of the region, the rainy season tends to start a little earlier over time. The start of the season evolves between the 13^{ème} (March 1-5) and 30^{ème} (May 26-31) pentad.




Figure 5: Variability of rainy season onset dates from 1981 to 2020

As for the end dates of the rainy season in the four departments that make up the Bounkani region, represented by Figure 6, it is apparent that the region as a whole is showing an upward trend with low slopes averaging 0.03. This reflects the fact that the end dates of the rainy seasons tend to occur later in the study area. However, the end dates vary differently across the departments. In Bouna and Tehini departments, the end of the rainy season varies between the 51^{ème} (Sept. 11-15) and 61^{ème} pentad (Nov. 1-5), while in Doropo and Nassian departments, the end of the rainy season fluctuates between the 49 (Sept. 1-5) and 61 (Nov. 1-5) pentads. In addition, early season ends (21-25 Aug.) were recorded in Doropo and Tehini departments in 1981.




Figure 6: Variability of rainy season end dates from 1981 to 2020

After determining the onset and end dates of the seasons, the analysis of rainfall amounts during the rainy season is presented in Figure 7. There is a decrease in rainfall amounts from 1981 to 2020 in the departments of the Bounkani region, with the exception of Doropo, which shows an increasing rainfall trend. The departments concerned by the decrease in rainfall are Bouna, Tehini, and Nassian with rainfall varying between 500 mm and 1100 mm during the rainy season. Rainfall in Doropo department varies between 600 mm and 1250 mm during the rainy season.




Figure 7: Variation in rainfall amounts in the rainy season from 1980 to 2020

Spatial variation in season onset and end dates⁶

Onset date of the rainy season

The spatial distribution of rainy season onset dates (Figure 8) shows a uniformity in the start⁷² of the season in the Bounkani region during the 1981-1985 five-year period. In this period, the start of the season was between April 11 and 15. However, the extreme south had an earlier start to the season than the north, which started a month later. The 1986-1990 five-year period shows that the start of the season varies between pentads 21 and 25 (11 Apr-5 May). This variation is observed in the east in Youndouo Sub-Prefecture and in the west in Comoé Park. In this first decade (1981-1990), a variation in season onset dates is observed. Initially, the region that had a season onset date (11-15 Apr) has now shifted to a date between 01 and 05 May, a shift of about 20 days.

The next three quinquennia 1991-1995, 1996-2000 and 2001-2005 show an earlier start of the season than the previous decade (1981-1990). The dates are located between pentads 17 and 19 (21 Mar-5 Apr). Furthermore, the 1991-1995 five-year period shows⁵⁹ that the rainy season tends to start earlier in the South, which generally begins in pentad 17 (21-25 Mar), and then progresses towards the North. From 2001 to 2005, the rainy season began more or less during pentad 19 (1-5 Apr) throughout the region, except for a few sub-prefectures in the departments of Bouna (Bonna, Ondefidouo) and Nassian (Sominassé), which experienced a delay of about one week.

The last three quinquennia (2006-2020) of the study period shows a shift in rainy season onset dates that varies between pentads 21 and 29 (11 Apr-25 May). The 2011-2015 quinquennium shows a strong shift in season onset dates in the time series (1981-2020). This shift²⁷ is more pronounced in the southern part of the region, particularly Nassian and the southern part of Bouna department. These areas have a late start to the season (21-25 May) in the 2011-2015 five-year period compared to the 1991-2000 decade, during which the season started between 21 and 25 March. This corresponds⁸⁰ to a delay in the start of the season of about 2 months or 60 days.




Figure 8: Map of the spatial and temporal distribution of the rainy season from 1981 to 2020

End of rainy season date

Figure 9 presents a spatial distribution of rainy season end dates in the Bounkani region. There is an increasing gradient from north to south in the date of the end of the season. In general, the end of the season occurs first in the north of the region and then gradually moves southward. We note that during the 1981-1985 five-year period, the region experienced shifts in the end of the season, particularly in Tehini Department and the sub-prefectures of Danoa and Bouko. In these localities, the rainy season ended between September 1 and 15. While the South of Nassian recorded the end of the season between October 6 and 20, one month after the rains stopped in the North.

The 2001-2005 and 2006-2010 five-year periods show a change in end dates along the South-North gradient. The northern half of the region experienced a season end in pentad 53 (Sept. 21-25) during the 2001-2005 quinquennium. While in the second part of the decade (2006-2010), the region observed a delay in the end of the rainy season. It thus shifted from pentad 53 to pentad 60 (26-31 Oct.) respectively in the south and east of the study area. With the exception of Tehini department, which maintained its end-of-season period.

The last decade 2011-2020, on the other hand, shows a decrease in the ending date trend. The southern part of the region that had recorded the end of the season during pentad 60 has regressed by 2 pentads corresponding to 10 days of lag.




Figure 9: Map of the spatial and temporal distribution of the end of the rainy season from 1981 to 2020

Statistical analysis of rainy days⁴¹

Obtaining the onset and end dates of the rainy season allowed for a statistical analysis of rainy days during the rainy season. Figure 10 shows the variations in the number of rainy days during the rainy season. There is a slight downward trend in the number of rainy days in Bouna and Tehini with slopes $a=-0.05$ and $a=-0.025$ respectively. In contrast to the previous departments, Doropo and Nassian experienced an increase in the number of rainy days with a steeper slope for Nassian (Figure 10).




Figure 10: Change in number of rainy days from 1981 to 2020³³

Discussion

In recent years, the rainy season has been confronted with anomalies arising from rainfall descriptors that are an important factor in rainfed agriculture. Indeed, the results obtained from statistical methods (Nicholson index, regression line) reveal that the Bounkani region has recorded a decrease in rainfall from 1981 to 2020. The chronicle shows deficit periods (the five-year periods 1981-1985 and 2011-2015 with rainfall of 930 mm and 950 mm respectively). According to Brou (2009), this is the most deficient area in terms of cumulative annual rainfall in Côte d'Ivoire. The year 1983 recorded a severe drought in the West African region, which was followed by a resumption of rainfall until 2013, which also recorded very little rainfall. This alternation of wet and dry periods has been illustrated by several previous studies (Noufè et al., 2015, Dekoula et al., 2018, etc.). These studies show the decline in rainfall at the end of the 1960s in the north of Côte d'Ivoire, which gradually spread

southwards (Brou, 1997 ; Bigot ⁸² *et al.* 2005 and Noufé *et al.* 2011). This disfunctioning of rainfall observed over the last few decades has had repercussions on the distribution of seasons.

Pentadal data were used for an accurate determination ⁷ of the onset and end dates of the rainy season to guard against the risks of false starts and early ends during the rainy seasons in the study area. The results show a shift in ⁵⁴ the onset and end dates of the rainy season. This shift is observed by a slight shift in ⁶ the onset and end dates of the rainy season in the departments of Bouna, Doropo and Tehini. This means ⁶ that the duration of the rainy season in these departments is almost stationary. While Nassian shows a decrease in ⁶ the trend in the onset dates of the wet season, the trend in the end dates varies little. These trends indicate ³⁵ a delay in the start of the rainy season in this department and little variation in the end of the season. This shows ⁵³ that there is a shortening of the length of the rainy season in this department. This difference between the northern and southern localities can be explained by their belonging to different climatic regimes. Indeed, the Bounkani region is located in the Sudanese climate regime (monomodal), except for the department of Nassian, which is located in the south in the ¹⁰ transition zone between the Sudanese climate and the equatorial climate with an attenuated transition and bimodal regime. This department straddles the gap between the monomodal and bimodal regimes. This corroborates the work of Noufé (2011) which reveals that the separation of these two climatic domains ⁸⁷ in the northeast quarter of Côte d'Ivoire follows the Nassian-Farako line south of Comoé National Park. In addition, there was a ⁶² decrease in the amount of rainfall during the rainy season throughout the region except for Doropo, which recorded a slight increase. Brou (2010) and Kanga and Assi Kaudjhis (2016) also noted a ¹⁰⁶ decrease in rainfall in the Bounkani area in their study on rainfall variation from 1951 to 2010 ⁹⁵ in the northeast of Côte d'Ivoire. ⁶⁰ The maps of spatio-temporal distribution of the dates of the beginning and end of the seasons show a progressive onset and end of the season in the Bounkani region. ⁴ The progressive start ⁷⁴ according to the South-North gradient is caused by the movement of the monsoon from the Atlantic and conversely the end that occurs ²⁷ according to the North-South gradient is generated by the migration of the harmattan from the continent. These phenomena are the basis ³³ for determining the number and duration of the rainy season (Eldin, 1971). ²¹ These results confirm those of Noufé *et al.* (2015) who presented a latitudinal ⁹ evolution of the start of the rainy season in eastern Côte d'Ivoire. ⁹⁸

Conclusion

This study has highlighted the high variability of rainfall from 1981 to 2020 in the Bounkani region of northeastern Côte d'Ivoire. This variation is characterized by an alternation of surplus and deficit years. The in-depth analysis indicates a seasonal variation, and the wet months are thus determined in order to assess the rainfall amounts during these periods for crops. A decrease in rainfall was observed in the region during the rainy season. In fact, the 1986-1995 decade received an average rainfall of about 1050 mm, whereas the 2011-2020 decade received 950 mm, or a loss of 100 mm. This rainfall recession is more pronounced in Bouna, which recorded an average rainfall of 730 mm during the 2011-2020 decade. In order to better understand seasonal variations for efficient rainfed crop planning, the onset and end dates as well as the number of rainy days (rainfall ≥ 1 mm) of the rainy season were determined. This increased trend reflects a shift in the season. The rainy season tends to start later and later in the Bounkani region. Nassian remains the most affected area with more than one month delay in the decade 2011-2020 compared to that of 1986-1995. In contrast, the trend in the variation of end dates in our study area is quasi-stationary over the study period. End dates vary on average between 1^{er} to 5 October in the North and 16 to 20 October in the South. Furthermore, South-North and North-South gradients are observed for the onset and end dates of the rainy season in the Bounkani region, respectively.

Library references

- Ahoussi, K. E., Koffi, Y. B., Kouassi, A. M., Soro, G., Soro, N., & Biémi, J. (2013). Étude de la variabilité hydroclimatique et de ses conséquences sur les ressources en eau du Sud forestier et agricole de la Côte d'Ivoire : cas de la région d'Abidjan-Agboville. *International Journal of Pure & Applied Bioscience*, 1(6), 30–50.
- Ardoïn-Bardin, S. (2004). Variabilité hydroclimatique et impacts sur les ressources en eau de grands bassins hydrographiques en zone soudano-sahélienne. Thèse de Doctorat, Université Montpellier II (France).
- Assani, A. A. (1999). Analyse de la variabilité temporelle des précipitations (1916-1996) à Lumbashi (Congo-Kinshasa) en relation avec certains indicateurs de la circulation atmosphérique (oscillation australe) et océanique (El Niño/La Niña). *Sécheresse*, 10(4), 245–252.
- Azpuruá, M., & Dos Ramos, K. (2010). Acomparision of spatial interpolation methods for estimation of average electromagnetic field magnitude. *Progress In Electromagnetics*

Research M, 14, 135–145.

- Balliet, R., Saley, M. B., Eba, E. L. A., Sorokoby, M. V., Bi, H. V. N., N'Dri, A. O., Djè, B. K., & Biémi, J. (2016). Évolution Des Extrêmes Pluviométriques Dans La Région Du Gôh (Centre-Ouest De La Côte d'Ivoire). *European Scientific Journal, ESJ*, 12(23), 74–74. <https://doi.org/10.19044/ESJ.2016.V12N23P74>
- Balme, M., Galle, S., & Lebel, T. (2005). Démarrage de la saison des pluies au Sahel : variabilité aux échelles hydrologique et agronomique, analysée à partir des données EPSAT-Niger. *Sécheresse*, 16(1), 15–22.
- Bigot, S., Brou, Y. T., Diédhiou, A., & Houndenou, C. (2005). Facteurs de la variabilité pluviométrique en Côte d'Ivoire et relations avec certaines modifications environnementales. *Sécheresse*, 16(1), 14–21.
- Brou, Y. T. (1997). Analyse et dynamique de la pluviométrie dans le Sud forestier ivoirien : recherche de corrélations entre les variables climatiques et les variables liées aux activités anthropiques. Thèse de Doctorat, Université de Cocody (Côte d'Ivoire).
- Brou, Y. T. (2009). Impacts des modifications bioclimatiques et des l'aménagement des terres forestières dans les paysanneries ivoiriennes: quelles solutions pour une agriculture durable en côte d'ivoire. *Cuadernos Geograficos*, 45, 13–29.
- Brou, Y. T. (2010). Variabilité climatique, déforestation et dynamique agrodémographique en Côte d'Ivoire. *Sécheresse*, 21(1), 1–6. <https://doi.org/doi: 10.1684/sec.2010.0277>
- Brou, Y. T., & Chaléard, J. L. (2007). Peasant visions of environmental in Ivory Coast. *Annales de Géographie*, 116(653), 65–87. <https://doi.org/10.3917/AG.653.0065>
- Camberlin, P., Okoola, R., Diop, M., & Valimba, P. (2003). Identification des dates de démarrage et de fin de saison des pluies : applications a l'Afrique de l'est et au Sénégal. *Publication de l'Association Internationale de Climatologie*, 15, 295–303.
- Dekoula, C. S., Kouame, B., Goran, K. E. N., Yao, G. F., Kassin, K. E., Kouakou, J. B., Guessan, E. B. N., & Soro, N. (2018). Variabilité des descripteurs pluviométriques intrasaisonniers à impact agricole dans le bassin cotonnier de Côte d'Ivoire : cas des zones de Boundiali, Korhogo et Ouangolodougou. *Journal of Applied Biosciences*, 130, 13199–13212. <https://doi.org/https://dx.doi.org/10.4314/jab.v130i1.7> RÉSUMÉ
- Dekoula, C. S., Kouame, B., N'goran, E. K., Yao, F. G., Ehounou, J.-N., & Soro, N. (2018). Impact De La Variabilité Pluviométrique Sur La Saison Culturelle Dans La Zone De Production Cotonnière En Côte d'Ivoire. *European Scientific Journal, ESJ*, 14(12), 143. <https://doi.org/10.19044/esj.2018.v14n12p143>
- Didi, S. R., Ly, M., Kouadio, K., Bichet, A., Diedhiou, A., Coulibaly, H. S. J., Kouadio, K. C.

- A., Coulibaly, T. J. H., Salomon, O., & Savané, I. (2020). Using the CHIRPS Dataset to Investigate Historical Changes in Precipitation Extremes in West Africa. *Climate*, 8(84), 1–26. <https://doi.org/10.3390/cli8070084>
- Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A., & Michaelsen, J. (2015). The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes. *Scientific Data*, 2(1), 1–21. <https://doi.org/10.1038/sdata.2015.66>
- Giráldez, L., Silva, Y., Zubieta, R., & Sulca, J. (2020). Change of the rainfall seasonality over central peruvian andes: Onset, End, Duration and its relationship with large-scale atmospheric circulation. *Climate*, 8(2), 23. <https://doi.org/10.3390/CLI8020023>
- Goula Bi, T. A., Soro, G. E., Dao, A., Kouassi, F. W., & Srohourou, B. (2010). Frequency analysis and new cartography of extremes daily rainfall events in Côte d'Ivoire. *Journal of Applied Sciences*, 10(16), 1684–1694.
- Hodam, S., Sarkar, S., Marak, A. G. R., Bandyopadhyay, A., & Bhadra, A. (2017). Spatial Interpolation of Reference Evapotranspiration in India: Comparison of IDW and Kriging Methods. *Journal of The Institution of Engineers (India): Series A*, 98(4), 511–524. <https://doi.org/10.1007/s40030-017-0241-z>
- INS (Institut National De la Statistique). (2014). *Statistique de la population du Recensement Général de la Population et de l'Habitat (RGPH 2014)*.
- Joseph, P. V., Eischeid, J. K., & Pyle, R. J. (1994). Interannual Variability of the Onset of the Indian Summer Monsoon and Its Association with Atmospheric Features, El Niño, and Sea Surface Temperature Anomalies. *Journal of Climate*, 7, 81–105.
- Kanga, H. K., & Assi Kaudjhis, J. P. (2016). La sécheresse dans le « Quart Nord-Est » de la Côte d'Ivoire: de la réalité climatique à la perception paysanne. *European Scientific Journal, ESJ*, 12(29), 214–231. <https://doi.org/10.19044/esj.2016.v12n29p214>
- Kouamé, B., Ehounou, J.-N., Kassin, K. E., Dekoula, C. S., Yao, G. F., N'goran, E. K., Kouakou, B. J., Koné, B., & Soro, N. (2018). Caractérisation Des Paramètres Agroclimatiques Clés De La Saison Culturelle En Zone De Contact ForêtSavane De Côte-d'Ivoire. *European Scientific Journal, ESJ*, 14(36), 243–259. <https://doi.org/10.19044/esj.2018.v14n36p243>
- Kouassi, A. M. (2007). Caractérisation d'une modification éventuelle de la relation pluie-débit et ses impacts sur les ressources en eau en Afrique de l'Ouest: cas du bassin versant du N'zi (Bandama) en Côte d'Ivoire. Thèse de Doctorat, Université de Cocody (Côte d'Ivoire).

- Lienou, G. (2007). Impacts de la variabilité climatique sur les ressources en eau et les transports de matières en suspension de quelques bassins versants représentatifs au Cameroun. Thèse de Doctorat, Université Yaounde I (Cameroun).
- Marengo, J. A., Liebmann, B., Kousky, V. E., Filizola, N. P., & Wainer, I. C. (2001). Onset and End of the rainy season in the Brazilian Amazon Basin. *Journal of Climate*, 14, 833–852.
- Marteau, R., Sultan, B., Moron, V., Baron, C., Traore, S. B., & Alhassane, A. (2010). Démarrage de la saison des pluies et date de semis du mil dans le sud-ouest du Niger. *23ième Colloque de l'Association Internationale de Climatologie*, 379–384.
- Ndong, J. B. (2003). Caractérisation De La Saison Des Pluies Dans Le Centre-Ouest Du Senegal. *Publication de l'Association Internationale de Climatologie*, 15 (figure 1), 326–332. http://www.climato.be/aic/colloques/actes/PubAIC/art_2003_vol15/Article_40_JB_Ndong.pdf
- Nicholson, S., Kim, J., & Hoopingarner, J. (1988). *Atlas of African rainfall and its interannual variability*. Florida State University Tallahassee.
- Noufé, D. (2011). Changements hydroclimatiques et transformations de l'agriculture : l'exemple des paysanneries de l'Est de la Côte d'Ivoire. Thèse de Doctorat, Université Paris 1 Panthéon-Sorbonne (France).
- Noufé, D. (2012). Changement climatique et agriculture : cas de l'Est ivoirien. (*Path*) https://www.researchgate.net/publication/285055834_Changement, 2–3.
- Noufé, D., Lidon, B., Mahé, G., Servat, E., Yao, T. B., Zueli, K. B., & Chaleard, J. L. (2011). Climate variability and rainfed maize production in the eastern Ivory Coast: Variabilité climatique et production de maïs en culture pluviale dans l'est ivoirien. *Hydrological Sciences Journal*, 56(1), 152–167. <https://doi.org/10.1080/02626667.2010.545247>
- Noufé, D., Mahé, G., Kamagaté, B., Servat, E., Goula, B. A., Tié, B., & Savané, I. (2015). Climate change impact on agricultural production: the case of Comoé River basin in Ivory Coast. *Hydrological Sciences Journal*, 60(11), 1972–1983. <https://doi.org/10.1080/02626667.2015.1032293>
- Ozer, P., & Erpicum, M. (1995). Méthodologie pour une meilleure représentation spatio-temporelle des fluctuations pluviométriques observées au Niger depuis 1905. *Sec*, 6(103), 8.
- Paturel, J.-E., Servat, E., Kouame, B., & Boyer, J.-F. (1995). Manifestation de la sécheresse en Afrique de l'Ouest non sahélienne. Cas de la Côte d'Ivoire, Togo et du Bénin. *Sécheresse*, 6(1), 95–102.

- Seregina, L. S., Fink, A. H., Linden, R. van der, & Elagib, N. A. (2018). A new and flexible rainy season definition : Validation for the Greater Horn of Africa and application to rainfall trends. *International Journal of Climatology*, 1–4. <https://doi.org/10.1002/joc.5856>
- Servat, E., Paturel, J.-E., Lubes-Niel, H., Kouame, B., Travaglio, M., & Marieu, B. (1997). Variabilité climatique en Afrique humide le long du golfe Guinée, Première partie : Analyse détaillée du phénomène en Côte d'Ivoire. *Journal of Hydrology*, 191, 1–15.
- Sie, K., Coulibaly, T. J. H., Coulibaly, N., Savane, I., Gone, L. D., Kouadio, K. C. A., Coulibaly, H. S. J. P., Cissé, S., Camara, I., & Sylla, G. (2021). Contribution of satellite imagery to the characterization of the relationship between climate and pyrological variables of bush fires in the savannah zone (case of the bounkani region). *Environment & Ecosystem Science*, 5(1), 64–72. <https://doi.org/10.26480/ees.01.2021.64.72>
- Sivakumar, M. V. K. (1988). Predicting rainy season potential from the onset of rains in Southern Sahelian and Sudanian climatic zones of West Africa. *Agricultural and Forest Meteorology*, 42(4), 295–305. [https://doi.org/10.1016/0168-1923\(88\)90039-1](https://doi.org/10.1016/0168-1923(88)90039-1)
- Soro, T. D., Soro, N., Oga, Y. M.-S., Lasm, T., Soro, G., Ahoussi, K. E., & Biémi, J. (2011). La variabilité climatique et son impact sur les ressources en eau dans le degré carré de Grand-Lahou (Sud-Ouest de la Côte d'Ivoire). *Physio-Géo*, 5, 55–73. <https://doi.org/10.4000/physio-geo.1581>
- Traore, S. B., Reyniers, F. N., Vaksman, M., Kone, B., Sidibé, A., Yorote, A., & Yattara, K. (2000). Adaptation à la sécheresse des ecotypes locaux de sorghos du Mali. *Science et Changements Planétaires/Sécheresse*, 11(4), 227–237.

Spatio-temporal variability of the onset and end dates of the rainy seasons in the Bounkani region (North-East of Côte d'Ivoire).

ORIGINALITY REPORT

31%	26%	28%	8%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	ijpsat.org Internet Source	3%
2	repositorio.igp.gob.pe Internet Source	2%
3	pdffox.com Internet Source	1%
4	link.springer.com Internet Source	1%
5	article.sciencepublishinggroup.com Internet Source	1%
6	S. H. Franchito, V. Brahmananda Rao, M. A. Gan. "Onset and end of the rainy season and corn yields in São Paulo State, Brazil", <i>Geofísica Internacional</i> , 2010 Publication	1%
7	vital.seals.ac.za:8080 Internet Source	1%
8	www2.mdpi.com Internet Source	1%
9	www.mdpi.com Internet Source	1%
10	"Agricultural Adaptation to Climate Change", Springer Nature, 2016 Publication	1%
11	"Handbook of Climate Change Resilience", Springer Science and Business Media LLC, 2020 Publication	1%
12	www.omicsonline.org Internet Source	<1%

13	m.scrip.org Internet Source	<1 %
14	M. González, V. Barros. "On the forecast of the onset and end of the convective season in the Amazon", Theoretical and Applied Climatology, 2002 Publication	<1 %
15	journalijdr.com Internet Source	<1 %
16	www.researchsquare.com Internet Source	<1 %
17	docplayer.net Internet Source	<1 %
18	Yaakov Aviad, Haim Kutiel, Hanoch Lavee. "Analysis of beginning, end, and length of the rainy season along a Mediterranean–arid climate transect for geomorphic purposes", Journal of Arid Environments, 2004 Publication	<1 %
19	ijcrar.com Internet Source	<1 %
20	Koffi Claude Alain Kouadio, Ernest Amoussou, Talnan Jean Honoré Coulibaly, Arona Diedhiou et al. "Analysis of hydrological dynamics and hydropower generation in a West African anthropized watershed in a context of climate change", Modeling Earth Systems and Environment, 2020 Publication	<1 %
21	eujournal.org Internet Source	<1 %
22	static1.1.sqspcdn.com Internet Source	<1 %
23	Submitted to Charles Sturt University Student Paper	<1 %
24	Cissé, Soukèye, Laurence Eymard, Catherine Otllé, Jacques Ndione, Amadou Gaye, and Françoise Pinsard. "Rainfall Intra-Seasonal	<1 %

Variability and Vegetation Growth in the Ferlo Basin (Senegal)", Remote Sensing, 2016.

Publication

25	ijoeear.com Internet Source	<1 %
26	meteoguinebissau.gw Internet Source	<1 %
27	www.climato.be Internet Source	<1 %
28	Atta Sanoussi, Ly Mohamed, Salack Seyni, Alan George David. "Adapting to climate variability and change in smallholder farming communities: A case study from Burkina Faso, Chad and Niger", Journal of Agricultural Extension and Rural Development, 2015 Publication	<1 %
29	mafiadoc.com Internet Source	<1 %
30	repository.pauwes-cop.net Internet Source	<1 %
31	Abdelaziz EL-BOUHALI, Mhamed AMYAY, Khadija EL OUAZANI ECH-CHAHDI. "Recent variations of water area in the Middle Atlas lakes (Morocco): A response to drought severity and land use changes", Research Square Platform LLC, 2023 Publication	<1 %
32	Pierre Camberlin, Joseph Boyard-Micheau, Nathalie Philippon, Christian Baron, Christian Leclerc, Caroline Mwongera. "Climatic gradients along the windward slopes of Mount Kenya and their implication for crop risks. Part 1: climate variability", International Journal of Climatology, 2014 Publication	<1 %
33	ijramr.com Internet Source	<1 %
34	www.ijarp.org Internet Source	<1 %

35 Kapoury Sanogo, Joachim Binam, Jules Bayala, Grace B. Villamor, Antoine Kalinganire, Soro Dodiomon. "Farmers' perceptions of climate change impacts on ecosystem services delivery of parklands in southern Mali", Agroforestry Systems, 2016

Publication

<1 %

36 Submitted to University of Energy and Natural Resources

Student Paper

<1 %

37 agritrop.cirad.fr

Internet Source

<1 %

38 A. Bodian, O. Ndiaye, H. Dacosta. "Evolution des caractéristiques des pluies journalières dans le bassin versant du fleuve Sénégal: Aavant et après rupture", Hydrological Sciences Journal, 2016

Publication

<1 %

39 Kokeb Zena Besha, Tamene Adugna Demessie, Fekadu Fufa Feyessa. "Spatiotemporal variability and trend detection of hydrological and climatic variables of Modjo catchment, central Ethiopia", Theoretical and Applied Climatology, 2023

Publication

<1 %

40 Dioumacor Faye, François Kaly, Abdou Lahat Dieng, Dahirou Wane et al. "Regionalization of the Onset and Offset of the Rainy Season in Senegal Using Kohonen Self-Organizing Maps", Atmosphere, 2024

Publication

<1 %

41 Fitsum Bekele, Diriba Korecha, Lisanework Negatu. "Demonstrating Effect of Rainfall Characteristics on Wheat Yield: Case of Sinana District, South Eastern Ethiopia", Agricultural Sciences, 2017

Publication

<1 %

42 Kwadwo Owusu, Peter R. Waylen. "The changing rainy season climatology of mid-

<1 %

Ghana", Theoretical and Applied Climatology, 2012

Publication

43	www.ijser.org Internet Source	<1 %
44	www.jmaterenvironsci.com Internet Source	<1 %
45	"Wadi Flash Floods", Springer Science and Business Media LLC, 2022 Publication	<1 %
46	123dok.net Internet Source	<1 %
47	Submitted to CVC Nigeria Consortium Student Paper	<1 %
48	J. V. Revadekar. "Statistical analysis of the relationship between summer monsoon precipitation extremes and foodgrain yield over India", International Journal of Climatology, 03/15/2012 Publication	<1 %
49	Kimmo Ruosteenoja, Jouni Räisänen, Pentti Pirinen. "Projected changes in thermal seasons and the growing season in Finland", International Journal of Climatology, 2011 Publication	<1 %
50	archiv.ub.uni-heidelberg.de Internet Source	<1 %
51	assets.researchsquare.com Internet Source	<1 %
52	cp.copernicus.org Internet Source	<1 %
53	geology-dnu.dp.ua Internet Source	<1 %
54	m.moam.info Internet Source	<1 %
55	Amine Tahiri, Fouad Amraoui, Mohamed Sinan, Lhoussaine Bouchaou, Faouzi Berrada, Khalid Benjmel. "Influence of climate	<1 %

variability on water resource availability in the upper basin of Oum-Er-Rabiaa, Morocco", Groundwater for Sustainable Development, 2022

Publication

-
- 56 Argemiro Teixeira Leite-Filho, Verônica Yameê Sousa Pontes, Marcos Heil Costa. "Effects of Deforestation on the Onset of the Rainy Season and the Duration of Dry Spells in Southern Amazonia", Journal of Geophysical Research: Atmospheres, 2019 <1 %

Publication

-
- 57 Brou Dibi, Arthur Brice Konan-Waidhet, Valérie Plagnes, Jean Biemi. "Development of a Model for Assessing Vulnerability to Pollution of Groundwater in Fissured Aquifers: The Case of the Ehania Watershed (South-Eastern Côte d'Ivoire)", Journal of Environmental Protection, 2021 <1 %

Publication

-
- 58 Daouda Nsangou, Zakari Mfonka, Amidou Kpoumié, Paulin Sainclair Kouassy Kalédjé et al. "Analysis of rainfall variability and impact on the start and end dates of rainy seasons in the urban humid tropical zone: a case of the Yaoundé Town, Cameroon (Central Africa)", Arabian Journal of Geosciences, 2024 <1 %

Publication

-
- 59 I. N. Smith, L. Wilson, R. Suppiah. "Characteristics of the Northern Australian Rainy Season", Journal of Climate, 2008 <1 %

Publication

-
- 60 Koffi Emmanuel KASSIN, Brou KOUAME, Klotioloma COULIBALY, Klotioloma COULIBALY et al. "Prospects for sustainable cocoa farming from the rainfall balance in the last thirty years at Lh-Djiboua and Gh post-pioneer regions, Cte d'Ivoire", Journal of Soil Science and Environmental Management, 2018 <1 %

Publication

61	Muhire, I., and F. Ahmed. "Spatio-temporal trend analysis of precipitation data over Rwanda", South African Geographical Journal, 2014. Publication	<1 %
62	Submitted to University of South Africa Student Paper	<1 %
63	markhamgeog336.blogspot.com Internet Source	<1 %
64	rjas.ro Internet Source	<1 %
65	www.europeanjournalofscientificresearch.com Internet Source	<1 %
66	www.ijsr.net Internet Source	<1 %
67	Daniel Berhanu, Tena Alamirew, Meron Teferi Taye, Degefie Tibebe, Solomon Gebrehiwot, Gete Zeleke. "Evaluation of CMIP6 models in reproducing observed rainfall over Ethiopia", Journal of Water and Climate Change, 2023 Publication	<1 %
68	Guan, Kaiyu, Eric F. Wood, David Medvigy, John Kimball, Ming Pan, Kelly K. Caylor, Justin Sheffield, Xiangtao Xu, and Matthew O. Jones. "Terrestrial hydrological controls on land surface phenology of African savannas and woodlands", Journal of Geophysical Research Biogeosciences, 2014. Publication	<1 %
69	Kolotioloma Alama Coulibaly, Pauline Agoh Dibi-Anoh, Bi Néné Jules Tah, Hervé Anoh et al. "Occurrence of Extreme Rainfall and Flood Risks in Yopougon, Abidjan, Southeast Côte d'Ivoire from 1971 to 2022", American Journal of Climate Change, 2024 Publication	<1 %
70	Piou Dobo Guilavogui, Ibrahima Kalil Kante, Magbini Tokpa Mamy. "Indicators of Spatio-Temporal Changes in Rainfall in Forest	<1 %

Guinea", Atmospheric and Climate Sciences, 2025

Publication

71	Valentin Brice Ebodé, Jean Guy Dzana, Elias Nkiaka, Bernadette Nka Nnomo, Jean Jacques Braun, Jean Riotte. "Effects of climate and anthropogenic changes on current and future variability in flows in the So'o River Basin (south of Cameroon)", Hydrology Research, 2022 Publication	<1 %
72	ageconsearch.umn.edu Internet Source	<1 %
73	ftp.academicjournals.org Internet Source	<1 %
74	hal-insu.archives-ouvertes.fr Internet Source	<1 %
75	hal.ird.fr Internet Source	<1 %
76	ijlrhss.com Internet Source	<1 %
77	larhyss.net Internet Source	<1 %
78	pubs.sciepub.com Internet Source	<1 %
79	scirp.org Internet Source	<1 %
80	spiral.imperial.ac.uk Internet Source	<1 %
81	www.innspub.net Internet Source	<1 %
82	"Climate Change and Water Resources in Africa", Springer Science and Business Media LLC, 2021 Publication	<1 %
83	"Congo Basin Hydrology, Climate, and Biogeochemistry", Wiley, 2022 Publication	<1 %

84 Adama Bamba, N'Datchoh E. Toure, Kouakou Kouadio, Stéphane A. A. Ahoua et al. "Contribution of Climate Scenarios RCP4.5 and RCP8.5 to the Study of Climate Change Impacts on Cocoa Farming in Côte d'Ivoire", Atmospheric and Climate Sciences, 2023
Publication

<1 %

85 Ha Pham-Thanh, Tan Phan-Van, Andreas H. Fink, Roderick Linden. " Rainy Season Onset Detection: A New Approach Based on Principal Component Analysis and its Application to Vietnam ", International Journal of Climatology, 2021
Publication

<1 %

86 Jean-Luc Kouassi, Amos Gyau, Lucien Diby, Yeboi Bene, Christophe Kouamé. "Assessing Land Use and Land Cover Change and Farmers' Perceptions of Deforestation and Land Degradation in South-West Côte d'Ivoire, West Africa", Land, 2021
Publication

<1 %

87 Jean-Noël Ehounou, Brou Kouamé, Mathias G. Tah, Emmanuel K. Kassim et al. "Global Warming Influence on Major Seasonal and Intra-seasonal Rainfall Indicators for Sustainable Cocoa Production in West-central Côte d'Ivoire", Journal of Experimental Agriculture International, 2019
Publication

<1 %

88 Konan Jean-Yves N'Guessan, Botou Moise Adahi, Arthur Brice Konan-Waidhet, Barbara Sidoine Abonoua Kouassi et al. "A vulnerability assessment of rice farmers to climate change in Yamoussoukro, central Côte d'Ivoire", Environmental Development, 2025
Publication

<1 %

89 Mohammed El Hafyani, Narjisse Essahlaoui, Ali Essahlaoui, Meriame Mohajane, Anton Van Rompaey. "Generation of climate change

<1 %

scenarios for rainfall and temperature using
SDSM in a Mediterranean environment: a
case study of Boufakrane river watershed,
Morocco", Journal of Umm Al-Qura University
for Applied Sciences, 2023

Publication

90	Namwinwelbere Dabire, Eugene Ezin, Adandedji Firmin. "Determination of Hydrological Flood Hazard Thresholds and Flood Frequency Analysis: Case Study of Nokoue Lake Watershed", Water, 2025 Publication	<1 %
91	climatologie.u-bourgogne.fr Internet Source	<1 %
92	horizon.documentation.ird.fr Internet Source	<1 %
93	innspub.net Internet Source	<1 %
94	iwaponline.com Internet Source	<1 %
95	orbi.uliege.be Internet Source	<1 %
96	wedocs.unep.org Internet Source	<1 %
97	www.journalijar.com Internet Source	<1 %
98	www.riges-uao.net Internet Source	<1 %
99	www.scielo.br Internet Source	<1 %
100	www.scirp.org Internet Source	<1 %
101	M. Beyer, M. Wallner, L. Bahlmann, V. Thiemig, J. Dietrich, M. Billib. "Rainfall characteristics and their implications for rain-fed agriculture: a case study in the Upper Zambezi River Basin", Hydrological Sciences Journal, 2015	<1 %

102 "Water Resources in Arid Areas: The Way Forward", Springer Science and Business Media LLC, 2017 <1 %
Publication

103 Artadji Attoumane, Dos Santos Stéphanie, Modeste Kacou, Alla Della André et al. "Individual perceptions on rainfall variations versus precipitation trends from satellite data: An interdisciplinary approach in two socio-economically and topographically contrasted districts in Abidjan, Côte d'Ivoire", International Journal of Disaster Risk Reduction, 2022 <1 %
Publication

104 Handbook of Climate Change Adaptation, 2015. <1 %
Publication

105 Jean-Noël Ehounou, Brou Kouamé, Mathias G. Tah, Emmanuel K. Kassin et al. "Impact of Local Global Warming on Rainfall and Annual Cocoa Water Requirements in the Regions of Loh-Djiboua and Goh in West-central Côte d'Ivoire", International Journal of Environment and Climate Change, 2019 <1 %
Publication

106 Kouamé Donald Kouman, Amos T. Kabobah, Boyossoro Hélène Kouadio, Komlavi Akpoti. "Spatio-Temporal Trends of Precipitation and Temperature Extremes across the North-East Region of Côte d'Ivoire over the Period 1981–2020", Climate, 2022 <1 %
Publication
