

An update on the current status of House Sparrow, *Passer domesticus*: A bird with uncertain future

Abstract

The house sparrow, commonly called *goraiya*, is a tiny brown bird with black streaks running down its back that is frequently seen in metropolitan settings. Due to their adaption to human settlements, they have spread over the world from their Middle Eastern origins. Nonetheless, their numbers have drastically decreased globally, especially in metropolitan suburbs. The sparrow population in India has plummeted, with declines of up to 70% in some locations. An ornithological survey conducted by the Indian Council of Agricultural Research (ICAR) unveiled a drastic decline of 80% in Andhra Pradesh alone, 20% in Kerala, Gujarat, and Rajasthan, and followed by 70-80% in coastal regions. House sparrow populations in Tamil Nadu, Andhra Pradesh, Karnataka, and Uttarakhand are declining due to environmental stressors, vegetation, and transition from automobiles to horse-drawn transportation. Reproductive potential is influenced by insectivorous food availability, while electromagnetic radiation and power lines pose health deterioration and microwave sickness.

Having status of least concern in IUCN, house sparrow is still experiencing a rapid decline in its population at global scale. To protect sparrow populations several national measures and conservation regulations are being implemented. House sparrows have been designated as the “State Bird of Delhi” and the “State Bird of Bihar”, making India the largest sparrow conservation initiative. The current review seeks suggestions for providing alternative nesting sites for House Sparrows, as this could potentially restore them to our lives and allow future generations to enjoy these tiny, singing birds in their homes.

Key words: population, reason for declining, conservation strategies.

Introduction

The house sparrow is a widespread and easily recognizable bird commonly found in metropolitan areas. Also known as the *Goraiya*, it is a small, stocky passerine measuring 14–16 cm in length, weighing 26–32 grams, and with a wingspan of 19–25 cm. belonging to the family Passeridae, it exhibits clear sexual dimorphism between males and females. Considered the world’s most abundant and widely distributed bird species (Anderson, 2006), the house sparrow originated in the Middle East and has since expanded across much of the globe. Its range includes large parts of Asia-particularly the Indian subcontinent-as well as Europe and North Africa (Figure 1). Within the Indian subcontinent, the species has also been introduced to the Andaman Islands, Pakistan, Bangladesh, Sri Lanka, and the Maldives (Ali & Ripley, 1983).

Of the 32 sparrow species recorded worldwide, five are found in India: the House, Spanish, Sind, European Tree, and Russet sparrows. Among these, *Passer domesticus* is the most widely distributed species across the country (Teotia et al., 2018). Its remarkable ability to adapt quickly to human-dominated landscapes has facilitated its widespread establishment (Dandapat et al., 2010). Despite its small size, the house sparrow holds great ecological significance, acting as an indicator of environmental balance. Closely associated with human settlements, it inhabits both urban and rural areas and thrives across diverse climates and habitats, though it generally avoids dense forests, vast grasslands, and deserts.

These adaptive traits make the house sparrow an important model species for studying biological questions such as evolutionary mechanisms, thermal metabolism, and pest control (Kumar et al., 2015). Its diet primarily consists of weed and cereal seeds, supplemented by insects and other small food items. However, sparrow populations have declined drastically in many parts of the world, particularly along the urban–suburban gradient (Chamberlain et al., 2007). In some regions of India, populations have decreased by nearly 70% (Dandapat et al., 2010). Once abundant in cities, towns, and villages, sparrows are now disappearing more rapidly from urban centers compared to rural areas. One proposed factor contributing to this decline is electromagnetic radiation in urban environments (Balmori&Hallberg, 2007).




Figure 1. House sparrow range map: green= native, yellow= introduced (map from Wikimedia Commons)

The genus passer

Summers-Smith's 1988 update on the genus *Passer*, also known as Old World sparrows, includes 20 species found primarily in Africa, Europe, and Asia, with his findings heavily cited in this narration (Table 1). Summers-Smith (1988) divided the house sparrow genus into two subgenera and three superspecies groups, including the Spanish sparrow (*P. hispaniolensis*), the Somali sparrow (*P. castanopterus*), and the desert sparrow (*P. simplex*), which are part of the same superspecies group (Anderson 2006).

Description

Compared to female house sparrows, males often weigh more and are bigger (Hanson et al., 2020b). Male *Passer* species have black stripes, gray crests, and white markings, while females have duller plumage with dark brown crests and light brown lines behind the eyes. Females lack the black marks on their heads and have gray-brown to light brown cheeks, beaks, and feathers. Their morphology resembles juveniles and females of other *Passer* species, making it difficult to identify them (Anderson, 2006). Subspecies varied in respect to bulk, size, and male plumage as well (Summers-Smith, 1988). The male sparrow's huge black neck patch is the most obvious physical difference between it and the female. The species' status as a model in behavioural ecology research may have been influenced by this characteristic marking (Sanchez-Tojar et al., 2018).

Diet and Scavenging

After hatching, fledgling house sparrows mostly eat insects during the first three days. They then often graze more on cereals following fledging, particularly in rural regions (Anderson, 2006). Adult house sparrows make most of their year-round food opportunistically, especially in urban and suburban regions where there is an abundance of human garbage (Summers-Smith, 1988). A potential cause for house sparrows' skill in taking advantage of a variety of meals might be the release of digestive enzymes with flexibility (Brzęk et al., 2009). The species is used as a model for range expansion due to behavioural reactions to feeding. House sparrows from an older invading population in New Jersey consume unusual foods less quickly than those from a 40-year-old Panama population. An inclination to explore novel meals may be advantageous for birds in circumstances where supplies are limited or unpredictable. But there are hazards associated with this activity as well, such as the immune system being triggered by tainted food or exposure to foreign poisons (Martin & Fitzgerald, 2005). It has been found that populations vary substantially in the composition of their immune systems and the parasites they harbor throughout their entire lives, lending credence to this idea (Kilvitis et al., 2019; Martin et al., 2015; Martin et al.,

87 2014; Coon and Martin, 2014; Coon et al., 2014),


88

89 **Table 2. House sparrow of India**

Name	Description	Distribution	References
<i>Passer hispaniolensis</i> (Spanish sparrow)	16cm, a chestnut and black sparrow with a stout bill. Sexes are dimorphic.	Erratic winter visitor to plains of Northwest India mainly east to Haryana and south to Rajasthan. Vagrant in Uttar Pradesh.	Modak, BK 2017
<i>Passer pyrrhonotus</i> (Sind sparrow)	12cm, slim, brown and grey sparrow with neat, rectangular black bib. Sexes are dimorphic.	Local breeding restricted to areas of Punjab and Haryana.	IrshadChandio et al. 2019.
<i>Passer rutilans</i> (Russet sparrow)	15cm, a bright chestnut sparrow with usually yellowish-buff underparts. Sexes are dimorphic.	Locally common breeding resident in northern mountains from north Pakistan to east Myanmar.	Naithani, A et al. 2012
<i>Passer montanus</i> (Eurasian Tree sparrow)	15cm, brown and buff sparrow with a rounded, chest-nut crown. Sexes alike.	Locally common breeding resident in Uttaranchal.	Kumar, A et al. 2015
<i>Passer domesticus</i> (house sparrow)	15cm, brown and grey sparrow with stout bill. Sexes are dimorphic.	Common breeding resident throughout plains and foothills. Rare in parts of the Northwest and the Northeast.	Swami, R and Kumar, M 2022

90

91



109

Figure 2. (A) Female and (B) Male House Sparrow(Images from Wikimedia commons)

Behaviour

Housesparrowsareoutgoingbirdsthatroostingroupsandarerecognizablethroughouttheyear.Their nest-buildingpreferencesarediverse;theyoftenchoosetoconstructtheirhomesincracksandcrevices in cliffs and banks, residential rooftops, and streetlights. Their food sources include insects, grains, seeds, flower nectar, and, in urban areas, leftover human food. Limited migration is conceivable in many parts of the world.

The breeding season in northern India extends from March to June. In central India, it is from September to October and in southern India, it is observed all year round (Dandapat et al., 2010). While not in the breeding season, they come together as massive flocks; otherwise,theyare seenincouples orsmall groups. Despite some

assistance from the female, the male chooses the nest site and builds the majority of the nest. March– April is when nesting starts (Dandapat et al., 2010). Male house sparrows fiercely defend their nesting grounds throughout the mating season, and both sexes actively assist in rearing the young. These birds live for twelve years on average. In order to find food on the ground and protect their nesting grounds, house sparrows employ wing flapping and skipping. To attract possible mates, fend off intruders, and notify others, they communicate through postures, gestures, and vocalizations, maintaining a safe and secure habitat (Paul, 2015).

Breeding Biology

House sparrows establish during matities and are monogamous. Although they have several broods, it is uncommon for more than two of them to succeed. Males select a nest site and remain there while they advertise by calling, which is a quiet chirrup that they repeat once every one to two seconds. They may also shiver their wings and raise their tails. If a falcon approaches and expresses interest in the nest, the male gets excited and jumps around, holding his head up and holding his wings out slightly. During this period, the male enters and exits the nest site, occasionally blocking the female's entry for two to three days. Males may offer nest material to females and even touch bills during the construction of the nest; bill contact also triggers copulations. Five to six eggs, profusely powdered, speckled, or blotched with black, brown, or ash-grey, are laid by the house sparrow on a creamy white or blue-tinted background. Every bird takes between 10 and 12 days to incubate. Older birds breed earlier in the season, which is the only way that their reproductive success rises with age. 4-5 eggs, ranging from 1 to 8, typically make up a clutch (Dandapat et al., 2010).



Figure 3. Potential nesting sites of the house sparrow in India (Images from Wikimedia commons)



Figure 4. (A) Birds having sunbath and rest (B) Birds perching on wire and Socializing

Status in world

Approximately $1/8^{\text{th}}$ of the world's bird species are under threat of going extinct, meaning that 1,226 of the 9,856 bird species are closer to going extinct. Worldwide, the population of sparrows is dropping (Crick et al., 2002, Summers-Smith 2006). According to recent studies, the House Sparrow population is drastically declining in several regions of the world. The population of most of Europe is dwindling dramatically. In fact, this species is listed as endangered in the Netherlands. In the United Kingdom, population declines in rural regions have been 47% and in urban and suburban areas 60% since the mid-1970s (Robinson et al., 2005). Reports of this species' population reduction also exist for northwest Europe (Prowse, 2002). The House Sparrow is now included on the European Conservation Concern species as a result (Shaw et al., 2008). Special conservation concern (Red List) in Britain (Baillie et al. 2010).

According to the British Trust for Ornithology, between 1994 and 2002, there was a 71% decrease in the sparrow population in London. Other European cities that have observed a fall in sparrow populations include Edinburgh, Dublin, Glasgow, Hamburg, and Berlin (Dandapat et al., 2010). Based on an investigation conducted four years ago at Stanford University by conservation scientist Cagan Sekercioglu, around 10% of all species are expected to become extinct and an additional 15% may be in danger of doing so by the year 2100. As a result, critical processes like seed distribution, pollination, and decomposition are probably going to diminish (Dandapat et al., 2010). The species is classified as least concern on the 2019 IUCN Red List, and the population trend is currently declining.

Status in India

Over the past few decades, house sparrow populations have shown a marked decline across several regions of India, including West Bengal, Bangalore, Punjab, Rajasthan, Delhi, and Haryana (Bandel, 2010). Limited studies from Uttarakhand, Karnataka, Tamil Nadu, and Andhra Pradesh have also documented reductions in population (Ghosh et al., 2010; Balaji et al., 2013; Hussain et al., 2014; Paul, 2015; Naik, 2018; Deepalakshmi & Salomi, 2019; Sharma & Binner, 2020; Renukadevi et al., 2021). Research from Kerala further suggests that house sparrows may already be locally extinct in Trivandrum City (Raju, 2015).

In Andhra Pradesh alone, sparrow numbers have fallen by nearly 80%, while states such as Kerala, Gujarat, and Rajasthan have recorded declines of about 20%. Coastal regions appear to be most affected, with reductions as high as 70–80%, according to an ornithological assessment by the Indian Council of Agricultural Research. Multiple surveys carried out in different parts of the country also confirm this downward trend (Rajashekar & Venkatesha, 2008; Daniels, 2008; Khera et al., 2010; Bhattacharya et al., 2010; Ghosh et al., 2010). The Bombay Natural History Society (BNHS) has reported decreases in both nest numbers and overall sparrow populations. Supporting this, a joint survey by the Central Ministry of Environment and Forests and

BNHS revealed a sharp decline in sparrow numbers in the Hyderabad–Ranga Reddy region beginning around 2005 (Sharma & Binner, 2020).

Causes of decline

Many theories have been developed to explain why House Sparrow populations in urban and suburban areas may be declining, including predation, ecological factors, competition, a lack of nesting places, disease, a shortage of food, and pollution (Summers-Smith, 1999; Daniels, 2008). Several theories have been proposed as potential reasons for the decrease in House Sparrow populations in rural, urban, and suburban areas are:

Predation: House sparrows are a common food source for owls and hawks. These include Eurasian Sparrow Hawks (*Accipiter nisus*) and domestic cats (*Felis catus*). However, the domestic cat is a significant predator of house sparrows. Several snakes, raccoons, and domestic dogs are other house sparrow predators. Predators including black kites, shikra, and crows also attack the chicks (Sharma & Binner, 2020).

The lack of food: Insects and grains are both consumed by house sparrows. The introduction of pre-cleaned food grains has reduced the availability of these foods; washing grains in a garden has become uncommon. Thus, the ensuing grain spillage that these birds were pecking at have vanished. The decorative plants and crotons that sustain less insect life the primary food source for sparrows have supplanted the vegetable gardens in old world homes (Kumar et al., 2015). Uncontrolled Pigeons The primary competitors for food in the two urban environments are *Columba livia* and gulls *Larus*, especially Lesser Black-backed Gulls *L. fuscus*. However, it is doubtful that either species would have a distinct impact (Sharma & Binner, 2020).

Absence of nest sites: Hundreds of shrubs and trees are being cut down to make way for large structures. The lack of cracks and holes in the terraced structures, which are favoured over tiled homes, has resulted in a sharp decrease in house sparrow nesting locations (Kumar et al., 2015).

Diseases: In urban and suburban environments, house sparrows can serve as reservoirs of disease (Juricova et al., 1998). Among free-ranging wild sparrows, salmonella infections are particularly common during the winter and spring seasons (Macdonald, 1978). Infected birds often exhibit enlarged and congested livers and spleens, along with abscesses on the skin, muscles, and liver. Communal feeding practices—such as the use of garden bird tables and feeders—can further facilitate the spread of infection (Macdonald, 1978). As the disease progresses, affected birds become weaker and increasingly susceptible to secondary infections. Immunocompromised individuals are also at greater risk of contracting a variety of illnesses from such pathogens. Disease is now recognized as being influenced not only by parasite infection but also by the host's overall health status. This makes it difficult to separate the effects of food scarcity and environmental stressors from those of disease. Consequently, disease-related mortality is generally considered unlikely to contribute significantly beyond other existing causes of sparrow mortality (Newton, 1998).

Environmental Pollutants: Pesticides, industrial effluents, and combustion emissions are among the many harmful substances continually released into the environment. Several of these chemicals are now recognized as major factors contributing to bird population declines worldwide, as they influence distribution and abundance patterns (Newton, 1998). A study conducted in urban Warsaw, Poland, showed that mortality in House Sparrow nestlings was linked to the presence of polychlorinated biphenyls (PCBs) and chlorinated hydrocarbons, which enter the environment through industry, sanitation, and agriculture (Karolewski et al., 1991). Similarly, Romanowski et al. (1991) reported that concentrations of zinc and lead in the livers of sick and dead sparrow nestlings were higher than in healthy ones. Nestlings with stunted growth (low body weight) also had elevated levels of zinc, cadmium, and lead compared to those with normal growth. Summers-Smith (2006) further noted that pollutants such as MTBE, emitted from cars, can reduce populations of invertebrates like aphids, indirectly threatening sparrow nestling survival. Although MTBE levels in the UK are not yet monitored, there is currently no conclusive evidence that pollution alone is responsible for the decline of House Sparrow

247 populations.

2486. **Owing to Shifts in Human Lifestyle:** Contemporary lifestyles have brought about significant modifications to
249 housing architecture. Because terraced constructions are chosen over tiled homes because they lack gaps and
250 cracks, house sparrow nesting grounds have drastically decreased (Kumar et al., 2015).

2517. **Use of chemicals:** When pesticides and insecticides are used excessively in gardens and
252 agricultural fields, insects are killed. The mortality rate for chicks and yearlings can be elevated because of a deficiency
253 of insects high in protein and contamination of grains and insects they ingest (Kumar et al., 2015).

254 **Conservation strategies**

255 House sparrows, being opportunistic in nature, can thrive wherever there is adequate food, suitable nesting sites,
256 and safe roosting places. However, their numbers have recently declined in India, raising concern for their
257 conservation. To prevent urbanization from driving them toward the category of Critically Endangered, various
258 initiatives have been launched across the country. Since 2006, Mohammed Dilawar, through the Nature Forever
259 Society (NFS), has been working to protect common species such as sparrows. The society has introduced
260 impactful programs like the *Adopt a Feeder and Nest Program*, *World Sparrow Day*, *Common Bird Monitoring*
261 *of India*, *Project Save Our Sparrows*, and the *Nature Forever Society Sparrow Award*. These initiatives have
262 contributed to noticeable improvements in sparrow presence in urban areas. In collaboration with the Uttar
263 Pradesh State Biodiversity Board, the Biodiversity and Wildlife Protection Lab at the University of Lucknow
264 has also taken steps to spread awareness on sparrow conservation. One of their unique approaches has been
265 gifting artificial nest boxes as souvenirs to dignitaries during awareness campaigns (Sharma & Binner, 2020).

266 To further emphasize its cultural and ecological importance, the House Sparrow was declared the "State Bird of
267 Delhi" in 2012 by then Chief Minister Sheila Dikshit under the *Rise for the Sparrows* campaign. Following this,
268 in 2013, Bihar's Chief Minister Nitish Kumar also recognized the sparrow as the state bird. The NFS's first
269 celebration of *World Sparrow Day* in 2009 marked a milestone in spreading awareness about protecting
270 common birds and biodiversity. The aim was to bring together nature lovers and sparrow enthusiasts to foster a
271 global conservation movement.

272 Ultimately, creating alternative nesting opportunities for sparrows is a simple yet effective way humans can
273 contribute to their survival, ensuring that future generations may continue to experience the charm of these
274 small, melodious birds (Sharma & Binner, 2020).

275 **Discussion**

276 The decline in the *Passer domestic* (house sparrow) population in India has alarmed environmentalists and
277 specialists alike. This phenomenon has been connected to several changes in the environment and human
278 impacts. A primary cause of this decline is the evolution of human behavior, particularly the modifications
279 made to house architecture. Modern construction lacks the openings and cracks that
280 sparrows need to construct their nests; thus, they opt for terraced homes rather than classic tiled homes. Furthermore, the
281 misuse of chemicals such as herbicides and insecticides in gardens and agricultural areas has resulted in a major
282 decline in the availability of protein-rich insects, which are vital for the
283 diet of sparrow chicks and yearlings. Another factor contributing to the decline in sparrow populations is the widespread
284 use of catapults, which pose a direct threat to their numbers. Urbanization and the associated increase in
285 electromagnetic radiation from communication towers have also been linked to the decline in population.
286 House sparrows are being protected via national programs. The Nature Forever Society (NFS), founded by
287 Mohammed Dilawar in 2006, has been in charge of these
288 conservation efforts. Initiatives such as "Adopt a Feeder and Nest Program," "World Sparrow Day," and "Project Save
289 Our Sparrows" have positively impacted the population of sparrows in urban areas. The importance of sparrow
290 conservation has been highlighted by state-level initiatives like the recognition of the house sparrow as the
291 "State Bird" of Delhi and Bihar.

292

Conclusion

Tackling the decline of house sparrow populations in India demands joint efforts from researchers, conservationists, and the wider community. The primary factors driving this decline include rapid urbanization, intensive chemical usage, and modifications in modern building designs. Studies indicate that initiatives by state governments and organizations such as the Nature Forever Society can play a significant role in stabilizing sparrow populations through targeted conservation programs. Nonetheless, consistent and long-term action remains crucial, as sparrows are important indicators of ecological balance. Future research should focus on identifying additional contributing factors and formulating comprehensive conservation strategies that can be applied on a broader scale. Promoting public awareness and restoring habitats represent effective steps toward protecting sparrows, ensuring their survival for future generations, and reversing current population declines.

References

- Ali, S., & Ripley, S.D. (1983). Handbook of the birds of India and Pakistan. Compact edition. Oxford University Press and BNHS, Mumbai. Ali, S. and S.D. Ripley (1995). The Pictorial Guide to the Birds of Indian Sub-continent. Oxford University Press and BNHS, Mumbai. Baskaran, S.T. (1992). Sighting of Dusky Horned Owl. Newsletter for Birdwatchers, 32(9), 10.
- Anderson, T. R. (2006). *Biology of the ubiquitous house sparrow: from genes to populations*. Oxford University Press.
- Baillie S.R., Marchant J.H., Leech D.I., Renwick A.R., Joys A.C., Noble D.G., Barimore C., Conway G.J., Downie I.S., Risely K. and Robinson R.A. (2010): Breeding birds in the wider countryside: their conservation status 2010. BTO Research Report no. 565. BTO, Thetford, www.bto.org/birdtrends.
- Balaji S., Baskaran S., Rajan M.K. and Pavaraj M. 2013. Investigating the causes for the decline and strategies to conserve house sparrow, *Passer domesticus* in Sivakasi taluk, Virudhunagar district, Tamil Nadu, India. *World Journal of Zoology*, 8(3), 278-284.
- Balmori, A., & Hallberg, Ö. (2007). The urban decline of the house sparrow (*Passer domesticus*): a possible link with electromagnetic radiation. *Electromagnetic biology and medicine*, 26(2), 141-151.
- Bandel H. (2010) Proceedings of National Seminar on Biodiversity, Water resource and Climate change issues, Department of Environmental Science, Kalyani University, pp: 147-152.
- Bhattacharya, R., R. Roy, S. Ghosh & A. Dey (2011). Observations on house sparrow at Bandel, Hoogly,
- BirdLife International (BirdLife International). (2016b, October 1). IUCN Red List of Threatened Species: *Passer domesticus*. IUCN Red List of Threatened Species. <https://www.iucnredlist.org/species/103818789/155522130>
- Birdlife International, *Passer domesticus* (amended version of 2017 assessment). The IUCN Red List of Threatened Species (2018).
- Brzęk, P., Kohl, K., Caviedes-Vidal, E., & Karasov, W.H. (2009). Developmental adjustments of house sparrow (*Passer domesticus*) nestlings to diet composition. *Journal of Experimental Biology*, 212(9), 1284-1293.
- Chamberlain, D. E., Toms, M. P., Cleary-McHarg, R., & Banks, A. N. (2007). House sparrow (*Passer domesticus*) habitat use in urbanized landscapes. *Journal of Ornithology*, 148(4), 453-462.
- Coon, C.A.C. and Martin, L.B. (2014). Patterns of haemosporidian prevalence along a range expansion in introduced Kenyan House Sparrows *Passer domesticus*. *J. Avian Biol.*, 45, 34-42
- Coon, Courtney A.C., "Host-Parasite Interactions in an Invasive Songbird" (2014). USF Tampa Graduate Theses and Dissertations. <https://digitalcommons.usf.edu/etd/5004>
- Crick H.Q.P., Robinson R.A., Appleton G.F., Clark N.A. and Rickard A.D., Investigation into the causes of the decline of starlings and house sparrows in Great Britain. British Trust for Ornithology, Thetford (2002).

344 Dandapat,A., Banerjee, D., &Chakraborty, D. (2010). The case of the Disappearing House Sparrow (*Passer*
345 *domesticusindicus*). *Veterinary World*, 3(2), 97.

346 Daniels,R.J.R.(2008).Canwesavethesparrow?*CurrentScience*,95,1527–1528.

347

348 DeepalakshmiS.andSalomiA.A.2019.Impactofurbanizationonhousesparrow(*Passerdomesticus*) diversity from
349 Erode and Namakkal districts, Tamilnadu, India. *International Journal of Advanced Research in Biological*
350 *Sciences*, 6(11), 22-27. <http://doi.org/10.22192/ijarbs.2019.06.11.004>.

351

352 Galib.,Hussain, S., Zaheer, K. et al., (2013). An Overview of Occurrence, Distribution, and Status of the Birds
353 of Khirthar Protected Area Complex (KPAC), Sindh. *Canadian Journal of Pure and Applied Sciences*. 7, 2515-
354 2532.

355

356 Gedeon, Kai &Cauldwell, Andrew &Ewnetu, Mihret&Regassa, Fekede&Schönbrodt, Robert &Töpfer, Till.
357 (2015). House Sparrow *Passer domesticus* and hybrids with Somali Sparrow *P. castanopterus* in Ethiopia. *The*
358 *Bulletin of the African Bird Club*. 22. 71-73.

359

360 Ghosh S., Ki-Hyun K. and Bhattacharya R. 2010. A survey on house sparrow population decline at Bandel,
361 West Bengal, India. *Journal of the Korean earth science society*, 31(5), 448-453.

362

363 Hanson,H.E.,Zolik,J.E.,&Martin,L.B.(2020).HouseSparrow(*Passerdomesticus*Linnaeus,1758). In *Invasive*
364 *Birds: Global trends and impacts* (pp. 85-96). Wallingford UK: CABI.

365

366 Higgins, P.J. ; Peter, J.M. & Cowling, S.J. (editors) 2006. Handbook of Australian, New Zealand & Antarctic
367 Birds. Volume 7, Boatbill to starlings; Part 78, Dunnock to starlings. Melbourne, Oxford University Press. [Vol.
368 5, pages 51-55] Vol. 78, pages 1067-1089; plate 31.

369

370 HussainA.,DasguptaS.andBargaliH.S.2014.Caseofhousesparrow(*Passerdomesticus*)population decline: role of
371 semi nomadic pastoralist community (Van Gujjar) in their community. *International Journal of Conservation*
372 *Science*, 5(4), 493-502.

373

374 IrshadChandio, Ali MurtazaDharejo, MUHAMMAD MUNIF KHAN, SAIMA NAZ, 2019. New host record for
375 the *Diplotrienamonticolae* (Filariidae: Nematoda) from the thoracic cavity of *Passer pyrrhonotus*, Blyth 1845,
376 (*Passeridae: Passeriformes*) in Larkana, Sindh, Pakistan. *Pure and Applied Biology*, 8(1), 580-584.

377

378 Juricova,Z.,Pinowski,J.,Literak,I.,Hahm,K.,andRomanowski,J.,1998,AntibodiestoAlphaviruses, Flavivirus, and
379 BunyavirusArboviruses in House Sparrows and Tree Sparrows in Warsaw. *Avian Diseases*, 42(1),182-5.

380

381 Karolewski,M.,Lukowski,A., Pinowski,J., and Trojanowski,J., 1991, The International Symposium
382 oftheWorkingGrouponGranivorousBirds.Chlorinatedhydrocarbonsineggsandnestlingsof*Passer montanus* and
383 *Passer domesticus* from urban and suburban Warsaw-Preliminary Report. In Pinowski,
384 J.,Kavanagh,B.,andGorski,W.(eds.),Nestlingmortalityofgranivorousbirdsduetomicroorganisms and toxic
385 substances. Polish Scientific Warsaw, Poland, 189-196.

386

387 Khera,N.,A.Das,S.Srivatsava&S.Jain(2010).Habitat-wisedistributonofthehousesparrow(*Passer domesticus*) in
388 Delhi, India. *Urban Ecosystems*, 13, 147–153.

389

390 Kilvitis,H.J.,Schrey,A.W.,Ragsdale,A.K.,Berrio,A.,Phelps,S.M.,& Martin,L.B.(2019).DNA
391 methylationpredictsimmunegeneexpressioninintroducedhousesparrows*Passerdomesticus*.*Journal of Avian*
392 *Biology*, 50(6).

393

394 Kumar, A., Kanaujia, A., Kushwaha, S., & Kumar, A. (2015). Strategies and efforts for conservation of House
Sparrow (*Passer domesticus*) in India. *Natural Environment*, 20, 58-66.

395

396 Macdonald,J.,1978,CutaneousSalmonellosisinaHouseSparrow.*BirdStudy*,25,p.59.

397

398 Martin LB, LieblAL, Kilvitis HJ. 2015. Covariation in stress and immune gene expression in a range expanding
bird. *General and Comparative Endocrinology* 211:14–19. DOI: <https://doi.org/10.1016/j.ygcen.2014.11.001>,
PMID: 25448257.

399 Martin, L. B., & Fitzgerald, L. (2005). A taste for novelty in invading house sparrows, *Passer domesticus*.
400 *Behavioral Ecology*, 16(4), 702-707.

401 Martin, L. B., Coon, C.A., Liebl, A. L., & Schrey, A.W. (2014). Surveillance for microbes and range expansion in
402 house sparrows. *Proceedings of the Royal Society B: Biological Sciences*, 281(1774), 20132690.

403 Modak, B.K. (2017). Impact of Urbanization on House Sparrow Distribution: A Case Study from Greater
404 Kolkata, India. *Proceedings of the Zoological Society*, 70, 21–27 <https://doi.org/10.1007/s12595-015-0157-4>
405

406 Naik S.P. 2018. Rapid study of house sparrows (*Passer Domesticus*) in Moodbidri. conference on conservation
407 and sustainable management of ecologically sensitive regions in Western Ghats: 290-293.
408 <http://ces.iisc.ernet.in/energy>
409

410 Naithani, A., Bhatt, D. (2012). Bird community structure in natural and urbanized habitats along an altitudinal
411 gradient in Pauri district (Garhwal Himalaya) of Uttarakhand state, India. *Biologia*, 67, 800–808
412 . <https://doi.org/10.2478/s11756-012-0068-z>
413

414 Newton, I., 1998, Population Limitation in Birds. Academic Press Limited, USA, 147 p.
415

416 Päckert M, Hering J, Belkacem AA, Sun Y-H, Hille S, Lkhagvasuren D, Islam S, Martens J (2021) A revised
417 multilocus phylogeny of OldWorld sparrows (Aves: Passeridae). *Vertebrate Zoology*, 71: 353-
418 366. <https://doi.org/10.3897/vz.71.e65952>
419

420 Paul, M. R. (2015). A review of house sparrow population decline in India. *Asia Pacific Journal of Research*
421 *Vol: I. Issue XXIX*. pp. 147–152. Proceedings of National Seminar on Biodiversity, Water Resource and Climate
422 change Issues. Kalyani University, West Bengal.
423

424 Prowse A. (2002) The urban decline of the house sparrow. *British Birds* 95(3), 144-145.
425

426 Rajashekhar, S. & M.G. Venkatesha (2008). Occurrence of House Sparrow, *Passer domesticus indicus* in and
427 around Bangalore. *Current Science*, 94, 446–449.
428

429 Raju S. 2015. Population trend of the common birds in a residential area of Thiruvananthapuram city, Kerala.
430 *Indian Birds*, 10(2), 40-45.
431

432 Renukadevi K., Maxmillion M.E., Angayarkanni P.S. and Rajendran P. 2021. Exploratory data analytics to
433 study the impact of parameters to enhance the population of house sparrow (*Passer domesticus*). *Research*
434 *Square*, 1-21. <https://doi.org/10.21203/rs.3.rs-440060/v1>.

435 Robinson, R. A., Siriwardena, G. M., & Crick, H. Q. (2005). Size and trends of the House Sparrow *Passer*
436 *domesticus* population in Great Britain. *Ibis*, 147(3), 552-562.

437 Romanowski, J., Pinowski, J., Sawicka-Kapusta, K. and Wlostowski, T, 1991, The effect of heavy metals upon
438 development and mortality of *Passer domesticus* and *Passer montanus* nestlings. In Pinowski, J., Kavanagh, B.,
439 and Gorski, W. (eds.), Nestling mortality of granivorous birds due to microorganisms and toxic substances. The
440 International Symposium of the Working Group on Granivorous Birds, 197-204.

441 Sanchez-Tojar, A., Nakagawa, S., Sanchez-Fortun, M., Martin, D. A., Ramani, S., Girndt, A., ...& Schroeder, J.
442 (2018). Meta-analysis challenges a textbook example of status signalling and demonstrates publication bias. *Elife*, 7,
443 e37385.

444 Sharma, P., & Binner, M. (2020). The decline of population of house sparrow in India. *International Journal of*
445 *Agricultural Science*, 5.
446

447 Shaw LM, Chamberlain D and Evans M. (2008) The House Sparrow *Passer domesticus* in urban areas: reviewing a
448 possible link between post-decline distribution and human socioeconomic status. *Journal of Ornithology*,
449 149(3), 293-299.
450

451 Sullivan, B.L., C.L. Wood, M.J. Iliff, R.E. Bonney, D. Fink, and S. Kelling. 2009. eBird: a citizen-based bird
 452 observation network in the biological sciences. *Biological Conservation*, 142: 2282-2292.
 453

454 Summers-Smith J.D., The Decline of the House Sparrow: a review, *Br. Birds*, 96, 439-446 (2003) 18. Chris
 455 Klok, Remko Holtkamp, Rob van Apeldoorn, Marcel E. Visser, Lia Hemerik, Analysing Population Numbers of
 456 the House Sparrow in the Netherlands With a Matrix Model and Suggestions for Conservation Measures,
 457 *Acta Biotheoretica*, 54(3), 161–178 (2006).
 458

459 Summers-Smith, D., 1999, Current status of the House Sparrow in Britain. *British Wildlife*, UK, 381- 386.

460 Summers-Smith, D., 2006, Decline of House Sparrows in Large Towns. *British Birds*, UK, 93p.

461 Summers-Smith, J.D. (1988). *The Sparrows: A Study of the Genus Passer*. T & A D Poyser. *Staffordshire, UK*.

462 Swami, R. and Kumar, M. Study on Ecology of House Sparrow, *Passer domesticus* (Linnaeus, 1758). *J Am Sci*
 463 2022;18(4), 107-110.

464 Teotia, D.S., Yadav, A., Kumar, V. and Kumar, A. (2018). Study of the declining population of common
 465 “house sparrow” (*Passer domesticus*) in urban and sub-urban areas of India. *Life Science Leaflets*, 101, 1-11.