

PLATELET-RICH FIBRIN (L-PRF), TREATMENT OPTION FOR CLOSING OROANTRAL COMMUNICATION IN AN ONCOLOGIC PATIENT: A REVIEW AND CASE REPORT.

by Jana Publication & Research

Submission date: 13-Oct-2025 07:14AM (UTC+0300)

Submission ID: 2769517919

File name: IJAR-54294.pdf (235.31K)

Word count: 2919

Character count: 15718

28
1 **PLATELET-RICH FIBRIN (L-PRF): TREATMENT OPTION FOR CLOSING**
2 **OROANTRAL COMMUNICATION IN AN ONCOLOGIC PATIENT: A REVIEW**
3 **AND CASE REPORT.**

9
5 **ABSTRACT**

6 Leukocyte- and Platelet-Rich Fibrin (L-PRF) is an autologous biomaterial used in grafting
7 procedures. It consists of a fibrin matrix enriched with leukocytes, platelets, leukocyte-
8 derived cytokines, and growth factors, all obtained from a simple blood sample. This case
9 report aims to present the clinical application of L-PRF in the treatment of oroantral
10 communication in an oncology patient. A 59-year-old female patient undergoing
11 chemotherapy presented with right maxillary sinusitis and a fractured maxillary first molar
12 (tooth #3), which exhibited a periapical lesion in continuity with the maxillary sinus. The
13 proposed treatment involved the extraction followed by closure of the resulting oroantral
14 fistula. To minimize invasiveness, the treatment strategy focused on obliterating the
15 communication between the oral cavity and the maxillary sinus by filling the extraction
16 socket with an autologous L-PRF membrane. After four months of follow-up, the patient
17 showed complete resolution of the maxillary and alveolar infection, with no evidence of
18 persistent communication between the oral cavity and the sinus. The use of L-PRF resulted in
19 satisfactory outcomes, including new bone formation and the prevention of maxillary
20 osteonecrosis. This case demonstrates that L-PRF can be a viable and effective option for the
21 management of oroantral communication, particularly in oncology patients where minimally
22 invasive approaches are preferred.

16
23 **Keywords:** Platelet-Rich Fibrin; Bone Regeneration; Oroantral Fistula.

14
24 **Introduction**

25 Due to the pneumatization of the maxillary sinus in the region of the upper molars and
26 the anatomical characteristics of this area, oroantral communications may occur during
27 extraction procedures of these teeth. Their resolution should be immediate or as early as
28 possible, given the risk of postoperative sinusitis and oroantral fistula, with treatment varying
29 according to the size of the communication¹. However, when oroantral communication occurs
30 in patients with systemic conditions that impair the healing process, treatment should be more
31 conservative.

32 One factor contributing to systemic impairment is the risk of osteonecrosis caused by
33 bisphosphonate-based chemotherapeutic agents in oncology patients. Bisphosphonates are
34 indicated for various conditions, including tumor-induced hypercalcemia, breast carcinoma,

multiple myeloma, heterotopic ossification, and stabilization of bone loss in cases of osteoporosis².

As members of a drug class that interacts with bone remodeling, prolonged use of bisphosphonates may compromise bone quality due to excessive suppression of bone repair. In recent years, osteonecrosis of the jaws has been reported as a significant complication of chronic bisphosphonate use. This complication frequently occurs following oral surgical procedures, primarily extractions³.

As a result, a series of studies have been initiated to explore adjunctive therapies that support the healing process in such procedures. Platelet aggregates have emerged as a promising biomaterial in oral surgery⁴⁻⁵. Among the types of concentrates, the slow and progressive polymerization of L-PRF enhances the incorporation of chemical mediators, leukocytes, platelets, leukocyte-derived cytokines, and growth factors into the fibrin mesh⁶⁻⁷. It is considered the closest to a natural clot, as it is obtained from pure blood and promotes satisfactory bone regeneration⁸⁻⁹.

The biological properties of L-PRF have the ability to accelerate physiological healing, and when combined with bone grafts, they enhance the bone regeneration process¹⁰. L-PRF can be obtained in its polymeric phase, forming a membrane, and in its monomeric phase, which initially presents as a liquid and becomes gelatinous upon polymerization¹¹. The aim of this study was to present a clinical case in which L-PRF was chosen as a treatment option for oroantral communication in an oncology patient.

Literature Review

Oroantral communications are defined as a passage between the oral cavity and the maxillary sinus, which may occur due to trauma, pathological conditions, or oral surgeries such as tooth extractions. If the perforation is small (up to 2 mm), it may close spontaneously¹². However, it is difficult to accurately estimate its size; therefore, once the perforation is detected through the Valsalva test, closure should always be performed immediately or as early as possible to prevent the formation of an oroantral fistula or chronic maxillary sinusitis¹³.

Various surgical techniques are available for treating these communications, with the most commonly used being buccal advancement flaps, palatal rotational flaps, and the Bichat

fat pad flap^{14 15}. However, in cases where rehabilitation is needed in patients with systemic compromise, a more conservative approach is required—such as in patients undergoing bisphosphonate-based treatment.

Bisphosphonates are drugs used in the treatment of cancer-related conditions, improving the quality of life for patients undergoing such therapies². However, studies have shown a correlation between the use of bisphosphonates and the development of medication-related osteonecrosis of the jaws (MRONJ)^{3 16}. This condition depends on three diagnostic criteria: current or previous treatment with bisphosphonates; exposed bone in the maxillofacial region persisting for more than eight weeks; and no history of radiation therapy to the jaws¹⁷. It may directly affect the maxillary sinus, leading to oroantral communications. Since bisphosphonates impair bone healing, in cases of oroantral communication in oncology patients treated with these drugs, a more conservative therapeutic adjunct is necessary to enhance healing—such as platelet-based biomaterials^{18 19 20}.

Different types of platelet concentrates are used in oral surgery, including platelet-rich plasma (PRP) and platelet-rich fibrin (PRF). Within this field, leukocyte- and platelet-rich fibrin (L-PRF) is a compound that includes, in addition to platelets and fibrin, leukocytes. Each of these components plays an important role in the healing process. The fibrin clot that forms acts as a provisional matrix in which platelets are trapped, responsible for releasing various growth factors, chemokines, and inflammatory mediators that directly influence the healing process⁷.

Leukocytes are divided into three groups: granular cells, lymphocytes, and monocytes. Among the granular cells are neutrophils, basophils, and eosinophils, which also include polymorphonuclear leukocytes (PMN and PML). Lymphocytes are divided into B and T cells, and monocytes migrate into tissues, maturing into macrophages. The primary role of these leukocytes is to produce inflammatory cytokines and various growth factors, promote wound repair, protect against bacterial infections, maintain tissue integrity, and enhance the effects of platelets^{6 21}.

L-PRF is produced by collecting approximately 9 ml of the patient's blood into glass tubes without anticoagulants, using standard venipuncture nursing techniques. Immediately after collection, the blood is placed into a centrifuge⁴. There are different protocols for

centrifugation, with rotation speeds ranging from 1500 to 3300 rpm for 10 to 12 minutes, depending on the centrifuge used²².

During centrifugation, the coagulation process is activated, forming a clot with a three-dimensional fibrin network where platelets and other cells are trapped²⁴. Unlike other platelet aggregates, this occurs without the addition of anticoagulants, which typically induce rapid fibrinogen polymerization with low mechanical strength. In L-PRF, fibrinogen polymerization occurs naturally, resulting in a membrane with good clinical handling and mechanical resistance^{4, 23}.

Case Report

The patient, a 58-year-old female undergoing chemotherapy for the management of metastatic tumors secondary to breast cancer, presented with complaints of right maxillary sinusitis. Clinical and radiographic examination revealed a root perforation of tooth #3, with a periapical lesion in continuity with the maxillary sinus, as well as opacification of the right maxillary sinus (Figures 1 and 2).

Following consultation with the oncologist responsible for the patient's treatment, the recommended approach was the extraction of tooth #3 and closure of the oroantral fistula to resolve the odontogenic infection. Considering the patient's systemic condition and the risk of osteonecrosis associated with her oncologic therapy, it was proposed obliteration of the communication using an autologous L-PRF membrane in its polymeric phase, combined with antibiotic therapy—Amoxicillin with Potassium Clavulanate 875mg administered preoperatively and continued for 14 days postoperatively.

At the beginning of the procedure, 18 ml of venous blood were collected into two 9 ml glass tubes containing an activator (silica). The tubes were placed in a centrifuge (KASVI - K14-0815®) at 2200 rpm for 10 minutes. At the end of the process, the L-PRF in its polymeric phase was decanted and placed into the extraction socket (Figure 3).

The procedure was performed with the patient conscious under local anesthesia. Following anesthetic administration, tooth #3 was extracted using a minimally traumatic technique to preserve the surrounding soft tissues. This was followed by vigorous curettage of the socket to remove all periapical lesion tissue, copious irrigation with 0.9% saline solution, and filling of the socket with folded L-PRF membranes in their polymeric phase. Tissue

125 closure was achieved using 5-0 nylon sutures to ensure stabilization of the L-PRF within the
126 socket (Figure 4).

127 The patient was monitored weekly, and no signs of surgical site infection were
128 observed (Figure 5). Sutures were removed three weeks postoperatively, and a Valsalva
129 maneuver was performed, yielding a negative result, confirming successful closure of the
130 oroantral communication.

131 After four ¹⁷months of follow-up, there was complete resolution of the infectious
132 condition affecting both the maxillary sinus and the alveolar region. No clinical signs of
133 alteration or continuity ⁵between the oral cavity and the maxillary sinus were observed.
134 ²⁰Radiographic evaluation revealed evidence of new bone formation in the area of the
135 extraction socket of tooth #3 (Figure 6).

136 Discussion

137 Oroantral communications ³require immediate or early intervention to ensure
138 successful resolution, as exposure of the maxillary sinus to the oral environment can lead to
139 chronic sinusitis¹³. Several treatment options have been proposed, with the most conventional
140 approaches involving vestibular or palatal flaps with suturing, or alternatively, obliteration of
141 the communication using the Bichat fat pad^{14 15}.

142 Given the compromised systemic condition of the patient described in this case, a
143 more conservative surgical approach was necessary, as her healing capacity was affected by
144 oncologic treatment involving bisphosphonate-based drugs, which have been widely reported
145 to carry a significant risk of maxillary osteonecrosis²⁰.

146 With this in mind, the chosen treatment was ²⁶the use of an L-PRF membrane, instead
147 of a flap, to close the oroantral communication and promote both soft tissue and bone healing
148 in the region. Studies have shown that the slow and progressive polymerization of L-PRF
149 leads to the aggregation of leukocytes, neutrophils, leukocyte-derived cytokines, and growth
150 factors within a fibrin network, which are released gradually and continuously at the site⁶.
151 This promotes healing and helps prevent local infections, making it effective in such cases²⁴
152 ²⁵.

153 It is important to note that the communication described in this case was of moderate
154 size (between 3 and 4 mm), making it suitable for closure using the L-PRF membrane. In

cases of larger communications, the L-PRF membrane may lack sufficient structural support, and additional materials such as autologous bone grafts or collagen membranes may be required to achieve satisfactory results²⁵.

Conclusion

After four months of follow-up, the patient showed complete resolution of the infectious condition in both the maxillary sinus and alveolar region, with no evidence of continuity between the oral cavity and the maxillary sinus. Thus, the use of L-PRF proved to be a satisfactory treatment option, promoting new bone formation and preventing the occurrence of maxillary osteonecrosis.

REFERENCES

1. Al-Juboori MJ, Al-Attas MA, Filho LCM. Treatment of chronic oroantral fistula with platelet-rich fibrin clot and collagen membrane: A case report. *Clin Cosmet Investig Dent*. 2018;10:245–9.
2. Ruggiero SL, Dodson TB, Assael L a, Landesberg R, Marx RE, Mehrotra B. Position paper AAOMS position paper 2009. *Aust Endod J*. 2009;35(3):119–30.
3. Al-Zoman KH, AlBazie S, Robert AA, Baskaradoss JK, Alsuywed AS, Ciancio S, et al. Surgical management of bisphosphonate-related osteonecrosis of the jaw: Report of three cases. *J Palliat Care*. 2013;29(1):52–7.
4. Madurantakam P, Yoganarasimha S, Hasan FK. Characterization of Leukocyte-platelet Rich Fibrin, A Novel Biomaterial. *J Vis Exp*. 2015;(103):1–8.
5. Orłowska A, Sader R, Kirkpatrick C, Lorenz J, Booms P, Choukroun J, et al. Advanced Platelet-Rich Fibrin: A New Concept for Cell-Based Tissue Engineering by Means of Inflammatory Cells. *J Oral Implantol*. 2014;40(6):679–89.
6. Bielecki T, M. Dohan Ehrenfest D, A. Everts P, Wiczowski A. The Role of Leukocytes from L-PRP/L-PRF in Wound Healing and Immune Defense: New Perspectives. *Curr Pharm Biotechnol*. 2012;13(7):1153–62.
7. Abukabda AB, Rutkowski JL, Clafshenkel WP, Davis VL, Radio NM, Witt-Enderby PA, et al. Platelet-Rich Preparations to Improve Healing. Part I: Workable Options for Every Size Practice. *J Oral Implantol*. 2012.
8. Vandessel J, Teughels W, Jacobs R, Castro A, Pinto N, Quirynen M, et al. The use of leucocyte and platelet-rich fibrin in socket management and ridge preservation: a split-mouth, randomized, controlled clinical trial. *J Clin Periodontol*. 2016;43(11):990–9.
9. Pinto GDDS, Pigossi SC, Pessoa T, Nicoli LG, Araújo RF de SB, Marcantonio C, et al. Successful Use of Leukocyte Platelet-Rich Fibrin in the Healing of Sinus Membrane Perforation: A Case Report. *Implant Dent*. 2018;27(3):375–80.
10. Nizam N, Eren G, Akcalı A, Donos N. Maxillary sinus augmentation with leukocyte and platelet-rich fibrin and deproteinized bovine bone mineral: A split-mouth

- 192 histological and histomorphometric study. Clin Oral Implants Res. 2018;29(1):67–75.
- 193 11. Cortellini S, Castro AB, Temmerman A, Van Dessel J, Pinto N, Jacobs R, et al.
194 Leucocyte- and platelet-rich fibrin block for bone augmentation procedure: A proof-of-
195 concept study. J Clin Periodontol. 2018;45(5):624–34.
- 196 12. Procacci P, Alfonsi AF, Tonelli P, Selvaggi F, Battista G, Fabris M, et al. Surgical
197 Treatment of Oroantral Communications. 2016;27(5):1190–6.
- 198 13. Akhlaghi F, Esmaeelinejad M, Safai P. Etiologies and Treatments of Odontogenic
199 Maxillary Sinusitis : A Systematic Review. 2015;17(12).
- 200 14. Visscher SH, Minnen B Van, Bos RRM. Closure of Oroantral Communications :
201 YJOMS. 2010;68(6):1384–91.
- 202 15. Hernando J, Gallego L, Junquera L, Villarreal P. Oroantral communications . A
203 retrospective analysis. 2010;15(3):1–5.
- 204 16. Marx RE, Sawatari Y, Fortin M, Broumand V. Bisphosphonate-induced exposed bone
205 (osteonecrosis/osteopetrosis) of the jaws: Risk factors, recognition, prevention, and
206 treatment. J Oral Maxillofac Surg. 2005;63(11):1567–75.
- 207 17. Colella G, Campisi G, Fusco V. American Association of Oral and Maxillofacial
208 Surgeons Position Paper: Bisphosphonate-Related Osteonecrosis of the Jaws-2009
209 Update: The Need to Refine the BRONJ Definition. J Oral Maxillofac Surg.
210 2009;67(12):2698–9.
- 211 18. Voss PJ, Vargas Soto G, Schmelzeisen R, Izumi K, Stricker A, Bittermann G, et al.
212 Sinusitis and oroantral fistula in patients with bisphosphonate-associated necrosis of
213 the maxilla. Head Face Med. 2016;12(1):1–6.
- 214 19. Mast G, Otto S, Mücke T, Schreyer C, Bissinger O, Kolk A, et al. Incidence of
215 maxillary sinusitis and oro-antral fistulae in bisphosphonate-related osteonecrosis of
216 the jaw. J Cranio-Maxillofacial Surg. 2012;40(7):568–71.
- 217 20. Weissman T, Yahalom R, Fliss DM, Yarom N. Involvement of the maxillary sinus in
218 bisphosphonate- related osteonecrosis of the jaw: Radiologic aspects. Am J Rhinol
219 Allergy. 2017;31(1):36–9.
- 220 21. Davis VL, Rutkowski JL, Cairone JV, Witt-Enderby PA, Abukabda AB, Radio NM, et
221 al. Platelet-Rich Preparations to Improve Healing. Part II: Platelet Activation and
222 Enrichment, Leukocyte Inclusion, and Other Selection Criteria. J Oral Implantol.
223 2014;40(4):511–21.
- 224 22. Dohan Ehrenfest DM, Pinto NR, Pereda A, Jiménez P, Corso M Del, Kang B-S, et al.
225 The impact of the centrifuge characteristics and centrifugation protocols on the cells,
226 growth factors, and fibrin architecture of a leukocyte- and platelet-rich fibrin (L-PRF)
227 clot and membrane. Platelets. 2018 Feb 17;29(2):171–84.
- 228 23. Khorshidi H, Raoofi S, Bagheri R, Banihashemi H. Comparison of the Mechanical
229 Properties of Early Leukocyte- and Platelet-Rich Fibrin versus PRGF/Endoret
230 Membranes. Int J Dent. 2016;2016:1–7.
- 231 24. Studies BC, Bilginaylar K. Comparison of the Clinical Outcomes of Buccal Platelet-
232 Rich Fibrin Application for the Immediate Closure of Acute Oroantral. J Craniofac
233 Surg. 2019;30(1):150–2.

- 234 25. Kapustecki M, Niedzielska I, Borgiel-marek H, Rózanowski B. Alternative method to
235 treat oroantral communication and fistula with autogenous bone graft and platelet rich
236 fibrin. 2016;21(5):3–8.

UNDER PEER REVIEW IN IJAR

PLATELET-RICH FIBRIN (L-PRF), TREATMENT OPTION FOR CLOSING OROANTRAL COMMUNICATION IN AN ONCOLOGIC PATIENT: A REVIEW AND CASE REPORT.

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1

www.ijmedrev.com

Internet Source

2%

2

Submitted to Trinity College Dublin

Student Paper

1%

3

Armelia Sari Widyarman, Muhammad Ihsan Rizal, Moehammad Orliando Roeslan, Carolina Damayanti Marpaung. "Quality Improvement in Dental and Medical Knowledge, Research, Skills and Ethics Facing Global Challenges", CRC Press, 2024

Publication

1%

4

Carolina Mendonça de Almeida Malzoni, Lélis Gustavo Nícoli, Gustavo da Col dos Santos Pinto, Suzane Cristina Pigossi et al. "The Effectiveness of L-PRF in the Treatment of Schneiderian Membrane Large Perforations: Long-Term Follow-Up of a Case Series", Journal of Oral Implantology, 2021

Publication

1%

5

Christian Bacci, Claudia Manera, Davide Meneghetti, Iris Dhelpra, Luca Sbricoli. "Efficacy of Bichat's Buccal Fat Pad Advancement in the Treatment of Odontogenic Maxillary Sinus Fistulas", Surgeries, 2025

Publication

1%

6

Submitted to University of Edinburgh

Student Paper

		1 %
7	lirias.kuleuven.be Internet Source	1 %
8	www.frontiersin.org Internet Source	1 %
9	www.ijcbr.in Internet Source	1 %
10	Guillermo Pardo-Zamora, Yanet Martínez, Jose Antonio Moreno, Antonio J. Ortiz-Ruíz. "Treatment of Stage 2 Medication-Induced Osteonecrosis of the Jaw: A Case Series", International Journal of Environmental Research and Public Health, 2021 Publication	1 %
11	Mohamed S. Elmaradny, Ahmed S. Elmahalawy, Hala R. Ragab. "THE USE OF BUCCAL ANTROSTOMY BONE AS A GRAFT FOR CLOSURE OF ORO- ANTRAL FISTULA", Alexandria Dental Journal, 2019 Publication	1 %
12	healthdocbox.com Internet Source	1 %
13	www.cro-pe.org.br Internet Source	1 %
14	link.springer.com Internet Source	<1 %
15	www.mdpi.com Internet Source	<1 %
16	Paulina Adamska, Magdalena Kaczoruk-Wieremczuk, Dorota Pylińska-Dąbrowska, Marcin Stasiak et al. "Treatment of Oroantral Communication and Fistulas with the Use of Blood-Derived Platelet-Rich Preparations Rich	<1 %

17 Shay Keren, Matthew Lapira, Ewan McCallum, Zuzana Sipkova et al. "Paediatric orbital lymphoma; a case series and review of the literature", Eye, 2022

Publication

18 poseido.info <1 %

Internet Source

19 worldwidescience.org <1 %

Internet Source

20 Tara L Aghaloo. "Periodontal disease and bisphosphonates induce osteonecrosis of the jaws in the rat", Journal of Bone and Mineral Research, 08/2011

Publication

21 clinicaterapeutica.it <1 %

Internet Source

22 cynbagley.blogspot.com <1 %

Internet Source

23 iris.unime.it <1 %

Internet Source

24 www.preprints.org <1 %

Internet Source

25 www.saudijournals.com <1 %

Internet Source

26 Ana B. Castro, Nastaran Meschi, Andy Temmerman, Nelson Pinto, Paul Lambrechts, Wim Teughels, Marc Quirynen. "Regenerative potential of leucocyte- and platelet-rich fibrin. Part B: sinus floor elevation, alveolar ridge preservation and implant therapy. A

systematic review", Journal of Clinical
Periodontology, 2017

Publication

- 27 Rodella, Luigi Fabrizio. "Platelet preparations in dentistry: How? Why? Where? When?", World Journal of Stomatology, 2015. <1 %

Publication

- 28 Ivo Lambrichts, Esther Wolfs, Annelies Bronckaers, Pascal Gervois, Tim Vangansewinkel. "The Effect of Leukocyte- and Platelet-Rich Fibrin on Central and Peripheral Nervous System Neurons—Implications for Biomaterial Applicability", International Journal of Molecular Sciences, 2023 <1 %

Publication

- 29 Sheeja S Varghese, Arunima Padmakumar Reshma, Lizymol P Pampadykandathil. "Comparative Evaluation of Platelet-derived Growth Factor Isoforms between Platelet-rich Fibrin Prepared with Three Different Centrifugation Protocols", World Journal of Dentistry, 2024 <1 %

Publication

- 30 hdl.handle.net <1 %

Internet Source

Exclude quotes On

Exclude matches Off

Exclude bibliography On