

TO EVALUATE AWARENESS AND ACCEPTANCE OF PERIODONTAL DISEASE AND THERAPY AMONG PATIENT VISITING AMC DENTAL COLLEGE: A QUESTIONNAIRE SURVEY

by Jana Publication & Research

Submission date: 14-Oct-2025 09:02AM (UTC+0300)

Submission ID: 2769515754

File name: IJAR-54331.pdf (1.19M)

Word count: 3055

Character count: 16443

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

TO EVALUATE AWARENESS AND ACCEPTANCE OF PERIODONTAL DISEASE AND THERAPY AMONG PATIENT VISITING AMC DENTAL COLLEGE: A QUESTIONNAIRE SURVEY

Manuscript Info

Manuscript History

Received: xxxxxxxxxxxxxxxx
Final Accepted: xxxxxxxxxxxxxxxx
Published: xxxxxxxxxxxxxxxx

Key words:-

Periodontal disease, Preventive dentistry,
Patient awareness, Acceptance of periodontal
therapy.

Abstract

Background: Periodontal diseases are among the most common oral health problems worldwide and are characterized by inflammation and destruction of the supporting structures of the teeth. They can lead to tooth loss, impaired function, and a negative impact on overall health and quality of life. This study aimed to assess patients' knowledge, understanding, and attitudes toward periodontal disease, and to evaluate their awareness and acceptance of periodontal treatment.

Materials and Methods: A cross-sectional, questionnaire-based study was conducted among 440 patients visiting the outpatient department of Periodontology at AMC Dental College, Ahmedabad, India. The study was carried out from January 2024 to June 2024. A structured questionnaire containing 13 questions was administered in the local language using Google Forms. Responses were recorded on a three-point scale (Yes, No, Maybe) and analysed using descriptive statistics.

Results: Most participants demonstrated awareness of basic aspects of periodontal health, including recognition of bleeding gums (60.5%) and tooth loss due to periodontal disease (59.8%). However, limited awareness was observed regarding diabetes as a risk factor (27.5%), medication-induced gingival swelling (27.5%), laser therapy (24.5%), bone grafting (17.5%), and local antibiotic delivery (16.6%). A common misconception was that scaling results in the loss of tooth substance (68.6%), indicating the need for improved patient education.

Conclusion: The findings highlight considerable gaps in knowledge and several misconceptions related to periodontal disease and its treatment. Targeted educational strategies are required to enhance patient understanding, increase treatment acceptance, and improve overall periodontal health outcomes.

Introduction:

Periodontal diseases are among the most prevalent oral health problems worldwide, affecting a large proportion of the adult population. Characterized by the inflammation and destruction of the supporting structures of the teeth, periodontal diseases can lead to tooth loss, impaired function, and significant impacts on general health and quality of life. In India, the burden of periodontal disease remains exceptionally high, with national data indicating that a large proportion of adults experience some form of periodontal breakdown (Bali et al., 2004)⁽¹⁾⁽²⁾⁽³⁾. Gingivitis and periodontitis not only compromise oral health but are also associated with systemic conditions such as diabetes mellitus and cardiovascular

47 disease⁽⁴⁾⁽⁵⁾. In India, periodontal disease is a major cause of tooth loss, particularly among adults over
48 40 years of age, accounting for approximately 50–70% of extractions in this group. (Bali et al., 2004;
49 Kumar et al., 2009; Sharma et al., 2016)⁽¹⁾⁽²⁾. Despite advances in preventive and therapeutic strategies,
50 many patients remain unaware of the causes, consequences, and treatment options related to periodontal
51 conditions.

52 Patient awareness and acceptance of periodontal therapy play a critical role in the prevention and
53 management of these diseases. Inadequate knowledge and negative perceptions can result in delayed
54 diagnosis, poor compliance with treatment recommendations, and suboptimal outcomes. In India, varying
55 literacy levels, socioeconomic factors, and cultural beliefs further influence attitudes toward oral health
56 care and utilization of dental services.⁽⁶⁾ AMC Dental College serves a diverse patient population,
57 providing an opportunity to assess current levels of understanding and acceptance regarding periodontal
58 disease and its management. Evaluating these factors is essential to identify knowledge gaps, improve
59 patient education, and design effective community-based interventions.

60 This questionnaire survey aims to assess the awareness and acceptance of periodontal disease and
61 therapy among patients attending AMC Dental College. The findings are expected to guide strategies to
62 enhance periodontal health literacy and promote positive attitudes toward periodontal care.

63 **Materials and Methods:**

64 This cross-sectional questionnaire-based study was conducted in the Outpatient Department of
65 Periodontology, AMC Dental College, Ahmedabad. Ethical clearance was obtained from the Institutional
66 Review Board (IRB) of AMC Dental College prior to the commencement of the study. A total of 440
67 patients visiting the department were recruited for the study. Data were collected using a pre-validated
68 structured questionnaire administered through Google Forms. To ensure better comprehension, the
69 questionnaire was provided in the participants' respective local language. The questionnaire comprised 13
70 questions, of which the first six questions assessed knowledge regarding periodontal disease, including its
71 symptoms and risk factors. The remaining seven questions evaluated participants' awareness and
72 acceptance toward periodontal treatment. Responses were recorded on a three-point Likert scale with the
73 options "Yes," "No," and "Maybe." (table no. 1) The collected data were compiled and analysed using
74 simple descriptive statistics to summarize the study variables.

75

76

77 **Table 1: Questionnaire to Assess Awareness and Acceptance of Periodontal Disease and**
 78 **Therapy Among Patients**

79

Sr. No.	Questions	Response
1	Do you feel bleeding gums while brushing or flossing is sign of gum disease?	Yes \ No \ May be
2	Do you feel tooth loss is due to periodontitis?	Yes \ No \ May be
3	Consulting dental specialist required when you notice a mobile tooth or bad breath?	Yes \ No \ May be
4	Do you feel that diabetes increase your risk of gum disease?	Yes \ No \ May be
5	Do you feel that swelling of gums can be caused by certain medications?	Yes \ No \ May be
6	Do you feel that smoking , tobacco chewing can aggravate gum disease?	Yes \ No \ May be
7	Do you feel that early gum therapy can help for reduction of tooth movement?	Yes \ No \ May be
8	Do you think scaling can contribute to tooth substance loss?	Yes \ No \ May be
9	Do you know that for dental procedure new technologies like laser are being use?	Yes \ No \ May be
10	Can smile design can be achieved by esthetic periodontal surgery?	Yes \ No \ May be
11	Do you know that bone graft material use for regeneration of bone?	Yes \ No \ May be
12	Are you aware that systemic antibiotic can be delivered locally in your gum?	Yes \ No \ May be
13	Are you aware that dental implants are used for replacement of your tooth?	Yes \ No \ May be

80

81

82

7

Results:

A total of 440 participants were included in this study, with ages ranging from 14 to 95 years (mean = 39.44 years, SD = 15.201), representing a diverse age group. Among the participants, 60.2% were male and 39.8% were female, indicating a higher participation rate among males.

Awareness of Periodontal Disease:

1. Yes, bleeding gums are a sign of periodontal disease. Among the respondents, 60.5% recognized bleeding gums as a sign of periodontal disease, while 35% did not consider it an indicator of the disease, demonstrating a substantial need for education (Figure 1).




Figure 1: Do you feel bleeding gums while brushing or flossing is sign of gum disease?

2. Yes, tooth loss is due to periodontitis. In this study, 59.8% of respondents were aware that periodontitis could lead to tooth loss, whereas 34.3% disagreed, highlighting a significant knowledge gap (Figure 2).




Figure 2: Do you feel tooth loss is due to periodontitis ?

3. Yes, consulting a specialist is required when bad breath or mobile teeth are noted. In this study, 63.2% of participants expressed a positive attitude toward seeking periodontal treatment, while 29.3% disagreed (Figure 3).




Figure 3: Consulting dental specialist required when you notice a mobile tooth or bad breath?

4. Yes, diabetes increases the risk of gum disease. Awareness regarding the link between diabetes and periodontal disease was poor only 27.5% of respondents identified diabetes as a risk factor, while 63% disagreed (Figure 4).




Figure 4 : Do you feel that diabetes increase your risk of gum disease?

5. Yes, swelling of the gums can be caused by medication. Knowledge about drug-induced gingival enlargement was limited, with only 27.5% of respondents reporting awareness, compared to 61.4% who were unaware (Figure 5).




Figure 5: Do you feel that swelling of gums can be caused by certain medications?

6. Yes, smoking and tobacco chewing aggravate gum disease. Awareness of the relationship between these habits and periodontal disease was moderate, with 61.4% of respondents recognizing the risk, while 28.4% disagreed (Figure 6).




Figure 6: Do you feel that smoking, tobacco chewing can aggravate gum disease?

Awareness of Periodontal Therapy:

7. Yes, early gum therapy helps to reduce tooth movement. When questioned about preventive periodontal care, only 29.5% of respondents agreed that it was essential, whereas 59.5% disagreed (Figure 7)




Figure 7: Do you feel that early gum therapy can help for reduction of tooth movement?

8.No, scaling does not contribute to tooth substance loss. A notable misconception was observed, with 68.6% of respondents believing that scaling leads to tooth loss, highlighting the persistence of myths related to routine dental procedures (Figure 8).




Figure 8:Do you think scaling can contribute to tooth substance loss?

9. Yes, lasers are used in dental procedures. Awareness of laser periodontal treatment options was generally poor, with only 24.5% of respondents reporting familiarity with laser therapy (Figure 9).






Figure 9:Do you know that for dental procedure new technologies like laser are being use?

157 **10.**Yes, smile design can be achieved through esthetic periodontal surgery. However, only 22.7% of
 158 respondents were aware of cosmetic periodontal procedures (Figure 10).




159
 160 **Figure 10 :Can smile design can be achieved by esthetic periodontal surgery?**

161 **11.**Yes, bone graft materials are used for bone regeneration. However, only 17.5% of respondents
 162 recognized that regenerative procedures can be used in periodontal therapy (Figure 11).



163
 164 **Figure 11:Do you know that bone graft material use for regeneration of bone?**

165
 166 **12.**Yes, systemic antibiotics can be delivered locally in the gums. However, only 16.6% of respondents
 167 were aware of the use of localized antibiotic delivery in periodontal therapy (Figure 12).



168
 169 **Figure 12:Are you aware that systemic antibiotic can be delivered locally in your gum?**

13. Yes, dental implants are used for the replacement of teeth. Among the respondents, 54.8% recognized the use of implants for tooth replacement, while 9.3% had no knowledge about dental implants.




Figure 13: Are you aware that dental implants are used for replacement of your tooth?

Comparative evaluation of awareness regarding periodontal disease and periodontal treatment:

On average, 50% of participants demonstrated correct knowledge regarding periodontal disease, while only 27% exhibited accurate knowledge about periodontal treatment modalities (Figure 14). Wrong knowledge is more common about periodontal therapy than about periodontal disease. Right knowledge is more common about periodontal disease than about periodontal therapy. This substantial gap underscores the need for improved patient education and counselling by dental professionals.




Figure 14: Comparative evaluation of awareness regarding periodontal disease and periodontal treatment:

Discussion :

The present survey highlights important insights into awareness and perceptions of periodontal disease and its management among patients visiting AMC Dental College. The findings show that while participants demonstrated good understanding of basic signs and consequences of periodontal disease, such as bleeding gums and tooth loss, their awareness about underlying risk factors and advanced treatment options was limited. In India, periodontitis is highly prevalent. According to the National Oral Health Survey and Fluoride Mapping 2002–2003, approximately 85% of adults over 35 years exhibited some form of periodontal disease. More recent estimates suggest that moderate to severe periodontitis affects 40–60% of Indian adults, with severe forms seen in about 10–15% (Sharma et al., 2016)⁽²⁾. Despite this high burden, public understanding remains inadequate.

A notable proportion of respondents in this survey recognized bleeding gums as an early sign of periodontal disease and were aware that untreated gum problems could lead to tooth mobility and eventual tooth loss. This finding is in agreement with previous studies reporting moderate awareness of visible symptoms (Prasad et al., 2014).⁽⁷⁾ However, widespread misconceptions persist, particularly the belief that scaling causes permanent damage to tooth structure—a myth endorsed by 68.6% of participants in this study. Similar misconceptions have been documented by Kumar et al. (2013)⁽⁸⁾ who found that nearly 70% of patients believed scaling would weaken teeth or remove enamel. Such beliefs are significant barriers to timely treatment and contribute to disease progression. Similar to the findings of previous studies conducted in North India and West Godavari, where over 70% of participants believed that scaling weakens teeth or causes mobility, this survey also revealed that 68.6% of patients shared this misconception, emphasizing the persistence of this myth across different populations in India and the urgent need for targeted educational interventions.⁽⁹⁾⁽¹⁰⁾

Additionally, awareness of systemic associations of periodontal disease was poor. Most participants did not recognize diabetes as a significant risk factor. This gap is concerning given the established bidirectional link between diabetes and periodontitis (Mealey & Oates, 2006)⁽¹²⁾. Patients with diabetes are

214 at higher risk of severe periodontal destruction, and poor periodontal health can, in turn, compromise
215 glycemic control. Knowledge about modern periodontal therapies, including laser treatment, bone
216 grafting, and localized antimicrobial delivery, was also found to be inadequate. These findings are in line
217 with previous reports indicating very low public awareness of advanced treatment modalities (Gupta et
218 al., 2013).⁽¹¹⁾ Limited exposure, perceived high cost, and lack of communication by dental professionals
219 contribute to this deficit.

220 Overall, the results of this survey underscore the urgent need for targeted educational interventions to
221 address misconceptions and improve knowledge about periodontal disease and its management. Dental
222 professionals play a critical role in counseling patients about the importance of preventive care, the safety
223 and necessity of scaling, the systemic impacts of periodontal infection, and the availability of effective
224 treatment options. Integrating oral health education into primary care, leveraging community outreach,
225 and using culturally appropriate, easy-to-understand materials may significantly improve awareness and
226 acceptance of periodontal therapy.

227 **Conclusion:**

228 These findings underscore the need for targeted oral health education initiatives to address specific
229 knowledge gaps and misconceptions. Improving patient understanding of periodontal disease and
230 available therapies will be essential to enhance prevention, encourage acceptance of treatment, and
231 ultimately improve periodontal health outcomes. To improve awareness, chairside education using simple
232 language and visuals should be provided, along with displaying posters in clinics and community centers,
233 implementing school-based oral health programs, and distributing local language videos and pamphlets
234 that emphasize the importance of early consultation.

235 **Acknowledgments:**

236 We sincerely thank our HOD, Dr. Bela Dave, for her guidance and support. We are also grateful to
237 interns, and all the patients who participated in this study for their cooperation.

238

239 **References:**

- 240 1) National Oral Health Survey and Fluoride Mapping 2002–2003 India; Bali RK, Mathur VB, Talwar PP,
241 Chanana HB. Dental Council of India. 2004.
- 242 2) Prevalence of periodontal disease in India; Sharma N, Gupta R, Tandon S. Journal of Indian Society of
243 Periodontology. 2016 Dec; 20(6):593–597.
- 244 3) Reasons for tooth extractions among adults: A global perspective; Chrysanthakopoulos NA.
245 International Dental Journal. 2011 Dec; 61(6):343–351. PMID: 22251317
- 246 4) Periodontal disease and systemic conditions: A bidirectional relationship; Kim J, Amar S.; Odontology
247 2006; 94:10–21. PMID: 17043864
- 248 5) Oral health: A window to general health; Saini R. Indian Journal of Dental Sciences. 2010 Jan; 2(1):1–2.
- 249 6) Factors influencing dental care services utilization in India using Andersen health behaviour model: a
250 systematic review; Singh S, Mathur A, Udhaya E, et al. BMC Health Services Research. 2025 Jul; 25:988.
251 PMID: 40730996

- 252 7)Awareness and knowledge of periodontal diseases among patients visiting dental colleges in India: A
253 questionnaire study; Prasad S, et al. Journal of Indian Society of Periodontology. 2014 Jul-Sep; 18(3):
254 329–334.
- 255 8)Assessment of knowledge and awareness of oral hygiene and scaling among North Indian patients;
256 Kumar S, Tadakamadla J, Johnson NW. Indian Soc Periodontology. 2013 Mar;17(2):221–225. PMID:
257 23946576XXX
- 258 9)Oral health knowledge, attitude, and behavior among rural population of West Godavari district,
259 Andhra Pradesh, India; Srinivas S, Ramesh A, Shashikanth ; Indian Soc Periodontology. 2019 Jul-
260 Aug;23(4):376–381. PMID: 31572031
- 261 10)Awareness about scaling and its effects among the population of Patna, India; Kumar S, et al. Journal
262 of International Oral Health. 2013 May; 5(3):44–47.
- 263 11)Awareness of laser dentistry among patients visiting dental hospitals: A survey; Gupta V, et al. Indian
264 Journal of Dental Sciences. 2013 May; 5(3):33–36.
- 265 12)Diabetes mellitus and periodontal diseases; Mealey BL, Oates TW. Journal of
266 Periodontology. 2006 Aug; 77(8):1289–1303. PMID: 16945027

267

268

269

270

271

272

273

274

275

276

277

278

279

TO EVALUATE AWARENESS AND ACCEPTANCE OF PERIODONTAL DISEASE AND THERAPY AMONG PATIENT VISITING AMC DENTAL COLLEGE: A QUESTIONNAIRE SURVEY

ORIGINALITY REPORT

6%

SIMILARITY INDEX

2%

INTERNET SOURCES

4%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|-----|
| 1 | Rajat Gajjar, Ashish Kaur, Sirsha Bhattacharjee, Sona Sheth, Krishna Shah, Heena Narsinghan. "To evaluate knowledge, awareness and attitude about periodontal disease among medical interns: A questionnaire based study", IP International Journal of Periodontology and Implantology, 2025
Publication | 2% |
| 2 | Submitted to Herzing University
Student Paper | 1% |
| 3 | www.jbclinpharm.org
Internet Source | 1% |
| 4 | Sami A. Zelkha. "Periodontal innate immune mechanisms relevant to atherosclerosis and obesity : Influence of periodontal disease on obesity and atherosclerosis", Periodontology 2000, 08/16/2010
Publication | <1% |
| 5 | jcdr.net
Internet Source | <1% |
| 6 | Sushil Anil Phansopkar, Anmol Mathur, Subhashree Mohapatra. "Root caries prevalence among older adults in Miraj, India: | <1% |

a cross-sectional study", Asian Journal of Gerontology and Geriatrics, 2024

Publication

7 Santhosh Kumar. "Oral hygiene status in relation to sociodemographic factors of children and adults who are hearing impaired, attending a special school", Special Care in Dentistry, 10/23/2008

Publication

8 Singh, Abhinav, Bharathi M. Purohit, Nitin Masih, and Praveen Kumar Kahndelwal. "Risk factors for oral diseases among workers with and without dental insurance in a national social security scheme in India", International Dental Journal, 2013.

Publication

9 assets-eu.researchsquare.com

Internet Source

10 www.researchgate.net

Internet Source

Exclude quotes On

Exclude matches Off

Exclude bibliography On