

An uncommon cause of infective endocarditis in a healthy 11-year-old girl with native heart valve

Abstract:

Infective endocarditis (IE) is a rare disease in children, and it can result in significant morbidity and mortality. The epidemiology of infective endocarditis in children has shifted in recent years with less rheumatic heart disease, more congenital heart disease survival, and increased use of central venous catheters in children with chronic illness. Less commonly, infective endocarditis occurs in children with no preexisting cardiac disease or other known risk factors (1).

IE due to anaerobic bacteria is an uncommon event, accounting for 2—16% of all cases of IE over the past three decades (2).

We present the case of an 11-year-old girl with no known cardiac disease or significant risk factors, who was diagnosed with infective endocarditis according to the modified Duke criteria. Her clinical presentation included a prolonged fever lasting 2.5 months, along with chest pain and easy fatiguability. Blood cultures identified *Anaerococcus prevoti*, a rare pathogen with limited available data and research.

This case underscores the importance of maintaining a high index of suspicion for infective endocarditis in children with prolonged fever, even in the absence of obvious sources of infection or predisposing factors such as cardiac anomalies. Prompt diagnosis is essential for effective treatment and improved patient outcomes.

Case report:

11-year-old previously healthy girl who presented to the emergency department of Al Jalila Children's Hospital (AJCH) in Dubai in Nov 2023 with a history of fever for 2.5 months, with temperatures reaching as high as 39.5°C. The fever was intermittent. She also reported occasional central chest pain with poor weight gain.

Her parents observed that she quickly became fatigued even with trivial exercise. For example, after walking around a mall for a few minutes, she experiences breathlessness and needs to sit down to rest. On a few occasions, her mother noticed central cyanosis, though there were no episodes of syncope.

For the past two months, her father mentioned that she has been complaining of lower back pain without any clear history of trauma or physical activity. They consulted several pediatricians, but symptomatic treatments were ineffective.

Upon further questioning, the father recalled that 10 days before her illness began, she had visited a dentist for a dental extraction. She has had no exposure to pets, no recent travel history, and has not consumed unpasteurized milk. There have been no changes in her appetite or any weight loss.

She was initially treated with a 10-day course of oral amoxicillin and clavulanic acid at a private hospital, but her symptoms did not improve. Due to the persistence of symptoms, she returned to the same hospital, where an echocardiogram revealed vegetation on the mitral valve (MV) along with mitral regurgitation (MR). There was no family history of congenital heart disease. She was referred to AJCH for further management

On admission to AJCH she had a temperature of 36.7°C, a pulse of 124 beats/minute, a blood pressure of 99/68 mmHg, a respiratory rate of 22 breaths/minute, and an oxygen saturation 98% in room air. Her physical examination she looked tired and fatigued. She didn't have any obvious tooth decay. She had no skin rash or petechiae. She was also noted to have clubbing in hand and feet. She had a regular heart rate and rhythm but had systolic murmur 3/6 more at apical and axillary area and much less at bases, no thrill no gallop. Her lungs were clear to auscultation bilaterally, and had soft abdomen without hepatosplenomegaly. The rest of her examination was unremarkable.

A complete blood count revealed a normal white blood cell count of $8.4 \times 10^3/\mu\text{L}$ (5.0 - $13.0 \times 10^3/\mu\text{L}$), haemoglobin 11.6 g/dL (11.5 - 15.5 g/dL) with platelets $293 \times 10^3/\mu\text{L}$ (170 - $450 \times 10^3/\mu\text{L}$). Electrolytes showed a sodium of 140 (136-145 mmol/L), a potassium of 4.2 (3.5-5.1 mmol/L), blood urea of 19 (15.6-40.6 mg/dL), a creatinine of 0.54 mg/dL (0.52-69 mg/dL), and a calcium of 9.3 mg/dL (8.8-10.8 mg/dL). Her C-reactive protein (CRP) was elevated at 8.1 mg/L (normal 0 - 5 mg/L, ESR 36 (Normal 0 - 20 mm/hr), Procalcitonin was 0.06 normal range < 0.5 ng/mL, ASO < 20 (Normal < 150 IU/mL), Troponin T was 4 (Normal < 11 ng/L), CKMB < 0.3 (Normal 0-3.1), GAS antigen throat negative. Urine analysis was normal.

Trans thoracic ECHO was done on the day of admission which showed large vegetation 8 x 14 mm mobile attached to posterior leaflet of the MV and posterior wall of LA. MV anterior leaflet full of Aschoff nodules, AM leaflet not prolapsing. Post leaflet looks prolapsing as vegetation is attached to it. Severe MV regurgitation. Dilated LA, and LV Good cardiac function EF 59% No pericardial effusion.

In the suspicion of IE three blood cultures were collected prior to starting antibiotics and patient was started on Lisinopril, Furosemide, Aspirin, Vancomycin and gentamicin. She became afebrile within 24 hours of admission.

Within 48 hours all 3 peripheral blood cultures came positive for *Anaerococcus prevotii*. After 48 hours of starting IV antibiotics blood culture was collected again which was also reported positive for *Anaerococcus prevotii*. Eye examination was done which was normal. After we got the culture sensitivity for the first culture the antibiotics were changed to IV ampicillin and clindamycin. Blood culture collected on day 5 of antibiotics which was reported negative and she was continued on total of IV clindamycin for 16 days and ampicillin for 7 weeks.

Table : 1

Susceptibility

	Anaerococcus prevotii AJCH SUSCEPTIBILITY
Ampicillin	Susceptible
Clindamycin	Susceptible
PIPERA./TAZOBACTAM	Susceptible
Vancomycin	Susceptible

ECHO was repeated after 2 weeks of admission showed large vegetation -Thread like measures 25-9 mm in length and 3 mm in width, friable mobile attached to atrial surface of posterior leaflet of the MV -Part of it is Mitral valve leaflet and Chordae. There were also two perforations across posterior leaflets one was big and the other is smallish causing Moderate Plus MR.


Following ECHO almost after 3 weeks of admissionshowed persistence of large vegetation, mobile attached to posterior leaflet of the MV and posterior wall of LA. Post leaflet looked prolapsing as vegetation is attached to it and leaflet torn at the level of P2. Severe MV regurgitation, Mild PR, 6 mmhg dilated left atrium and left ventricle. In view of the ECHO findings cardiac surgeons decided to repair the mitral valve and excise the vegetations. Prior to surgery 2 peripheral blood cultures were noted to be negative and patient was given antibiotics for almost 3 weeks prior to surgery.

During surgery it was found the main reason of the MV regurgitation was due to a floating segment of the MV posterior leaflet which was covered by an elongated vegetation. There was also a generalized picture of post IE changes on the rest of the mitral valve. These were mostly healed changes with good quality tissues. No vegetations were identified elsewhere. The posterior leaflet had an abnormal short secondary attachment which was also tethering segments of the valve leading to significant degree of regurgitation.The Vegetation was removed fully with the damaged scallop of the MV. Excision of all infected or suspected rims of tissues from the LA side of the MV. The resultant quadrangular defect on the posterior leaflet was closed. The incised rims of tissue were sent for culture which was reported negative. Following day of surgery, ECHO was repeated which showed successful surgical results,MV well repaired, no MR,no vegetations, no effusion, good - improved cardiac function.She received almost 4 weeks of IV ampicillin after surgery and was continued on aspirin and lisinopril with follow up with pediatric cardiology team. She was discharged home in stable condition.

109

110

111 Figure 1 :



112

113

114 Figure 2:

UNDER PTL





Figure 3:



Figure 4:




119

120 Figure 1- 4 :It shows the echocardiography before the surgery. Showing Large vegetation. It is
121 mobile attached to posterior leaflet of the mitral valve and posterior wall of Left atrium .Mitral
122 valve anterior leaflet full of Aschoff nodules,Post leaflet looks prolapsing as vegetation is
123 attached to it. Severe mitral valve regurgitation.


124 Figure 5:

UNDER PETER



125

126 Figure 6:



127

Figure 5-6 shows echocardiography post surgery . It shows mitral valve well repaired, no mitral regurgitation no mitral stenosis, no vegetations,no effusion

Discussion:

Infective endocarditis refers to the inflammation of the endocardium and the valves of the heart. It is mainly caused by bacterial infections and can present with a variety of symptoms and complications. If not diagnosed and treated promptly, numerous intracardiac and distant extracardiac issues can arise. Thus, a comprehensive evaluation including an in-depth medical history, as pointed out in our case and physical examination, is essential for diagnosing the condition and guiding treatment to reduce both mortality and morbidity.

Infectious endocarditis is a rare condition with an estimated yearly incidence of 3 to 10 cases per 100,000 people.[3] Historically, this disease process has demonstrated a predilection for males, with a male to female ratio of nearly 2 to 1. The average age of infectious endocarditis patients is now greater than 65 years old. This preponderance for the elderly likely corresponds to the increased prevalence of predisposing factors such as prosthetic valves, indwelling cardiac devices, acquired valvular disease, hemodialysis, and diabetes mellitus within this demographic.[4] Although previously a major risk factor, rheumatic heart disease now underlies less than 5% of all cases in the modern antibiotic era. Recreational intravenous drug use represents a growing risk factor that now accounts for about 10% of all infectious endocarditis cases.[5]

Awareness among the family physician and primary physician about this entity is very crucial in early diagnosis and appropriate management.

The vast majority of infectious endocarditis cases stem from gram-positive streptococci, staphylococci, and enterococci infection. Together, these three groups account for 80% to 90% of all cases, with *Staphylococcus aureus* specifically responsible for around 30% of cases in the developed world.[6]

Infections due to anaerobic bacteria are common, and can be serious and life-threatening. The recent increase in the recovery of these organisms from all infectious sites [7], including bacteremia [8], has led to greater appreciation of the role anaerobes play in infections at all body sites, including infective endocarditis (IE). Most cases of anaerobic IE are caused by the anaerobic and microaerophilic streptococci, *Propionibacterium acnes* and *B. fragilis* [9], [10].

The mortality rate for patients with anaerobic IE is 21–43% [11], [12] *Anaerococcus prevotii* is a gram-positive, strictly anaerobic bacteria that belongs to genus *Anaerococcus*. It is part of the commensal human microbiota. However, it has been associated with various infections, including ovarian abscesses, chronic wounds, vaginal discharge, foot ulcers, knee arthritis, urinary tract infections, pleural empyema, blood infections, and soft tissue infections. This bacterium is non-motile and does not form spores.

In adults there have been some case reports of *Anaerococcus prevotii* infections in various clinical cases, including injuries from road traffic accidents (RTAs) and odontogenic brain abscesses.

There are no documented pediatric cases of IE due to *Anaerococcus prevotii* or any other *Anaerococcus* species in the medical literature. Our patient represents what appears to be the first reported case in a child, reinforcing the need to consider rare and unusual pathogens in febrile children without classic risk factors for endocarditis.

This case further highlights the importance of:

- Maintaining a high index of suspicion for IE in children with prolonged fever.
- Utilizing echocardiography early in the evaluation.
- Ensuring thorough microbiologic workup including multiple blood cultures.
- Considering anaerobic coverage when clinical suspicion is high.

Advanced molecular techniques such as metagenomic sequencing may further assist in diagnosing culture-negative IE, especially in the setting of fastidious or slow-growing organisms such as *Anaerococcus* spp.

Conclusion:

This case highlights the complex presentation and management of infective endocarditis (IE) caused by *Anaerococcus prevotii* in a previously healthy pediatric patient. While *Anaerococcus prevotii* is part of the commensal microbiota, it is rarely reported as a causative agent of endocarditis, making this case particularly unusual. To our knowledge, there have been no reported cases, particularly in pediatric patients with native valves. Infective endocarditis should always be considered in febrile children, even in the absence of known heart disease or other obvious risk factors. It is important to remain vigilant for even the rarest causative organisms, such as *Anaerococcus prevotii*, which was identified after nearly 48 hours of incubation in our case.

References:

- 1- Salloum S, Bugnitz CJ. A case report of infective endocarditis in a 10-year-old girl. Clin Pract. 2018 Jul 10;8(3):1070. doi: 10.4081/cp.2018.1070. PMID: 30090217; PMCID: PMC6060482.
- 2- Brook I. Infective endocarditis caused by anaerobic bacteria. Arch Cardiovasc Dis. 2008 Oct;101(10):665-76. doi: 10.1016/j.acvd.2008.08.008. Epub 2008 Nov 11. PMID: 19056073.
- 3- Cahill TJ, Prendergast BD. Infective endocarditis. Lancet. 2016 Feb 27;387(10021):882-93. [PubMed]
- 4- Correa de Sa DD, Tleyjeh IM, Anavekar NS, Schultz JC, Thomas JM, Lahr BD, Bachuwar A, Pazdernik M, Steckelberg JM, Wilson WR, Baddour LM. Epidemiological trends of infective endocarditis: a population-based study in Olmsted County, Minnesota. Mayo Clin Proc. 2010 May;85(5):422-6. [PMC free article] [PubMed]

- 207 5- Murdoch DR, Corey GR, Hoen B, Miró JM, Fowler VG, Bayer AS, Karchmer AW,
208 Olaison L, Pappas PA, Moreillon P, Chambers ST, Chu VH, Falcó V, Holland DJ, Jones
209 P, Klein JL, Raymond NJ, Read KM, Tripodi MF, Utili R, Wang A, Woods CW, Cabell
210 CH., International Collaboration on Endocarditis-Prospective Cohort Study (ICE-PCS)
211 Investigators. Clinical presentation, etiology, and outcome of infective endocarditis in the
212 21st century: the International Collaboration on Endocarditis-Prospective Cohort
213 Study. *Arch Intern Med*. 2009 Mar 09;169(5):463-73. [PMC free article] [PubMed]
214 6- Barnett R. Infective endocarditis. *Lancet*. 2016 Sep 17;388(10050):1148. [PubMed]
215 7- I. Brook Anaerobic infections in children *Microbes Infect*, 4 (2002), pp. 1271-1280 View
216 PDF View article View in Scopus Google Scholar
217 8- B. Lassmann, D.R. Gustafson, C.M. Wood, J.E. Rosenblatt Reemergence of anaerobic
218 bacteremia *Clin Infect Dis*, 44 (2007), pp. 895-900 View at publisher Crossref Google
219 Scholar
220 9- C.E. Nord Anaerobic bacteria in septicemia and endocarditis *Scand J Infect Dis* (1982),
221 pp. 95-104 View in Scopus Google Scholar
222 10- D. Kiani, E.L. Quinn, K.H. Burch, T. Madhavan, L.D. Saravolatz, T.R. Neblett. The
223 increasing importance of polymicrobial bacteremia. *J Am Med Assoc*, 242 (1979),
224 pp. 1044-1047. View at publisher Crossref View in Scopus Google Scholar
225 11- J.M. Felner, V.R. Dowell Jr. Anaerobic bacterial endocarditis *N Engl J Med*, 283 (1970),
226 pp. 1188-1192 View in Scopus Google Scholar
227 12- M.D. Sans, J.G. Crowder Subacute bacterial endocarditis caused by *Eubacterium*
228 *acrofaciens*. Report of a case *Am J Clin Pathol*, 59 (1973), pp. 576-580 View at
229 publisher Crossref View in Scopus Google Scholar