

The impact of varying bulb sizes, plant spacing, and planting dates on the yield and quality of onion seeds var.

Taherpuri

by Jana Publication & Research

Submission date: 24-Oct-2025 12:23PM (UTC+0300)

Submission ID: 2769518898

File name: IJAR-54485.pdf (1.44M)

Word count: 11914

Character count: 51093

¹³
The impact of varying bulb sizes, plant spacing, and planting dates on the yield and quality of onion seeds

var. Taherpuri

⁷³
Abstract

³
Background and objective: Onion is a spice crops, which are particularly important because it uses in daily life. The crop is also appreciated as condiments for flavoring foods. It can be used as a stimulant, promotes the flow of urine, uses against tuberculosis, venereal disease, typhoid fever, chest and lungs problem and it contains vitamin B and trace of vitamin C and traces of iron and calcium. The objective to study to improve the yield and quality of onion seeds (var. Taherpuri) through management practices.

²⁸
Methodology: The experiment was designed using the Randomized Complete Block Design (RCBD) with three replications and included three parameters. The unit plot was randomly assigned to the experiment's treatments. In every trial, the unit plot size was the same (2 m × 1 m). Plots were separated by 0.5 meters and blocks by 1 meter.

⁷
Main Results: The tallest plant height (48.50, 57.96 and 62.11 cm) and the maximum number of leaves per plant (15.52, 23.25 and 24.66) at 45, 60 and 75 DAP was found from the large bulb size (16-20 g) × widest plant spacing (25 cm x 20 cm) × early planting on 25th October treatments interaction, respectively. The highest seed yield of 1075.00 kg/ha, maximum weight of 1000 seeds (3.75 g), the highest germination percentage (88.33%), the highest shoot length (8.73 cm), highest root length (3.23 cm) and the maximum seed vigor index (1057.10) was obtained from the treatment interaction of the largest bulb (16-20 g) × widest plant spacing (25 cm x 20 cm) × early planting on 25th October treatments interaction.

Conclusion: The integrated use of large mother bulbs, optimal plant spacing, and early planting significantly enhances onion seed yield and quality. Specifically, the combination of largest bulbs (16–20 g) and wider plant spacing (25 × 20 cm) with early planting dates (25th October) consistently results in the highest seed yield up to 1,075 kg/ha along with superior quality traits: heavier 1,000-seed weight, higher germination rates, longer shoots and roots, and elevated vigor index.

Key words: Onion, *Allium cepa*, spacing, planting date, bulb, yield and quality

Introduction

The onion (*Allium cepa* L.), a member of the Alliaceae family, is the most widely grown and significant spice crop worldwide including in Bangladesh. For the manufacture of bulbs, it has been grown annually for more than 4700 years. Additionally, it is a vegetable in many parts of the world (Brewster, 1994). According to McCollum (1976), the Mediterranean region is the second-largest origin center for onions, with Central Asia

33 being the principal center. According to FAO (2022), China is the world's largest producer of onions, followed
34 by India, the United States, and then Pakistan, Korea, the Netherlands, Israel, Japan, Turkey, Syria, Egypt,
35 Lebanon, and so on. Bangladesh generates 2% of global production annually, while China and India account for
36 26% and 21% of global production, respectively (Quaiyum, 2020).

37

38 The onion is referred to be the "Queen of the kitchen" because of its highly prized flavor, scent, and distinct
39 taste as well as the therapeutic qualities of its flavor compounds (Selvaraj, 1976). Of all the spice crops, onions
40 are particularly important because of their many applications in daily life. The crop is also appreciated as
41 condiments for flavoring foods. It can be used as a stimulant, promotes the flow of urine, uses against
42 tuberculosis, venereal disease, typhoid fever, chest and lungs problem. Onion contains vitamin B and trace of
43 vitamin C and traces of iron and calcium. In onions, allyl-propyl-disulphide, a volatile oil, is the cause of the
44 onion's notable pungency. According to Ahmed (2020), the renowned physician Hippocrates recommended
45 onions as emmenagogues, laxatives, and diuretics. Additionally, he utilized onions to cure pneumonia and to
46 topically heal septic wounds. The most prevalent vitamin in onion bulbs is ascorbic acid, which has a
47 concentration of 1 mg/g dry weight. It also contains steroidal saponins, which stop the intestines from absorbing
48 cholesterol. In 2023, the top 20 nations that produced the most onions generated 80.04 million tons annually,
49 with China, India, and the United States accounting for 24.97, 22.82, and 3.17 million tons, respectively
50 (FAOSTAT, 2023). In BBS, 2023, the other 17 nations each produced over 1.36 million tons. Egypt is the
51 world's fourth-largest producer, with 3.08 million tons produced annually in Africa (FAO, 2023). Additionally,
52 according to FAOSTAT (2023), 93.23 million metric tons of onions are produced annually worldwide. With
53 24.04 million metric tons produced annually, China leads the globe in onion production, followed by India with
54 21.42 million metric tons. In the larger districts of Pabna, Faridpur, Rajbari, Manikganj, Kushtia, Rajshahi,
55 Jhenaidah, Magura, Meherpur, Madaripur, Shariatpur, Gopalganj, Narayanganj, Dinajpur, and Rangpur in
56 Bangladesh, onions are grown extensively. Out of all the 64 districts, Pabna produces the most onions. The
57 country's total onion acreage was 1.87 lakh hectares, of which 42566.40 hectares were in Pabna and 2617
58 hectares were in Faridpur (Aker et al., 2023). Roughly 75.79% of the nation's onion production comes from
59 Pabna, Faridpur, Rajbari, Rajshahi, Kushtia, and Jhenaidah (BBS, 2023). Bangladesh's 3.5 million metric tons of
60 annual demand for winter onions are primarily fulfilled by domestically grown winter onions. Still, 237,000
61 hectares of land are used to grow the current production, which is between 2.5 and 2.6 million tons. At a cost of
62 almost BDT 40,000 million, the nation must import one million tons of onions annually to meet the present
63 demand (ACI Seed Limited, 2023). Multi-crop cultivation methods, the challenges of increasing cultivation
64 acreage in the winter, and the lack of adequate land all make output shortfalls and storage losses worse.

65 According to ¹⁹ Rahim et al. (1993), however, its seeds are grown in the districts of Faridpur, Natore, Pabna,
66 Rajshahi, Dinajpur, and Thakurgaon.

67

68 Farmers employ a variety of seed types, including their own seed, enhanced seed, local quality seed, HYV and
69 hybrid seeds, which result in varying onion bulb sizes. About 1245 metric tons of seeds were needed in 2018–19
70 to cultivate 1.73 lakh hectares of land (Quaiyum, 2020). It is estimated that farmers produce 90% of the seeds,
71 with BADC and other private companies producing the remaining 10% (Quaiyum, 2020). Thus, a market crisis
72 among Bangladeshi onion growers during the growing season is caused by a shortage of edible onion bulbs and
73 an enormous amount of onion seed used to produce bulbs annually. Practically speaking, compared to bulb
74 crops, onion plants grown from the bulb for seed production must be maintained in the field for a month as a
75 standing crop. In addition to impacting onion producers and the nation's total onion crop, lower seed production
76 makes the seed market susceptible. Because of some environmental constraints, such as Bangladesh's brief winter
77 season, plants grown from seed typically do not produce quality seed in the same season. The bulb-to-bulb
78 method is generally suitable for Bangladeshi onion seed production. In Bangladesh, because the winter season is
79 brief and mild, onions are typically planted when the season begins. This crop, which is favored by short days,
80 aids in seed production. Climate constraints prevent Bangladeshi farmers from using early planting. Due to late
81 rainfall or floods, illnesses frequently have a significant impact on early-planted onion seed crops (Ahmed,
82 2020). Normal seed growth and fruit setting are also impacted by high temperatures. In nature, onions are cross-
83 pollinated by flies, bees, and other insects. To maximize the potential of onion seed, a significant population of
84 pollination insects must be present.

85

86 In Bangladesh, more and more land is being planted with onions each year, but farmers are having trouble
87 finding high-quality seeds. Small farmers continue to supply high-quality onion seeds despite their lack of
88 expertise in bulb grade selection. Our nation's farmers always produce onion seeds using tiny bulbs.
89 Furthermore, the production of high-quality onion seeds is significantly impacted by the size of the mother bulb.
90 Low quality and low seed production are caused by our nation's onion seed growers' ignorance of bulb size.
91 Ahmed et al. (2020) discovered that for the Taherpuri variety, large bulbs outperformed small bulbs. Brewster
92 (1994) noted that high-quality seeds and ideal plant spacing were thought to be crucial for excellent production,
93 quality, and plant growth. Numerous studies have demonstrated the importance of plant and row spacing for the
94 optimal plant population, which may be reflected in the best production and quality (Kumar et al. 2018). Onions
95 are a crop that is sensitive to photo thermoregulation (Davis and Jones, 1944). Temperature, humidity, rainfall,
96 and length of day are some of the environmental elements that affect onion seed production. These elements are

essential for seed growth and setting (Mishra and Mishra, 1991). For the largest yield and highest-quality onion seeds, Mondal (1980) believed that the latter week of October was the ideal time to plant.

Even with the right planting time and spacing between bulbs, small farmers still distribute high-quality onion seed because they lack the information necessary to choose the right bulb grade. Therefore, the experiment was conducted to find the ideal planting date, plant spacing, and bulb size for a high yield and excellent quality of onion seeds of var. Taherpuri in order to obtain detailed information about the aforementioned characteristics.

Materials and methods

Experimental site

During the 2020–21 rabi season, an experiment was conducted to examine the impact of planting date, plant spacing, and bulb size on the yield and quality of onion seeds of the Taherpuri variety. The laboratory work was overseen at the MNT seed testing lab in Bashon, Gazipur, while the research was carried out at the Lal Teer Research Farm. The experimental site is 8.4 meters above sea level and lies between 23°53' and 24°21' N latitudes and 90°09' and 90°39' E longitudes. The soil on the experimental site is clay loam, moderately acidic in reaction (soil pH 6–6.5, Haider et al. 1991), and has poor fertility status. It is part of the Modhupur tract, which is classified as being in Agro Ecological Zone-28 (FAO, 1988).

Climatic conditions

The most significant determinant of onion seed production is the agroclimatic component. The experimental area had a sub-tropical climate, which was marked by scarce rainfall, low humidity, low temperatures, and short days during the "Rabi" season (October–March) and high rainfall from April to September. Additionally, this climate is distinguished by several seasons, such as the per-monsoon period or hot season from March to April, the winter or dry season from November to February, and the monsoon, which lasts from May to October (Edris et al., 1979).

Experimental material

The experimental material was the bulbs of "Taherpuri or BARI Piaz-1 (HRC, 1996), a cherished local open pollinated cultivar. This variety is well grown in Taherpur area at Bagmara Upazila of Rajshahi district according to the name of Taherpur area, it is called Taherpuri.

General characteristics of onion variety (var. Taherpuri)

It is selected from the local variety Taherpuri, selected by HRC, BARI, Joydebpur, Gazipur and registered by the NSB, MOA, and GOB in 1996 to grow all over the country. The variety is characterized by 50-55cm plant height, 10-12 leaves, round-cum-flat shape and compact single bulb, thin necked, reddish-pink bulb, highly pungent, good shelf life (8-9 months), takes 130-140 days for bulb maturation, yields 12-16 tons/ha of fresh bulbs. The picking of matured umbels is made within 145-155 days of planting mother bulbs. The yield of seed is 950-1050 kg/ha. The diseases such as purple (*Alternaria porri*) blotch, leaf blight and insect like thrips and cutworms may attack the plants.

Source of the seed quality test

The healthy, disease free and desired sizes of seeds were obtained from the store of (at ambient temperature) Lal Teer R&D farm, Bashedon, Gazipur. Seeds obtained from the respective experiments conducted for the seed production, were used for the quality test such as germination, 1000 seed weight and vigor test.

Preparing the land

On October 15, 2020, the experimental ground was first ploughed with a tractor-drawn disc plough. Four cross-ploughs were then performed using a power tiller and ladder. The earth was then let to dry out. They spaded the land's corners. For the soil to be in good tilth condition, it was then harrowed. A ladder then leveled the land completely. We cleared the field of stubble and weeds. Every clod was fragmented into tiny bits. Before planting, the unit plots were also easily prepared with a spade.

Design and treatments of experiments

Three replications of the experiment were set up using a randomized complete block design (RCBD). Various numbers of unit plots were created for each block based on the amount of treatments. The unit plot's experimental treatments were assigned at random. The experiment's bulb size, plant spacing, and planting date varied, but the unit plot size remained the same at 2 m × 1 m. Plots were separated by 0.5 meters and blocks by 1.0 meters. The experiment involved planting three different bulb sizes (11, 12, 15, and 16 g) at four different

spacing's (25 cm × 20 cm, 25 cm × 15 cm, 20 cm × 20 cm, and 20 cm × 15 cm) on three separate dates (October 25, November 5, and November 15).

Observation data

During the study period, morphological data were recorded, including seed germination, stalk length, plant height, and number of leaves per plant, as well as yield contributing characteristics, such as flower per umbel, fruit set percentage, seed yield, seed weight per hill, 1000 seed weight, etc.

Measurement and data collection

At various growth phases, information on various growth, seed yield, and seed quality metrics was gathered from each plot's middle two rows. Data were collected on plant height (cm), leaves per plant (no), tillers per plant (no), length (cm) of the flowering stalk, umbels per plant (no), umbel diameter (cm), flowers per umbel (no), days to 80% maturity, seeded fruits per umbel (no), seed yield per plant (g), seed yield per plot (g), and seed yield per hectare by randomly selecting ten plants from each plot's two central rows. It was represented in kg/ha and was derived from the corresponding seed yield per plot. At Desire Seed Laboratory, however, seed quality contributing factors such seed weight (g), seed germination (%), and seed vigor test were carried out in accordance with ISTA (1985).

Records of growth and yield parameters

Except for the percentage of bulbs that emerged, data on growth and development were gathered from ten randomly chosen plants for each treatment.

Statistical analyses

To determine the statistical significance of the experimental results, statistical analysis was performed on the gathered data on various growth, yield, and quality contributing parameters. Using Microsoft Excel, all treatments were computed, analyzed, and tabulated. The F-variance test was used to analyze the variance of the data for each parameter using Statistix 10 and RStudio software. The least significant difference (LSD) test was used to assess the significance between treatment means at 1% and 5% of provability for result interpretation (Gomez and Gomez, 1994).

Results and discussion

Growth parameters

Plant height and leaf count

In terms of plant height and the number of leaves per plant at 45, 60, and 75 DAP, the interaction impact of bulb size, plant spacing, and planting date was shown to be statistically significant. The findings showed that the bulb size (16–20 g) × spacing (25 cm × 20 cm) × planting date at 25 October were used to determine the tallest plant height (48.50, 57.96, and 62.11 cm) and the largest number of leaves per plant (15.52, 23.25, and 24.66) (Figure 1 and 2). The harvest from the smallest bulbs had the shortest plants at every stage of growth. This could be because large bulbs store a comparatively larger amount of internal food stores than small bulbs, which promotes robust vegetative growth and increased plant height. The number of leaves dropped as the size of the bulb shrank. The reason why larger bulbs produce more leaves than smaller ones may be because they store more food material and have more sprouting leaf initials (Brewster, 1994).

The current results are in good accord with the findings of Islam et al. (2008), Haque (2011), Verlag (2013), and Mukhtadir et al. (2001), who also observed a substantial influence of bulb size on the augmentation of plant height. The findings of Kumar et al. (2015) also complement these findings. According to reports by Ali et al. (2015) and SK-Phor et al. (2019), the greatest mother bulb size resulted in the highest plant height. These findings corroborated those of Phor et al. (2019), who found that crops planted with the widest spacing generated noticeably taller plants than crops planted with the narrowest spacing. In contrast to the last eight plantings, the early planting may have given the plants a comparatively cooler period, which could have contributed to the rise in plant height. The cooler weather promotes the buildup of gibberellin and cytokines, which alters the hormonal balance and causes the plant to grow more. It is also the cause of the flower stalk's elongation. According to research by Ashagrie et al. (2014), onions planted in October grew to their maximum height when compared to those planted in November. These findings are also consistent with those of other authors who found that larger bulbs generated more leaves per plant since there were more food supplies stored, which led to more robust plants. Haque (2011), Asaduzzaman et al. (2012), and Hussain et al. (2001). According to a study by Mollah et al. (2015), onion plants planted in November had more leaves per plant and were longer than those planted in October. The study site's varying climate (temperature) may be the cause of this discrepancy.




Fig 1. Interaction effects of bulb size, plant spacing and planting date on plant height (cm) of onion seeds var. Taherpuri




Fig 2. Interaction effects of bulb size, plant spacing and planting date on leaves/plant of onion seeds var. Taherpuri

Note: B₁ = ~ 11 g size bulb, B₂ = 12-15 g size of bulb and B₃ = 16-20 g size of bulb, D₁ = Planting distance 25 cm × 20 cm, D₂ = 25 cm × 15 cm, D₃ = 20 cm × 20 cm and planting distance D₄ = 20 cm × 15 cm, T₁ = Planting date 25 October, T₂ = 5 November and T₃ = 15 November.

14

72

Number of flowers and fruits per plant, number of tillers, length of stalk, number of umbels, and diameter of umbels

umbels

1

Statistically significant effects were observed in the number of tillers, stalk length, umbel number, umbel diameter, and number of flowers and fruits per plant when varying bulb size, plant spacing, and planting date were combined. According to Table 1, the interaction effect of bulb size (16–20 g) × plant spacing (25 cm × 20 cm) × planting date (25 October) resulted in the highest number of tillers per plant (3.46), the longest flowering stalk (95.03 cm), the number of umbels per plant (3.50), the maximum umbel diameter (7.50 cm), the highest number of flowers per umbel (486.43), and the maximum number of seeded fruits per umbel (374.0). The current findings closely align with those of other researchers. The findings of Ashagrie (2021) corroborate the considerable influence of bulb size on improving plant growth indices, as reported by Muktadir et al. (2000), Hussain et al. (2001), Haque (2011), Asaduzzaman et al. (2012), and El-Helaly and Karam (2012). Additionally, Singh and Sachan (1999) discovered that the plants with the widest spacing had the most tillers per plant. Asaduzzaman et al. (2012) and Haque (2011) also discovered that the plants with the widest spacing had more tillers per plant. Because there was less competition for nutrients, light, space, and moisture, plants with the widest spacing produced more green leaves. In the end, these leaves promoted the production of new flowering stalks by accumulating photosynthate. The widest spacing also produced the greatest number of blooming stalks per plant, according to Singh and Sachan (1999). The current findings were nearly identical to those of Haque (2011) and Asaduzzaman et al. (2012), who reported that the length of the flowering stalk rose as plant separation increased. The reason for this could be that the plants with the widest spacing helped to develop more green leaves and extra nourishment, which could later promote the length of the flowering stalk.

255

The number of umbels per plant and the width of the umbels were found to be superior when bulbs were planted with a larger spacing of 25 cm by 20 cm, followed by another treatment, and a smaller umbel was planted with a spacing of 20 cm by 15 cm. The quantity of umbels and their diameter per plant, however, rose as planting spacing increased. With the wider plant spacing, there may be less competition amongst the plants for nutrients, light, and moisture. Significantly larger umbel was the outcome of the greater nitrogen rate at wider spacing (Nehra et al., 1988; Pandey et al., 1992). These results can be explained by the fact that plants with wider spacing absorb more water and nutrients, particularly nitrogen, than plants with closer spacing, which may account for the larger and more numerous umbels. These results are similarly consistent with those of Khan (2003), Singh and Sachan (1999a), and Hoque (2010) and Haque (2011), who also found that greater spacing led to a bigger umbel diameter. Both Phor et al. (2019) and Asaduzzaman et al. (2012) concurred that wider-spaced crops result in higher umbel diameters. Wider spacing was seen to result in a significant increase in the

number of flowers per umbel. Bulbs spaced at the widest spacing of 25 cm × 20 cm produced the most blooms per umbel (350.14), while bulbs spaced at the closest spacing of 20 cm × 15 cm produced the fewest 311.80. It's possible that this is because the widest spacing resulted in the largest umbel diameter and length by absorbing more nutrients, which in turn promoted the production of the most flowers per umbel. Begum (1998) also made a comparable observation.

While the number of seeded fruits gradually decreased with the closest spacing, the maximum number of seeded fruits per umbel (263.76) was recorded from the wider spacing of 25 cm × 20 cm. The preceding aspects mentioned in the earlier criteria may be to blame for this. Begum (1995) found that the greater spacing resulted in the greatest quantity of seeded fruits per umbel. These findings also supported those of Haque (2011) and Asaduzzaman et al. (2012), who found that crops planted with the widest spacing produced considerably more seeded fruits per umbel than crops planted with the narrowest spacing. Regarding growth parameters such as the number of umbels, the diameter of the umbels per plant, and the length of the stalks, there was a notable fluctuation in the planting time. Out of the three planting dates, the outcomes were best when done early (on October 25) as opposed to late (on November 15). Larger bulbs had enough reserve food to support plant growth, which led to the maximum plant growth attributes. Other possible explanations for the increase in plant growth parameters include early planting, which may have given plants a comparatively cooler period compared to later plantings. This outcome is consistent with research by Haque (2011) and Nag and Oishimaya (2017), which found that planting date may have an impact on plant growth parameters. Furthermore, Khodadadi (2012) found that the date of onion bulb planting significantly impacted plant growth characteristics. The current investigation's findings are consistent with those of Healy and Karam (2012), as well as with those of Kumar et al. (2018), Thalkari et al. (2019), and Ashagrie et al. (2021). Because of the planting timing, there was a substantial (5% level) variation in the amount of flowers and seeded fruits available per umbel. The crop was planted early on October 25th, which produced the most flowers and seeded fruits per umbel, and late on November 15th, which produced the fewest flowers and seeded fruits per umbel. This could be because of the crop's poor growth and delayed bolting in November, as well as the high temperatures during the umbel-forming stage, which may have decreased the quantity of flowers and seeds per umbel. The current findings are in good agreement with those of Mishra (1986), Nehra et al. (1989), and Singh and Singh (1985). Furthermore, comparable findings were observed by Ashagrie et al. (2014) and Thalkar (2019), who found that crops planted early generated the most flowers and seeds per umbel.

298 **Table 1.** The number of tillers, stalk length, umbel count, umbel diameter, and number of flowers and fruits per
 299 onion seed plant are all impacted by the interaction of bulb size, plant spacing, and planting date.

Interaction	Number of	Length of	No. of	Umbel	No. of flowers	No. of fruits		
effect	tillers/plant	flowering	umbels/ plant	diameter (cm)	/umbel	/umbel		
stalks (cm)								
B	D	T1	2.43k-o	89.40a-e	2.57h-k	6.20f-k	297.87mnopq	231.60f-l
		T2	2.37l-o	88.23a-f	2.53h-k	6.17f-l	293.53opqr	224.77g-m
		T3	2.27m-p	86.23def	2.43i-l	6.07g-n	290.20pqr	218.07i-m
	D	T1	2.50j-n	87.90b-f	2.53h-k	6.10f-m	294.30nopq	224.97g-m
		T2	2.33mno	87.00c-f	2.37i-l	5.90i-o	284.30qrs	151.83n
		T3	2.20nop	86.20def	2.23jkl	5.70mno	274.30s	213.33klm
	D	T1	2.47k-n	88.43a-f	2.43i-l	5.87j-o	289.53pqr	226.00g-m
		T2	2.37l-o	87.13b-f	2.37i-l	5.77l-o	274.27s	216.00j-m
		T3	2.27m-p	86.93c-f	2.23jkl	5.67no	274.20s	209.33lm
	D	T1	2.37l-o	83.70ef	2.30i-l	6.03h-n	285.67qrs	232.63e-l
		T2	2.10op	83.90ef	2.20kl	5.87j-o	289.00pqrs	215.97j-m
		T3	1.93p	82.03f	2.10l	5.60o	279.00rs	205.97m
B	D	T1	2.97d-h	88.27a-f	3.10b-f	6.97bc	332.53hij	241.27d-i
		T2	2.83f-j	88.10b-f	3.07b-f	6.90cd	312.87l	239.27d-j
		T3	2.77g-k	87.00c-f	3.03c-f	6.87cde	308.93lmn	233.53e-l
	D	T1	2.87f-i	93.93ab	2.90e-h	6.50def	360.97de	253.17c-f
		T2	2.60i-m	92.60a-d	2.63ghi	6.33fgh	347.63efg	237.17d-k
		T3	2.50j-n	88.93a-f	2.60g-j	5.80k-o	317.63jkl	229.17f-m
	D	T1	2.90e-i	92.57a-d	2.97d-g	5.97h-o	339.53fghi	235.03e-k
		T2	2.83f-j	91.60a-d	2.90e-h	6.03h-n	332.87ghi	223.97g-m
		T3	2.70h-l	91.17a-d	2.87fgh	5.97h-o	302.87lmnop	218.37i-m
	D	T1	3.00c-h	91.23a-d	3.10b-f	6.46efg	342.00fghi	248.17c-g
		T2	2.43k-o	91.23a-d	2.40i-l	6.20f-k	312.00lm	222.50h-m
		T3	2.37l-o	89.67a-e	2.30i-l	5.93h-o	306.33lmno	205.83m
B	D	T1	3.47a	95.03a	3.50a	7.50a	486.43a	347.00a
3	1	T2	3.33abc	94.03ab	3.37abc	7.37ab	419.43b	327.50ab

	T3	3.27a-d	93.70abc	3.30a-d	7.17abc	409.43b	310.83b
D	T1	3.40ab	93.23abc	3.43ab	6.30f-i	381.00c	268.80c
	T2	3.30a-d	90.57a-e	3.33a-d	6.20f-k	364.00d	260.13cd
	T3	3.17a-f	89.57a-e	3.20a-f	6.03h-n	345.67fgh	246.80c-h
D	T1	3.37ab	93.70abc	3.40abc	6.50def	332.60hi	256.97cde
	T2	3.23a-e	92.37a-d	3.27a-e	6.30f-i	349.27def	247.63c-g
	T3	3.17a-f	90.97a-d	3.27a-e	6.17f-l	332.60hi	236.63d-k
D	T1	3.30a-d	93.00a-d	3.20a-f	6.23f-j	346.93efgh	256.87cde
	T2	3.17a-f	92.00a-d	3.17a-f	6.13f-l	328.30ijk	230.87f-l
	T3	3.10b-g	90.37a-e	3.07b-f	6.13f-l	316.93kl	226.50g-m
	LSD (0.05)	0.25	5.20	0.29	0.31	11.25	18.39
	LSD (0.01)	0.33	6.90	0.39	0.42	14.948	24.42
	Level of sig.	**	**	**	**	**	**
	CV (%)	5.59	3.56	6.50	3.12	2.12	4.74

Note: ** = Significant at 1% level of probability. B₁ = ~ 11 g size bulb, B₂ = 12-15 g size of bulb and B₃ = 16-20 g size of bulb, D₁ = Planting distance 25 cm × 20 cm, D₂ = 25 cm × 15 cm, D₃ = 20 cm × 20 cm and planting distance D₄ = 20 cm × 15 cm, T₁ = Planting date 25 October, T₂ = 5 November and T₃ = 15 November.

Seed yield (kg/ha)

The seed yield had a considerable impact on the interactions between bulb size, spacing, and planting date. When the data on seed yield per plot was converted to yield (kg/ha), the smallest bulb, widest spacing, and early planting all showed the same upward trend. The largest bulb (B₃) × widest plant spacing (D₁) × early planting (T₁) treatments produced the highest calculated seed yield of 1075.00 kg/ha, while the bulb planted of the smallest bulb (B₁) × closest spacing (D₄) × late planting of (T₁) interaction produced the lowest performance for the same parameter, 700.0 kg/ha (Table 2). As crops grew from larger bulbs, the seed production per hectare increased, whereas the small bulb treatment yielded a lower yield. As the size of the bulb expanded, so did the seed weight. The disparity in seed output may have resulted from the comparatively large food reserves kept in large bulbs, which increased the formation of robust, healthy plants with abundant seed heads and, as a result,

increased seed yield kg/ha (Table 2). This outcome supports the findings of Haque (2011), who found that the size of the mother bulb could have an impact on the amount of onion seed produced per hectare, with large mother bulbs producing the maximum harvest and small mother bulbs producing the lowest. The findings of Healy and Karam (2012), Ashagrie et al. (2014), Ali et al. (2015), Mollah et al. (2015), Debashis et al. (2016), Khan (2016), Thalkari (2019), Ahmed et al. (2020), and Ashagrie et al. (2021) that the larger bulb produced the higher seed yield per hectare are also consistent with this result.

The highest seed output was obtained from bulbs planted at the tightest spacing (20 cm × 15 cm) and the widest spacing (25 cm × 20 cm). As plant spacing increased, there was a discernible rise in seed yield per plot and per hectare (Table 2). This result validates the findings of Kanwar et al. (2000) and Cuocolo and Barbieri (1988). Similar outcomes were also reported by Haque (2011), who achieved higher seed yields per hectare and per plot with wider plant spacing. Early planting on October 25 produced the highest seed yield, which may have been caused by pollination-friendly weather conditions that resulted in more seed sets per umbel and per plant. Late planting on November 15 produced the lowest seed yield per unit area, which may have been caused by weather conditions that negatively affected pollination and seed set, resulting in fewer seed sets per plant (Table 2). According to Healy and Karam (2012), flower abortion may potentially be the cause of the decreased seed output in crops that are planted later. Furthermore, our findings are consistent with those of Haque (2011), Asaduzzaman et al. (2012), and Thalkari (2019), who found that the highest seed production was observed on an early planting calendar. The findings of Ashagrie et al. (2021), Debashis et al. (2016), Ali et al. (2015), and Mollah et al. (2015) are all consistent with this outcome.

Quality parameters

The interaction effect of large bulb size (16-20 g) × the widest spacing (25 cm × 20 cm) × early planting (25 October) produced the highest values for the 1000-seed weight (3.75 g), the highest germination percentage (88.33%), the maximum root length (3.23 cm), the maximum shoot length (8.73 cm), and the lowest seed vigor index (1057.10). This result is consistent with larger onion bulbs increasing the weight of 1000 seeds, as reported by Ali et al. (1998), Muktadir (2000), Asaduzzaman et al. (2012), Khan (2016), Ahmed et al. (2020), and Ashagrie et al. (2021). The germination rate was higher for seeds harvested from large bulbs and lower for seeds harvested from small bulbs. The medium-sized bulbs were used to produce seeds with germination percentages in the middle. Asaduzzaman et al. (2012), Islam (2008), Hoque (2010), El-Helaly and Karam (2012), Khan (2016), and Ashagrie (2021) also found higher germination percentages of seed from larger bulbs, which was in keeping with the aforementioned findings. The bulb size significantly increased the seed vigor

index. The findings showed that as bulb size increased, the vigor index rose as well. The larger bulbs may have a higher vigor index because their roots and shoots are longer and their seeds germinate at a higher rate than those of smaller bulbs. This result is consistent with that of Ashagrie et al. (2014) and Badawi et al. (2010). Ashagrie et al. (2014) discovered that the big bulbs had the highest seed vigor index.

Additionally, the results of Badawi et al. (2010) were nearly identical to the current findings, indicating that the quality of onion seed as measured by the percentage of germination shoot and root lengths, varied significantly across the three mother bulb sizes. These outcomes also supported Khan's (2016) findings, which showed that the maximum SVI from large bulbs was much higher than that from smaller ones. Ali et al. (1998) also discovered that the wider spacing resulted in a much greater onion seed weight of 1000. Moreover, Haque (2011) and Asaduzzaman et al. (2012) concurred that the onion's 1000-seed weight was noticeably greater at the wider spacing. The plants that are closer together may have less access to water and nutrients. According to Khan (2003a) and Hogue (2010), who found a higher percentage of seed germination at wider spacing, the current result is consistent with their findings. According to Teshome et al. (2015), a wider spacing and fewer fungal pathogen incidences often resulted in a higher percentage of seeds germinating. This outcome is similar to that of Asaduzzaman et al. (2012), who found that the maximum spacing resulted in the highest germination percentage (92.64%) while the smallest spacing resulted in the lowest germination percentage (83.85%). When compared to seeds with the closest intra-row spacing, the widest plant spacing produced the highest seed vigor index, which may be explained by the maximum root and shoot length and germination % of seeds. This could be because, in contrast to plants with closer spacing, the plants with wider spacing had greater access to food materials for the developing seeds, while the plants with closer spacing had less access to water and nutrients. According to Teshome et al. (2015), this may also be because plants with small populations (wide spacing) have less rivalry for assimilation and seed distribution, which could lead to vigor growth.

Similar findings were made by Singh et al. (1985), who showed that a wider spacing improved the seedling vigor index more than a narrower spacing did. In general, when space rose (plant population density decreased), the seed vigor index increased as well. This may be explained by less rivalry amongst plants for growth components, which promotes greater development and the accumulation of food sources in the seed for the following generation. In their study, Sarker et al. (2021) discovered that the wider spacing outperformed the closing ones in terms of germination percentage, seedling length, and seedling vigor index. They did this with spacing of 30 cm × 20 cm (S1), 25 cm × 20 cm (S2), and 20 cm × 20 cm (S3). Significant differences in seed germination were noted depending on the planting date. According to the results, seeds originating from the 25

October planting date had the highest seed germination rate (78.78%), while seeds originating from the 15 November planting date had the lowest seed germination rate (70.92%). According to the following authors, this outcome is consistent. Early planting produced the highest percentage of seed germination, according to El-Helaly and Karam (2012) and Ashagrie et al. (2021), who also found that planting date had a major impact on seed germination. Malik et al. (1999) found that early planting produced the highest seed vigor index, which was connected to this finding. According to Ashagrie et al. (2021), early planting resulted in a significantly higher seed vigor index after the planting date.

Table 2. Effects of planting date, plant spacing, and bulb size on the yield and quality characteristics of onion seed varieties. Taherpuri,

Interaction	Seed yield	1000 seed weight	Germination	Shoot length	Root	Seed vigor
effect	(kg/ha)	(g)	(%)	(cm)	length (cm)	Index
T1	970.0d	3.46c-g	79.00d-h	7.96b-f	2.53d-g	829.8e-i
D1 T2	891.7ij	3.36e-i	78.00e-k	7.93b-g	2.66c-f	827.0e-i
T3	850.0mn	3.16i-m	71.00n-qr	7.06i-n	2.60d-g	686.8l-p
T1	915.0gh	3.35e-i	78.33e-j	7.63d-j	2.90a-d	825.2e-j
D2 T2	841.7n	3.30f-k	77.00f-l	7.60e-j	2.63c-f	788.3g-k
T3	795.0p	3.10klm	70.33o-s	6.70mno	2.56d-g	652.0n-q
T1	895.0ij	3.36e-i	74.33j-o	7.20g-m	2.56d-g	726.2j-o
D3 T2	816.7o	3.26g-l	73.00l-p	7.13h-m	2.23g	684.0m-p
T3	775.0q	3.06lmn	66.33st	6.33nop	2.23g	568.6qr
T1	820.0o	3.15i-m	68.00q-t	6.96j-n	2.56d-g	648.5opq
D4 T2	741.7r	3.06lmn	67.33rst	6.73l-o	2.33fg	610.5pq
T3	700.0t	2.85n	61.00u	6.06op	2.23g	507.1r
T1	995.0c	3.65abc	79.00d-h	8.50ab	3.13ab	919.4b-e
D1 T2	916.7g	3.55a-e	82.66bcd	7.60e-j	2.90a-d	868.7c-h
T3	875.0kl	3.35e-i	75.66h-m	7.56e-k	2.80b-e	784.7h-l
T1	940.0ef	3.60a-d	82.66bcd	8.16a-e	3.13ab	934.8bcd
D2 T2	861.7lm	3.50b-f	81.33cde	8.13a-e	2.90a-d	897.6c-f
T3	820.0o	3.30f-k	74.66i-n	7.26f-m	2.80b-e	751.9i-n

	T1	920.0g	3.55a-e	78.66d-i	7.73c-i	2.80b-e	828.7e-i
D3	T2	841.7n	3.45c-h	77.66e-k	7.70c-j	2.56d-g	797.7f-k
	T3	800.0p	3.25g-m	70.00p-s	6.83k-n	2.46efg	651.7opq
	T1	845.0n	3.33e-j	73.00l-p	7.50e-k	2.80b-e	752.9i-m
D4	T2	766.7q	3.23h-m	72.00m-q	7.46e-l	2.56d-g	722.5k-o
	T3	725.0s	3.03mn	64.66tu	6.53m-p	2.46efg	582.3qr
	T1	1075.0a	3.75a	88.33a	8.73a	3.23a	1057.1a
D1	T2	996.7c	3.65abc	87.33a	7.83b-h	3.00abc	946.0bcd
	T3	955.0de	3.45c-h	80.33d-g	7.83b-h	2.90a-d	862.7d-h
	T1	1020.0b	3.70ab	86.00ab	8.43abc	3.23a	1003.5ab
D2	T2	941.7e	3.60a-d	85.00abc	8.36a-d	3.00abc	966.8abc
	T3	900.0hi	3.40d-h	74.66i-n	7.50e-k	2.90a-d	776.9h-m
B3	T1	998.3c	3.65abc	81.66cde	7.96b-f	2.90a-d	888.3c-g
D3	T2	921.5g	3.45c-h	81.00c-f	8.06a-e	2.66c-f	869.7c-h
	T3	880.0jk	3.35e-i	74.00k-p	7.06i-n	2.56d-g	713.3k-o
	T1	925.0fg	3.43c-h	76.33g-l	7.70c-j	2.90a-d	809.5f-k
D4	T2	846.7mn	3.35e-i	75.33h-m	7.70c-j	2.66c-f	781.0h-m
	T3	805.0op	3.11j-m	68.33q-t	5.80p	2.53d-g	568.9qr
LSD _{0.05}		11.62	0.16	3.21	0.57	0.28	75.32
LSD _{0.01}		15.43	0.22	4.26	0.75	0.37	100.00
Level of sig.		**	**	**	**	**	**
CV (%)		0.81	2.98	2.60	4.68	6.42	5.93

** = Significant at 1% level of probability. B₁ = ~ 11 g size bulb, B₂ = 12-15 g size of bulb and B₃ = 16-20 g size of bulb, D₁ = Planting distance 25 cm × 20 cm, D₂ = 25 cm × 15 cm, D₃ = 20 cm × 20 cm and planting distance D₄ = 20 cm × 15 cm, T₁ = Planting date 25 October, T₂ = 5 November and T₃ = 15 November.

Economic analysis

The findings from economic analysis in Table 3 demonstrated that the larger the seed bulbs (16–20 g), the higher the gross return, net return, and gross margin. However, the smallest net return, gross margin, and benefit-cost ratio (BCR) were obtained from the tiny seed bulbs (~11 g). Because of the greater seed rate, the larger bulbs had the highest possible overall cost of production. Accordingly, the larger seed rate and larger

yield from the large seed bulbs may be the cause of the larger net-return, higher gross margin, and lower BCR from the large bulbs (Table 3).

Accordingly, differences in production costs, net returns, and benefit-cost ratios were noted for various bulb sizes. Although the benefit cost ratio (2.17) was higher for small size bulbs because of their relatively lower production costs and smaller quantity, the cost of production (TK. 584410/ha) and net return (TK 681050/ha) were highest for large size bulbs. The benefit cost ratio (1.49) was lowest for large bulbs because of the relatively bigger quantity of bulbs and higher production costs, while the net return (TK. 440652/ha) and production cost (TK. 780335/ha) were lower for tiny bulbs.

Table 3. The impact of bulb size and plant spacing on the cost of production, net profit, gross return, and benefit cost ratio (BCR) of Taherpuri onion seeds

Interaction		Cost of production	Gross return	Net return	BCR
effect		(tk ha ⁻¹)	(tk ha ⁻¹)	(tk ha ⁻¹)	
B1	D1	604410	1265460	681050	2.09
	D2	619159	1169575	550416	1.89
	D3	611140	1119015	507875	1.83
	D4	651235	998918	347682	1.53
B2	D1	614110	1311683	709572	2.14
	D2	648859	1223460	574601	1.89
	D3	640840	1174113	533272	1.83
	D4	680935	1051515	370580	1.54
B3	D1	605550	1432905	789395	2.37
	D2	708259	1359308	651048	1.92
	D3	700240	1306620	606380	1.87
	D4	780335	1160988	440652	1.49

Conclusion

The integrated use of large mother bulbs, optimal plant spacing, and early planting significantly enhances onion seed yield and quality. Specifically, the combination of largestbulbs(16–20 g) and wider plant spacing (25 × 20 cm) with early planting dates (around late October)consistently results in the highest seed yield up to

1,075 kg/ha along with superior quality traits: heavier 1,000-seed weight, higher germination rates, longer shoots and roots, and elevated vigor index.

Acknowledgments

The authors acknowledge the support of the Lal Teer Seed Company Limited, Bangladesh for providing the necessary leave of deputation, all research inputs and bearing the cost of field experiment. The authors would like to thank Md Rezaul Karim, Professor, Bangladesh Agricultural University, Mymensingh-2202 and Dr. Md Jahangir Alam, Cotton Development Board, Ministry of Agriculture, Bangladesh for their help and support.

Funding

Not funding is available.

Authors and Affiliations

Lal Teer Seed Company Limited, Bangladesh

Md Asaduzzaman

Professor, Bangladesh Agricultural University, Mymensingh-2202

Md Rezaul Karim

Professor, Bangladesh Agricultural University, Mymensingh-2202

Md Abdur Rahim

Cotton Development Board, Ministry of Agriculture, Bangladesh

Md. Jahangir Alam

Professor, Bangladesh Agricultural University, Mymensingh-2202

Md Shafiqul Islam

Professor, Bangladesh Agricultural University, Mymensingh 2202, Bangladesh

Ahmed Kairul Hasan

Author Contribution

Conceptualization: [MA]; Methodology: [JA, SI, AKH]; Formal analysis and investigation: [JA, SI, AKH];

Writing-original draft preparation: [MA, MRK, JA]; Writing-review and editing: [JA, MAR, SI, AKH]; Funding

acquisition: [MA]; Supervision: [SI, AKH]

Ethics declarations

Competing Interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

References

- Ahmed S, Rahim MA, Moniruzzaman M, Khatun MA, Jahan FN, Akter R 2020: Effect of bulb sizes on seed yield of two onion (*Allium Cepa* L.) varieties. *SAARC Journal of Agriculture* **18** 51 - 65.
- Akter S, Rahman KT, Anik AR and Amin MR 2023: Growth and trend analysis of onion production in Bangladesh. *Annals of Bangladesh Agriculture* **27** 201 - 210.
- Ali AM, Hossain MM, Zakaria M, Naznin A, Islam MM 2015: Effect of bulb size on quality seed production of onion. *International Journal of Agronomy and Agricultural Research* **4** 174 - 180.
- Ali M, Quadir MA, Chowdhury AR, Alam MZ 1989: Effect of positive selection of mother bulbs on seed and bulb yields of onion. *Bangladesh Journal of Agriculture* **14** 57-63
- Asaduzzaman MD, Hasan MM, Hasan MM, Moniruzzaman MD, Howlader KHM 2012: Effect of bulb size and plant spacing on seed production of onion (*Allium cepa* L.) *Bangladesh Journal of Agricultural Research* **37** 405 - 414.
- Ashagrie TD, Belew S, Alamerew Y, Getachew 2014: Effects of planting time and mother bulb size on onion (*Allium cepa* L.) seed yield and quality at Kobo Woreda Northern Ethiopia. *International Journal of Agriculture Research* ISSN 1816 - 4897.
- Ashagrie T, Belew D, Nebiyu A and Moral MT (2021): Influence of planting date and bulb size on yield and quality of onion (*Allium cepa* L.) seed production. *Journal of Cogent Food and Agriculture* **7** 2382.
- Badawi MA, Seadth SE, El-Emery MI and Shalaby AEM 2010: Onion seed yield and its quality as influenced by storage methods mother bulb size and harvesting time. Mansoura University, Egypt, *Journal of Plant Production* **1** 239 - 249.
- BBS 2023: Year Book of Agricultural Statistics of Bangladesh, Bangladesh Bureau of Statistics, Statistics Division, Ministry of Planning, Government of the People's Republic of Bangladesh.
- Begum A 1995: Effects of planting time, irrigation, set size and spacing on the seed production of some onion cultivars. M. S Thesis, Department of Horticulture, BAU, Mymensingh p80.

481 Begum A, Rahim MA, Haider MA 1998: Effect of set size and spacing on the seed production of onion.
 482 *Bangladesh Journal of Seed Science Technology* **2** 91-100.
 483 Brewster JL 1994: Onion and other vegetable Alliums. *CAB International* UK 236.
 484 Brewster JI 1994: Onions and other Vegetable Alliums. *Horticulture Research International* Wellesbourne UK
 485 p236.
 486 Cuocolo L and Barbieri G 1988: Effect of nitrogen fertilization and plant density on seed yield of onion (*Allium*
 487 *cepa* L.). *Rivista – di – Agronomia* **22** 195 - 202.
 488 Davis GN and Jones AA 1944: Experiment with the transplant onion crop in California. Bull. California
 489 Agricultural Experiment Station, 682 p20.
 490 Debashis M, Santra P, Maity TK and Basu AK 2016: Quality seed production of onion (*Allium cepa* L.) cv.
 491 Sukhsagar as influenced by bulb size and date of planting. *Bangladesh Journal of Agricultural Research*
 492 *and Technology* 1–6.
 493 Edris KM, Islam AMT, Chowdhury MS and Haque AKMM 1979: Detailed Soil Survey of Bangladesh, Dept.
 494 Soil Survey, BAU and Govt. Peoples Republic of Bangladesh. p118.
 495 FAO 2022: Food and Agricultural Quarterly Bulletin of Statistics, Rome, Italy **10** 91-94.
 496 FAO 1988: Land Resources Appraisal of Bangladesh. Report 6, FAO, Rome, Italy.
 497 FAOSTAT 2023: Statistical Year Book of Food and Agricultural Organization of the United
 498 Nations, Rome, Italy 356.
 499 Gomez KA, Gomez AA 1984: *Statistical Procedure for Agricultural Research*. John Wiley and Sons, New
 500 York 680.
 501 Haider R, Rahman AA and Huq S (eds.) 1991: "Cyclone '91: An Environmental and Perceptual Study",
 502 Bangladesh Centre for Advanced Studies, Dhaka, 91 pp.
 503 Healy MA, Karam SS 2012: Influence of planting date on the production and quality of onion seeds.
 504 *Journal of Horticulture Science & Ornamental Plants* **4** 275-279.
 505 Haque MR 2011: Development of package of technology for quality seed production of onion var.
 506 Taherpuri. Ph.D. Thesis, Department of Horticulture, Patuakhali Science and Technology
 507 University, Patuakhali, Bangladesh.
 508 Hoque OM 2010: Manipulation of growth cycle in summer onion seed crop with specific reference to
 509 high yield and quality seed production in Bangladesh. Ph.D. Thesis, Dept. of Hort., Hajee
 510 Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh 233.
 511 HRC 1996 BARI Onion-1: A high yielding variety of onion. A Leaflet, published by Horticulture
 512 Research Centre, Bangladesh agricultural Research Institute, Joydebpur, Gazipur.

513 Hussain SW, Ishtiaq M, and Hussain, SA 2001: Effects of different bulb sizes and planting dates on green leaf
514 production of onion (*Allium cepa* L.). *Journal of Biological Science* **8** 433-439.

515 ISTA 1985: International rules for seed testing. International Seed Testing Association, *Seed Science and*
516 *Technology* **13** 299-355.

517 Islam ZM, Rahim MA, Rahman MM, Alam MJ 2008: Effect of planting time, bulb size and shading on yield
518 and quality of onion seed. *Bangladesh Journal of Seed Science and Technology* **12** 23-28.

519 Khan MA, Rahim MA 2003a: Effect of bulb size and plant spacing on yield and quality of onion seed.
520 *Bangladesh Journal of Seed Science Technology* **7** 149-154.

521 Khan MA 2016: Integrated crop management for quality seed production of onion. Ph.D. Thesis, Department
522 of Horticulture, Hajee Mohammad Danesh Science and Technology University, Dinajpur,
523 Bangladesh.

524 Kanwar JS, Gill BS and SS 2000: Response of planting time and density to onion seed yield and quality. *Seed*
525 *Research* **28** 212-214.

526 Khodadadi M, 2012: The effects of planting date and mother bulb size on quantitative and qualitative seed traits of
527 onion Red Rey Variety. *International Journal of Agriculture Research and Review* **2** 324-327.

528 Kumar P, Savita, Kumar S, Thakur Y, Kaur D and Kamboj A 2018: Effect of planting density and inorganic
529 fertilizers on growth and yield of onion. Lovely Professional University, Panjab. *International Journal*
530 *of Current Microbiology and Applied Science* **7** 2319-7706.

531 Kumar S, Tomar BS, Jain SK, Singh N, Prasad R, and Munshi AD 2015: Effect of planting time and density on plant
532 growth, seed yield and quality attributes in onion (*Allium cepa*) cv. *Pusa Riddhi*. *Indian Journal of Agricultural*
533 *Sciences* **85** 1578-1585.

534 Malik YS, Singh N, Nehra BK 1999: Effect of planting time, bulb cut pinching of bolt treatment on yield and
535 quality of onion seed. *Vegetable Science* **26** 143-145.

536 Mc Collum GD 1976: Evaluation of Crop Plants. Ed. N.W. Simmonds, Logman, London and New York 186-
537 190.

538 Mishra HP and Mishra B 1991: Effect of time of planting and bulb size on the yield and yield components of
539 onion seed crop. *Indian Journal of .Agricultural Research* **25** 107-112.

540 Mishra HP 1986: Effect of time of planting and size of mother bulbs on onion seed production. *Indian Journal of*
541 *Horticulture Science* **43** 140-143.

542 Mondal MF and Husain A 1980: Effect of time of planting onion bulbs on the yield and quality of seeds.
543 *Bangladesh Journal of Agriculture* **5** 131-134.

544 Mollah, MRA, Ali MA, Ahmad M, Hassan MK, and Alam MJ 2015: Effect of bulb size on the yield and
545 quality of true seeds of onion. *European Journal of Biotechnology and Bioscience* **3** 23–27.

546 Muktadir MS 2000: Effect of planting time, bulb size and vernalization on the yield and quality of onion seed.
547 MS Thesis, Department of Horticulture BAU, Mymeningh 124.

548 Muktadir MS, Farooque AM, Rahim MA and Hossain MM 2001: Yield and quality of onion seed as influenced
549 by the planting time and bulb size. *Bangladesh Journal of Seed Science Technology* **5** 47-52.

550 Nag S, and Oishimaya 2017: The top onion producing countries in the World. Worldatlas.com/articles/the - top -
551 onion - producing-countries - in-the world.
552 html.<https://www.agriculturenigeria.com/production/cropproduction/horticulture/onion/>

553 Nehra BK, Pandita M.L, Singh K 1989: Cultural and nutritional studies in relation to seed production in onion.
554 *Haryana Journal of Horticulture Science* **17**106-110.

555 Nehra BK, Pandita M.L, Singh K 1988: Cultural and nutritional studies in relation to seed production in onion.
556 *Haryana Journal of Horticulture Science* **17**106-110.

557 Pandey UC, Dhingra RP, Singh K and Mangal JL 1992: Effect of N fertilizer, spacing and their combined seed
558 yield of onion var. Hissar 2. *Progressive Horticulture* **14** 75-77.

559 Pandey UB, Panwar DS, Sarma VP 1992: Effect of spacing and nitrogen on growth and seed yield of kharif
560 onion. *Seed Research* **20** 147-148.

561 Phor SK 2019: Effect of spacing and bulb size on growth, yield and economics of onion seed crop in Haryana.
562 *Journal of Pharmacognosy Phytochemistry* **5** 59-62.

563 Quaiyum A 2020: Demand and supply of onion in Bangladesh. The Daily Observer, Thursday, 29 October,
564 2020.

565 Rahim MA, Amin MMU and Haider MA 1993: Onion seed production technologies in Bangladesh. Allium
566 Improvement Newsletter (USA) **3** 33.

567 Sarker Pc, Kaiser MO, Mozumder SN, Hossain MA, Sen R and Choudhury AK 2021: Effect of plant spacing
568 and fertilizer dose on seed yield and quality of onion. A Scientific Research, Regional Agricultural
569 Research Station (RARS), BARI, Cumilla, Bangladesh.

570 Selvaraj S 1976: Onion: queen of the kitchen. *Kisan World* **3** 32-34.

571 Singh KP, Malik YS, and Lal S 1985: Effect of stickling, planting and spacing in radish seed crop. *Seed Research* **13**
572 176-179.

573 Singh SR, Sachan BP 1999: Evaluation of different bulb size, spacing and varieties for higher seed yield and
574 yield attributing traits on onion (*Allium cepa* L.). *Crop Research Hissar* **17** 351-355.

575 Teshome H, Woldeeslassie A, Simon T 2015: Seed Yield and Quality of Onion (*Allium Cepa* Var. *Cepa*).
576 Seed as influenced by bulb treatment and spacing patterns at Larena, Southern Ethiopia, *Journal*
577 *of Natural Sciences Research* 5 2225-0921.
578 Thalkari GN, BM Kallalbandi BM, RD Baghele RD and MM Maind MM 2019: Effect of spacing, bulb size and
579 time of planting on onion seed yield (*Allium cepa* L.) *International Journal of Chemical Studies* 73383-3386.
580 Verlag G, 2013: The Effect of Bulb Size and Plant Density on Yield and Quality of Onion (*Allium cepa* L.)
581 seed, at Ziway, Central Ethiopia. *Google-eBook Technology and Engineering* 70.

582

583

584 **Table 1.** The number of tillers, stalk length, umbel count, umbel diameter, and number of flowers and fruits per
585 onion seed plant are all impacted by the interaction of bulb size, plant spacing, and planting date.

Interaction	Number of	Length of	No. of	Umbel	No. of flowers	No. of fruits	
effect	tillers/plant	flowering	umbels/ plant	diameter (cm)	/umbel	/umbel	
stalks (cm)							
D 1	T1	2.43k-o	89.40a-e	2.57h-k	6.20f-k	297.87mnopq	231.60f-l
	T2	2.37l-o	88.23a-f	2.53h-k	6.17f-l	293.53opqr	224.77g-m
	T3	2.27m-p	86.23def	2.43i-l	6.07g-n	290.20pqr	218.07i-m
D 2	T1	2.50j-n	87.90b-f	2.53h-k	6.10f-m	294.30nopq	224.97g-m
	T2	2.33mno	87.00c-f	2.37i-l	5.90i-o	284.30qrs	151.83n
	T3	2.20nop	86.20def	2.23jkl	5.70mno	274.30s	213.33klm
B 1	T1	2.47k-n	88.43a-f	2.43i-l	5.87j-o	289.53pqr	226.00g-m
	T2	2.37l-o	87.13b-f	2.37i-l	5.77l-o	274.27s	216.00j-m
	T3	2.27m-p	86.93c-f	2.23jkl	5.67no	274.20s	209.33lm
D 4	T1	2.37l-o	83.70ef	2.30i-l	6.03h-n	285.67qrs	232.63e-l
	T2	2.10op	83.90ef	2.20kl	5.87j-o	289.00pqrs	215.97j-m
	T3	1.93p	82.03f	2.10l	5.60o	279.00rs	205.97m
B 1	T1	2.97d-h	88.27a-f	3.10b-f	6.97bc	332.53hij	241.27d-i
	T2	2.83f-j	88.10b-f	3.07b-f	6.90cd	312.87l	239.27d-j
	T3	2.77g-k	87.00c-f	3.03c-f	6.87cde	308.93lmn	233.53e-l
D 2	T1	2.87f-i	93.93ab	2.90e-h	6.50def	360.97de	253.17c-f
	T2	2.60i-m	92.60a-d	2.63ghi	6.33fgh	347.63efg	237.17d-k

	T3	2.50j-n	88.93a-f	2.60g-j	5.80k-o	317.63jkl	229.17f-m
D 3	T1	2.90e-i	92.57a-d	2.97d-g	5.97h-o	339.53fghi	235.03e-k
	T2	2.83f-j	91.60a-d	2.90e-h	6.03h-n	332.87ghi	223.97g-m
	T3	2.70h-l	91.17a-d	2.87fgh	5.97h-o	302.87lmnop	218.37i-m
D 4	T1	3.00c-h	91.23a-d	3.10b-f	6.46efg	342.00fghi	248.17c-g
	T2	2.43k-o	91.23a-d	2.40i-l	6.20f-k	312.00lm	222.50h-m
	T3	2.37l-o	89.67a-e	2.30i-l	5.93h-o	306.33lmno	205.83m
D 1	T1	3.47a	95.03a	3.50a	7.50a	486.43a	347.00a
	T2	3.33abc	94.03ab	3.37abc	7.37ab	419.43b	327.50ab
	T3	3.27a-d	93.70abc	3.30a-d	7.17abc	409.43b	310.83b
D 2	T1	3.40ab	93.23abc	3.43ab	6.30f-i	381.00c	268.80c
	T2	3.30a-d	90.57a-e	3.33a-d	6.20f-k	364.00d	260.13cd
	T3	3.17a-f	89.57a-e	3.20a-f	6.03h-n	345.67fgh	246.80c-h
D 3	T1	3.37ab	93.70abc	3.40abc	6.50def	332.60hi	256.97cde
	T2	3.23a-e	92.37a-d	3.27a-e	6.30f-i	349.27def	247.63c-g
	T3	3.17a-f	90.97a-d	3.27a-e	6.17f-l	332.60hi	236.63d-k
D 4	T1	3.30a-d	93.00a-d	3.20a-f	6.23f-j	346.93efgh	256.87cde
	T2	3.17a-f	92.00a-d	3.17a-f	6.13f-l	328.30ijk	230.87f-l
	T3	3.10b-g	90.37a-e	3.07b-f	6.13f-l	316.93kl	226.50g-m
	LSD (0.05)	0.25	5.20	0.29	0.31	11.25	18.39
	LSD (0.01)	0.33	6.90	0.39	0.42	14.948	24.42
	Level of sig.	**	**	**	**	**	**
	CV (%)	5.59	3.56	6.50	3.12	2.12	4.74

Note: ** = Significant at 1% level of probability, B₁ = ~ 11 g size bulb, B₂ = 12-15 g size of bulb and B₃ = 16-20 g size of bulb, D₁ = Planting distance 25 cm × 20 cm, D₂ = 25 cm × 15 cm, D₃ = 20 cm × 20 cm and planting distance D₄ = 20 cm × 15 cm, T₁ = Planting date 25 October, T₂ = 5 November and T₃ = 15 November.

594

595

596 **Table 2.** Effects of planting date, plant spacing, and bulb size on the yield and quality characteristics of onion
 597 seed varieties. Taherpuri,

Interaction		Seed yield	1000 seed weight	Germination	Shoot length	Root	Seed vigor
effect		(kg/ha)	(g)	(%)	(cm)	length	Index
		(cm)					
B1	T1	970.0d	3.46c-g	79.00d-h	7.96b-f	2.53d-g	829.8e-i
	D1 T2	891.7ij	3.36e-i	78.00e-k	7.93b-g	2.66c-f	827.0e-i
	T3	850.0mn	3.16i-m	71.00n-qr	7.06i-n	2.60d-g	686.8l-p
	T1	915.0gh	3.35e-i	78.33e-j	7.63d-j	2.90a-d	825.2e-j
	D2 T2	841.7n	3.30f-k	77.00f-l	7.60e-j	2.63c-f	788.3g-k
	T3	795.0p	3.10klm	70.33o-s	6.70mno	2.56d-g	652.0n-q
	T1	895.0ij	3.36e-i	74.33j-o	7.20g-m	2.56d-g	726.2j-o
	D3 T2	816.7o	3.26g-l	73.00l-p	7.13h-m	2.23g	684.0m-p
	T3	775.0q	3.06lmn	66.33st	6.33nop	2.23g	568.6qr
B2	T1	820.0o	3.15i-m	68.00q-t	6.96j-n	2.56d-g	648.5opq
	D4 T2	741.7r	3.06lmn	67.33rst	6.73l-o	2.33fg	610.5pq
	T3	700.0t	2.85n	61.00u	6.06op	2.23g	507.1r
	T1	995.0c	3.65abc	79.00d-h	8.50ab	3.13ab	919.4b-e
	D1 T2	916.7g	3.55a-e	82.66bcd	7.60e-j	2.90a-d	868.7c-h
	T3	875.0kl	3.35e-i	75.66h-m	7.56e-k	2.80b-e	784.7h-l
	T1	940.0ef	3.60a-d	82.66bcd	8.16a-e	3.13ab	934.8bcd
	D2 T2	861.7lm	3.50b-f	81.33cde	8.13a-e	2.90a-d	897.6c-f
	T3	820.0o	3.30f-k	74.66i-n	7.26f-m	2.80b-e	751.9i-n
B2	T1	920.0g	3.55a-e	78.66d-i	7.73c-i	2.80b-e	828.7e-i
	D3 T2	841.7n	3.45c-h	77.66e-k	7.70c-j	2.56d-g	797.7f-k
	T3	800.0p	3.25g-m	70.00p-s	6.83k-n	2.46efg	651.7opq
	T1	845.0n	3.33e-j	73.00l-p	7.50e-k	2.80b-e	752.9i-m
	D4 T2	766.7q	3.23h-m	72.00m-q	7.46e-l	2.56d-g	722.5k-o
	T3	725.0s	3.03mn	64.66tu	6.53m-p	2.46efg	582.3qr

	T1	1075.0a	3.75a	88.33a	8.73a	3.23a	1057.1a
D1	T2	996.7c	3.65abc	87.33a	7.83b-h	3.00abc	946.0bcd
	T3	955.0de	3.45c-h	80.33d-g	7.83b-h	2.90a-d	862.7d-h
	T1	1020.0b	3.70ab	86.00ab	8.43abc	3.23a	1003.5ab
D2	T2	941.7c	3.60a-d	85.00abc	8.36a-d	3.00abc	966.8abc
	T3	900.0hi	3.40d-h	74.66i-n	7.50e-k	2.90a-d	776.9h-m
B3	T1	998.3c	3.65abc	81.66cde	7.96b-f	2.90a-d	888.3c-g
D3	T2	921.5g	3.45c-h	81.00c-f	8.06a-e	2.66c-f	869.7c-h
	T3	880.0jk	3.35e-i	74.00k-p	7.06i-n	2.56d-g	713.3k-o
	T1	925.0fg	3.43c-h	76.33g-l	7.70c-j	2.90a-d	809.5f-k
D4	T2	846.7mn	3.35e-i	75.33h-m	7.70c-j	2.66c-f	781.0h-m
	T3	805.0op	3.11j-m	68.33q-t	5.80p	2.53d-g	568.9qr
LSD _{0.05}		11.62	0.16	3.21	0.57	0.28	75.32
LSD _{0.01}		15.43	0.22	4.26	0.75	0.37	100.00
Level of sig.		**	**	**	**	**	**
CV (%)		0.81	2.98	2.60	4.68	6.42	5.93

** = Significant at 1% level of probability. B₁ = ~ 11 g size bulb, B₂ = 12-15 g size of bulb and B₃ = 16-20 g size of bulb, D₁ = Planting distance 25 cm × 20 cm, D₂ = 25 cm × 15 cm, D₃ = 20 cm × 20 cm and planting distance D₄ = 20 cm × 15 cm, T₁ = Planting date 25 October, T₂ = 5 November and T₃ = 15 November.

Table 3. The impact of bulb size and plant spacing on the cost of production, net profit, gross return, and benefit cost ratio (BCR) of Taherpuri onion seeds

Interaction effect		Cost of production (tk ha ⁻¹)	Gross return (tk ha ⁻¹)	Net return (tk ha ⁻¹)	BCR
B1	D1	604410	1265460	681050	2.09
	D2	619159	1169575	550416	1.89
	D3	611140	1119015	507875	1.83
	D4	651235	998918	347682	1.53
B2	D1	614110	1311683	709572	2.14
	D2	648859	1223460	574601	1.89
	D3	640840	1174113	533272	1.83

	D4	680935	1051515	370580	1.54
B3	D1	605550	1432905	789395	2.37
	D2	708259	1359308	651048	1.92
	D3	700240	1306620	606380	1.87
	D4	780335	1160988	440652	1.49

Fig 1.Interaction effects of bulb size, plant spacing and planting date on plant height (cm) of onion seeds var. Taherpuri





Fig 2.Interaction effects of bulb size, plant spacing and planting date on leaves/plant of onion seeds var. Taherpuri



The impact of varying bulb sizes, plant spacing, and planting dates on the yield and quality of onion seeds var. Taherpuri

ORIGINALITY REPORT

21 %

SIMILARITY INDEX

18 %

INTERNET SOURCES

17 %

PUBLICATIONS

5 %

STUDENT PAPERS

PRIMARY SOURCES

1

www.researchgate.net

Internet Source

2 %

2

archive.saulibrary.edu.bd:8080

Internet Source

2 %

3

www.sac.org.bd

Internet Source

2 %

4

www.banglajol.info

Internet Source

1 %

5

www.bowl.com

Internet Source

1 %

6

Teshome Ashagrie, Derbew Belew, Amsalu Nebiyu. " Influence of planting date and bulb size on yield and quality of onion (L.) seed production ", Cogent Food & Agriculture, 2021

Publication

1 %

7

www.saulibrary.edu.bd

Internet Source

1 %

8

Submitted to Northeast Community College

Student Paper

1 %

9

ageconsearch.umn.edu

Internet Source

<1 %

10

www.chemijournal.com

Internet Source

<1 %

11

www.scialert.net

Internet Source

<1 %

12	agriculturalsocietynigeria.org Internet Source	<1 %
13	indianecologicalsociety.com Internet Source	<1 %
14	krishikosh.egranth.ac.in Internet Source	<1 %
15	www.iiste.org Internet Source	<1 %
16	"Phytochemical Drug Discovery for Central Nervous System Disorders", Wiley, 2023 Publication	<1 %
17	hal.science Internet Source	<1 %
18	agrosym.ues.rs.ba Internet Source	<1 %
19	banglajol.info Internet Source	<1 %
20	cgspace.cgiar.org Internet Source	<1 %
21	docslib.org Internet Source	<1 %
22	Fikre Limeneh Demis, Mohammed Beshir Hussien, Gebretensay Mengistu Fekadu. "Influence of Nitrogen and Phosphorus fertilizer Rates on Seed Yield, Yield Components and Quality of Onion (<i>Allium cepa</i> L.) at Kulumsa in Arsi Zone, South East Ethiopia", African Journal of Agricultural Research, 2019 Publication	<1 %
23	Sharmin Akter, Kazi Tamim Rahman, Asif Reza Anik, Md Ruhul Amin. "Growth and trend	<1 %

analysis of onion production in Bangladesh",
Annals of Bangladesh Agriculture, 2023

Publication

24	bari.portal.gov.bd Internet Source	<1 %
25	downloads.hindawi.com Internet Source	<1 %
26	pharmchoices.com Internet Source	<1 %
27	(8-31-14) http://67.225.160.147/profiles/profiles/onion-onion-products/z,,63,0,a/index.html Internet Source	<1 %
28	Submitted to QEC LUAWMS Student Paper	<1 %
29	T. AKTER, A. K. M. A. ISLAM, M. G. RASUL, S. KUNDU, KHALEQUZZAMAN, J. U. AHMED. "Evaluation of genetic diversity in short duration cotton (Gossypium hirsutum L.)", Journal of Cotton Research, 2019 Publication	<1 %
30	hspublishing.org Internet Source	<1 %
31	www.mdpi.com Internet Source	<1 %
32	M. Badawi, S. Seadh, M. El-Emery, A. Shalaby. "ONION SEED YIELD AND ITS QUALITY AS INFLUENCED BY STORAGE METHODS, MOTHER BULB SIZE AND HARVESTING TIME", Journal of Plant Production, 2010 Publication	<1 %
33	Submitted to University of Duhok Student Paper	<1 %

34 Asaduzzaman, Md., Md. Mainul Hasan, Md. Mahmudul Hasan, Md. Moniruzzaman, and Mohammad Humayun Kabir Howlander. "Effect of bulb size and plant spacing on seed production of onion (*Allium cepa* L.)", Bangladesh Journal of Agricultural Research, 2012.
Publication

35 Teshome Ashagrie, Derbew Belew, Sentayehu Alamerew, Yehenew Getachew. "Effects of Planting Time and Mother Bulb Size on Onion (*Allium cepa* L.) Seed Yield and Quality at Kobo Woreda, Northern Ethiopia", International Journal of Agricultural Research, 2014
Publication

36 innspub.net
Internet Source

37 isws.org.in
Internet Source

38 www.baripmis.org
Internet Source

39 www.f2ffoundation.org
Internet Source

40 T. K. Hazarika, Boris Aheibam. " Soil nutrient status, yield and quality of lemon (*Burm.*) cv. 'Assam lemon' as influenced by bio-fertilizers, organics and inorganic fertilizers ", Journal of Plant Nutrition, 2019
Publication

41 impactfactor.org
Internet Source

42	"Asian Digital Libraries. Looking Back 10 Years and Forging New Frontiers", Springer Science and Business Media LLC, 2007 Publication	<1 %
43	M. Ashrafuzzaman, M. N. Millat, M. Razi Ismail, S. M. Shahidullah. " Influence of paclobutrazol and bulb sizes on seed yield and yield attributing traits of onion (L) cv Taherpuri ", Archives of Agronomy and Soil Science, 2009 Publication	<1 %
44	barcapps.gov.bd Internet Source	<1 %
45	www.plantarchives.org Internet Source	<1 %
46	Submitted to University of Rajshahi Student Paper	<1 %
47	1library.net Internet Source	<1 %
48	Hossain, Md Shakhawat, MAR Sarkar, M Jahiruddin, AK Chaki, and ASM MR Khan. "Productivity and partial budget analysis in wheat-rice sequences as influenced by integrated plant nutrition system and legume crops inclusion", Bangladesh Journal of Agricultural Research, 2016. Publication	<1 %
49	OA Fakir, MK Alam, MJ Alam, M Jahiruddin, MR Islam. "Effects of different methods and time of boron application on the nutrient concentration and uptake by wheat (Triticumaestivum L.)", Bangladesh Journal of Agricultural Research, 2018 Publication	<1 %

50	S. K. Dingre, D. D. Pawar, K. D. Kale, M. M. Kadam. "Onion seed productivity, nutrient use, and quality response to drip fertigation in semi-arid India", Journal of Plant Nutrition, 2016 Publication	<1 %
51	hdl.handle.net Internet Source	<1 %
52	juniperpublishers.com Internet Source	<1 %
53	nib.portal.gov.bd Internet Source	<1 %
54	uilspace.unilorin.edu.ng Internet Source	<1 %
55	www.phytojournal.com Internet Source	<1 %
56	Carmine Amalfitano, Nadezhda A. Golubkina, Laura Del Vacchio, Giuseppe Russo et al. "Yield, Antioxidant Components, Oil Content, and Composition of Onion Seeds Are Influenced by Planting Time and Density", Plants, 2019 Publication	<1 %
57	Hayrettin Kıraç, Akife Dalda Şekerci, Ömer Faruk Coşkun, Osman Gülşen. "Morphological and molecular characterization of garlic (Allium sativum L.) genotypes sampled from Turkey", Genetic Resources and Crop Evolution, 2022 Publication	<1 %
58	Sunday E. Obalum, Ifeoma G. Edeh, Odirichi N. Imoh, Ogechi M. Njoku et al. "Agronomic evaluation of seedbed and mulching	<1 %

alternatives with plant spacing for dry-season
fluted pumpkin in coarse-textured tropical
soil", Food and Energy Security, 2017

Publication

59	arkgroup.arkgroup.co.in Internet Source	<1 %
60	bmcvetres.biomedcentral.com Internet Source	<1 %
61	docsdrive.com Internet Source	<1 %
62	journals.hu.edu.et Internet Source	<1 %
63	prh.hec.gov.pk Internet Source	<1 %
64	sfna.org.my Internet Source	<1 %
65	updatepublishing.com Internet Source	<1 %
66	Ghulam Khaliq, Sajid Ali. "Postharvest Physiological Disorders of Fruits", Apple Academic Press, 2025 Publication	<1 %
67	M R Ara, M M H Masud, K Akter, M R Islam. "Evaluation of different field treatments on seed and seedling quality of onion against purple blotch disease", Bangladesh Journal of Agriculture, 2023 Publication	<1 %
68	S Ahmed, MA Rahim, M Moniruzzaman, MA Khatun, FN Jahan, R Akter. "Effect of Bulb Sizes on the Seed Yield of Two Onion (Allium Cepa L.) Varieties", SAARC Journal of Agriculture, 2021	<1 %

69

doaj.org

Internet Source

<1 %

70

Changxing Fu, Jiayan Fan, Gaohua Fan, Heqi Wang, Congwen Wang, Wei Qiang, Dafu Yu, Yingxin Huang. "Soil and plant communities co-regulating plant biomass allocation patterns along a saline-alkali gradient, case study of *Allium ramosum* in Songnen Grassland, Northeast China", *Frontiers in Plant Science*, 2025

Publication

<1 %

71

Haile Ashenafi, Tesfaye Bizuayehu, Worku Walelign. "Seed yield of onion (*Allium cepa* L.) as affected by bulb size and intra-row spacing", *African Journal of Agricultural Research*, 2017

Publication

<1 %

72

eprints.utas.edu.au

Internet Source

<1 %

73

scialert.net

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On