Jana Publication & Research

ANAESTHESIA FOR INTERVENTIONAL RADIOLOGICAL PROCEDURES A REVIEW ARTICLE

VRC34

Document Details

Submission ID
trn:oid:::2945:338686598

Submission Date
Dec 22, 2025, 10:42 AM GMT+5:30

Download Date
Dec 22, 2025, 10:44 AM GMT+5:30

File Name
IJAR-55314.pdf

File Size
1.1 MB

20 Pages

5,248 Words

31,302 Characters

6% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- Bibliography
- Quoted Text

Match Groups

24 Not Cited or Quoted 6%
Matches with neither in-text citation nor quotation marks

0 Missing Quotations 0%
Matches that are still very similar to source material

0 Missing Citation 0%
Matches that have quotation marks, but no in-text citation

0 Cited and Quoted 0%
Matches with in-text citation present, but no quotation marks

Top Sources

3% Internet sources
4% Publications
4% Submitted works (Student Papers)

Match Groups

- **24** Not Cited or Quoted 6%
  Matches with neither in-text citation nor quotation marks
- **0** Missing Quotations 0%
  Matches that are still very similar to source material
- **0** Missing Citation 0%
  Matches that have quotation marks, but no in-text citation
- **0** Cited and Quoted 0%
  Matches with in-text citation present, but no quotation marks

Top Sources

3% Internet sources
4% Publications
4% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Student papers	Clarkson College on 2025-08-01 <1%
2	Student papers	hcpc on 2024-04-17 <1%
3	Publication	Kealan Westby, Daniel Westby, Kevin McKevitt, Brian M. Moloney. "Artificial Intell... <1%
4	Publication	Nathan David P. Concepcion, Bernard F. Laya, Edward Y. Lee. "Current updates in ... <1%
5	Student papers	University Of Tasmania on 2025-08-30 <1%
6	Internet	scholar.archive.org <1%
7	Publication	"CIRSE 2006, Rome, Italy, 9–13 September, Abstracts & Index", CardioVascular an... <1%
8	Student papers	Sinclair Community College on 2025-12-01 <1%
9	Internet	archive.org <1%
10	Internet	pmc.ncbi.nlm.nih.gov <1%

11 Publication

Didolkar, Manjiri M., Megan E. Anderson, Mary G. Hochman, Julia G. Rissmiller, Jef... <1%

12 Student papers

University of Bristol on 2019-04-03 <1%

13 Internet

ia801803.us.archive.org <1%

14 Internet

www.ahcmedia.com <1%

15 Publication

"CIRSE 2007, Athens, Greece, 8–12 September, Abstracts & Author Index", CardioV... <1%

16 Publication

"CIRSE 2015", CardioVascular and Interventional Radiology, 2015 <1%

17 Publication

David C. Madoff, Ron C. Gaba, Charles N. Weber, Timothy W. I. Clark, Wael E. Saad... <1%

18 Publication

Elizabeth von Ende, Sean Ryan, Matthew A. Crain, Mina S. Makary. "Artificial Intell... <1%

19 Publication

Image-Guided Cancer Therapy, 2013. <1%

20 Student papers

Universidad Tecnica De Ambato- Direccion de Investigacion y Desarrollo , DIDE o... <1%

ANAESTHESIA FOR INTERVENTIONAL RADIOLOGICAL PROCEDURES: A REVIEW ARTICLE

ABSTRACT: Interventional Radiology (IR) is a rapidly advancing subspecialty of medicine that utilizes real-time imaging techniques such as fluoroscopy, ultrasound, computed tomography and magnetic resonance imaging to perform minimally invasive diagnostic and therapeutic procedures. These procedures span a wide range of medical disciplines including vascular, neuro, hepatobiliary, musculoskeletal, genitourinary, and oncologic interventions. Anaesthesia plays a vital role in ensuring procedural success, patient safety and comfort. Anaesthetic management in IR is challenging due to diverse procedures, patient comorbidities, and the remote location of IR suites. Approaches must be carefully tailored, ranging from local anaesthesia with sedation to general anaesthesia, based on procedure complexity and patient factors. Monitored anaesthesia care is common for minor procedures, while general anaesthesia for more complex or high-risk interventions likeTrans-arterial chemoembolization, Radiofrequency ablation, and neuro-interventions. Special considerations must be needed for paediatric, geriatric, and critically ill patients, Effective collaboration between the anaesthesiology and radiology teams is essential to overcome challenges such as limited access, radiation exposure, and the constraints of working in a non-operating room environment. Ensuring effective communication and planning enhances both procedural success and patient safety.This review aims to provide an in-depth overview of anaesthetic considerations for interventional radiology procedures, including indications, anaesthetic techniques, patient-specific challenges, and future developments in this continually evolving field.

ANAESTHESIA MANAGEMENT IN INTERVENTIONAL RADIOLOGY: CURRENT PERSPECTIVES

Keywords: anaesthesia, Interventional radiology, Image-guided interventions, IR suite challenges, non-operating room anaesthesia, patient safety, Sedation,

Introduction:

The field of Interventional radiology (IR) was introduced in the 1960s by Dr. Charles Dotter, known as the "father of interventional radiology," who introduced percutaneous transluminal angioplasty. Since then, IR has rapidly evolved into a vital component of modern medicine.IR is a rapidly expanding field that utilizes image guidance to perform minimally invasive diagnostic and therapeutic procedures across multiple medical specialties. Since its inception in the 1960s, IR has become essential in managing vascular, oncological, hepatobiliary, neurological, and musculoskeletal conditions. The spectrum of procedures ranges from simple image-guided biopsies to complex interventions such as trans-arterial chemoembolization (TACE), radiofrequency ablation (RFA), endovascular aneurysm repair (EVAR), and cerebral thrombectomy. [1,2]The increasing complexity, length, and invasiveness of IR procedures necessitate the development and implementation of robust anaesthetic strategies tailored to individual patient needs. Anaesthesiologists play a critical role in ensuring patient safety, comfort, immobility, and optimal physiological conditions throughout these procedures. Moreover, many IR procedures are performed in remote locations, outside the conventionaloperating room (OR), often posing challenges in termsof equipment, monitoring, staffing, and emergency preparedness. [3,4] Traditionally, IR procedures were performed under local anaesthesia or moderate sedation. However, with the advent of more invasive and prolonged techniques, there is a growing demand for monitored anaesthesia care (MAC), regional anaesthesia, and general anaesthesia (GA), especially in paediatric, geriatric, and critically ill populations.[5] The choice of anaesthetic technique depends on several factors including the type and duration of the procedure, patient comorbidities, cooperation level, pain expected, and the need for absolute immobility.[6]Effective anaesthesia management in IR requires thorough pre-procedural evaluation, risk stratification, and multidisciplinary collaboration between the anaesthesia, radiology, and nursing teams. In addition, the anaesthesiologist must be proficient in managing sedation-related complications, airway emergencies, and haemodynamic instability, especially in patients with limited physiological reserve.[7]

Materials and Methods

The data for this review were compiled from a wide range of articles published between 1982 and 2025, sourced from multiple academic journals. These papers were carefully selected and reviewed to extract relevant information applicable to the focus of this study.

Key Principles:

- **Minimally Invasive Techniques:** IR procedures are done through small incisions,resulting in reduced pain, quicker recovery, and shorter hospital stays.[8]
- **Image Guidance:** Real-time imaging ensures high precision and safety throughout the procedure.[9]
- **Therapeutic and Diagnostic:** IR offers both diagnostic interventions (e.g., biopsies) and therapeutic interventions (e.g., tumour ablation, embolization).[10]
    - **Patient-Focused Anaesthesia Planning:**local, sedation, MAC, or GA is selected based on the procedure's complexity, duration, and patient-specific needs.[11]
    - **Pre-Procedure Risk Assessment:** A comprehensive evaluation should include review of comorbidities (cardiac, hepatic, renal), coagulation profile, allergies (especially to contrast media), and airway evaluation.[12]
    - **Monitoring and Safety in Remote Environments:** Standard ASA monitoring (ECG, NIBP, SpO_2, $EtCO_2$) must be ensured even in off-site IR settings, with readiness for emergencies despite spatial limitations.[13]
    - **Effective Airway Management:** prepared for difficult airway scenarios due to limited patient access during procedures.[14]
    - **Radiation Safety:** Minimize radiation exposure to staff and patients; use protective equipment (lead aprons, thyroid shields) and comply with ALARA principles (As Low AsReasonably Achievable).[15]
    - **Communication and Teamwork:** Maintain continuous communication with the radiology team for timing, contrast use, patient positioning, and any procedural complications.[16]
    - **Contrast Allergy Management:** Identify contrast sensitivities early, premedicate at-risk individuals, and have emergency drugs and airway equipment readily available.[17]

- **Post-Procedure Monitoring:** Provide proper recovery and post-anaesthesia care, monitoring for issues such as prolonged sedation, bleeding, or contrast-induced nephropathy.[18]
- **Use of Sedation Protocols:** Administer sedatives like propofol, fentanyl, or midazolam in titrated doses, with the ability to escalate to GA if required.[19]
- **Multidisciplinary Collaboration:** Optimal outcomes require collaboration between anaesthesiologists, interventional radiologists, nurses, and technologists.[20]
- **Role in Modern Medicine:** IR is now a first-line treatment option for many conditions and is integrated into multidisciplinary care pathways. It continues to expand with innovations in catheter technology, imaging resolution, and targeted therapies.[21]

Common Procedures:

Vascular

1. **Angiography and Angioplasty**
   Diagnostic imaging of blood vessels using iodinated contrast to identify stenosis, occlusion, or vascular anomalies.
   Commonly performed in cerebral, coronary, renal, and peripheral arteries.[22]
2. **Angiography:** catheter-based visualization of blood vessels using contrast dye.
   To assess vascular pathology and guide interventions.[23]
3. **Percutaneous transluminal angioplasty (PTA):** minimally invasive procedureusing a balloon catheter to dilate narrowed vessels, commonly for peripheral arterialdisease(PAD).[24]
4. **Balloon Angioplasty:**Treats vessel stenosis in PAD, renal artery stenosis, and coronary artery disease (CAD).[25]
5. **Stent Placement:** Stents are deployed in blood vessels to maintain patency post-angioplasty, commonly used in renal arteries, iliac arteries, and carotid arteries.
6. **Embolization:** Embolization is the intentional occlusion of blood vessels to treat aneurysms, arteriovenous malformations, gastrointestinal bleeding, or to reduce tumour vascularity (pre-operative embolization).

- Uterine Artery Embolization (UAE) for fibroids.
- Bronchial Artery Embolization (BAE) for haemoptysis.
- Trauma-related embolization for internal bleeding.
- Gastrointestinal Bleeding Control: Embolization of mesenteric arteries.
- Agents include coils, polyvinyl alcohol (PVA) particles, glue, or gelfoam.[26]

7. **Inferior Vena Cava (IVC) Filter Placement:** Used to prevent pulmonary embolism in patients with contraindications to anticoagulation therapy.[27]

Oncologic:

1. **Radiofrequency Ablation (RFA) / Microwave Ablation (MWA)**
   - Local thermal destruction of solid tumors (commonly liver, lung, kidney, bone).
   - Image-guided (usually CT or ultrasound), minimally invasive procedure using either high-frequency alternating current (RFA) or electromagnetic waves (MWA) to induce coagulative necrosis. [28,29]

2. Cryoablation: Induces cell death via repeated freeze-thaw cycles, causing intracellular ice crystal formation and vascular stasis.

   - Renal cell carcinoma (especially in high-risk surgical patients), bone metastases, and soft tissue tumors.[30,31]

3. **Trans-arterialChemoembolization:**A palliative procedure for hepatocellular carcinoma where chemotherapy drugs and embolic agents are injected directly into the tumour's blood supply.[32,33]
4. **Selective Internal Radiation Therapy:** Involves injection of radioactive microspheres into liver tumours via hepatic artery for localized radiation therapy.[34,35]

Hepatobiliary and pancreatic

1. **Percutaneous Transhepatic Biliary Drainage:** Relief of malignant or benign biliary obstruction causing obstructive jaundice (e.g., cholangiocarcinoma, pancreatic cancer).

Insertion of a catheter into dilated intrahepatic ducts under ultrasound and fluoroscopy guidance. [36,37]

2. **Biliary Stenting**

To maintain bile flow and relieve obstruction

Malignant biliary obstruction (e.g., pancreatic cancer, cholangiocarcinoma), benign strictures, or post-surgical leaks. [38,39]

3. **Trans-jugular Intrahepatic Portosystemic Shunt**

Refractory ascites, variceal bleeding, Budd-Chiari syndrome.

A stent is placed between the portal and hepatic veins via the internal jugular vein to reduce portal hypertension.[40,41]

4. **Cholecystostomy**: Temporary decompression of the gallbladder in critically ill or surgical high-risk patients with acute cholecystitis.

Percutaneous catheter placement into gallbladder under ultrasound or CT guidance.[42,43]

Genitourinary

1) **Percutaneous Nephrostomy:** Drainage of obstructed urinary system.
2) **Ureteral Stenting:** Relieves ureteric obstruction (e.g., stones, malignancy).
3) **Varicocele Embolization:** Minimally invasive treatment for varicocele related infertility.
4) **Renal biopsy:**Image-guided diagnostic procedure for glomerular and parenchymal disease.[44,45,46,47]

Neurointerventional radiology

1) **Cerebral Angiography:** for diagnosis of aneurysms, AVMs, and stroke.
2) **Intracranial Aneurysm Coiling:** Endovascular technique using platinum coils to prevent rupture.
3) **Mechanical Thrombectomy**: in acute ischemic stroke has become standard care for large vessel occlusions.
4) **Carotid Artery Stenting:** Used for significant carotid stenosis as an alternative to endarterectomy.

5) **Balloon-assisted coiling and flow diverters**: are employed for aneurysm treatment.[48,49,50,51,52]

Musculoskeletal

1) **Vertebroplasty and Kyphoplasty:** for osteoporotic vertebral fractures.
2) **Joint Aspiration and Injection:** for diagnosis or corticosteroid delivery.
3) **Bone and Soft Tissue Biopsies:** Image-guided core biopsies of bone or soft tissue lesions.[53,54,55]

Gastrointestinal

1) **Percutaneous Gastrostomy/Jejunostomy:**Feeding tube placement under fluoroscopic or CT guidance**.**
2) Percutaneous drainageof abscesses, hematomas, and pseudocysts using image guidance is standard practice.
3) It reduces the need for surgical drainage and supports antibiotic therapy.[56,57]

Pulmonary

1) **Pulmonary Angiography:** Diagnoses pulmonary embolism (PE).
2) **Pulmonary AVM Embolization**: Embolization of arteriovenous malformations to prevent paradoxical emboli.[58,59]

Central venous access:

1) Peripherally Inserted Central Catheter (PICC)
2) Tunneled Dialysis Catheters
3) Implantable Ports (Port-a-Cath): Used for long-term IV access in oncology, dialysis, or chronic diseases.[60,61]

Imaging Modalities Used in IR

1. **Ultrasound:** Provides real-time, radiation-free imaging. Commonly used for vascular access, biopsies, and regional anaesthesia. Its portability and safety make it ideal for bedside procedures.

2. **Computed Tomography:** Offers **high-resolution anatomic detail**, especially useful for accessing **deep organs** like the **lungs**, **liver**, and **adrenal glands**. Enables precise planning and targeting in complex procedures.
3. **Fluoroscopy:** Provides **dynamic, real-time X-ray imaging**. Widely used in **vascular**, **biliary**, and **gastrointestinal interventions** (e.g., angioplasty, stent placement, cholangiography).
4. **Magnetic Resonance Imaging:** Offers superior **soft tissue contrast** without ionizing radiation. Used selectively due to equipment limitations and patient constraints. Primarily applied in **neuro-interventions** and **musculoskeletal procedures**.[62]

Applications and Benefits of Interventional Radiology (IR)

- Lower risk of infection
- Less postoperative pain and discomfort
- Reduced overall healthcare costs
- High procedural accuracy through real-time imaging guidance (e.g., fluoroscopy, ultrasound, CT)
- Suitable for patients unfit for conventional surgery due to comorbid conditions
- Lower complication and morbidity rates compared to open surgery
- Cost-effective management of complex medical conditions
- Improved patient safety in high-risk groups
- Valuable in treating high-risk surgical patients
- Broad applications in:Oncology, Vascular surgery, Neurology, Nephrology, Gastroenterology
- Faster patient recovery and shorter hospital stays
- Enhanced patient comfort and cooperation due to minimally invasive approach

- Ideal for cases requiring **prolonged patient immobility** (e.g., embolization procedures)
- Essential in managing:
    - Paediatric or uncooperative patients
    - Lengthy or technically complex procedures
    - Airway risk or high aspiration risk
    - Patients with significant coexisting medical diseases
    - Interventions with high bleeding or embolism risk
    - Claustrophobic or highly anxious individuals
    - Patients needing controlled ventilation or deep sedation.[63,64,65]

Anaesthetic considerations:

Pre-Procedural Assessment: A detailed assessment includes ASA classification, airway evaluation, fasting status, comorbidities (renal, hepatic, cardiovascular), and risk of contrast-induced nephropathy. Attention must be given to anticoagulation, allergies (especially to iodinated contrast), and patient cooperation.[66,67]

Local Anaesthesia: Suitable for minor procedures such as superficial biopsies and vascular access.[68]

Moderate Sedation: Involves benzodiazepines and opioids, with monitoring of oxygenation and ventilation; appropriate for cooperative patients.[69]

Monitored Anaesthesia Care: Enables deeper sedation under anaesthesiologist supervision using agents like propofol or dexmedetomidine.[70,71]

General Anaesthesia: Indicated for complex, long, or painful procedures and for uncooperative or high-risk patients; includes airway control and advanced monitoring.[72]

Regional Anaesthesia: Ultrasound- or fluoroscopy-guided nerve blocks provide site-specific analgesia, reduce opioid use, and enhance recovery.[73,74]

Patient-Specific Considerations

- **Paediatric Patients:** Typically require GA due to poor cooperation; regional anaesthesia may reduce opioid use.[75,76]
- **Elderly Patients:** More sensitive to sedatives; prefer lighter sedation or regional techniques to avoid delirium and cardiovascular instability.[77]
- **Critically Ill Patients:** Require advanced haemodynamic monitoring and careful selection of anaesthetic technique to maintain organ perfusion.[78]

Monitoring and Safety: Standard ASA monitoring includes ECG, pulse oximetry, non-invasive blood pressure, and capnography. For high-risk patients, arterial lines, central venous pressure, and urine output monitoring are necessary. Equipment for airway management and resuscitation must be readily available.[79,80]

Challenges in IR Anaesthesia:

- Remote location from main operating theatres
- Radiation exposure risks
- Limited space and staff availability
- Sudden conversion from sedation to GA in emergencies
- Complex comorbidities in patients undergoing IR.

Communication and Team Dynamics

- Need for **close coordination** with radiologists and nurses.
- Often poor **pre-procedure planning** or incomplete patient information.
- **Communication barriers** due to noise or lead shields.

Patient Factors

- **High-risk** (ASA III/IV) with multiple comorbidities.
- **Haemodynamically unstable**, septic, or coagulopathic.
- **Non-fasting**, increasing aspiration risk.
- **Paediatric, geriatric, or pregnant**, requiring special considerations.

Procedure-Related Challenges

- **Lengthy and complex procedures**
- **Use of contrast media** risk of allergic reactions or nephropathy.
- **Painful or anxiety-provoking** (e.g., ablations, biopsies).
- **Risk of sudden complications**: bleeding, embolism, pneumothorax.[81,82]

Discussion and Future Directions:

1. Integration of artificial intelligence (AI) for image interpretation, procedural planning, and real-time guidance, improving accuracy and efficiency
2. Adoption of predictive analytics from AI to aid patient selection, outcome prediction, and complication prevention.[83,84]
3. Fusion imaging (e.g., CT/MRI with ultrasound) and intraoperative 3D/AR visualization for enhanced anatomical guidance
4. Advancements in intraoperative imaging such as cone-beam CT to allow dynamic assessment during procedures.[85]
5. Focus on radiation dose reduction through AI-optimized fluoroscopy and dose monitoring.[86]
6. Deployment of robotic-assisted IR systems to enhance precision, reduce radiation, and enable tele-robotic interventions.
7. Development of minimally invasive devices — smaller catheters, bioabsorbable materials, and drug-eluting technologies.[87]
8. Implementation of radio-genomics to tailor treatments using imaging and biomarker data.[88]
9. Real-time monitoring systems and automated documentation to enhance patient safety and infection control.[89]

10. Use of high-fidelity simulators, VR, and AR for structured training to address IR workforce gaps
11. Transition to outpatient/day-care IR with ambulatory procedures and enhanced recovery protocols.
12. Increased use of federated learning to share AI model training across institutions while preserving data privacy
13. Development of explainable AI (XAI) frameworks to improve clinician trust and meet regulatory standards
14. Stronger multidisciplinary collaboration among IR, anaesthesia, surgery, oncology, and critical care in tumour boards and complex cases
15. Expansion of therapeutic applications across oncology, vascular, neuro, GI, and pain management with novel targeted delivery systems.
16. Growth of multimodal data integration using foundation models to support autonomy in pre-, peri-, and post-procedural settings.
17. Establishment of standardized regulatory guidelines and frameworks for AI/robotics integration, including safety, liability, and ethics.
18. Movement towards partially or fully autonomous robotic navigation in endovascular procedures, leveraging reinforcement and imitation learning.[90]

Conclusion: IR stands at the forefront of minimally invasive medicine, offering high-precision diagnostic and therapeutic options across diverse specialties. As IR procedures become increasingly complex, prolonged, and patient-specific, anaesthesia assumes a critical role not only in ensuring procedural success but also in enhancing patient safety, comfort, and recovery. Anaesthesiologists are now indispensable members of the IR team, tasked with managing a wide spectrum of clinical scenarios from stable outpatients to critically ill individuals in non-operating room environments. This demands a tailored approach to anaesthetic planning, incorporating meticulous pre-procedural assessment, vigilant intra-procedural monitoring, and well-coordinated postoperative care.The evolution of technology such as advanced imaging, robotic assistance, and AI-driven procedural planning has further redefined the expectations from anaesthetic services in IR. These innovations call for adaptability, continual upskilling, and interdisciplinary collaboration. As IR continues to

expand its therapeutic reach into oncology, neurology, hepatobiliary, and musculoskeletal domains, the role of anaesthesia will also broaden to match the growing demands in precision, safety, and efficiency. Looking ahead, the integration of data-driven decision-making, personalized sedation protocols, and real-time analytics will be pivotal in enhancing outcomes. Thus, anaesthesia in IR is not merely a supportive element but a cornerstone of patient-centered, technology-enabled, and outcome-driven interventional care.

REFERENCES

1. Vari A, Gangi A. Anaesthesia practices for interventional radiology in Europe. Cardiovasc InterventRadiol. 2017;40(6):803–13.
2. Olsen JW, Barger RL Jr, Doshi SK. Moderate sedation: What radiologists need to know. AJR Am J Roentgenol. 2013;201(5):941–6.
3. Martin ML, Lennox PH. Sedation and analgesia in the interventional radiology department. J VascIntervRadiol. 2003;14(9 Pt 2):1119–26.
4. Royal College of Radiologists. Sedation and Pain Management in Interventional Radiology: Audit Tool. London: RCR; 2018.
5. Whalin MK, Halenda KM, Haussen DC, et al. Dexmedetomidine: A safe alternative to general anesthesia for endovascular stroke treatment. J Neurointerv Surg. 2014;6(3):270–5.
6. ASA Committee on Pain Medicine. Statement on Anesthetic Care During Interventional Pain Procedures for Adults. Schaumburg: American Society of Anesthesiologists; 2021.

7. Takahashi CE, DiNapoli V, Mlynash M, et al. Intraprocedural blood pressure and end-tidal CO_2 are associated with outcomes in acute stroke intervention. Neurocrit Care. 2014;21(3):448–56.

8. Kaufman JA, Lee MJ. Vascular and Interventional Radiology: The Requisites. 2nd ed. Philadelphia: Elsevier; 2013.

9. Baerlocher MO, Asch MR. Image-guided procedures: diagnostic and therapeutic. Can Assoc Radiol J. 2010;61(5):254–61.

10. Wallace MJ, Kuo MD, Glaiberman C, et al. Clinical practice: quality improvement guidelines for percutaneous needle biopsy. J VascIntervRadiol. 2010;21(7):969–75.

11. Hemming AE, Shere-Wolfe KD. Anesthesia for interventional radiology. Anesthesiol Clin. 2014;32(3):663–79.

12. Bhatia PK, Bhandari S, Tulsiani KL. Preanesthetic medical evaluation: what to look for. Indian J Anaesth. 2009;53(6):696–703.

13. American Society of Anesthesiologists Task Force. Standards for Basic Anesthetic Monitoring. Anesthesiology. 2020;132(1):1–3.

14. Apfelbaum JL, Hagberg CA, Caplan RA, et al. Practice guidelines for management of the difficult airway. Anesthesiology. 2013;118(2):251–70.

15. Dauer LT, Thornton RH, Miller DL, et al. Occupational radiation protection in interventional procedures. J VascIntervRadiol. 2010;21(11):1615–22.

16. Haskal ZJ, Kundu S, Kopecky KK, et al. Quality improvement guidelines for interventional radiology. J VascIntervRadiol. 2003;14(9 Suppl):S199–202.

17. Greenberger PA, Patterson R. Radiocontrast media hypersensitivity. J Allergy Clin Immunol. 2011;127(3):586–92.

18. Bettmann MA. Frequently asked questions: iodinated contrast agents. Radiographics. 2004;24 Suppl1:S3–10.

19. Coulter FL, Anderson HL. Anesthesia for interventional radiology. Anesthesiol Clin. 2002;20(1):163–79.

20. Koelemay MJ, Legemate DA, Elgersma OE. Multidisciplinary approach in interventional radiology. Cardiovasc InterventRadiol. 2009;32(Suppl 2):S223–6.

21. Baerlocher MO, Asch MR. The future of interventional radiology. Can Assoc Radiol J. 2012;63(1):6–8.

22. Kaufman JA, Lee MJ. Vascular and Interventional Radiology: The Requisites. 2nd ed. Philadelphia: Elsevier; 2014.

23. Suri R, Gupta AK. Diagnostic catheter angiography: techniques and applications. Indian J Radiol Imaging. 2008;18(1):16–25.

24. Norgren L, Hiatt WR, Dormandy JA, et al. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). J Vasc Surg. 2007;45(1 Suppl):S5–67.

25. Safian RD, Freed MI. The role of balloon angioplasty in peripheral vascular disease. Circulation. 1991;83(6 Suppl):I97–107.

26. Marin ML, Veith FJ, Panetta TF, et al. Stent graft repair of abdominal aortic aneurysms: long-term outcomes. J Vasc Surg. 1995;21(3):465–77.

27. Kaufman JA, Kinney TB, Streiff MB, et al. Guidelines for the use of retrievable and convertible vena cava filters. J VascIntervRadiol. 2006;17(3):449–59.

28. Livraghi T, Meloni F, Solbiati L, et al. Treatment of focal liver tumors with percutaneous radio-frequency ablation: complications encountered in a multicenter study. Radiology. 2003;226(2):441–51.

29. Shibata T, Niinobu T, Ogata N, et al. Microwave coagulation therapy for multiple hepatic metastases from colorectal carcinoma. Cancer. 2000;89(2):276–84.

30. Littrup PJ, Jallad B, Vorugu V, et al. Lethal isotherms of cryoablation in a phantom study. J VascIntervRadiol. 2009;20(10):1343–51.

31. Atwell TD, Callstrom MR, Farrell MA, et al. Percutaneous cryoablation of large renal masses. J Endourol. 2008;22(12):2461–5.

32. Llovet JM, Real MI, Montaña X, et al. Arterial embolisation or chemoembolisation vs symptomatic treatment in hepatocellular carcinoma. Lancet. 2002;359(9319):1734–9.

33. Vogl TJ, Naguib NN, Nour-Eldin NE, et al. Transarterial chemoembolization in hepatic tumors. Expert Rev Anticancer Ther. 2011;11(3):405–20.

34. Salem R, Lewandowski RJ, Mulcahy MF, et al. Radioembolization for hepatocellular carcinoma using Yttrium-90 microspheres. Gastroenterology. 2010;138(1):52–64.

35. Kennedy AS, Coldwell D, Sangro B, et al. Radioembolization for the treatment of liver tumors. Am J Clin Oncol. 2012;35(1):91–9.

36. Mueller PR, Ferrucci JT Jr, Harbin WP, et al. Percutaneous biliary drainage: technical and catheter-related problems. AJR Am J Roentgenol. 1982;138(1):17–23.

37. Uberoi R, Das N, Moss J, et al. Biliary drainage and stenting registry (BDSR). Cardiovasc InterventRadiol. 2012;35(1):127–38.

38. Baron TH, Harewood GC. Endoscopic stenting for pancreatic cancer. Cancer Control. 2004;11(1):32–7.

39. Moss AC, Morris E, Leyden J, et al. Metal stents in malignant biliary obstruction. Eur J Gastroenterol Hepatol. 2007;19(12):1119–24.

40. Boyer TD, Haskal ZJ. TIPS in the management of portal hypertension: update 2009. Hepatology. 2010;51(1):306.

41. Garcia-Tsao G, Sanyal AJ, Grace ND, et al. Management of gastroesophageal varices. Hepatology. 2007;46(3):922–38.

42. Melin AA, Brethauer SA, Kroh M, et al. Percutaneous cholecystostomy for acute cholecystitis. Am J Surg. 2014;208(6):1021–4.

43. Barak O, Dreznik Y, Cohen SM, et al. Percutaneous cholecystostomy in critically ill patients. Isr Med Assoc J. 2000;2(10):757–60.

44. Farrell TA, Hicks ME. Percutaneous nephrostomy. Tech VascIntervRadiol. 2001;4(4):183–94.

45. Wah TM, Weston MJ, Irving HC. Percutaneous nephrostomy insertion. Clin Radiol. 2004;59(3):255–61.

46. Nabi G, Asterlings S, Greene DR, et al. Percutaneous embolization of varicoceles. Urology. 2004;63(2):359–63.

47. Maya ID, Maddela P, Barker J, et al. Percutaneous renal biopsy: blind vs real-time ultrasound technique. Semin Dial. 2007;20(4):355–8.

48. Cloft HJ, Joseph GJ, Dion JE. Risk of cerebral angiography: a meta-analysis. Stroke. 1999;30(2):317–20.

49. Molyneux A, Kerr R, Stratton I, et al. ISAT of clipping vs coiling. Lancet. 2002;360(9342):1267–74.

50. Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy for ischemic stroke: a meta-analysis. Lancet. 2016;387(10029):1723–31.

51. Brott TG, Hobson RW 2nd, Howard G, et al. Stenting versus endarterectomy for treatment of carotid-artery stenosis. N Engl J Med. 2010;363(1):11–23. doi:10.1056/NEJMoa0912321

52. Fiorella D, Lylyk P, Szikora I, et al. Curative cerebrovascular reconstruction with the Pipeline Embolization Device. AJNR Am J Neuroradiol. 2009;30(6):1130–6. doi:10.3174/ajnr.A1531

53. Lieberman IH, Dudeney S, Reinhardt MK, et al. Initial outcome and efficacy of kyphoplasty in the treatment of painful osteoporotic vertebral compression fractures. Spine. 2001;26(14):1631–8. doi:10.1097/00007632-200107150-00019

54. Roberts WW, Sandhu GS, Lobash SA, et al. Image-guided corticosteroid injections in joint disease. Radiol Clin North Am. 1999;37(3):617–30. doi:10.1016/S0033-8389(05)70119-3

55. Welker JA, Henshaw RM, Jelinek J, et al. The percutaneous needle biopsy is safe and effective for diagnosing musculoskeletal lesions. Clin OrthopRelat Res. 2000;(375):104–13. doi:10.1097/00003086-200004000-00014

56. Pugash RA, Lim R, Temple ME, et al. Radiologic percutaneous gastrostomy: incidence of major and minor complications. Can Assoc Radiol J. 1997;48(5):339–42. PMID:9350108

57. van Sonnenberg E, Mueller PR, Ferrucci JT. Percutaneous drainage of 250 abdominal abscesses and fluid collections. Part I. AJR Am J Roentgenol. 1984;142(1):1–6. doi:10.2214/ajr.142.1.1

58. Remy-Jardin M, Remy J. Spiral CT angiography of the pulmonary circulation. Radiology. 1999;212(3):615–36. doi:10.1148/radiology.212.3.r99se05615

59. Letourneau-Guillon L, Faughnan ME, Soulez G, et al. Embolization of pulmonary arteriovenous malformations with Amplatzer vascular plugs: safety and midterm effectiveness. J VascIntervRadiol. 2010;21(5):649–56. doi:10.1016/j.jvir.2010.01.010

60. Vesely TM. Central venous catheter tip position: a continuing controversy. J VascIntervRadiol. 2003;14(5):527–34. doi:10.1097/01.RVI.0000062023.05893.D6

61. Funaki B. Central venous access: a primer for the diagnostic radiologist. AJR Am J Roentgenol. 2002;179(2):309–18. doi:10.2214/ajr.179.2.1790309

62. Wallace MJ, Kuo MD, Glaiberman C, Binkert CA, Orth RC, Soulez G. Quality improvement guidelines for percutaneous image-guided epidural steroid injections. J VascIntervRadiol. 2010;21(6):879–91. doi:10.1016/j.jvir.2010.02.009

63. Patel PJ, Patel NM, Patel R. Interventional radiology: physical principles and clinical applications. Cureus. 2020;12(8):e9698. doi:10.7759/cureus.9698

64. Covey AM, Brody LA, Getrajdman GI, Sofocleous CT, Brown KT. Safety and efficacy of interventional radiologic procedures in high-risk patients. Radiology. 2000;214(3):694–700. doi:10.1148/radiology.214.3.r00mr28694

65. Kessel DO, Robertson I. Interventional radiology: a guide to clinical practice. 2nd ed. London: Springer; 2010. p. 7–35.

66. Bhatia PK, Bhandari S, Tulsiani KL, et al. Preanesthetic medical evaluation: what to look for. Indian J Anaesth. 2009;53(6):696–703.

67. Bettmann MA. Frequently asked questions: iodinated contrast agents. Radiographics. 2004;24 Suppl1:S3–10.

68. Kaufman JA, Lee MJ. Vascular and Interventional Radiology: The Requisites. 2nd ed. Philadelphia: Elsevier; 2013.

69. American Society of Anesthesiologists. Practice guidelines for sedation and analgesia by non-anesthesiologists. Anesthesiology. 2002;96(4):1004–17.

70. Hemming AE, Shere-Wolfe KD. Anesthesia for interventional radiology. Anesthesiol Clin. 2014;32(3):663–79.

71. Jalowiecki P, Rudner R, Goniewicz M, et al. Use of dexmedetomidine in anaesthesia and intensive care. Anaesthesiol Intensive Ther. 2014;46(4):332–41.

72. Coulter FL, Anderson HL. Anesthesia for interventional radiology. Anesthesiol Clin. 2002;20(1):163–79.

73. Neal JM, Bernards CM, Butterworth JF 4th, et al. The ASRA evidence-based guidelines for regional anesthesia in patients on antithrombotic therapy. Reg Anesth Pain Med. 2010;35(1):64–101.

74. Mariano ER, Loland VJ, Ilfeld BM. Ultrasound-guided regional anesthesia and acute pain management. Curr OpinAnaesthesiol. 2010;23(5):632–8.

75. Mason KP. Sedation trends in children undergoing procedures outside the operating room. Curr OpinAnaesthesiol. 2007;20(4):365–72.

76. Ecoffey C. Pediatric regional anesthesia – update. Curr OpinAnaesthesiol. 2007;20(3):232–5.

77. Sieber FE. Postoperative delirium in the elderly surgical patient. Anesthesiol Clin. 2009;27(3):451–64.

78. Barash PG, Cullen BF, Stoelting RK, et al. Clinical Anesthesia. 8th ed. Philadelphia: Wolters Kluwer; 2017.

79. American Society of Anesthesiologists. Standards for basic anesthetic monitoring. Anesthesiology. 2020;132(1):1–3.

528 80. Ehrenfeld JM, Urman RD, Segal S. Anesthesiology. 7th ed. Philadelphia: Elsevier;
529 2017.

530 81. Kaye AD, Urman RD, Vadivelu N. Essentials of interventional radiology anesthesia.
531 Best Pract Res Clin Anaesthesiol. 2016;30(1):59–71. doi:10.1016/j.bpa.2016.01.004

532 82. Yang S, Hemming AE, Falco-Walter JJ, et al. Anesthesia for interventional radiology. J
533 Anesth. 2014;28(6):882–9. doi:10.1007/s00540-014-1848-6

534 83. 83. Glielmo P, Fusco S, Zantonelli G, et al. *Artificial intelligence in interventional*
535 *radiology: state of the art*. EurRadiol Exp. 2024;8:62.

536 84. 84. Mastriani C, et al. *AI and Interventional Radiology: narrative review*. Diagnostics.
537 2023;15(7):893.

538 85. Seah J, Boeken T, Sapoval M, Goh GS. *Artificial Intelligence, Augmented Reality, and*
539 *Virtual Reality Advances and Applications in Interventional Radiology*. Tech
540 VascIntervRadiol. 2023;26(3):100919.

541 86. Zimmermann S, Bang J, et al. *AI-enabled fluoroscopy reduces radiation exposure in*
542 *IR*. DiagnInterv Imaging. 2020;.

543 87. Lanza C, Carriero S, Buijs EF M, et al. *Robotics in Interventional Radiology: Review of*
544 *Current and Future Applications*. J Med Syst. 2023;

545 88. [PMC] *Applications and challenges of artificial intelligence in diagnostic and*
546 *interventional radiology*. 2022.

547 89. Seah J, et al. *Artificial intelligence in interventional radiology: Current concepts and*
548 *future trends*. DiagnInterv Imaging. 2025 Jan;106(1):5-10.

549 90. Use of Artificial Intelligence in Non-Oncologic Interventional Radiology: Current
550 State and Future Directions. PMC.

551

552