

# **Knowledge -Practice gap in children's nutrition among mothers of youngchildren: An Urban–Rural Comparative Study**

## Abstract

Child nutrition is a fundamental determinant of growth, development, and overall health during early childhood. Inadequate nutrition during the under-five period can lead to long-term physical, cognitive, and developmental impairments that may persist into adulthood. Mothers, as primary caregivers, play a pivotal role in ensuring appropriate feeding practices and nutritional care for young children. Therefore, assessing maternal knowledge and practices regarding child nutrition is essential, particularly across diverse socio-cultural settings such as urban and rural communities. The present study aimed to assess and compare the knowledge and practices related to child nutrition among mothers of under-five children residing in selected urban and rural areas of District Patiala, Punjab, and to examine the association between knowledge and practices with selected socio-demographic variables. A descriptive comparative research design was adopted for the study. The research was conducted in June 2025, and a total of 70 mothers of under-five children were selected using purposive sampling, comprising 35 mothers from urban areas and 35 from rural areas. Data were collected using a self-structured questionnaire that included socio-demographic characteristics and items assessing maternal knowledge and practices related to child nutrition. Statistical analysis was carried out using descriptive statistics such as mean and standard deviation, along with inferential statistics including the t-test. Findings revealed that the socio-demographic characteristics of urban and rural mothers were largely similar, with no statistically significant differences observed except for religion ( $p=0.001$ ). The majority of mothers demonstrated average knowledge regarding child nutrition in both urban (54.3%) and rural (45.7%) areas, and no mothers were found to have poor knowledge. In terms of practices, good nutritional practices were more commonly observed among rural mothers (56.5%) compared to urban mothers (43.5%). A statistically significant difference was found in practice scores between urban and rural mothers ( $p=0.001$ ), whereas no significant difference was noted in knowledge scores ( $p=0.933$ ). Educational status and occupation showed a significant association with knowledge and practice scores among rural mothers ( $p<0.05$ ). Although urban mothers exhibited slightly higher mean knowledge scores, rural mothers consistently demonstrated better nutritional practices. This highlights a gap between knowledge and practice, particularly in urban settings. The study underscores the need for targeted educational interventions and community-based nutrition programs to bridge this gap and promote optimal child nutrition practices, thereby improving health outcomes among under-five children.

Keywords: Child nutrition, maternal knowledge, feeding practices, urban mothers, rural mothers

45

46 **Introduction**

47 Early childhood nutrition plays a critical role in supporting optimal growth, cognitive  
48 development, immunity, and long-term health outcomes.<sup>1</sup> Malnutrition—both  
49 undernutrition and overnutrition—remains one of the leading causes of childhood  
50 morbidity and mortality worldwide.<sup>2</sup> The first 1,000 days of life are especially crucial,  
51 as nutritional deficiencies during this period can lead to stunting, impaired cognitive  
52 performance, and increased susceptibility to infections.<sup>3</sup> Parents play a central role in  
53 safeguarding child nutrition, as many common childhood health issues can be  
54 prevented through healthy feeding practices.<sup>4</sup> Nutrition is strongly influenced by  
55 socioeconomic status, maternal education, cultural norms, and access to healthcare  
56 services.<sup>5</sup> Children of educated mothers have been shown to have better dietary intake  
57 and lower risk of malnutrition.<sup>6</sup> Inadequate nutrition not only affects individual child  
58 potential but also undermines community health and long-term economic  
59 development.<sup>7</sup> According to the World Health Organization, malnutrition includes  
60 deficiencies, excesses, or imbalances in energy or nutrient intake.<sup>8</sup> Rural children are  
61 more prone to undernutrition due to poverty, limited healthcare access, and poor  
62 sanitation, while urban children increasingly face obesity risks due to processed food  
63 consumption and sedentary lifestyles.<sup>9 10</sup> Feeding practices among mothers vary  
64 widely between urban and rural communities, influenced by cultural beliefs,  
65 availability of nutrition information, and access to health services.<sup>11 12</sup> India continues  
66 to bear a substantial burden of malnutrition, despite economic growth and national  
67 nutrition programs. NFHS-5 reports high prevalence of stunting, wasting, and  
68 underweight among children under five.<sup>13</sup> Punjab, although relatively developed,  
69 exhibits concerning rates of childhood malnutrition, with 25.7% stunted, 21.6%  
70 underweight, and 17.2% wasted children.<sup>14</sup> Poor maternal knowledge regarding  
71 breastfeeding, complementary feeding, dietary diversity, and appropriate portion sizes  
72 further contributes to nutritional deficiencies among young children.<sup>15 16</sup> Significant  
73 differences between rural and urban maternal feeding practices highlight the need for  
74 comparative research.<sup>10</sup>

75 **Need of study**

76 According to WHO, nearly 45% of deaths among children under five years are  
77 attributable to malnutrition, particularly in low- and middle-income countries.<sup>16</sup>  
78 Malnutrition includes not only undernutrition but also micronutrient deficiencies and  
79 the rising burden of childhood overweight.<sup>17</sup> India contributes significantly to global  
80 malnutrition levels. NFHS-5 reported that 35.5% of children under five are stunted,  
81 19.3% wasted, and 32.1% underweight.<sup>13</sup> Punjab also shows concerning malnutrition  
82 levels, with 25.7% stunted, 21.6% underweight, and 17.2% wasted.<sup>14</sup> These figures  
83 highlight the continued challenges in child nutrition even in states considered  
84 relatively developed. Poor maternal knowledge regarding breastfeeding,  
85 complementary feeding, and dietary diversity is a major factor contributing to  
86 malnutrition.<sup>15</sup> Incorrect timing of complementary feeding, inadequate portion sizes,  
87 and low intake of micronutrient-rich foods are common concerns.<sup>18</sup> Maternal practices  
88 are also influenced by socioeconomic status, cultural norms, food availability, and  
89 access to health services. Rural mothers often follow traditional methods and may  
90 lack nutrition-related information, whereas urban mothers may rely more on packaged  
91 foods due to lifestyle influences.<sup>10</sup>

92 **Material and Methods**

93 The study adopted a comparative, non-experimental research design to assess and  
94 compare the knowledge and practices regarding child nutrition among mothers of  
95 under-five children residing in urban and rural areas. A total of 70 mothers were  
96 selected, with 35 from urban areas and 35 from rural areas, using a non-probability  
97 purposive sampling technique. A structured questionnaire was used to collect data on  
98 mothers' knowledge and practices related to nutrition. The key research variable was  
99 used to assess the knowledge and practice regarding child nutrition among mother of  
100 under five at urban and rural community area.

101 **Results**

102 The findings of the study indicated that the urban and rural groups were largely  
103 homogeneous in terms of baseline demographic characteristics. Religion was the only  
104 variable that showed a statistically significant difference between the two groups ( $p <$   
105 0.05), with Hindu mothers predominantly belonging to urban areas and Sikh mothers  
106 mainly representing rural areas. The data were normally distributed; therefore,  
107 parametric statistical tests were applied for further analysis. Assessment of maternal  
108 knowledge regarding child nutrition revealed no statistically significant difference  
109 between urban and rural mothers ( $p > 0.05$ ), with both groups demonstrating similar  
110 distributions of average and good knowledge levels, indicating that area of residence  
111 did not influence knowledge scores. In contrast, maternal practices related to child  
112 nutrition differed significantly between the two groups. Rural mothers exhibited  
113 significantly better practices compared to urban mothers, with a higher proportion  
114 demonstrating good practices (56.5% versus 43.5%), and the mean practice scores  
115 were significantly higher among rural participants ( $p < 0.001$ ), suggesting a strong  
116 influence of residence on maternal practices. Further analysis showed that educational  
117 status and occupation were the only demographic variables that had a significant  
118 influence on maternal knowledge scores ( $p < 0.05$ ), whereas other variables such as  
119 age, number of children, gender, religion, family type, and family income did not  
120 show a significant association. Between-group comparisons also revealed significant  
121 differences in knowledge scores based on educational status and occupation across  
122 urban and rural areas. With respect to maternal practices, the age of the youngest child  
123 was the only demographic variable that showed a significant association with practice  
124 scores ( $p < 0.05$ ), while other demographic factors did not demonstrate a significant  
125 effect. However, independent samples analysis revealed highly significant differences  
126 in practice scores across all demographic variables ( $p < 0.001$ ), with rural mothers  
127 consistently scoring higher than urban mothers. Overall, the results highlight that  
128 although maternal knowledge levels were comparable between urban and rural  
129 communities, maternal practices differed markedly, with rural mothers demonstrating  
130 more favorable practices related to child nutrition, underscoring the significant role of  
131 area of residence in influencing maternal practices.

132 **Table 1: Comparison of level of knowledge regarding child nutrition among**  
133 **mothers**

134

**N=70**

<b>Level of</b>	<b>N</b>	<b>Urban (n=35)</b>	<b>Rural (n=35)</b>	<b>p-value</b>
-----------------	----------	-------------------------	-------------------------	----------------

knowledge		F	%	F	%	
<b>Poor</b>	0	0	0.0	0	0.0	0.473 <sup>NS</sup>
<b>Average</b>	35	19	54.3	16	45.7	
<b>Good</b>	35	16	45.7	19	54.3	

135 **Fisher's Exact test** NS=Non-significant at  $p>0.05$ ; \*= Significant at  $p<0.05$

136

137 Table 1 reveals no statistically significant difference between the two groups ( $p > 0.05$ ),  
 138 with both urban and rural subjects having a similar distribution of average and good  
 139 knowledge levels.

140

141 **Table 2:Comparison of level of practices regarding child nutrition among**  
 142 **mothers**

143

N=70

Practices	N	Urban (n=35)		Rural (n=35)		p-value
		F	%	F	%	
<b>Poor</b>	0	0	0.0	0	0.0	0.005**
<b>Average</b>	8	8	100.0	0	0.0	
<b>Good</b>	62	27	43.5	35	56.5	

144 **Fisher's Exact test** NS=Non-significant at  $p>0.05$ ; \*\*=Significant at  $p<0.01$

145 Table 2 showed a statistically significant difference between urban and rural mothers  
 146 ( $p = 0.005$ ,  $p < 0.05$ ), with a higher proportion of rural mothers demonstrating good  
 147 practices (56.5%) compared to urban mothers (43.5%).

148 **Table 3:Comparison of mean and standard deviation of knowledge scores**  
 149 **regarding child nutrition between respondents of Urban and rural community**  
 150 **areas**

151

N=70

Residence	N	Mean $\pm$ SD	Mean difference	df	t	p-value
<b>Urban</b>	35	20.20 $\pm$ 4.16	0.086	68	8.36	0.933 <sup>NS</sup>
<b>Rural</b>	35	20.29 $\pm$ 4.28				

152 t=Independent Samples t-test NS=Non-significant at  $p>0.05$ ; \*= Significant at  $p<0.05$

153 Table 3 revealed that no statistically significant difference between the groups ( $p >$   
 154 0.05) suggesting that area residence had no effect on the knowledge scores of  
 155 mothers, and both groups had similar levels of knowledge.

156 **Table 4:Comparison of mean and standard deviation of practice scoresregarding**  
 157 **child nutritionbetween respondents of Urban and rural community areas**

158

**N=70**

Residence	N	Mean± SD	Mean difference	df	t	p-value
Urban	35	15.31± 1.87	3.257	68	8.36	0.001***
Rural	35	18.57± 1.35				

159 **t=Independent Samples t-testNS=Non-significant at p>0.05; \*\*\*=Significantat p<0.001**

160

161 Table 4showed statistically significant difference in mean practice scores between  
 162 urban and rural participants ( $p <0.001$ ) indicating that residence has a significant  
 163 impact on maternal practice scores. Rural participants demonstrated significantly  
 164 better practices compared to their urban counterparts

165

166 **Table5:Comparison of mean knowledge scores and demographic variables in**  
 167 **urban and rural areasrespondents**

168

**N=70**

Variable	Urban (n=35)					Rural (n=35)								
	N	Mean	SD	F	p-value	N	Mean	SD	F	p-value				
<b>Age(years)</b>														
18–25	7	21.57	2.5	0.63	0.540 <sup>NS</sup>	12	20.08	3.91	0.83	0.448 <sup>NS</sup>				
26–32	26	19.73	4.49			20	19.95	4.66						
33–39	2	21.5	4.95			3	23.33	2.51						
<b>t</b>	<b>0.12</b>	<b>p- value</b>	<b>0.90<sup>NS</sup></b>											
<b>Number of children</b>														
1	11	21	3.43	0.570	0.570 <sup>NS</sup>	7	21.43	4.57	1.208	0.298 <sup>NS</sup>				
2	16	19.38	4.41			24	20.08	4.23						
3	8	20.75	4.77			3	17.33	3.05						
≥ 4	0	–	–			1	26	–						
<b>t</b>	<b>-1.02</b>	<b>p- value</b>	<b>0.32<sup>NS</sup></b>											
<b>Gender of child</b>														
Male	17	19.65	4.78	0.453	0.453 <sup>NS</sup>	21	19.95	4.33	0.320	0.581 <sup>NS</sup>				
Female	18	20.72	3.54			14	20.79	4.33						
<b>t</b>	<b>-0.065</b>	<b>p- value</b>	<b>0.95<sup>NS</sup></b>											
<b>Age of youngest child(years)</b>														
0–1	0	–	–	0.33	0.72 <sup>NS</sup>	4	16	3.83	1.360	0.260 <sup>NS</sup>				
1–2	18	20.28	4.39			14	20.43	3.81						
2–3	12	19.75	4.35			12	20.67	4.67						
3–4	4	21.75	3.5			3	22.33	2.51						
4–5	1	18	–			2	22.5	6.36						

<b>t -0.49p - value 0.63<sup>NS</sup></b>														
<b>Educational status</b>														
No formal education	0	—	—	0.87	0.684 <sup>NS</sup>	2	14.5	2.12	5.07	0.028*				
Primary	6	18.33	2.06			5	16.6	4.39						
Matriculation	14	20.5	5.06			15	21.2	4.07						
Senior Sec.	11	20.45	4.29			8	20.63	4.06						
Graduation>	4	21.25	2.75			5	23	1.58						
<b>t -0.009p - value 0.99<sup>NS</sup></b>														
<b>Religion</b>														
Hindu	17	20.06	3.68	0.31	0.245 <sup>NS</sup>	1	24	—	0.77	0.387 <sup>NS</sup>				
Muslim	2	25	1.41			0	—	—						
Sikh	16	19.75	4.62			34	20.18	4.3						
<b>t -0.17p- value 0.86<sup>NS</sup></b>														
<b>Occupation</b>														
Homemaker	29	19.62	4.28	6.45	0.016 <sup>S</sup>	28	20.68	4.07	3.42	0.036*				
Govt. employee	0	—	—			2	12	1.41						
Private employee	6	23	2			3	21	3.6						
Self employed	0	—	—			2	22	0						
<b>t 0.20 p- value 0.84<sup>NS</sup></b>														
<b>Family income (in Rs.)</b>														
< 10,000	3	19.33	1.15	0.09	0.964 <sup>NS</sup>	4	15.75	4.42	1.87	0.162 <sup>NS</sup>				
10,000–20,000	13	20.54	4.52			16	20.69	4.19						
20,000–30,000	18	20.17	4.44			12	21.08	4.07						
> 30,000	1	19	—			3	21	3.6						
<b>t 0.044p- value 0.97<sup>NS</sup></b>														
<b>Type of family</b>														
Nuclear	21	21.29	4.12	3.96	0.058 <sup>NS</sup>	13	20.92	3.4	0.37	0.507 <sup>NS</sup>				
Joint	14	18.57	3.79			22	19.91	4.77						
<b>t 1.99p- value 0.055<sup>NS</sup></b>														

169 F =One-way ANNOVA    t=Independent Samples t-test    NS=Non-significant at p>0.05

170 \*=Significant at p<0.05

171 Table 5 revealed that a significant mean difference (p<0.05) was observed only for  
172 educational status and occupation, indicating their influence on knowledge scores. No  
173 significant mean differences (p>0.05) were found across other demographic variables,  
174 suggesting that variables such as age, number of children, gender, family type,  
175 religion, and income did not affect the knowledge scores. Between groups,  
176 independent samples t-tests revealed significant mean differences (p<0.05) in  
177 educational status and occupation, suggesting these factors influenced maternal  
178 knowledge scores across urban and rural communities.

179 **Table 6: Comparison of mean practice scores and demographic variables in**  
180 **urban and rural areasrespondents**

181

N=70

Variable	Urban(n=35)					Rural (n=35)				
	N	Mean	SD	F	p-value	N	Mean	SD	F	p-value
<b>Age(years)</b>										
18–25	7	15.43	1.98	2.09	0.15 <sup>NS</sup>	12	18.33	1.43	0.19	0.83 <sup>NS</sup>

26-32	26	15.42	1.83			20	18.75	1.33		
33-39	2	13.5	2.12			3	18.33	1.52		

**t =-7.32; p 0.001\*\*\***

**Number of children**

1	11	15.45	1.63	0.11	0.935 <sup>NS</sup>	7	19	1.15	2.11	0.233 <sup>NS</sup>
2	16	15.19	1.87			24	18.54	1.35		
3	8	15.38	2.38			3	17.33	1.52		
≥ 4	0	—	—			1	20	—		

**t =-6.63; p 0.001\*\*\***

**Gender of child**

Male	17	14.94	1.56	1.42	0.259 <sup>NS</sup>	21	18.62	1.28	0.067	0.804 <sup>NS</sup>
Female	18	15.67	2.11			14	18.5	1.5		

**t =-8.49; p 0.001\*\*\***

**Age of youngest child(years)**

0-1	0	—	—	0.13	0.981 <sup>NS</sup>	4	18	1.15	3.05	0.047*
1-2	18	15.22	2.07			14	18.29	1.26		
2-3	12	15.5	1.38			12	18.83	1.33		
3-4	4	15.25	2.87			3	20	0		
4-5	1	15	—			2	18	2.82		

**t = -8.21 p 0.001\*\*\***

**Educational status**

No formal education	0	—	—	3.14	0.039 <sup>NS</sup>	2	19.5	0.7	2.42	0.110 <sup>NS</sup>
Primary	6	14.33	1.5			5	17.2	1.09		
Matriculation	14	15.14	1.99			15	18.53	1.3		
Senior Sec.	11	15.82	1.88			8	19	1.3		

**t = -8.35 p 0.001\*\*\***

**Religion**

Hindu	17	15.24	2.01	0.78	0.667 <sup>NS</sup>	1	20	—	0.73	0.292 <sup>NS</sup>
Muslim	2	16.5	0.7			0	—	—		
Sikh	16	15.25	1.84			34	18.53	1.35		
Hindu	17	15.24	2.01			1	20	—		
Muslim	2	16.5	0.7			0	—	—		

**t = -8.35 p 0.001\*\*\***

**Occupation**

Homemaker	29	15.21	1.91	0.55	0.465 <sup>NS</sup>	28	18.57	1.31	0.32	0.912 <sup>NS</sup>
Govt. employee	0	—	—			2	18	1.41		
Private employee	6	15.83	1.72			3	18.67	2.3		
Self employed	0	—	—			2	19	1.41		

**t = -8.4 p < 0.001\*\*\***

**Family income (in Rs.)**

< 10,000	3	16.33	2.3	0.607	0.745 <sup>NS</sup>	4	17.75	1.5	1.64	0.20 <sup>NS</sup>
----------	---	-------	-----	-------	---------------------	---	-------	-----	------	--------------------

10,000–20,000	13	15	1.78			16	18.5	1.31						
20,000–30,000	18	15.39	1.97			12	19	1.2						
> 30,000	1	15	–			3	18.33	2.08						
<b>t = -8.43 p 0.001***</b>														
<b>Type of family</b>														
Nuclear	21	15.67	1.9			1.96	0.177 <sup>NS</sup>	13	18.69	1.25				
Joint	14	14.79	1.76					22	18.5	1.43				
<b>t = -8.59 p 0.001***</b>														

182 **F=One-way ANNOVA    t=Independent Samples t-test    NS=Non-significant at p>0.05**

183 **\*\*\*=Significant at p<0.001**

184

185 Table 6 revealed that a significant mean difference( $p<0.05$ ) was observed for the age  
186 of the youngest child, indicating its influence on practice scores. No significant mean  
187 differences ( $p>0.05$ ) were found across other demographic variables, suggesting that  
188 variables such as age, number of children, gender, educational status, religion,  
189 occupation, family income, and type of family did not affect the practice scores.  
190 Between groups, independent samples t-tests revealed highly significant mean  
191 differences across all variables ( $p<0.001$ ), with rural participants consistently scoring  
192 higher than urban participants implying area of residence significantly influenced the  
193 measured practicescores, with rural participants showing better outcomes than urban  
194 participants.

195 The findings revealed that religion was the only demographic variable showing  
196 significant differences, with Hindu participants mainly from urban areas and Sikh  
197 participants mainly from rural areas. Overall knowledge scores were similar across  
198 residences, but rural mothers demonstrated significantly better child care and nutrition  
199 practices. Within-group comparisons showed minimal variation across demographic  
200 subgroups, whereas between-group comparisons indicated that educational status and  
201 occupation influenced knowledge scores, with rural participants generally scoring  
202 higher. Additionally, the age of the youngest child significantly affected practice  
203 scores, while other demographic variables had no significant effect. Overall, rural  
204 participants consistently outperformed urban participants, highlighting residence as a  
205 key factor influencing maternal knowledge and child care practices.

## 206 **Discussion**

207 The findings revealed that both urban and rural mothers had almost equal levels of  
208 knowledge regarding child nutrition, and no statistically significant difference was  
209 observed between the two groups ( $p > 0.05$ ). A study in Maharashtra also found that  
210 health education initiatives improved complementary feeding practices more  
211 effectively among rural mothers, supporting the current finding that rural mothers  
212 showed better practices than urban mothers.<sup>19</sup>A comparative study in Mysuru,  
213 Karnataka, reported that urban mothers had somewhat better practices compared to  
214 rural mothers, though both groups demonstrated gaps in correct feeding  
215 practices.<sup>20</sup>Similarly, research in Belgaum, Karnataka, found that urban mothers  
216 initiated complementary feeding more appropriately than rural mothers.<sup>21</sup>Studies from  
217 Thrissur, Kerala, also indicated that urban mothers adhered better to weaning  
218 guidelines, whereas rural mothers often started earlier than recommended.<sup>22</sup>

219 Despite these variations, the role of maternal education has been consistently  
220 emphasized. In the present study, education was significantly associated with  
221 knowledge among rural mothers. This is supported by a multi-state study in India  
222 which showed that higher maternal health literacy significantly reduced the risk of  
223 underweight and stunting among children.<sup>23</sup> Regarding practices, demographic factors  
224 such as occupation, type of family, and age of the youngest child were significantly  
225 associated with better nutrition-related practices. Overall, while the present study  
226 highlighted that rural mothers demonstrated better practices despite similar  
227 knowledge levels, evidence from other regions shows mixed trends.

## 228 **Conclusion**

229 The study concluded that while both urban and rural mothers had comparable levels  
230 of knowledge regarding child nutrition, rural mothers demonstrated significantly  
231 better practices. Education, occupation, and family type influenced mothers'  
232 knowledge and practices. The results highlight a gap between awareness and  
233 implementation, particularly among urban mothers.

234 Effective community-based health education, practical demonstrations, and active  
235 involvement of family members are necessary to bridge this gap. Strengthening  
236 government nutrition programs and involving nurses and health workers in continuous  
237 support can promote optimal feeding practices. A holistic, family-centered approach  
238 is essential to improve child nutrition and support the growth and development of  
239 under-five children.

## 240 **Acknowledgement**

241 The authors thank their institution for granting permission to conduct the study. We  
242 extend our heartfelt thanks to the academic experts and faculty members for  
243 validating the tool and providing continuous guidance. We also extend appreciation to  
244 mothers of under five children who willingly participate and cooperate with us during  
245 the data collection.

## 246 **Reference**

- 247 1. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al.  
248 Maternal and child undernutrition and overweight in low-income and middle-income  
249 countries. *Lancet.* 2013 [Internet];382(9890):427–51. Available  
250 from: [https://doi.org/10.1016/S0140-6736\(13\)60937-X2](https://doi.org/10.1016/S0140-6736(13)60937-X2).
- 251 2. UNICEF, World Health Organization, World Bank Group. Levels and trends in  
252 child malnutrition: key findings of the 2021 edition. Geneva: World Health  
253 Organization; 2021 [Internet]. Available from:  
254 <https://www.who.int/publications/i/item/9789240025257>
- 255 3. Victora CG, de Onis M, Hallal PC, Blössner M, Shrimpton R. Worldwide timing of  
256 growth faltering: revisiting implications for interventions. *Pediatrics.* 2010  
257 [Internet];125(3):e473–80. Available from: <https://doi.org/10.1542/peds.2009-1519>
- 258 4. Bhutta ZA, Das JK, Rizvi A, Gaffey MF, Walker N, Horton S, et al. Evidence-based  
259 interventions for improvement of maternal and child nutrition: what can be done and  
260 at what cost? *Lancet.* 2013 [Internet];382(9890):452–77. Available  
261 from: [https://doi.org/10.1016/S0140-6736\(13\)60996-4](https://doi.org/10.1016/S0140-6736(13)60996-4)

262 5.Alderman H, Headey DD. How important is parental education for child nutrition?  
263 World Dev. 2017 [Internet];94:448–64. Available  
264 from:<https://doi.org/10.1016/j.worlddev.2017.02.016>

265 6.Hoddinott J, Alderman H, Behrman JR, Haddad L, Horton S. The economic  
266 rationale for investing in stunting reduction. Matern Child Nutr. 2013  
267 [Internet];9(Suppl 2):69–82. Available from:<https://doi.org/10.1111/mcn.12080>

268 7.Yirga AA, Mwambi HG, Ayele DG, Melesse SF. Factors affecting child  
269 malnutrition in Ethiopia. Afr Health Sci. 2019 [Internet];19(3):1897–1909. Available  
270 from:<https://doi.org/10.4314/ahs.v19i3.38>

271 8.World Health Organization, UNICEF. Infant and young child feeding counselling:  
272 an integrated course. Geneva: WHO; 2006 [Internet]. Available  
273 from:<https://apps.who.int/iris/handle/10665/43599>

274 9.Menon P, Headey D, Avula R, Nguyen PH. Understanding the geographical burden  
275 of stunting in India: a regression-decomposition analysis of district-level data from  
276 2015–16. Matern Child Nutr. 2018 [Internet];14(4):e12620. Available  
277 from:<https://doi.org/10.1111/mcn.12620>

278 10.Srivastava S, Kumar S. Urban–rural differentials in child nutrition in India: a  
279 cross-sectional study. Clin Epidemiol Glob Health. 2021 [Internet];12:100871.  
280 Available from:<https://doi.org/10.1016/j.cegh.2021.100871>

281 11.Victora CG, Bahl R, Barros AJD, França GVA, Horton S, Krusevec J, et al.  
282 Breastfeeding in the 21<sup>st</sup> century: epidemiology, mechanisms, and lifelong effect.  
283 Lancet. 2016 [Internet];387(10017):475–90. Available  
284 from:[https://doi.org/10.1016/S0140-6736\(15\)01024-7](https://doi.org/10.1016/S0140-6736(15)01024-7)

285 12.Ministry of Women and Child Development. Poshan Abhiyaan: National Nutrition  
286 Mission. Government of India; 2018 [Internet]. Available from:<https://icds-wcd.nic.in/nnm/>

288 13.Kavle JA, Landry M. Addressing barriers to optimal breastfeeding: an integrative  
289 review. Matern Child Nutr. 2018 [Internet];14(3):e12544. Available  
290 from:<https://doi.org/10.1111/mcn.12544>

291 14.International Institute for Population Sciences (IIPS), ICF. National Family Health  
292 Survey (NFHS-5), 2019–21: Punjab. Mumbai: IIPS; 2021 [Internet]. Available  
293 from:<http://rchiips.org/nfhs/>

294 15.Aguayo VM, Menon P. Stop stunting: improving child feeding, women’s nutrition  
295 and household sanitation in South Asia. Matern Child Nutr. 2016 [Internet];12(Suppl  
296 1):3–11. Available from:<https://doi.org/10.1111/mcn.12283>

297 16.World Health Organization. Malnutrition. Geneva: WHO; 2021 [Internet].  
298 Available from:<https://www.who.int/news-room/fact-sheets/detail/malnutrition>

299 17.International Institute for Population Sciences (IIPS), ICF. National Family Health  
300 Survey (NFHS-5), 2019–21: India. Mumbai: IIPS; 2021 [Internet]. Available  
301 from:<http://rchiips.org/nfhs/>

302 18.Gupta N, Goel K, Shah P, Misra A. Childhood obesity in developing countries:  
303 epidemiology, determinants, and prevention. Endocr Rev. 2012 [Internet];33(1):48–  
304 70. Available from:<https://doi.org/10.1210/er.2010-0028>

305 19.Pawar PA, Katkade VB. A study to assess the knowledge and practice of mothers  
306 regarding complementary feeding in rural and urban areas of Maharashtra.  
307 ShodhKosh J Arts Humanit Soc Sci. 2023;5(2):123–130. Available  
308 from:<https://www.granthaalayahpublication.org/Arts-Journal/ShodhKosh/article/view/6088>

310 20.Girish T, Venkatesh G. Comparison of feeding practices among children under two  
311 years of age in urban and rural areas of Mysuru, Karnataka. Int J Community Med  
312 Public Health. 2023;10(3):1220–1226. Available  
313 from:<https://www.ijcmph.com/index.php/ijcmph/article/view/11359>

314 21.Patil SS, Angadi MM. A cross-sectional study on complementary feeding practices  
315 among mothers of under-five children in Belgaum. J Evid Based Med Healthc.  
316 2015;2(70):1037–1043. Available  
317 from:<https://jebmh.com/abstract/658c3be291384525835df9ede19ec4cb>

318 22.Nair MK, Augustine LF. Complementary feeding practices in Thrissur District,  
319 Kerala. Natl J Community Med. 2012;3(2):222–227. Available  
320 from:<https://njcmindia.com/index.php/file/article/view/336>

321 23.V, Kashyap S, Seth V, Agarwal S. Maternal health literacy in India: association  
322 with child nutritional status. Matern Child Health J. 2016;20(5):1001–1011. Available  
323 from:<https://pubmed.ncbi.nlm.nih.gov/27306895/>

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

UNDER PEER REVIEW IN IJAR