

1 Unusual absence of musculocutaneous nerve along with median nerve formation in mid arm

2

3 Abstract:

4 The lateral root of the brachial plexus issues three branches namely musculocutaneous nerve,
5 lateral root and lateral pectoral nerve. The musculocutaneous nerve innervates all the muscles of
6 anterior compartment of the arm. The median nerve is formed by union of lateral root of lateral
7 cord and medial root of medial cord in axilla.

8 But we found in one male cadaver, the absence of musculocutaneous nerve. The median nerve
9 was formed by lateral root of lateral cord and medial root of medial cord as usual but in mid arm.
10 In this case, the coracobrachialis is supplied by branch from lateral cord and biceps brachii and
11 brachialis muscles were innervated by median nerve.

12 The comprehension of these variations of musculocutaneous and median nerves are essential for
13 clinicians for detecting weakness or paralysis of flexor muscles of arm due to injury to median
14 nerve along with loss of flexion and supination of forearm, loss of sensation in the skin of lateral
15 portion of the forearm and motor loss along the median nerve distribution

16 Key words: Brachial plexus, median nerve, musculocutaneous nerve, biceps brachii,
17 coracobrachialis

18

19

20

21

22

23

24

25

26

27

28 Introduction:

29 Brachial plexus is a complex network of nerves formed from anterior primary rami of spinal
30 nerves C5-T1. The spinal nerves C5 and C6 join to form superior trunk, C7 continues as middle
31 trunk and inferior trunk is formed by union of spinal nerves C8-T1. Each of the above mentioned
32 trunks bifurcate into anterior and posterior divisions. Divisions join to form cords. Three
33 cords namely lateral, medial and posterior cord are formed by union of divisions (Dutta, 2017).

34 The lateral cord gives of three branches: lateral pectoral nerve, musculocutaneous nerve and
35 lateral root of median nerve. The musculocutaneous nerve enters the arm by piercing the
36 coracobrachialis muscle after which it courses between the biceps brachii and brachialis brachii
37 muscles supplying the muscles of anterior compartment of arm. At the elbow, it continues as the
38 lateral cutaneous nerve of forearm (Standring, 2005).

39 The median nerve is formed by union of lateral root from the lateral cord and medial root from
40 the medial cord. These two roots join together anterior to third part of axillary artery in axilla.
41 The lateral root contains C8 and T1 fibers while medial root is comprised of C5, C6, and C7
42 fibers (Dutta, 2017). The median nerve continues in the forearm anterior to the brachial artery
43 and in the upper arm, it crosses the brachial artery and comes to lie medial to brachial artery. The
44 median nerve does not give any branch in the arm (Standring, 2005).

45 The anatomical variations of brachial plexus and its branches are well documented. However, the
46 musculocutaneous nerve may pass deep to coracobrachialis or cling to the median nerve then
47 course deep to biceps or a branch to musculocutaneous nerve arises from the median nerve
48 (Standring, 2005). The musculocutaneous nerve may occur in a pair or may be absent. But
49 incidence of absence of musculocutaneous nerve is not exactly known. The musculocutaneous
50 nerve is observed to arise from the lateral cord in 90.5%, from posterior cord and medial cord in
51 4% and 2% of population respectively and separately but simultaneously from medial and lateral
52 cords in 1.4% of individuals (Arora and Dhingra), 2005.

53 Here we describe a case of absence of musculocutaneous nerve along with variant formation of
54 median nerve in the middle of arm in place of axilla.

55 Case presentation:

56 The axilla and anterior compartment of arm of right side of formalin fixed male cadaver of 60
57 year old was dissected using standard dissection technique described in Cunningham's manual
58 for teaching to first year MBBS student of 2025-2026 batch. The musculocutaneous nerve was
59 found to be absent and median nerve was observed to be formed by union of lateral root of
60 lateral cord and median root of medial cord in the middle of arm anterior to brachial artery
61 (Figure 1). It was also found that the median nerve crossed the brachial artery from anterior to
62 medial side of brachial artery. The median nerve issued separate muscular branches to biceps
63 brachii and brachialis muscles of the anterior compartment of the arm but nerve to
64 coracobrachialis arose from lateral root (Figure 1). Rest of the course of median was normal as
65 described in standard text books of Anatomy. On contralateral limb, the configuration of
66 musculocutaneous and median nerve was normal.

67 Discussion:

68 The variation of brachial plexus formation along with its branches are well cited in literature. We
69 observed absence of musculocutaneous nerve along with formation of median nerve by lateral
70 root of lateral cord and medial root of medial cord anterior to brachial artery. But the formation
71 of median nerve was in the middle of arm. The combination of aforementioned anomalies is rare.

72 The variable course of the musculocutaneous and median nerve is described by various authors
73 (Le Minor, 1992; Fregnani et al., 2008; Pacholczak et al, 2011; Parchand and Patil, 2013).

74 Some authors classified the course of musculocutaneous nerve and median nerve into five types
75 (Le Minor, 1992). The type five of this classification coincided with our case report but with
76 difference that median nerve was formed in mid arm in our case.

77 In addition, the absence of musculocutaneous nerve is also reported by various authors (Fregnani
78 et al., 2008; Pacholczak et al, 2011; Parchand and Patil, 2013; Malukar O, Rathva), 2011; Rao
79 and Chaudhary, 2001; Nasr, 2012). In all aforementioned cases, the place of musculocutaneous
80 was taken by the median nerve and the median nerve supplied all muscles of arm normally

81 innervated by musculocutaneous nerve. Prasad Rao and Chaudhary found the absence of
82 musculocutaneous nerve in 8% of upper limb dissected. In our case also, the musculocutaneous
83 nerve was absent and biceps brachii and brachialis muscles supplied by
84 musculocutaneous muscles are innervated by median nerve. But, in addition to above
85 observations, in our case the median nerve was formed in the mid arm which normally is
86 observed to form in axilla.

87 Conclusion:

88 Variations of musculocutaneous and median nerve are of utmost use for surgeons and clinicians
89 during shoulder and upper arm reconstruction as these nerves are likely to be injured during
90 these procedures. Moreover, detailed information of these variations are important to understand
91 weakness or paralysis of flexor muscles of arm due to injury to median nerve. When place of
92 musculocutaneous nerve is taken by median nerve, injury to median nerve may cause loss of
93 flexion and adduction of arm, flexion and supination of forearm, loss of sensation in the skin of
94 lateral portion of the forearm and motor loss along the median nerve distribution.

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110 References:

- 111 1. Dutta A.K. (2017) Essentials of Human Anatomy, part-3, 5th edition, Lenin Saranee,
112 Kolkata, p41-43
- 113 2. Shetty Arpitha. (2024) Unilateral Absence of the Musculocutaneous Nerve - A Case
114 Report. J Ayurveda Integr Med Sci;9:332-335. <http://dx.doi.org/10.21760/jaims.9.9.54>
- 115 3. Standring S. Grey's anatomy: The anatomical basis of clinical practice, 39th ed.
116 Edinburgh: Churchill Livingstone.2005:848-57.
- 117 4. Arora L, Dhingra R. (2005) Absence of Musculocutaneous nerve and accessory head of
118 biceps brachii: a case report. IJPS.;38(2):144-6.
- 119 5. Le Minor JM. (1992) A rare variant of the median and Musculocutaneous nerves in man.
120 Arch Anat Histo Embryo.; 73:33-42.
- 121 6. Fregnani JH, Macéa MI, Pereira CS, Barros MD and Macéa JR. (2008) Absence of the
122 Musculocutaneous nerve: a rare anatomical variation with possible clinical surgical
123 implications. São Paulo Med J; 126(5):288-90
- 124 7. Pacholczak R, Klimek-Piotrowska W and Walocha J. (2011) Absence of the
125 Musculocutaneous nerve associated with a supernumerary head of biceps brachii: a case
126 report. Surg Radiol Anat; 33(6):551-4.
- 127 8. Parchand MP, Patil ST. (2013) Absence of Musculocutaneous nerve with variations in
128 course and distribution of the median nerve. Anat. Sci. Int;88(1):58-60.
- 129 9. Malukar O, Rathva A. (2011) A study of 100 cases of Brachial plexus. Nat J Comm
130 Med;2(1):66 70.
- 131 10. Rao PV, Chaudhary SC. (2001) Absence of Musculocutaneous Classification of
132 communication between the Musculocutaneous and median nerves. two case reports.
133 Clin. Anat.14(1):31-5.
- 134 11. Nasr AY. (2012) Morphology and clinical significance of the distribution of the median
135 nerve within the arm of human cadavers. Neurosciences;17(4): 336-44.

136

137

138


139 Figure legend:

140 Figure 1: displaying absence of musculocutaneous nerve with formation median nerve in midarm

141 LR: lateral root, MR: medial root, NC: nerve to coracobrachialis muscle, NB: nerve to brachialis

142 muscle, BB: biceps brachii muscle, NBB: nerve to biceps brachii muscle, MN: median nerve,

143 BA: brachial artery



UNDE.