



ISSN NO. 2320-5407

Journal homepage: <http://www.journalijar.com>

INTERNATIONAL JOURNAL
OF ADVANCED RESEARCH

RESEARCH ARTICLE

Occupancy of gastropod shells by the hermit Crab *Clibanarius erythropus* (Crustacea, Anomura) inhabiting the Mediterranean Sea coast of Libya

Fatma El-Zahraa A. Abd El-Aziz, Khaleid F. Abd El-Wakeil, Ahmad H. Obuid-Allah, and Faraj O. Ramadan

Department of Zoology, Faculty of Science, Assiut University, Egypt. E-mail: F_abdelhameed @yahoo.com

Manuscript Info

Manuscript History:

Received: 12 October 2015

Final Accepted: 25 November 2015

Published Online: December 2015

Key words:

Hermit crab, *Clibanarius erythropus*, Shell occupation, Mediterranean Sea coast, Libya.

*Corresponding Author

Fatma El-Zahraa A. Abd El-Aziz

Abstract

The present study deals with the use of gastropod shells by the hermit crab *Clibanarius erythropus* (Decapoda: Anomura) inhabiting the Mediterranean Sea coast of Libya. Specimens of the crab were collected on 21 June 2014 from Misrata city, western Mediterranean Sea coast of Libya. A total of 260 individuals were collected which included 63 males (24.23%), 74 non-ovigerous females (28.46%) and 123 ovigerous females (47.31%). The study revealed that the crab occupied 11 gastropod shell species. The most common occupied shell species was *Pisania striata* (36%) followed by *Cerithium lividulum* (26%), *Cerithium vulgatum* (15%) and *Phorcus turbinatus* (8%). Shell utilization patterns appear to be determined by respective sex, size of hermit crabs and shell species. Male *C. erythropus* occupies the relatively large and heavy gastropod shell species (*Pisania striata* and *Phorcus turbinatus*). Small crabs occupy relatively larger shell species than large crabs. Positive relationships were recorded between the variables of the shells occupied and the hermit crabs size variables. The present study referred that the shell length and shell aperture width were the most important choice factors for the studied hermit crab.

Copy Right, IJAR, 2015,. All rights reserved

INTRODUCTION

Empty shells of marine mollusks are often inhabited by the anomuran decapod crustaceans. Hermit crabs generally utilize gastropod shells because their abdominal exoskeleton lacks calcification. So gastropod shells are an indispensable resource for hermit crabs (Hasegawa et al., 2009). Shells of gastropods provide protection against predators (Vance, 1972 and Hazlett, 1990) and physical stress (Reese, 1969). Habitat partitioning has been demonstrated since closely related species show variable use of gastropod shells depending on the shell size, shape and availability (Teoh and Chong, 2014), while a more heterogeneous habitat provides more niches and ways to exploit the available resources (Bazzaz, 1975). The life cycle of hermit crabs depends mostly on the processes that make suitable gastropod shells (Hazlett, 1981). Studies on the utilization of gastropod shells by hermit crabs have been conducted by many authors like Fotheringham, (1976); Koutsoubas et al., (1993); Manjón-Cabeza and García Raso (1999); Martinelli and Mantelatto (1999); Botelho and Costa (2000); Mantelatto and Garcia (2000) Ateş et al. (2007) and Rodrigues and Rodrigo, (2009). No such study was done on the shell utilization by hermit crabs of the Mediterranean Sea coast of Libya.

The intertidal hermit crab *Clibanarius erythropus* (Latreille, 1818) is a common species of Mediterranean shores and is also present along the Atlantic coasts from Brittany to the Azores (Zariquiey Alvarez, 1968 and Ingle, 1993). The present study was carried out to evaluate the utilization patterns of the gastropod shells by males, females and ovigerous females of *C. erythropus*; the most common species of hermit crabs found at Misrata city which lies on the western Mediterranean Sea coast of Libya.

MATERIALS AND METHODS

Samples were collected by hand from the shallow intertidal water from the public beach region of Misrata city (North $32^{\circ} 25'$, East $15^{\circ} 02'$) Fig.1 on June 21th, 2014. This area is characterized by rocky shores and by a mixture of sandy and rocky soils. A total of two hundred and sixty specimens were collected and preserved in 10% seawater formalin, then transported to the laboratory. In the laboratory; crabs were removed carefully from their shells. Shells were labeled for identification. Crab specimens were classified into males, non ovigerous females and ovigerous females. They were measured for shield length (SL: measured from the tip of the rostrum to the midpoint of the posterior margin of the shield, providing an indication of animal size), carapace length (CaL: measured from the tip of the rostrum to the midpoint of the posterior margin of the carapace).

Size of the gastropod shells occupied by these hermit crabs were measured, including shell aperture length (SAL), aperture width (SAW) and shell length (ShL), by using calipers (precision 0.01mm). Also shell weight (ShW) was estimated after removing the crab. SPSS software package (version 21) (SYSTAT statistical program) and MS Excel were used for statistical analysis of the data.

RESULTS

Sorting of the collected 260 individuals of *Clibanarius erythropus*; the study revealed the presence of 63 males constituting (24.15%), 74 non-ovigerous females, constituting (28.46%) and 123 ovigerous females constituting (47.33%). Sizes of *Clibanarius erythropus* (based on cephalothoracic shield length) ranged from 2.5 mm to 7mm (fig. 5). They occupied shells of 11 gastropod species (Fig 2-3) (Table1). The most commonly occupied shell species was *Pisania striata* which included 86 individuals constituting (33%) followed by *Cerithium lividulum* 75 (28%), *Cerithium vulgatum* 40 (15%), *Phorcus turbinatus* 22(8%), *Columbella rustica* 9 (3.5%), *Conus mediterraneus* 7(2.7%), *Gibbula divaricata* 7 (2.7%), *Gibbula varia* 6(2.3%), *Muricopsis cristata* 5(1.9%), *Gyroscala lamellose* 2(0.8%) and *Fasciolaria lignaria* 1(0.4%) (Table1) (Fig 3)). Males (33%), non-ovigerous females (45%) and ovigerous females (75%) occupied the shells of *Pisania striata* at higher percentages (Fig 4)..



Fig (1): Showing the location of Misrata city on the Mediterranean Sea coast of Libya and the sampling site at Misrata ($32^{\circ} 25'N$, $15^{\circ} 02'E$).



Figure (2): Shells used by *Clibanarius erythropus*. (a) *Pisania striata*. (b) *Muricopsis cristata*. (c) *Columbella rustica*. (d) *Cerithium vulgtatum*. (e) *Cerithium lividulum*. (f) *Gibbula varia*. (g) *Gyroscala lamellose*. (h) *Gibbula divaricata*. (i) *Fasciolaria lignaria*. (j) *Phorcus articulates* and (k) *Conus mediterraneus*.




Figure (3): Number of specimens of each gastropod shell species occupied by *Clibanarius erythropus* at Misrata; western Mediterranean Sea coast.




Fig. (4): Numbers of males, non ovigerous females, ovigerous females, in the different occupied shell species.





Fig. (5): Numbers of the (2.5 to 7 mm) sized crabs in the different occupied shell species.

Table. (1): Frequency and percentage of the different eleven species of gastropod shells occupied by Clibanarius erythropus.

No.	Name of shell species	Frequency	% of the total occupied shells
1	Cerithium vulgatum	40	15.4
2	Pisania striata	86	33.1
3	Cerithium lividulum	75	28.8
4	Phorcus turbinatus	22	8.5
5	Gibbula divaricata	7	2.7
6	Fasciolaria lignaria	1	0.4
7	Conus mediterraneus	7	2.7

8	Gibbula varia	6	2.3
9	Gyroscaia lamellosa	2	.8
10	Columbella rustica	9	3.5
11	Muricopsis cristata	5	1.9
Total		260	100

Table 2: Correlation coefficients for association between Clibanarius erythropus crab sizes variables and occupied shell size variables (r= correlation coefficient, Sig.=significant, **= statistically significant at 0.01, N=260).

variables	Shield length		Carapace length	
	r	Sig.	r	Sig.
Shell length	0.335	**	0.352	**
Aperture length	0.496	**	0.529	**
Aperture width	0.768	**	0.761	**
Shell weight	0.744	**	0.716	**

Table 3: Regression equations for the relations between Clibanarius erythropus crab sizes variables and occupied shell size variables (SL = Shield length, CaL = Carapace length, ShL = Shell length, SAL = shell aperture length, SAW= SAL = shell aperture width, ShW = Shell weight, N=260).

Dependent Variable	Selected variable	P value	R	Regression equations
Shield Length	Constant	< 0.001	0.808	SL=2.333 + 0.107 ShL +0.533SAW+0.279ShW
	Shell Length	0.019		
	Aperture Width	< 0.001		
	Shell weight	< 0.001		
Carapace Length	Constant	< 0.001	0.801	CaL=2.294+0.187ShL+ 0.141SAL+ 0.660SAW
	Shell Length	< 0.001		
	Aperture Length	.002		
	Aperture Width	< 0.001		

DISCUSSION

The present study revealed that Clibanarius erythropus inhabiting the Mediterranean Sea coast of Libya at Misrata city occupied 11 gastropod shell species. The most commonly occupied shells included: Pisania striata, Cerithium vulgatum, Cerithium lividulum and Phorcus turbinatus. Similar previous studies showed that hermit crabs and closely tied to a high proportion of the total gastropod shells present eg. Lewinsohn (1982),

Reay and Haig, (1990) Gherardi and McLaughlin (1994), Hogarth et al. (1998), Botelho and Costa (2000), Barnes (2003), Terossi et al. (2006), Michelle (2010), Khaleid et al. (2010) and Tahir et al. (2013).

The present study reported that there was a close relationship between the crab size and the size of the occupied shell. Similar relationship was recorded for other *Clibanarius* species from different areas. *C. erythropus* from São Miguel (Azores) (Botelho and Costa, 2000) and *C. virescens* from South Africa (Reddy and Biseswar, 1993). The small hermit crabs occupy larger shells to allow their future growth (Dominciano and Mantelatto, 2004). This might give the chance for smaller crab to choose from a wide range of gastropod shells between adequate shell size and larger shell for future growth.

In the present study *C. erythropus* size variables were positively correlated with all of the measured shell variables. This indicated the effect of the measured variables of shell size on crab size, this means that shells affect growth of the crab. Many authors reported the effects of shell on crab growth (Markham, 1968; Fotheringham, 1976b; Bertness, 1981; Turra and Leite, 2003; 2004; Khaleid et al., 2010). It is also known that hermit crab growth is slowed by occupation of a small shell (Hazlett, 1981; Lancaster, 1988; Botelho and Costa, 2000).

Multiple regressions for the present data show that shell length and shell aperture width were the selected shell features characterized the association between crab size (shield length) and the occupied shells.

The degree of importance for several morphological shell characteristics was always debatable (Floeter et al., 2000). Previous studies have shown that the most important variables in shell occupation by hermit crabs are shell aperture width, shell weight and internal volume (Botelho and Costa, 2000; Floeter et al., 2000; Garcia and Mantelatto, 2000; Mantelatto and Meireles, 2004; Dominciano and Mantelatto, 2004; Terossi, et al. 2006 Khaleid et al., 2010).

The above-mentioned results mean that hermit crabs are searching for adequate shell, which has suitable shell aperture and shell weight. Since the shell aperture is the main entrance for the crab, it should provide enough space to accommodate crab size and can close well to confer protection against desiccation. While the shell should be heavy enough to increase the crab resistance against the water current (Garcia and Mantelatto, 2001) and predators (Floeter et al., 2000), its weight should be suitable to minimize the energetic cost of carriage (Osorno et al., 1998). Therefore, it can be concluded that the shell weight and shell aperture are the most important choice factors for hermit crab.

Shell use in nature was also demonstrated to be dependent on hermit crab species. Also, shell use is dependent on a combination of site-specific features such as wave energy, predators, habitat complexity, shell availability, and shell use history, which may or may not differ among areas (Dominciano et al., 2009).

As a conclusion, gastropod shells preferred by hermit crabs depend on the region and site features. It was found out that exotic gastropod shells are mostly preferred along the Libyan Sea coast. Monthly or seasonal studies are required to be done along this coast determined whether there is a variation in utilized of this shells by juvenile or adult hermit crabs and whether exotic gastropod shells are occupied by hermit crabs through their lives only for certain stages or their life.

Acknowledgments

Our deep sincere thanks to D. Sherif Ramadan and Dr. Khaled Ahmad; (Institute of Oceanography, Alexandria, Egypt) and Dr. Fabio crocetta; (Laboratory of Cellular and Developmental Biology, Stazione Zoologica Anton Dohrn, Napoli, Italy) for cooperation in identification of gastropod shell species.

References

- Abd El-Wakeil, K.F., Ahmed, E. S., Obuid-Allah , A.H. and El-Shimy,N.A. (2010):** Gastropod shell occupation by hermit crab *Clibanarius signatus* (Crustacea: Decapoda) on the red sea coast, Egypt. Assiut Univ. J. of Zoology , **39(1), P-P. 1-10..**
- Ateş A.S., K atağan T., and Kocataş , A. (2007):** Gastropod Shell Species occupied by Hermit Crabs (Anomura: Decapoda) along the Turkish Coasts of the Aegean Sea. Turkish Journal of Zoology, 31: 13-18
- Barnes D.K.A. (2003):** Local, regional and global patterns of resource use in ecology: hermit crabs and gastropod shells as an example. Mar. Ecol. Prog. Ser. 246: 211-223.
- Bazzaz F.A. (1975):** Plant species diversity in old-field successional ecosystems in southern Illinois. Ecology 56:485–488.
- Bertness M.D. (1981):** The influence of shell type on the growth rates and clutch sizes of hermit crabs. Crustaceana, 40: 197-205.
- Botelho AZ, Costa AC. (2000):** Shell occupancy of the intertidal hermit crab *Clibanarius erythropus* (Decapoda, Diogenidae) on São Miguel (Azores). Hydrobiologia, 440: 111-117

- Dominciano L.C.C. and Mantelatto F.L.M. (2004)** The influence of shell species and size on the shell selection pattern of *Paguristes tortugae* (Decapoda, Diogenidae) from Anchieta Island (Ubatuba, Brazil) *Iheringia, Sér. Zool., Porto Alegre*, 94(4):425-428.
- Dominciano LCC, Sant'anna BS, and Turra A. (2009):** Are the preference and selection patterns of hermit crabs for gastropod shells species- or site-specific? *Journal of Experimental Marine Biology and Ecology*, 378(1-2): 15-21.
- Floeter S.R., Nalesso R.C., Rodrigues, M.P. and Turra A. (2000):** Patterns of shell utilization and selection in two sympatric hermit crabs (Anomura: Diogenidae) in south-eastern Brazil. *J. Mar. Biol. Ass. U. K.*, 80: 1053-1059.
- Fotheringham N. (1976):** Population consequences of shell utilization by hermit crabs. *Ecology*, 57: 570-578
- Fotheringham, N. (1976b):** Population consequences of shell utilization by hermit crabs. *Ecology*, 57:570–578.
- Garcia R.B. and Mantelatto F.L. (2000):** Variability of shell occupation by intertidal and infralittoral *Calcinus tibicen* (Anomura: Diogenidae) populations. *Nauplius* 8:99-105.
- Garcia R.B. and Mantelatto F.L. (2001):** Shell selection by the tropical hermit crab *Calcinus tibicen* (Herbst, 1791) (Anomura, Diogenidae) from Southern Brazil. *Journal of Experimental Marine Biology and Ecology*, 265(1):1-14.
- Gherardi F and McLaughlin PA (1994) Shallow water hermit crabs (Crustacea: Decapoda: Anomura: Paguridea) from Mauritius and Rodrigues Island, with description of a new species of *Calcinus*. *Raffles Bull. Zool.* 42:613-656.
- Hasegawa H., Wada S, Aoki M, & Wada K. (2009):** Regional variation in shell utilization patterns of the hermit crab *Pagurus filholi*. *Plankton and Benthos Research*, 4(2): 72-76.
- Hazlett B.A. (1981):** The behavioral ecology of hermit crab. *Annual Review of Ecology and Systematics*, 12: 1-22.
- Hazlett B.A. (1981):** The behavioural ecology of hermit crabs. *Annual Review of Ecology and Systematics*, 12: 1-22.
- Hazlett B.A. (1990):** Shell Exchange in Hawaiian hermit crabs. *Pacific Science*, 44(4): 401-406.
- Hogarth PJ, Gherardi F and McLaughlin PA (1998):** Hermit crabs (Crustacea Decapoda Anomura) of the Maldives with the description of a new species of *Catapagurus* A. Milne Edwards 1880. *Trop. Zool.* 11:149-175.
- Ingle, R. (1993):** Hermit crabs of the northeastern Atlantic Ocean and Mediterranean Sea. *Journal of Chapman & Hall*. Pp.495.
- Koutsoubas D, Labadariou N, and Koukouras A. (1993):** Gastropod shells inhabited by Anomura Decapoda in the North Aegean Sea. *Bios*, 1: 247-249
- Lancaster, I. (1988):** *Pagurus bernhardus* (L.) An introduction to the natural history of the hermit crabs. *Field Studies* 7:189-238.
- Lewinsohn CH (1982):** Researches on the coast of Somalia. The shore and the dune of Sar Uanle. 33. Diogenidae, Paguridae and Coenobitidae (Crustacea, Decapoda, Paguridea). *Monit. Zool. Ital* 2:35-68
- Manjón-Cabeza ME, Garcia Raso JE. (1999):** Shell utilization by the hermit crabs *Diogenes pugilator* (Roux, 1829), *Paguristes eremita* (Linnaeus, 1767) and *Pagurus forbesii* Bell, 1845 (Crustacea: Decapoda: Anomura), in a shallow-water community from Southern Spain. *Bulletin of Marine Science*, 65: 391-405
- Mantelatto F.L. and Garcia R.B. (2000):** Shell utilization patterns of the hermit crab *Calcinus tibicen* (Diogenidae) from southern Brazil. *Journal of Crustacean Biology* 20:460–467.
- Mantelatto FLM, Garcia RB. (2000):** Shell utilization pattern of the hermit crab *Calcinus tibicen* (Diogenidae) from southern Brazil. *Journal of Crustacean Biology*, 20: 460-467
- Markham J. (1968):** Notes on the growth patterns and shell utilization of the hermit crab *Pagurus bernhardus*. *Ophelia*, 5:189-205.
- Martinelli JM, & Mantelatto F.L.M. (1999):** Shell utilization by hermit crab *Loxopagurus loxochelis* (Diogenidae) in the Ubatuba Bay, Brazil. In: *Crustaceans and the Biodiversity Crisis* (Schram FR, Vaupel Klein JC, eds.) (719-731, Brill, Leiden, Netherlands
- Michelle w. (2010):** Resource utilization and reproduction of the hermit crab *Clibanarius virescens* (Crustacea: Decapoda: Anomura) in South Africa. Nelson Mandela Metropolitan University.
- Osorno JJ, Fernandez-Casillas L and Rodriguez-Juarez C (1998):** Are hermit crabs looking for light and large shells? Evidence from natural and field induced shell exchanges. *Journal of Experimental Marine Biology and Ecology*, 222:163-173.
- Reay PJ and Haig J (1990):** Coastal hermit crabs (Decapoda: Anomura) from Kenya, with a review and key to East African species. *Bull. Mar. Sci.* 46:578-589.
- Reese E.S. (1969):** Behavioral adaptations of intertidal hermit crabs. *American Zoologist*, 9: 343-355.
- Rodrigues P, Rodrigo RK. (2009):** Shell occupancy by the Hermit Crab *Clibanarius erythropus* (Crustacea) on the south coast of São Miguel, Açores. *Açoreana, Sup.* 6: 211-216

Tahir Ö, Bilal Ö, Tuncer K, Banu B. (2013): Gastropod shell species utilized by hermit crabs (Decapoda: Anomura) along the Turkish coast of the Levantine Sea, Fisheries Faculty, Ege University, 35100 Bornova-Izmir, Turkey, *Arthropods*, 2013, 2(2): 45-52.

Teoh HW, and Chong VC (2014): Shell use and partitioning of two sympatric species of hermit crabs on a tropical mudflat. *J Sea Res* 86:13–22.

Terossi M, Espósito DLA, Meireles AL, Biagi R and Mantelatto FL (2006): Pattern of shell occupation by the hermit crab *Pagurus exilis* (Anomura, Paguridae) on the northern coast of São Paulo State, Brazil. *Journal of Natural History*,; 40(1–2): 77-87.

Terossi M., Espósito D.L.A., Meireles A.L., Biagi R. and Mantelatto F.L. (2006): Pattern of shell occupation by the hermit crab *Pagurus exilis* (Anomura, Paguridae) on the northern coast of São Paulo State, Brazil. *Journal of Natural History*,; 40(1–2): 77-87.

Turra A. and Leite F.P. (2003): The molding hypothesis: linking shell use with hermit crab growth, morphology and shell-species selection. *Mar. Ecol. Prog Ser*, 265:155-163.

Turra A. and Leite F.P. (2004): Shell-size selection by intertidal sympatric hermit crabs. *Marine Biology*, 145: 251-257.

Vance,R.R. (1972): The Role of shell adequacy in behavioral interactions involving Hermit Crabs. *Ecology*, 53(6): 1075-1083.

Zariquiey A.L. and Arez. R. (1968): Crustáceos decápodes ibéricos. *litvest. Pcsq. Tomo 32. Barcehoua.*