

 ISSN NO. 2320-5407	Journal Homepage: - www.journalijar.com INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR) Article DOI: 10.21474/IJAR01 DOI URL: http://dx.doi.org/10.21474/IJAR01	 INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR) ISSN 2320-5407 Journal homepage: http://www.journalijar.com Journal DOI: 10.21474/IJAR01
---	---	--

RESEARCH ARTICLE

**International Conference on Recent Advances in Biotechnology, Biomolecules and Pharmacy RABBP – 2020
(Organized in Virtual Mode due to COVID-19 Pandemic) during 17th to 19th December 2020 at KL
University, Vijayawada, Andhra Pradesh, India.**

DOCKING STUDIES OF BCL-2 WITH *HEMIDESMUS WALLICHII* COMPOUNDS FOR ANTI-CANCER STUDIES

**Kavya Sri Arumilli, Roshni Mocharla, Likhita Ventrapragada, Thelma Aradhya Mudda, Chaturya
Ankamma Pamarthi and Himavarshini Kadiyala**

Department of Biotechnology, Koneru Lakshmaiah Education Foundation, Guntur, Andhra Pradesh, India.

Manuscript Info

Key words:-

Hemidesmus wallichii,
Diabetic, Anticancer, Docking
Alpha Glucosidase

Anti-
studies,

Abstract

Medicinal plants have wide spread properties due to the presence of phytochemicals and are the alternative medicines available for those who cannot be helped by conventional medicine. In this work we have selected bioactive compounds from *Hemidesmus wallichii* medicinal plant extracts. Gas chromatography and Mass Spectrum studies were studied to identify the compounds present in the ethanolic extracts based on the retention time and area. The identified compounds were used for anti-cancer activity by *in silico* method with BCL-2 which plays prominent role in causing cancer. Out of twenty selected compounds, docking results showed Methyl-1-Cyclo Hexane carboxylate and 1,2-diacetoxy-5-ido hexane as best docked to the BCL-2.

Copy Right, IJAR, 2020., All rights reserved.

Introduction:-

Cancer is a global health problem with high morbidity and mortality and poses both economic and psychological challenges. Cancer cure and prevention therefore remain a high priority for the scientific community across the world [1]. Insight gained into the etiology of cancer through various epidemiological studies encompassing various parameters such as geographical location, ethnicity, sex, age, and trans-migratory populations have collectively revealed that lifestyle is one of the major influencing factors [2]. Other factors include environmental aspects such as automobile exhaust pollutants, solar UV radiation, occupational exposure to carcinogens and mutagens, bacterial/viral infection, and genetic susceptibility. Lifestyle factors are usually classified as modifiable risk factors and include diet intake, smoking, caspase-3 activity, alcohol consumption, and physical activity and body mass. In general, physical activity instead of inactivity, abstinence from smoking and alcohol consumption, low body mass, and diets low in fat/calories are usually recommended for overall good health and have a positive influence on reducing the risk of cancer, especially breast and colorectal cancers [3]. Because all these factors can be modified, they also provide us with leverage to use them as interventional/preventive measures. Accordingly, the American Cancer Society has suggested guidelines on nutrition and physical activity for the prevention of cancer and early detection/screening for cancers of certain sites [4]. From the epidemiological data indicating that dietary habits influence cancer risk, considerable scientific interest has been generated in developing various preventive measures.

Corresponding Author:- Kavya Sri Arumilli

Address:- Department of Biotechnology, Koneru Lakshmaiah Education Foundation, Guntur, Andhra Pradesh, India.

based on diet, especially those involving fruits and vegetables. Fruits and vegetables, along with probiotics belonging to plant kingdom, represent a vast source of phytochemicals of varied chemical structure; many of them have already been studied extensively for their potential anticancer or chemopreventive efficacy[5]. As such, interventions based on fruits and vegetables are not only “more natural” in lowering cancer risk without posing “any side effects” but also in maintaining good general health based on the fact that they are major sources of vitamins, minerals, and fiber.

Methodology:-

GC-MS method for identification of compounds

GC-MS analysis was carried out on a GC CLARUS 550 PerkinElmer system comprising a gas chromatograph interfaced to a mass spectrometer (GC-MS) instrument employing the following conditions: Column Elite-1 fused silica capillary column (30×0.25 mm ID×1EM df, composed of 100% Dimethyl poly siloxane), operating in electron impact mode at 70 eV[6]; helium (99.999%) was used as carrier gas at a constant flow of 1ml/min and an injection volume of 0.5 EI was employed with split ratio of 10:1 injector temperature 250°C; ion-source temperature 280°C[7]. The oven temperature was fixed from 110°C (isothermal for 2 min), with an increase of 10°C/min, to 200°C, then 5°C/min to 280°C, ending with a 9 min isothermal at 280°C. Mass spectra were taken at 70 eV; a scan interval of 0.5 s and fragments from 40 to 550 Da.

BCL-2 Active site Identification

The structure of BCL-2 (PDB: 1GJH) was retrieved from PDB database and unnecessary chains, hetero atoms were removed using SPDBV software, hydrogens were added to the protein and used for active site identification. Active site of BCL-2 of Homo sapiens was identified using CASTp server[8]. A new program, CASTp, for automatically locating and measuring protein pockets and cavities, is based on precise computational geometry methods, including alpha shape and discrete flow theory. CASTp identifies and measures pockets and pocket mouth openings, as well as cavities[9]. The program specifies the atoms lining pockets, pocket openings, and buried cavities; the volume and area of pockets and cavities; and the area and circumference of mouth openings.

Docking method


Docking was carried out using GOLD (Genetic Optimization of Ligand Docking) software which is based on genetic algorithm which allows as partial flexibility of protein and full flexibility of ligand[10]. The compounds identified in GC-MS are docked to the active site of the BCL-2 of Homo sapiens[11]. The interaction of the compounds with the active site residues are thoroughly studied using molecular mechanics calculations. The parameters used for GA were population size (100), selection pressure (1.1), number of operations (10,000), number of island (1) and niche size (2). Operator parameters for crossover, mutation and migration were set to 100, 100 and 10 respectively[12]. Default cutoff values of 3.0 Å (dH-X) for hydrogen bonds and 6.0 Å for vanderwaals were employed. During docking, the default algorithm speed was selected and the ligand binding site in the targets were defined within a 10 Å radius with the centroid as CE atom of active residues. The number of poses for each inhibitor was set 100, and early termination was allowed if the top three bound conformations of a ligand were within 1.5 Å RMSD[13]. After docking, the individual binding poses of compounds were observed and their interactions with the protein were studied. The best and most energetically favorable conformation of ligands was selected.

Gold Score fitness function:

Gold Score performs a force field based scoring function and is made up of four components: 1. Protein-ligand hydrogen bond energy (external H-bond); 2. Protein-ligand vander Waals energy (external vdw); 3. Ligand internal vander Waals energy (internal vdw); 4. Ligand intramolecular hydrogen bond energy (internal- H- bond)[14]. The external vdw score is multiplied by a factor of 1.375 when the total fitness score is computed. This is an empirical correction to encourage protein-ligand hydrophobic contact. The fitness function has been optimized for the prediction of ligand binding positions. $\text{GoldScore} = S(\text{hb_ext}) + S(\text{vdw_ext}) + S(\text{hb_int}) + S(\text{vdw_int})$, Where $S(\text{hb_ext})$ is the protein-ligand hydrogen bond score, $S(\text{vdw_ext})$ is the protein-ligand van der Waals score, $S(\text{hb_int})$ is the score from intramolecular hydrogen bond in the ligand and $S(\text{vdw_int})$ is the score from intramolecular strain in the ligand[15].

Results And Discussion:-

From the PDB databank, the PDB files were collected and the final stable structure of the BCL-2 of Homo sapiens obtained is shown in Figure 1. The ligands present in the crystal structure were removed along with hetero atoms for docking studies.



Structure of BCL-2 retrieved from Protein data bank with seven helices

Active site Identification

After the final model was built, the possible binding sites of BCL-2 was searched based on the structural comparison of template and the model build and also with CASTP server and was shown in Figure 2. Infact from the final refined model of BCL-2 domain using SPDBV program, it was found that secondary structures are highly conserved and the residues shown below




Fig 2:- Amino acids in the active site region (red colour) of the BCL-2 protein.

Docking of inhibitors with the active site

Docking of the compounds with BCL-2 was performed using GOLD 3.0.1, which is based on genetic algorithm[16]. This program generates an ensemble of different rigid body orientations (poses) for each compound conformer within the binding pocket and then passes each molecule against a negative image of the binding site. Poses clashing with this 'bump map' are eliminated. Poses surviving the bump test are then scored and ranked with a Gaussian shape function. We defined the binding pocket using the ligand-free protein structure and a box enclosing the binding site. This box was defined by extending the size of a cocrystallized ligand by 4Å. This dimension was considered here appropriate to allow, for instance, compounds larger than the cocrystallized ones to fit into the binding site. One unique pose for each of the best-scored compounds was saved for the subsequent steps. The compounds used for docking was converted in 3D with SILVER. To this set, the substrate corresponding to the

protein was added. Docking of best inhibitor with the active site of protein showed the activity of the molecule on protein function[17].




Fig 3:- Methyl-1-Cyclo Hexane carboxylate docked to BCL-2 active site.

In docking studies, Methyl-1-Cyclo Hexane carboxylate showed a docking energy of 22.56K.cal/mol with BCL-2. In docking, Methyl-1-Cyclo Hexane carboxylate (O21) docked to LEU414 with a bond length of 1.733Å[18].




Fig 4:- 1,2-diacetoxy-5-idohexanedocked to BCL-2 active site.

In docking studies, 1,2-diacetoxy-5-idohexane showed a docking energy of 20.24K.cal/mol with BCL-2. In docking, 1,2-diacetoxy-5-idohexane (H33) docked to LEU414 with a bond length of 2.559Å[19].

Conclusion:-

From the studies, we can conclude that GC-MS analysis identified twenty phytocompounds from *Hemidesmus wallichii* root ethanolic extract. The identified phytocompounds were checked for their anti-cancer

activity using insilico method. BCL-2 protein was retrieved from the database and its active site was identified using CASTp server[20]. The twenty phytocompounds were docked to the BCL-2 for their anti-cancer activity. Among the phytocompounds docked, Methyl-1-Cyclo Hexane carboxylate showed a docking energy of 22.56K.cal/mol and 1,2-diacetoxy-5-idohexane showed 20.24K.cal/mol with BCL-2. From these docking studies we can conclude that among the phytocompounds identified, these two compounds have good BCL-2 inhibitory activity.

References:-

1. Vineis P, Wild CP. Global cancer patterns: causes and prevention. *The Lancet*. 2014 Feb 8;383(9916):549-57.
2. Kaur M, Agarwal C, Agarwal R. Anticancer and cancer chemopreventive potential of grape seed extract and other grape-based products. *The Journal of nutrition*. 2009 Sep 1;139(9):1806S-12S.
3. Hayes SC, Spence RR, Galvão DA, Newton RU. Australian Association for Exercise and Sport Science position stand: optimising cancer outcomes through exercise. *Journal of science and medicine in sport*. 2009 Jul 1;12(4):428-34.
4. Smith RA, Cokkinides V, Eyre HJ. American Cancer Society guidelines for the early detection of cancer, 2003. *CA: a cancer journal for clinicians*. 2003 Jan;53(1):27-43.
5. Nerurkar P, Ray RB. Bitter melon: antagonist to cancer. *Pharmaceutical research*. 2010 Jun 1;27(6):1049-53.
6. Selvamangai G, Bhaskar A. GC-MS analysis of phytocomponents in the methanolic extract of *Eupatorium triplinerve*. *Asian Pacific Journal of Tropical Biomedicine*. 2012 Jan 1;2(3):S1329-32.
7. Kifayatullah M, Sarker MM, Mustapha MS. Phytochemical investigation of ethanolic extract of *Pericampylus glaucus* leaves from Malaysia by GC-MS analytical technique. *International Journal of PharmTech Research*. 2017;10:234-40.
8. Gautam B, Singh G, Wadhwa G, Farmer R, Singh S, Singh AK, Jain PA, Yadav PK. Metabolic pathway analysis and molecular docking analysis for identification of putative drug targets in *Toxoplasma gondii*: novel approach. *Bioinformatics*. 2012;8(3):134.
9. Liang J, Woodward C, Edelsbrunner H. Anatomy of protein pockets and cavities: measurement of binding site geometry and implications for ligand design. *Protein science*. 1998 Sep;7(9):1884-97.
10. Jones G, Willett P, Glen RC, Leach AR, Taylor R. Development and validation of a genetic algorithm for flexible docking. *Journal of molecular biology*. 1997 Apr 4;267(3):727-48.
11. Sentürk M, Ercan F, Yalcin S. The secondary metabolites produced by *Lactobacillus plantarum* downregulate BCL-2 and *BUFFY* genes on breast cancer cell line and model organism *Drosophila melanogaster*: molecular docking approach. *Cancer Chemotherapy and Pharmacology*. 2020 Jan 1;85(1):33-45.
12. Park BJ, Choi HR, Kim HS. A hybrid genetic algorithm for the job shop scheduling problems. *Computers & industrial engineering*. 2003 Dec 1;45(4):597-613.
13. Lorber DM, Shoichet BK. Flexible ligand docking using conformational ensembles. *Protein Science*. 1998 Apr;7(4):938-50.
14. Kuhn B, Mohr P, Stahl M. Intramolecular hydrogen bonding in medicinal chemistry. *Journal of medicinal chemistry*. 2010 Mar 25;53(6):2601-11.
15. Annamala MK, Inampudi KK, Guruprasad L. Docking of phosphonate and trehalose analog inhibitors into *M. tuberculosis* mycolyltransferase Ag85C: Comparison of the two scoring fitness functions GoldScore and ChemScore, in the GOLD software. *Bioinformatics*. 2007;1(9):339.
16. Kavitha K, Srikrishna D, Aparna P. An efficient one-pot three-component synthesis of 2-(4-(2-oxo-2H-chromen-3-yl)thiazol-2-yl)-3-arylacrylonitriles and their cytotoxic activity evaluation with molecular docking. *Journal of Saudi Chemical Society*. 2019 Mar 1;23(3):325-37.
17. Remenyi A, Good MC, Lim WA. Docking interactions in protein kinase and phosphatase networks. *Current opinion in structural biology*. 2006 Dec 1;16(6):676-85.
18. Crane L, Anastassiadou M, El Hage S, Stigliani JL, Baziard-Mouysset G, Payard M, Leger JM, Bizot-Espiard JG, Ktorza A, Caignard DH, Renard P. Design and synthesis of novel imidazoline derivatives with potent antihyperglycemic activity in a rat model of type 2 diabetes. *Bioorganic & medicinal chemistry*. 2006 Nov 15;14(22):7419-33.
19. Huang N, Shoichet BK, Irwin JJ. Benchmarking sets for molecular docking. *Journal of medicinal chemistry*. 2006 Nov 16;49(23):6789-801.
20. Singh A, Pal TK. In silico sequence analysis, structure prediction and function annotation of human Bcl-X beta protein. *International Journal Bioautomation*. 2014;18(1):23-30.