

 ISSN NO. 2320-5407	Journal Homepage: - www.journalijar.com INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR) Article DOI: 10.21474/IJAR01/18229 DOI URL: http://dx.doi.org/10.21474/IJAR01/18229	 INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR) ISSN 2320-5407 Journal Homepage: http://www.journalijar.com Journal DOI: 10.21474/IJAR01
---	--	---

RESEARCH ARTICLE

PREVALENCE OF VISUAL ACUITY DISORDERS AMONG INTERMEDIATE SCHOOL STUDENTS IN RIYADH, SAUDI ARABIA 2022

Mohanad Abdulaziz Aljetaily

Manuscript Info

Manuscript History

Received: 31 October 2022

Final Accepted: 30 November 2022

Published: December 2022

Key words:-

Demographics/Epidemiology, Visual Acuity, Myopia, Hyperopia

Abstract

Our objectives in this study are to determine the prevalence of visual acuity disorders and to identify the important risk factors that associated with visual acuity disorders among Intermediate school students in Riyadh, 2022. Design: a cross sectional study was performed the prevalence of visual acuity has been measured in Riyadh schools district. a sample size of 580 students throw a Multi-stage random sampling technic was conducted. The tools that has been used to collect data in this study are self administered questioner and the Snellen chart through placing it on 6 meter distance from the student, it contains C-letter-like shapes in different direction and sizes, which should be identified by the student.

Study Sample: All Male intermediate school students in governmental schools in Riyadh city, 2022. which included a total of 580 students.

Results: The prevalence of visual acuity impairment in our study was 14.3%, with a mean age of 14.14 ± 5.497 years old. 44.58 % of the students who are visually impaired didn't wear glasses. Mean hours spent on watching TV of visually impaired students is 4.02 ± 2.437 and mean hours spent on reading 1.76 ± 1.061 . Mean number of students who feel tired after time watching TV is 15.70%. There is increase in visual impairment among students as age increased. We observed that 60% of visually impaired students don't follow up with ophthalmologists.

Conclusion: We found that there is a significant relationship between visual acuity impairment and inheritance. Also one of the risk factors among students who suffer from visual acuity impairment is that they don't regularly follow up with an ophthalmologist, and 44.58 % of the students who are visually impaired didn't wear glasses.

Copy Right, IJAR, 2022., All rights reserved.

Introduction:-

Visual disorders are a prominent health problem. There was an estimation of 161 million people around the world with visual disorder. Most of them are in developing countries, for instance Saudi Arabia. Visual acuity disorders are still considered a major setback that has a negative effect on the community. (2) Furthermore, visual acuity disorders are commonly found with difficulties in handling daily life issues, as it is considered the main sense of communication. The World Health Organization concluded that community-derived data on the incidence of visual acuity disorders are instantly desired as they are being important for detecting requirements for therapy and rehabilitation, development, and conducting screening programs, and assessing main concerns for different

communities(4).Data regarding visual acuitydisorders in Saudi Arabia are uncommon. Only minorities of researches wereimplementedin the last twenty years. Yet, the urge for additionalresearches to assess the requirements for implementing, evaluating and screening interventional programs are needed. We are implementing thisresearchfor the purpose ofassessing the magnitude of visual acuitydisordersamong school students in Riyadh(capital of Saudi Arabia). Our objectiveswereto determine the prevalence of visual acuity disorders and to identify the significant risk factors that associated with visual acuity disorders among Intermediate school students.

Materials and Methods:-

Study area Riyadh city, Saudi Arabia. Study populationAll school students in governmental schools in Riyadh city in 2022.

A cross-sectional study with a sample size of 580 students throw aMulti-stage random sampling technic was done. Riyadh has been divided to five regions and one school was chosen from each region by using simple random sampling technique. Using stratified random sampling with proportional allocation one class from each school year was chosen. All students present at the time of the study in the chosen classes were included in the study.

Data collection technique:

Upon visiting the selected schools we randomly chose one class from each grade, the randomly selected students were then gathered in the study room and a self administered questionnaire with standard sequence and pre-coded answers (Structured Technique) was distributed to the students targeted for gathering of statistics (nationality, level of education, age) Additional significant information associated with visual disorders and its risk factors, like diabetes, hypertension or family history of similar diseases , time spent on reading or in front of TV, frequency of visiting an ophthalmologist, then history of eye disorders were documented. The visual acuity of the students was tested by using a Snellen chart (which developed by the Dutch ophthalmologist Herman Snellen), through placing it on 6 meter distance from the student, and contain E-letter-like shapes in different directions and sizes, which should be identified by the student. World health organization defined visual impairment as a visual acuity of less than 6/18 in the better. A scale was used to measure weight and height. Then filled questionnaires were collected.

Statistical analysis:

Mean + SD was used to describe each quantitative variable :(Age, Age start using glasses, Family member who wears glasses, Times a year visiting the ophthalmologist(number). Time spent watching TV (hours)

Frequency (percentage): was used to describe each qualitative variable (Nationality, Family hypertension, Diabetes, Wear glasses)

T- test: was used to find if there is any difference between quantitative variables and one qualitative variable with only two categories. ANOVA will be used to find if there is any difference between quantitative variables and one qualitative variable with more than two categories. Chi-Square test: was used to find any association between two qualitative variables. Multivariate analysis: Regression analysis was used to predict visual acuity impairment (dependent variable) from one or more risk factor(s) (independent variables). P- value was set to be less than 0.05 throughout the study.

Ethical considerations:

Permission from school health Director was obtained prior to collecting data.

Results:-

Mean age of 580 students how participated in the study(SD) was $14.14 \pm (5.497)$. Totalnumber of students in each phase of the selected schools are Grade one 190 (32.8%). Grade two 202 (34.8%). Grade three 188 (32.4%). ThePrevalence of Visual acuity disorders Among Intermediate School Students was (14.3%). Total of students whom visually impaired are 83 students .we can see the number of students increases gradually with grade (age). And the percentages are grade 1 (13.15%), grade 2 (14.36%), and grade 3 (16.48%).Mean and (SD) of the Hours Spent on Reading $1.43 \pm (1.226)$, and Hours Spent on TV $4.02 \pm (2.402)$.Most of thestudents (85.7%) had visual acuity that consider normal according to world health organization.Distribution of both eyes Visual Acuity Value Among Intermediate School Students in Riyadh 2022:

Visual Acuity	R Eye		L Eye	
	N	%	N	%
6	196	33.8	223	38.4
9	207	35.7	198	34.1
12	64	11.0	57	9.8
18	43	7.4	28	4.8
24	29	5.0	25	4.3
36	25	4.3	34	5.9
60	16	2.8	15	2.6
Total	580	100.0	580	100.0

-as we see there is no big difference between right and left eye visual acuity according to WHO (ICD 10) worse than 18/6 in the better eye is visually impaired

Visual impairment	Times visiting an Ophthalmologist per Year								Total
	0	1	2	3					
	N	%	N	%	N	%	N	%	
Yes	50	60.24	15	18.07	14	16.86	4	4.81	83
Total	498	85.86	44	7.58	25	4.31	13	2.24	580

*in this table we can see that 60.24% of visually impaired students did not visit an ophthalmologist regularly

Relationship Between Wearing Glasses and Family Member who Wears Glasses

	Wearing glasses				Total		P
	Yes	No					
Family members who wear glasses	N.	%	N.	%	N.	%	<.001
Yes	66	17.27	316	82.73	382	65.86	
No	13	6.56	185	93.44	198	34.14	
Total	79	13.62	501	86.38	580	100	

-in this table we can see that there is a significant relation between visual impairment and family history.

Discussion:-

The study aimed to determine the prevalence of visual acuity disorders and associated risk factors among intermediate school students in Riyadh in 2022. The prevalence of visual acuity disorders was 14.3%, which agrees with a study done by Al-Shaalin in al jouf that found the prevalence of visual acuity disorders was 13.9% (1). A much lower prevalence was reported in another study done by Khalid F. Tabbara in Saudi Arabia In 2011 and on contrast disagrees with findings that stated the prevalence of visual acuity disorders was 7.8% (3). This disagreement could be explained by the difference of the participant's age groups in both studies and place. Our study revealed that refractive error is the primary cause of visual acuity disorders and its increases with the age. This finding was coherent with other researches that indicated that older persons tended to have lower rates of myopia than younger persons.(5) This remarkable decline could be explained by the fact that the estimated occurrences for causes of visual impairments in our study were based on the proportion of each cause in relation to all other causes, we found that there is a significant relationship between visual acuity disorder and inheritance. 17.27% of the students who wear glasses and suffer from visual acuity disorder, found to have a family members how wear glasses, which may suggest genetic predisposition among patients suffering from visual acuity disorders.

Acknowledgements:-

Our deep appreciation to those who participated in our study including, students, organizers and School principles.

References:-

- 1)Al-Shaalin FF, Bakrman MA, Ibrahim AM, Aljouidi AS Source Directorate of Primary Health Care, Aljouf, Saudi Arabia.
- 2)M F Al Faran, A A Al-Rajhi, O M Al-Omar, S A Al-Ghamdi, M Jabak International Ophthalmology (1993)Volume: 17, Issue: 3, Pages: 161-165.

- 3) Reprint requests to Research Department, King Khaled Eye Specialist Hospital, PO Box 7191, Riyadh 11462, Saudi Arabia (Dr Tabbara). Khalid F. Tabbara MD; Dennis Ross-Degnan MSPH.
- 4) Behrman, R, Kligman, R, Jenson, H. Nelson Textbook of Pediatrics 16th edition, Philadelphia.
- 5) Broderic P. Pediatric Vision Screening for family Physician, American Family Physician, Clinical ophthalmology.