



RESEARCH ARTICLE

EXOSOMES FOR FACIAL REPAIR, REGENERATION AND REJUVENATION: COMPARISON BETWEEN DIFFERENT COMMERCIAL PRODUCTS

Antonio Salvaggio¹, Anna Privitera^{1,2}, Greta Ferruggia¹, Massimo Zimbone³ and Maria Violetta Brundo¹

1. Department of Biological, Geological and Environmental Sciences, University of Catania.
2. Department of Biomedical and Biotechnological Sciences, University of Catania.
3. Institute for Microelectronics and Microsystems, National Research Council of Italy (CNR-IMM), 95123 Catania, Italy.

Manuscript Info

Manuscript History

Received: 22 October 2024

Final Accepted: 25 November 2024

Published: December 2024

Key words:-

Plant Nanovesicles, Bovine Colostrum Derived Exosomes, *Rosa Damascena* Callus, *Centella Asiatica* Callus, *Euphorbia Supina* Stem, MTT Test, Wound Healing

Abstract

Exosomes have been studied as a potential therapeutic option for improving skin texture. The mode of action of exosomes for skin texture likely involves several mechanisms, including stimulation of collagen production, promotion of skin cell proliferation; reduction of oxidative stress; regulation of inflammatory and antioxidant responses; extracellular matrix remodeling, which is important for maintaining skin structure and function. Aim of this study is to compare the *in vitro* effects of some commercial products that use new technologies containing plant derived nanovesicles from *Rosa damascena* callus, *Centella asiatica* callus, *Euphorbia supina* stem and exosomes from bovine colostrum passively loaded with growth factors and cytokines purified from bovine colostrum, for facial repair, regeneration and rejuvenation. Of all the products tested, only product containing exosomes purified from colostrum gave very encouraging results in terms of effects on proliferation, cell viability and wound repair. The other products tested gave results comparable to the untreated samples.

Copyright, IJAR, 2024. All rights reserved.

Introduction:-

In the last decade, interest in extracellular vesicles has grown considerably, since the key role of exosomes in intercellular communication has been demonstrated [1,2]. Exosomes are spherical extracellular nanovesicles with an endosomal origin with sizes ranging from 30 to 100 nm [3], as transporters of different bioactive molecules including proteins and lipids, can take part in different physiological mechanisms by transferring their contents to recipient cells, thus acting as extracellular messengers in cell-to-cell communication, but they are also attracting attention as potential therapeutic tools and positive effects on the regeneration of many tissues [3,4]. In this context, exosomes are attracting growing interest thanks to their relevance in the modulation of cellular processes (physiological and pathological), as well as for their potential beneficial effects on the organism, especially anti-inflammatory, anti-tumor, tissue regeneration and modulation of commensal microbiota. Furthermore, it is hypothesized to also use them as transporters of active molecules, to exploit them to improve the absorption and effectiveness of drugs thanks to their low immunogenicity and high stability in the gastrointestinal tract [5]. It is widely accepted that the therapeutic potential of several components purified from bovine colostrum may be mediated largely by paracrine factors, and exosomes might be the main components of such paracrine factors [6]. Thus, exosomes derived from bovine colostrum, as they induce the repair of damaged tissues, represent a

Corresponding Author:-Maria Violetta Brundo

Address:-Department of Biological, Geological and Environmental Sciences, University of Catania.

relevant therapeutic option in regenerative medicine and have a more promising future [7-11], especially when compared to the results obtained with exosomes purified from stem cells [12] or plant nanovesicles [13]. Among the soluble molecules that derive from colostrum we find cytokines and growth factors, which are involved in immunomodulation[14]; in fact, as demonstrated with a quantitative proteomics analysis, exosomes from bovine colostrum are significantly enriched with proteins that can potentially regulate the immune response and growth [14]. Also, wound healing, a dynamic and complex process involving a sequence of coordinated cellular events through which the organism attempts to restore the structural integrity and complete or partial functionality of the tissue, can be improved using products based on exosomes purified from bovine colostrum [8]. The healing of skin wounds begins at the very moment of the wound and involves cells present at the site of the lesion, cells migrated from different areas of the organism (such as keratinocytes, endothelial cells, fibroblasts, inflammatory cells) and inflammatory mediators (such as cytokines, chemokines, growth factors). Much research agrees that interleukin-6 (IL-6) [15] and transforming factor beta-1 (TGF- β 1) [16] play key roles in these processes. It is therefore clear that to promote the regeneration and healing processes of the skin it is useful to stimulate cell vitality, promote healing, stimulate the proliferation and migration of fibroblasts, endothelial cells and keratinocytes, reduce the inflammatory state and oxidative damage [17]. Similar processes are necessary for tissue rejuvenation and skin tightening [17].

The aim of this study is to compare the *in vitro* effects of some commercial products that use new technologies containing plant-derived nanovesicles or bovine colostrum-derived exosomes for facial repair, regeneration and rejuvenation.

Materials and Methods:-

For our research we purchased commercial products that utilize new technologies containing among the ingredient's extracellular vesicles. One product contains plant extracellular vesicles extract from *Rosa damascena* callus (indicated with Rosa-NVs). The second product contains plant extracellular vesicles extract from *Centella asiatica* callus (indicated with Centella-NVs). The third product contains plant extracellular vesicles extract from *Euphorbia supina* stem (indicated with Euphorbia-NVs). The fourth compared product contains exosomes purified from bovine colostrum passively loaded with growth factors and cytokines purified from bovine colostrum (indicated with Colostrum-Exo). This latest technology is called AMPLEX plus technology.

Propagation and maintenance of cells

Normal adult primary human epidermal keratinocytes (HEKa) (Invitrogen) were maintained in a humidified incubator with 5% CO₂ at 37°C in Epilife Medium containing Epilife defined growth supplement according to the manufacturer's instructions and split every 2–3 days depending on cell confluence.

Normal adult human dermal fibroblasts (HDFa) (ThermoFisher Scientific) were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% FBS, streptomycin (0.3 mg mL⁻¹) and penicillin (50 IU mL⁻¹), and GlutaMAX (1 mM) using 75 cm² polystyrene culture flasks. Cells were maintained in a humidified environment (37 °C and 5% CO₂) and split every 2–3 days depending on cell confluence.

Analysis of cell proliferation/metabolic status

Keratinocytes and fibroblasts were harvested by using a trypsin-EDTA solution, counted with a hemocytometer and plated in 96-well plates (5×10⁵ keratinocytes/well; 4×10⁴ fibroblasts/well). The following day, cells were treated with four products at three different concentrations (0.5%, 1% and 2%) and incubated for 24 in a humidified environment (37 °C and 5% CO₂). At the end of the treatment, MTT solution (1 mg/mL) was added to each well and cells were incubated for 2 hours in a humidified environment (37°C and 5% CO₂). Epoch Microplate Spectrophotometer (Bio Tec) was used to read the absorbance at 569 nm. Values were normalized with respect to control untreated cells and were expressed as the percent variation of cell proliferation/metabolic activity.

Scratch-wound assay

Human non immortalized fibroblasts were plated into six-well plates at the density of 2.5 x 10⁶ cells/well allowed to incubate until confluence. Once the confluence was reached, cells were scraped by using a 1 mL sterile pipette tip to form a wound. Cells were washed by using PBS to remove detached cells before adding the medium, in absence or presence of tested products used at 2%. Untreated cells were considered as control. Images were captured at 0, 1 and 24 hours after scratching.

Quantification and characterization of plant-derived nanovesicles and colostrum-derived exosomes

The characterization of plant-derived nanovesicles and colostrum-derived exosomes was performed with a light scattering according to [18]. Measurements were performed with a homemade apparatus using a quartz scattering cell, confocal collecting optics, a Hamamatsu photomultiplier mounted on a rotating arm, a BI-9100AT hardware correlator (Brookhaven Instruments Corporation) and illuminating the sample with a 660 nm laser. The power ranged between 5 and 15mW. Low power intensity was used to avoid convective motions due to local heating.

Statistical analysis

The statistical analysis was carried out by using Graphpad Prism software (version 8.0) (Graphpad software, San Diego, CA, USA). Two-way analysis of variance (ANOVA), followed by Tukey's post hoc test, was used for multiple comparisons. The statistical significance was set at p-values < 0.01. Data were reported as the mean $\pm$ SD of at least 2 independent experiments.

Results and Discussion:-

The size distribution of extracellular vesicles was determined using the Dynamic Light Scattering technique. In this measurement technique, a monochromatic radiation with wavelength λ and wavevector k_i , generally obtained via a laser, is passed through a polarizing filter and then made to impact on a sample of the colloidal dispersion of nanoparticles. When light radiation impacts a particle, scattering occurs where the light is scattered in all directions with a wave vector k_f which depends on the shape, dimensions and optical properties of the particle. Since the suspension is made up of a very large number N of particles, the intensity of the total scattering radiation that reaches the detector and is measured by it will be the sum (due to interference) of the scattering contributions of the individual particles. If we consider a colloidal dispersion, whose particles move independently with Brownian-type motion because of thermal agitation, the position of the particles themselves varies randomly over time. Therefore, if the light radiation emitted by scattering is collected, after being filtered by a second polarizer, by a detector, the measured intensity will fluctuate around an average value (random walking of nanoparticles) [19,20]. In table 1 are indicated the diameter size and the concentration of the plant-derived nanovesicles and colostrum-derived exosomes present in tested products. The dynamic light scattering analysis has revealed that the diameter size of Rosa-NVs (Fig 1A), Centella-NVs (Fig 1B) and Euphorbia-NVs (Fig 1C) are 565 nm, 677 nm and 667 nm respectively and that diameter size of colostrum-derived exosomes is 113 nm (Fig 1D). Exosomes are a class of cell-derived extracellular vesicles of endosomal origin and are typically 30-150 nm in diameter. They are the smallest type of extracellular vesicles, and their role is closely linked to size. Plant nanovesicles called also exosome-like are nano-sized vesicles of 50-1000 nm diameter [13]. Recent data have suggested that plant exosome-like nanovesicles (PENVs) and PENV-derived artificial nanocarriers (APNVs) may have possible clinical applications, but there are many obstacles to overcome [13]. Currently extracted plant nanovesicles are highly heterogeneous, containing multiple vesicle types, and of unknown origin. originates in plant cells [13]. Another problem to overcome depends on the fact that, because PENVs and APNVs do not come from mammalian cells, they may have poor targeting ability in some targeted tissues [13].




Figure 1:- DLS analysis used to characterize the size distributions of exosomes within four products: (A) product with plant extracellular vesicles from *Rosa damascena* callus; (B) product with plant extracellular vesicles from *Centella asiatica* callus; (C) product with plant extracellular vesicles from *Euphorbia supina* stem; (D) product with exosomes from colostrum bovine loaded with growth factors and cytokines purified from bovine colostrum.

In the four products analyzed the concentration of nanovesicles was also different (Table 1), lower for plant-derived nanovesicles (2^6 part/ml, 2^6 part/ml and 2^5 part/ml for Rosa-NVs, Centella-NVs and Euphorbia-NVs respectively) and higher for colostrum-derived exosomes (3.8^9 part/ml).

Table 1:- Diameter size and concentration of extracellular vesicles analyzed with DLS.

Product	Diameter size (nm)	Concentration (particles/ml)
Rosa-NVs	565	2^6
Centella-NVs	677	2^6
Euphorbia-NVs	667	2^5
Colostrum-Exo	113	3.8^9

The viability cell assay was conducted for both, human keratinocytes and dermal fibroblasts, and the results are presented in Figures 2 and 3, respectively. The results showed a statistically significant increase in the level of cell viability after treatment with product containing colostrum exosomes at different doses ($p < 0.001$) compared to cells

treatment with plant-derived nanovesicles, whose results were lower than the negative control (untreated cells) at all concentrations tested.




Figure 2:-The cell survival of human keratinocytes at 24h following treatment with the four products tested to different concentrations (0.5%, 1% and 2%). NC (Negative Control, cells without treatment); PC (Positive Control, 80 μ M H₂O₂). The asterisks denote the degree of significance between results: **p<0.001. Errors bars represent the Standard Deviation of the mean (experiment was repeated 2 times).




Figure 3:- The cell survival of human fibroblasts at 24h following treatment with the four products tested to different concentrations (0.5%, 1% and 2%). NC (Negative Control, cells without treatment); PC (Positive Control, 80 μ M H₂O₂). The asterisks denote the degree of significance between results: **p<0.001. Errors bars represent the Standard Deviation of the mean (experiment was repeated 2 times).

The experimental results obtained with the scratch test have demonstrated a significant increase in the level of cell proliferation following a 1 hour and 24 hours treatment with product containing colostrum-derived exosomes loaded with colostrum biomolecules (growth factors and cytokines) at 2% concentrations. Figure 4, in fact, shows a statistically significant increase in the level of cell viability after treatment with product containing colostrum exosomes ($p < 0.001$) compared to cells treatment with plant-derived nanovesicles and negative control (untreated cells).




Figure 3:- Scratch assay on human fibroblasts treated with the four products to 2% concentration at 0-, 1- and 24-hours' time points. Untreated cells were considered as a control (NC). The asterisks denote the degree of significance between results: ** $p < 0.001$. Errors bars represent the Standard Deviation of the mean (experiment was repeated 3 times). The box shows the images comparing the treatment with colostrum-derived exosomes and plant-derived nanovesicles (the results were comparable for the three products containing plant NVs).

Wound healing, as well as tissue rejuvenation and skin tightening, are a complex process that involves multiple stages, hemostasis, inflammation, proliferation, and tissue regeneration being the most critical steps[21]. The focus of this study was to examine the *in vitro* effects of some commercial products that use new technologies containing plant derived nanovesicles or exosomes for facial repair, regeneration and rejuvenation. The results obtained have demonstrated that the product containing colostrum-derived exosomes passively loaded with growth factors and cytokines from colostrum (AMPLEX plus technology) promote cell vitality and wound healing, determining a statistically significant increase in the level of cell viability after treatment at different doses (0.5%, 1% and 2%) compared to cells treatment with plant-derived nanovesicles. Also, the results obtained with the scratch test have demonstrated a significant increase in the level of cell proliferation already after a 1 hour treatment (78%) with product containing colostrum-derived exosomes loaded with colostrum biomolecules (growth factors and cytokines) at 2% concentrations. Our results are in accordance with other studies published in which the effectiveness of plant-derived nanovesicles was analyzed on facial repair, regeneration and rejuvenation. The results obtained in these studies have showed a low efficacy of plant nanovesicles, almost always comparable to untreated cells or even lower [13, 22-26]. Some Authors, for example, hypothesized that *Rose damascena* stem cell in culture can release exosome-like particles that may have biological function in cells relevant to skin, such as hair papilla cells, fibroblasts and melanocytes [23]. The Authors have found that the *Rose Stem Cell* cultured supernatant reduce viability in human dermal papilla cells, demonstrating its cytotoxic effect. However, the *Rose damascena* stem cell nanovesicles did not show any detrimental effect on viability of Human Dermal Papilla cells, but they have not even demonstrated a positive effect. Furthermore, the Authors declare that the *Rose damascena* stem cell nanovesicles also stimulate skin fibroblast proliferation and collagen production, as well as *in vitro* wound healing, but their results are just higher than those of untreated cells and after 24 hours there was a closure of just 27.5% [23], much

lower than obtained in our experiments with bovine colostrum exosomes-derived after only one hour of treatment, equal to 78%. Other Authors to confirm the cellular toxicity of *Centella asiatica* extract and *Centella asiatica* exosomes, have performed the MTT assay with a keratinocyte cell line (KEKa cells)[24]. *Centella a.* extract and *Centella a.* exosomes were nontoxic to the cells at all concentrations and induced cell proliferation compared with the control. Moreover, *Centella a.* extract induced higher proliferation than *Centella a.* exosomes; in particular, *Centella a.* exosomes has a maximum percentage of cell viability equal to $123.7 \pm 4.8 \%$ with a tested concentration of 1×10^5 particles/ml. In literature there is no research on the effects of exosomes purified from *Euphorbia asiatica*, but some authors have studied the antioxidant and skin-whitening effects of a 70% ethanol extract of *Euphorbia* extract. Authors have showed that *Euphorbia* extract significantly reduced tyrosinase activity and melanin content in a dose-dependent manner and MTT test it showed that the *Euphorbia* extract was not toxic as the percentage cell viability was the same to that of untreated cells[25]. Exosomes from colostrum bovine instead have an excellent structural and have a great potential as natural therapeutic agents to repair UV-irradiated skin aging and damage[27]. Authors showed that treatment with colostrum exosomes prevented the UV-induced generation of intracellular reactive oxygen species in epidermal keratinocytes. In UV-stimulated melanocytes, have reduced significantly the production of the protective skin-darkening pigment melanin; in the human dermal fibroblasts treated with colostrum exosomes, the expression of matrix metalloproteinases was suppressed, whereas increased cell proliferation was accompanied by enhanced production of collagen, a major extracellular matrix component of skin[27]. Other Authors suggested its potential antiaging properties. These effects are attributed to its capacity for moisturization and wrinkle reduction, which correlate with improvements in the expression levels of FLG, CD44 in keratinocytes, and HAS2 in fibroblasts[28].

Conclusion:-

Extracellular vesicles, including exosomes, are produced via the endosomal pathway and released in the extracellular space upon fusion with the plasma membrane[29]. Several studies show that these extracellular vesicles play a key role in cell-to-cell communication, transporting bioactive molecules[30-32]. Exosomes were reported to regulate also all phases of skin repair and rejuvenation[33, 36]. Exosomes from colostrum they appear to have a high potential for the applications in regenerative medicine[27, 28], as also demonstrated by our research. Of all the products tested, in fact, only product containing exosomes purified from colostrum passively loaded with growth factors and cytokines derived purified from colostrum (AMPLEX plus technology) gave very encouraging results in terms of effects on proliferation, cell viability and wound repair. The other products tested gave results comparable to the untreated samples, or even lower.

Author Contributions:

Conceptualization, A.S and M.V.B.; methodology, M.V.B.; software, G.F. and M.Z.; formal analysis, G.F. and M.Z.; investigation, G.F.; data curation, A.P., G.F. and M.Z.; writing original draft preparation, A.S. and M.V.B.; Writing – Review & Editing A.S. and A.P.; supervision, M.V.B. All authors have read and agreed to the published version of the manuscript.

Funding:

This research received no external funding.

Institutional Review Board Statement:

This study was performed in line with the principles of the Declaration of Helsinki and does not require approval from the Ethics Committee of University of Catania.

Acknowledgments:-

A.P. thanks the International PhD Program in Neuroscience and G.F. thanks the Ph.D. program FSE Notice 1/2021, at the University of Catania.

Conflicts of Interest:

The authors declare that they have no conflict of interest regarding the contents of this article.

References:-

1. Bischoff JP, Schulz A, Morrison H. The role of exosomes in intercellular and inter-organ communication of the peripheral nervous system. *FEBS Lett.* 2022, 596(5):655-664. doi: 10.1002/1873-3468.14274.

2. Schneider A, Simons M. Exosomes: vesicular carriers for intercellular communication in neurodegenerative disorders. *Cell Tissue Res.* 2013, 352(1):33-47. doi: 10.1007/s00441-012-1428-2.
3. Poinot V, Pizzinat N, Ong-Meang V. Engineered and Mimicked Extracellular Nanovesicles for Therapeutic Delivery. *Nanomaterials (Basel).* 2024, 14(7):639. doi: 10.3390/nano14070639.
4. Basu J, Ludlow JW. Exosomes for repair, regeneration and rejuvenation. *Expert Opinion on Biological Therapy*, 2016, 16(4): 489–506. <https://doi.org/10.1517/14712598.2016.1131976>.
5. Zeng H, Guo S, Ren X, Wu Z, Liu S, Yao X. Current Strategies for Exosome Cargo Loading and Targeting Delivery. *Cells.* 2023, 12(10):1416. doi: 10.3390/cells12101416.
6. Yağıntaş YM, Duman H, López JMM, Portocarrero ACM, Lombardo M, Khallouki F, Koch W, Bordiga M, El-Seedi H, Raposo A, Alves JLB, Karav S. Revealing the Potency of Growth Factors in Bovine Colostrum. *Nutrients.* 2024, 16(14):2359. doi: 10.3390/nu16142359.
7. Dos Santos PR, Kraus RB, da Silva Nascente P. Exploring the potential of bovine colostrum as a bioactive agent in human tissue regeneration: A comprehensive analysis of mechanisms of action and challenges to be overcome. *Cell BiochemFunct.* 2024, 42(4):e4021. doi: 10.1002/cbf.4021.
8. Jafarinejad S, Baumgartner A, Elis GJ and Najafzadeh, M. The Efficacy of a Novel Biological Formulation from Bovine Milk Colostrum Exosomes and its Growth Factors in Enhancing the Process of Wound Healing. *Research & Reviews: Journal of Biology* 2024, 12 (1).doi: 10.4172/2322-0066.12.1.001.
9. Privitera A, Ferruggia G, Contino M, Maugeri S, Zimbone M, Cardile V, Caruso G, Brundo MV. Nutri Complex 150+: a new and effective approach to facial rejuvenation. *bioRxiv* 2024.doi: 10.1101/2024.10.05.616805.
10. Mandloi V, Banerjee T, Sharma A, Pratap A, Ansari MA, Srivastava V. Role of Bovine Colostrum Dressing on Chronic Non-Healing Wounds in Comparison to Conventional Dressing: A Case-Control Study. *Int J Low Extrem Wounds.* 2024;15347346241241578. doi: 10.1177/15347346241241578.
11. Kim H, Kim DE, Han G, Lim NR, Kim EH, Jang Y, Cho H, Jang H, Kim KH, Kim SH, Yang Y. Harnessing the Natural Healing Power of Colostrum: Bovine Milk-Derived Extracellular Vesicles from Colostrum Facilitating the Transition from Inflammation to Tissue Regeneration for Accelerating Cutaneous Wound Healing. *Adv Healthc Mater.* 2022, 11(6):e2102027. doi: 10.1002/adhm.202102027.
12. Rezabakhsh A, Sokullu E, Rahbarghazi R. Applications, challenges and prospects of mesenchymal stem cell exosomes in regenerative medicine. *Stem Cell Res Ther.* 2021, 12(1):521. doi: 10.1186/s13287-021-02596-z.
13. Chen X, Xing X, Lin S. et al. Plant-derived nanovesicles: harnessing nature's power for tissue protection and repair. *J Nanobiotechnol* 2023, 21: 445. <https://doi.org/10.1186/s12951-023-02193-7>.
14. Samuel M, Chisanga D, Liem M. et al. Bovine milk-derived exosomes from colostrum are enriched with proteins implicated in immune response and growth. *Sci Rep* 2017, 7: 5933.doi: 10.1038/s41598-017-06288-8.
15. Johnson BZ, Stevenson AW, Prêle CM, Fear MW, Wood FM. The Role of IL-6 in Skin Fibrosis and Cutaneous Wound Healing. *Biomedicines.* 2020, 8(5):101. doi: 10.3390/biomedicines8050101.
16. Ramirez H, Patel SB, Pastar I. The Role of TGFβ Signaling in Wound Epithelialization. *Adv Wound Care (New Rochelle).* 2014, 3(7):482-491. doi: 10.1089/wound.2013.0466.
17. Boen M, Vanaman Wilson MJ, Fabi S. Skin tightening-does it really exist? *Semin Cutan Med Surg.* 2017, 36(4):202-206. doi: 10.12788/j.sder.2017.034.
18. Ferruggia G, Contino M, Zimbone M, Brundo MV. Effectiveness of a Novel Compound HAIR & SCALP COMPLEX on Hair Follicle Regeneration. *Cosmetics* 2024, 11(1): 10. doi: 10.3390/cosmetics11010010.
19. Zimbone M, Musumeci P, Baeri P, Messina E, Boninelli S, Compagnini G, Calcagno L. Rotational dynamics of gold nanoparticle chains in water solution. *J. Nanopart Res.* 2012, 14, 1308. doi: 10.1007/s11051-012-1308-4
20. Zimbone M, Baeri P, Calcagno L, Musumeci P, Contino A, Barcellona ML, Bonaventura G. Dynamic light scattering on bioconjugated laser generated gold nanoparticles. *PLoS ONE* 2014, 9, e89048. doi: 10.1371/journal.pone.0089048
21. Tottoli EM, Dorati R, Genta I, Chiesa E, Pisani S, Conti B. Skin Wound Healing Process and New Emerging Technologies for Skin Wound Care and Regeneration. *Pharmaceutics.* 2020, 12(8): 735. doi: 10.3390/pharmaceutics12080735.
22. Wu W, Zhang B, Wang W, Bu Q, Li Y, Zhang P, Zeng L. Plant-Derived Exosome-Like Nanovesicles in Chronic Wound Healing. *Int J Nanomedicine.* 2024, 19:11293-11303. doi: 10.2147/IJN.S485441.
23. YJ Won, E Lee, SY Min, BS Cho. Biological Function of Exosome-like Particles Isolated from Rose (Rosa Damascena) Stem Cell Culture Supernatant. *bioRxiv* 2023. doi: <https://doi.org/10.1101/2023.10.17.562840>.
24. Kim M, Yoon EJ, Kim JS, Park SJ, Lee H. Comparative analysis of the transcriptome and efficacy of bioactive Centella asiatica exosomes on skin cells. Pre-print 2023. doi: 10.21203/rs.3.rs-2787704/v1.

25. Kang SH, Jeon YD, Cha JY, Hwang SW, Lee HY, Park M, Lee BR, Shin MK, Kim SJ, Shin SM, Kim DK, Jin JS, Lee YM. Antioxidant and skin-whitening effects of aerial part of *Euphorbia supina* Raf. Extract. BMC Complement Altern Med. 2018, 18(1):256. doi: 10.1186/s12906-018-2323-5.
26. Pintus F, Floris S, Fais A, Era B, Porcedda C, Tuberose CIG, Caddeo C. *Euphorbia characias* Extract: Inhibition of Skin Aging-Related Enzymes and Nanoformulation. Plants 2022; 11(14):1849. doi: 10.3390/plants11141849.
27. Han G, Kim H, Kim DE, Ahn Y, Kim J, Jang YJ, Kim K, Yang Y, Kim SH. The Potential of Bovine Colostrum-Derived Exosomes to Repair Aged and Damaged Skin Cells. Pharmaceutics. 2022, 14(2):307. doi: 10.3390/pharmaceutics14020307.
28. Lu L, Bai W, Wang M, et al. Novel roles of bovine milk-derived exosomes in skin antiaging. J Cosmet Dermatol. 2024, 23:1374-1385. doi:10.1111/jocd.16112.
29. Zhang, Y., Liu, Y., Liu, H. et al. Exosomes: biogenesis, biologic function and clinical potential. Cell Biosci 9, 19 (2019). doi: 10.1186/s13578-019-0282-2.
30. Record M. Intercellular communication by exosomes in placenta: a possible role in cell fusion? Placenta. 2014;35(5):297–302. doi: 10.1016/j.placenta.2014.02.009.
31. Simpson RJ, Lim JW, Moritz RL, Mathivanan S. Exosomes: proteomic insights and diagnostic potential. Expert Rev Proteom. 2009;6(3):267–83. doi: 10.1586/epr.09.17.
32. Camussi G, Deregibus MC, Bruno S, Cantaluppi V, Biancone L. Exosomes/microvesicles as a mechanism of cell-to-cell communication. Kidney Int. 2010;78(9):838–48. doi: 10.1038/ki.2010.278.
33. Prasai A, Jay JW, Jupiter D, Wolf SE, El Ayadi A. Role of Exosomes in Dermal Wound Healing: A Systematic Review. J Invest Dermatol. 2022, 142(3 Pt A):662-678.e8. doi: 10.1016/j.jid.2021.07.167.
34. Hajialiasgari Najafabadi A, Soheilifar MH, Masoudi-Khoram N. Exosomes in skin photoaging: biological functions and therapeutic opportunity. Cell Commun Signal. 2024, 22(1):32. doi: 10.1186/s12964-023-01451-3.
35. Sreeraj H, AnuKiruthika R, Tamilselvi KS, Subha D. Exosomes for skin treatment: Therapeutic and cosmetic applications, Nano TransMed 2024, 3:100048. doi: 10.1016/j.ntm.2024.100048.