



ISSN NO. 2320-5407

Journal Homepage: -[www.journalijar.com](http://www.journalijar.com)

## INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/20863  
DOI URL: <http://dx.doi.org/10.21474/IJAR01/20863>



INTERNATIONAL JOURNAL OF  
ADVANCED RESEARCH (IJAR)  
ISSN 2320-5407  
Journal Homepage: <http://www.journalijar.com>  
Journal DOI:10.21474/IJAR01

### RESEARCH ARTICLE

## STUDY OF BACTERIAL PATHOGENS ON HIGH TOUCH SURFACES AND THEIR ANTIMICROBIAL SUSCEPTIBILITY PATTERN IN A TERTIARY CARE HOSPITAL

Mounika S<sup>1</sup> and Pavani G<sup>2</sup>

1. PG Student, Department of Microbiology, Apollo Institute of Medical Sciences and Research, Hyderabad.
2. Professor&HOD, Department of Microbiology, Apollo Institute of Medical Sciences and Research, Hyderabad.

### Manuscript Info

#### Manuscript History

Received: 20 February 2025

Final Accepted: 23 March 2025

Published: April 2025

#### Key words:-

Hospital Environment, High touch areas,  
Bacterial Pathogens

### Abstract

**Background:** The hospital environment surfaces plays an important role in causing Hospital acquired infections. This study conducted to assess the surface contamination in four different wards at Apollo General Hospital, Hyderabad which are frequently touched by health care professionals. Hence we conducted a study to assess the bacterial pathogens on frequently touched areas in the hospital and to determine antibiotic susceptibility of these pathogens

**Materials and methods:** A Cross-sectional study was conducted in Apollo General Hospital from April to May 2024. A total of 48 swabs collected from surfaces of floor, bed, wall, doorhandles, stethoscope, thermometer, IV stand, examination table, sink tap, bed side trolley and drug trolley from Medicine, OBG, Surgery and Paediatric wards using sterile moist swabs. Gram staining and culture on blood agar and MacConkey agar was done. Identification of the isolates was done as per standard protocol. Antimicrobial susceptibility testing was done as per CLSI guidelines

**Results:** Out of the 48 samples processed, 52% showed bacterial growth. *CoNS* (68%) was the predominant isolate followed by *Staphylococcus* (16%), *Acinetobacter spp* (4%) and *Proteus* (4%). The other isolates were *Enterococcus* (4%) and *Burkholderia* (4%). Majority of the *CoNS* were isolated from floor, bed and wall. Most of the isolates were sensitive to ciprofloxacin and gentamicin and resistant to penicillin

**Conclusion:** These results in our study suggests that there should be increased emphasis on frequent cleaning and disinfection of frequently touched surfaces in hospital to reduce bacterial contamination.

"© 2025 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

### INTRODUCTION

Bacterial contamination of high touch surfaces in health care facilities is a significant concern as they harbour potential pathogens and act as source of hospital acquired infections [1]. These bacteria persist in hospital environments through formation of biofilm and can resist disinfection [2]. High touch surfaces such as bed, floor, wall, tables and many other surfaces in hospital environment can become reservoirs of bacterial pathogens [3,4]. The

**Corresponding Author:-Mounika S**

Address:-PG Student, Department of Microbiology, Apollo Institute of Medical Sciences and Research, Hyderabad.

most common organisms associated with health care associated infections are both gram positive and gram-negative bacteria such as *S. aureus*, *CoNS*, *Pseudomonas*, *Proteus species* [5]. This study is planned to assess the distribution of bacterial contamination of frequently touched surfaces shared by healthcare workers, patients and visitors. Identification of these sites and bacterial pathogens help to reduce transmission of pathogens.

**MATERIALS AND METHODS**

A hospital based Cross-sectional study was conducted in Apollo General Hospital from April to May 2024. A total of 48 swabs collected from surfaces of floor, bed, wall, doorhandles, stethoscope, thermometer, IV stand, examination table, sink tap, bed side trolley and drug trolley from Medicine, OBG, Surgery and Paediatric wards using sterile moist swabs. Gram staining and culture on blood agar and MacConkey agar was done. Identification of the isolates was done as per standard protocol. Antimicrobial susceptibility testing was done as per CLSI guidelines.

**RESULTS AND DISCUSSION**

Out of the 48 samples processed, 52% showed bacterial growth. *CoNS* (68%) was the predominant isolate followed by *Staphylococcus* (16%), *Acinetobacter spp* (4%) and *Proteus* (4%). The other isolates were *Enterococcus* (4%) and *Burkholderia* (4%). Majority of the *CoNS* were isolated from floor, bed and wall. Most of the isolates were sensitive to ciprofloxacin and gentamicin and resistant to penicillin

Chart 1: Profile of bacterial isolates identified:





Chart 2: Percentage of organisms isolated:



In the present study out of the total isolates identified, the predominant isolate was *CoNS* (68%) followed by *Staphylococcus* (16%), *Acinetobacter spp* (4%) and *Proteus spp* (4%). The other isolates were *Enterococcus* (4%) and *Burkholderia* (4%). This was similar to a study conducted by Atsedewoyn et al [3]. This suggests that *CoNS* and *S. aureus* are the most common pathogens on frequently touched areas. The higher prevalence of these organisms as they are present as part of normal flora in human body.

Areas from which organisms isolated	Our Study	Teshale et al [5]
Floor	CoNS 50% <i>S. aureus</i> 16.6% <i>Acinetobacter</i> 16.6% <i>Burkholderia</i> 16.6%	CoNS 26.3% <i>S. aureus</i> 42.1% <i>Proteus</i> 5% <i>Serratia</i> 5%
Wall	CoNS 75% <i>S. aureus</i> 25%	<i>E. coli</i> 27.2% <i>Klebsiella</i> 27.2%
Bed	CoNS 75% <i>S. aureus</i> 25%	CoNS 33.3% <i>S. aureus</i> 33.3%
Bedside trolley	CoNS 100%	CoNS 7% <i>S. aureus</i> 7%
Drug trolley	<i>S. aureus</i> 100%	
Examination table	CoNS 100%	
Door handle	CoNS 100%	CoNS 15.3% <i>S. aureus</i> 30.7%
Stethoscope	CoNS 100%	CoNS 50%
Thermometer	CoNS 100%	CoNS 5.2% <i>S. aureus</i> 16.6%
Light switches	CoNS 100%	
Sink tap	<i>Enterococcus</i> 100% <i>Proteus</i> 100%	

In our study most contaminated surfaces were floor and bed. The other surfaces were wall, bedside trolley, drug trolley, examination table, stethoscope, light switches, thermometer, sink tap have also presented with bacterial contamination. This suggests that these surfaces are commonly used by health care staff, patients and visitors and secondly it is related that these surfaces are not adequately cleaned or disinfected

Isolates	CoNS		<i>S.aureus</i>		<i>Enterococcus</i>		<i>Acinetobacter</i>		<i>Proteus</i>		<i>Burkholderia</i>	
	Our study	Teshale et al study	Our study	Teshale et al study	Our study	Atsedewoyn et al study	Our Study	Bhatta et al study	Our study	Kalu MU et al study	Our study	Atsedewoyn et al study
Abx												
AMP	-	-	-	-	100%	77.7%	-	-	0	18%	-	-
PEN	23.5%	29.4%	-	-	-	-	-	-	-	-	-	-
CIP	75%	-	75%	-	-	-	0	29.8%	100%	45%	-	-
CTR		-			-	-	100%	68.7%	0	27%	-	-
GEN	64.7	76.4	75%	-	100%	66.7%	0	28.5%	100%	54%	-	-
COT	52.9%	47.6%	50%	53.7%	-	-	100%	49.8%	100%	100%	0	0
ERY	35.25	28.9%	33.35%	49.3%	-	-	-	-	-	-	-	-
CD	64.7	73.8	50%	69.5%	-	-	-	-	-	-	-	-
VAN	76.4	-	75%	-	100%	-	-	-	-	-	-	-

Abx- Antibiotics-AMP-Ampicillin, PEN-Penicillin, CIP-Ciprofloxacin, CTR-Ceftriaxone, GEN-Gentamicin, COT-Cotrimoxazole, ERY-Erythromycin, CD-Clindamicin, VAN-vancomycin

In our study, most of the isolates of *CoNS* showed resistance to Penicillin and Erythromycin. This correlated with the study conducted by Teshale et al [5] also showed resistance to Penicillin, Erythromycin and Amoxicillin. This suggests that *CoNS* isolated from frequently touched objects in hospital are resistant to Penicillin and Macrolides.

The isolates of *S.aureus* showed resistant to Penicillin. In accordance study conducted by Atsedewoyn et al [4] showed resistant to Penicillin. Also showed resistance to Clindamycin, Cefoxitin. Our study showed susceptibility to ciprofloxacin, gentamycin, ceftriaxone.

The isolates of *Acinetobacter* showed resistance to ciprofloxacin, gentamycin. It correlates with the study conducted by Bhatta et al [1] showed resistance to ciprofloxacin and gentamycin

The isolates of *Enterococcus* from Emergency department showed susceptibility to ampicillin and vancomycin, resistant to penicillin which agrees with the findings reported by Atsedewoyn et al showed susceptibility to vancomycin and ampicillin.

The isolates of *Proteus* isolated from sink tap of surgery ward showed resistance to ampicillin, ceftriaxone which correlates with study conducted by Mary Uche Kalu et al[6] showed susceptibility to ampicillin and ceftriaxone.

*Burkholderia* isolated from floor of labour room showed resistance to cotrimoxazole which agrees with the findings of study conducted by Atsedewoyn et al also showed resistance to cotrimoxazole.

All these studies have shown the presence of pathogenic bacteria on high touch surface areas of the hospital, which can cause serious infections for patients, health care and visitors.

## CONCLUSION

In our study *CoNS* and *S.aureus* are the predominant isolates on frequently touched surfaces. This study emphasizes on infection control practices like regular hand hygiene, cleaning and disinfection of the hospital environment

## REFERENCES

1. Dharm Raj Bhatta, et al. Bacterial contamination of frequently touched objects in a tertiary care hospital of Pokhara, Nepal: how safe are our hands? *Antimicrobial Resistance and Infection Control* (2018) 7:97
2. Alwari SK, Assi MS, et al. Medical devices, Tools and Equipment Surfaces Contamination in Three Departments at a Tertiary Hospital in Baghdad. *Annals of Tropical Medicine & Public Health*. 2020; (S11): SP231130. DOI: <http://doi.org/10.36295/ASRO.2020.231130>.
3. Atsedewoyn, et al. Bacterial profile of high-touch surfaces, leftover drugs and antiseptics together with antimicrobial susceptibility patterns at University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia. *BMC Microbiology* (2021) 21: 309. 1-13
4. Erick Odoyo et al. Environmental contamination across multiple hospital departments with multidrug-resistant bacteria pose an elevated risk of healthcare-associated infections in Kenyan hospitals. *Antimicrobial Resistance & Infection Control* (2023) 12:22. 1-9
5. Teshaleworku, et al. Bacterial Profile and Antimicrobial Susceptibility Pattern of the Isolates from Stethoscope, Thermometer, and Inanimate Surfaces of Mizan-Tepi University Teaching Hospital, Southwest Ethiopia. *International Journal Of Microbiology* 2018: (1-7)
6. Kalu MU, et al. Bacterial Contamination of Labor Wards and Delivery Rooms from Selected Primary Healthcare Facilities in Abia State. *Suan Science and Technology Journal*. 2023.
7. Sebre S, et al. Bacterial Profiles and Antimicrobial Susceptibility Pattern of Isolates from Inanimate Hospital Environments at Tikur Anbessa Specialized Teaching Hospital, Addis Ababa, Ethiopia. *Infectious and Drug Resistance* 2020;13: 4439-4448.
8. Elouakfaoui Aziz, et al. Microbiological Study of Surfaces in the Hospital Environment Case of the Provincial Hospital of Sidi Kacem, Morocco. *Indian Journal of Forensic Medicine & Toxicology*, January- March 2022, Vol 16, No.1: 419-426