



RESEARCH ARTICLE

BEYOND FIELD SUCCESS: INSIGHTS INTO FUNCTIONAL AND NUMERICAL RESPONSE OF THE PREDATORY MITE *TYPHLODROMALUS ARIPO* AND THE ROLE OF CASSAVA EXUDATE IN PREDATORY PERFORMANCE

Nazer Famah Sourassou^{1*}, Rachid Hanna^{2*}, Komi Agboka^{3*}, Koffi Negloh^{4*} and Atti Tchabi^{1*}

1. Centre d'Excellence en Protection Durable des Cultures, Laboratoire des Sciences Agronomiques et Biologiques Appliquées, Institut Supérieur des Métiers de l'Agriculture, Université de Kara, B.P. 404 Kara, Togo.
 2. Center for Tropical Research, University of California, 1300 Franz Hall, Box 951496, Los Angeles, CA 90095-1496, United States.
 3. Laboratoire de Recherche sur les Agroressources et la Santé Environnementale, Ecole Supérieure d'Agronomie, Université de Lomé; BP. 1515 Lomé, Togo.
 4. Institut Togolais de Recherche Agronomique, Kara, B.P. 169, Kara, Togo.
- *Formerly International Institute of Tropical Agriculture, 08, BP 0932, Cotonou, Benin, West Africa

Manuscript Info

Manuscript History

Received: 13 June 2025

Final Accepted: 15 July 2025

Published: August 2025

Key words:-

Predation, oviposition, efficiency of conversion, prey stage, biological control, cassava

Abstract

The predatory mite *Typhlodromalus aripo* is a key natural enemy of cassava green mite *Mononychellus tanajoa* on cassava in Sub-Saharan Africa, yet its functional and numerical responses remain poorly understood. We evaluated, under controlled laboratory conditions (25 °C, 70–90% RH, 12L:12D photoperiod), the functional and numerical responses of *T. aripo* to four immature stages of its prey *M. tanajoa* (eggs, larvae, protonymphs, deutonymphs) offered at four densities (5, 15, 30, and 60 individuals), in the presence or absence of cassava exudate. The predator *T. aripo* exhibited a Type II functional response across all prey stages, with highest attack rates on larvae and protonymphs, and shortest handling times on eggs and larvae. Predation rates were consistently higher on early prey stages, but cassava exudate reduced attack rates and increased handling times, except on larvae where both parameters increased. Oviposition and efficiency of conversion of ingested food (ECI) rose with prey stage, peaking on deutonymphs, and exudate enhanced reproduction and ECI particularly on early stages. These results demonstrate the effectiveness of *T. aripo* as biological control agent against cassava green mite, capable of exploiting a range of stages. Cassava exudate, while influencing predatory behavior, can enhance reproductive performance, thereby demonstrating the predator's long-term persistence in cassava fields. The functional and numerical response data presented here provide valuable reference information for selecting and developing new predatory mite species for use as biological control agents in similar agroecological contexts.

"© 2025 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Corresponding Author:-Nazer Famah Sourassou

Address:-Centre d'Excellence en Protection Durable des Cultures, Laboratoire des Sciences Agronomiques et Biologiques Appliquées, Institut Supérieur des Métiers de l'Agriculture, Université de Kara, B.P. 404 Kara, Togo.

Introduction

The functional and numerical responses are among the most studied attributes of the predator-prey relationships in the context of biological control of arthropod pests (i.e., Dalir et al., 2020; Fathipour et al., 2017). Functional response describes how the rate of prey consumption by a predator changes with prey density, while numerical response refers to changes in predator reproduction based on prey availability. These metrics are fundamental for predicting the impact of a predator on pest populations and are increasingly used as trait-based tools for selecting and comparing candidate biological control agents. Holling (1959, 1961) described three types of functional responses that characterize how a predator's rate of prey consumption changes with prey density. A type I describes a linear increase in prey consumption with increasing prey density, until a maximum is reached (due to predator satiation), a type II describes an accelerating curve – prey consumption increases with prey density but at a decreasing rate due to handling time, and a Type III describing a sigmoidal curve – slow increase at low prey densities, rapid increase at intermediate densities, and saturation at high densities.

Predatory mites of the family Phytoseiidae have been by far the most extensively studied around the world, including Africa, because of their potential as biological control agents of important pests such as spider mites, thrips, and whiteflies in agricultural crops (Gerson et al. 2003; McMurtry et al. 2013). They have been used in biological control programs with two success stories; the classical biological control of the citrus and avocado mites by *Euseius stipulatus* (Athias-Henriot) in California and the classical biological control of the cassava green mite by *Typhlodromalus aripo* (DeLeon) in Africa (Yaninek and Hanna, 2003; McMurtry et al., 2015). In the latter case – biological control of the cassava green mite *Mononychellus tanajoa* (Bondar) (Acari: Tetranychidae) in Africa – several species of predatory mites were introduced from Brazil, origin of the pest, for release in Africa (Yaninek and Hanna, 2003).

The predator *T. aripo* was the most successfully established predator (along with followed by *T. manihoti*) due to its high dispersal rate and its ability to persist in cassava fields, hence effectively controlling the pest *M. Tanajoa* (Yaninek and Hanna, 2003). A series of ecological studies were conducted prior and post-release of *T. aripo* in cassava fields for a better understanding of the host-predator and predator-prey interactions, as well as predator-predator interactions (i.e., Gnanvossou et al., 2003; Onzo et al., 2003; Hanna and Onzo, 2009; Onzo et al., 2012; Onzo et al., 2013; Onzo et al., 2014;). However, despite the several ecological studies, a good understanding of the functional response of *T. aripo* is still missing. Therefore, it is essential to provide data on the functional response of this effective biocontrol agent, which can be important when developing a new species for use as a biological control agent in a similar context.

Moreover, functional response is not static, but rather influenced by biotic factors, including the predator's age (Fathipour et al., 2017; Dalir et al., 2020), the prey's developmental stage (Dalir et al., 2020), and the availability of alternative non-prey food sources (Fathipour et al., 2020). The availability of alternative plant-based foods – such as pollen, plant exudates – can modulate predator predation and enhance the biological control of target pest (McMurtry and Scriven, 1966; Nomikou et al., 2010; Konstantinos Samaras, 2021). In the context of cassava agroecosystems, cassava plants produce large quantities of exudates, which have been identified as important alternative food sources for predatory mites, allowing their survival during periods of prey scarcity (Klein, 1990; Bakker and Klein, 1990; Toko et al., 1994; Gnanvossou et al., 2005). These findings suggest that cassava exudate may play a critical role in modulating the predator–prey dynamics between *T. aripo* and *M. tanajo*. Therefore, evaluating the influence of prey stage and cassava exudate on the functional and numerical responses of *T. aripo* provides a better insight into its effectiveness as a robust and adaptable biological control agent in sub-Saharan Africa cassava agroecosystems. This study was conducted to provide a comprehensive assessment of the functional and numerical responses of *T. aripo* to its prey, *M. tanajoa*, under controlled laboratory conditions, and evaluate the influence of additional cassava foliar exudate on predatory performance.

Material and methods

Mite sources and rearing

The predators (*T. aripo*) used in this study were obtained from a laboratory colony originally established from individuals collected in July 1998 from cassava fields near Sè, Cotonou, Benin (Gnanvossou et al. (2003). The colony has been maintained under controlled environmental conditions: $25 \pm 1^\circ\text{C}$, 70–90% relative humidity, and a 12:12 h light–dark photoperiod. The rearing method followed the procedure described by McMurtry and Scriven (1965) and Friese et al. (1987), with minor adaptations. Predators were maintained on black PVC tiles (4 x 4 cm) placed atop a 1 cm-thick foam pad of the same dimensions inside a 14.5 cm diameter Petri dish filled with water. To

prevent mite escape and contamination, absorbent paper bands were arranged along the edges of the tile, with one edge immersed in water to serve as a continuous moisture source. An oviposition site was prepared using a square of hydrophobic cotton wool, lightly stretched to form an open central platform, and covered with a transparent plastic sheet of matching shape. Fresh cassava leaves infested with all developmental stages of *M. tanajoa* were provided as a food source. These leaves were collected from greenhouse-grown cassava plants reared in pots. For the experiments, specific developmental stages of *M. tanajoa* were obtained from colonies maintained in a screenhouse on 2- to 3-week-old potted cassava plants. These colonies were initiated with cohorts of eggs laid by *M. tanajoa* females collected from cassava fields in southwestern Benin.

Functional response bioassay

To initiate the bioassay, a cohort of 1-day-old *T. aripo* eggs was obtained by placing 100 mated females from the mother colony onto 2.5 cm diameter black PVC disks and allowing them to oviposit for 24 hours. The eggs laid were then carefully transferred to new rearing units and reared to adulthood, following the procedure described by Mégevand et al. (1993). Upon emergence, adult mites were sexed, and females were isolated individually for experimentation. Each experimental unit consisted of a 2.5 cm diameter cassava leaf disc, placed on a layer of solidified agar-agar (1% solution) at the bottom of a Petri dish of similar dimensions. To allow adequate ventilation, two 2 mm-diameter holes were made on opposite sides of each dish and covered with fine nylon mesh.

All experiments were conducted under controlled conditions: $25 \pm 1^\circ\text{C}$, 70–90% relative humidity, and a 12:12 h light:dark photoperiod. To assess the functional and numerical responses of *T. aripo* to different prey stages, newly emerged adult females were transferred singly to the prepared experimental units and offered one of the four immature stages (egg, larvae, protonymph, and deutonymph) of *M. tanajoa* at densities of 5, 15, 30, and 60 individuals per unit. Only immature stages of *M. tanajoa* were used, as these are typically the stages most targeted by phytoseiid mites (Reis et al., 2000; Forero et al., 2008; Marafeli et al., 2011). Each prey stage and density combination was tested in both the presence and absence of cassava foliar exudate, which was provided ad libitum within each unit. A total of 20 replicates (females) were tested per prey stage and density combination. Prey consumption and oviposition was recorded daily over a 7-day period, with both prey and cassava exudate replenished every 24 hours.

Data analysis

Data on functional response were analyzed in two steps (Juliano, 2001). First, a logistic regression model of the proportion of prey consumed (N_e/N_0) as a function of prey density (N_0) was used to determine the type of the functional response of the predator to individual immature stage (equation 1):

$$N_e / N_0 = \frac{\exp(P_0 + P_1 N_0 + P_2 N_0^2 + P_3 N_0^3)}{1 + \exp(P_0 + P_1 N_0 + P_2 N_0^2 + P_3 N_0^3)} \quad (\text{Equation 1})$$

where N_0 is the initial density of prey offered, N_e is the number eaten, P_0 , P_1 , P_2 , and P_3 are the intercept, linear, quadratic, and cubic coefficients, respectively, estimated using the maximum likelihood method. The type of the functional response curve is determined based on the sign of P_1 and P_2 from equation 1 (Juliano, 2001): if $P_1 < 0$, it implies Type II functional response. If $P_1 > 0$ and $P_2 < 0$, it is Type III functional response. After the shape of functional response was determined (Type II), the next step was to estimate the handling time (T_h) and the attack rate (a) of parameters using the random predator equation (equation 2; Roger, 1972):

$$N_e = N_0 \{1 - \exp[a(T_h N_e - T)]\} \quad (\text{Equation 2})$$

where T is the total time available for the predator for search during the experiment (24 h), a is the attack rate (searching efficiency), T_h is the handling time of prey by the predator, and N_e and N_0 as described above. A nonlinear least-squares regression (PROC NLIN, SAS Institute, 2004) was used to fit the random predator equation to data and to estimate the parameters describing the type II response.

Numerical response

To assess the numerical response of *T. aripo*, the number of eggs laid by the predator was recorded concurrently with the number of prey individuals consumed, using the same experimental units over a 7-day observation period. The predator's daily oviposition rate was then regressed against the initial prey densities, following the approach of Fathipour and Maleknia (2016). This regression analysis quantified the relationship between prey density and predator fecundity, offering insights into the predator's reproductive potential in response to increasing prey availability.

The efficiency of conversion of ingested food (ECI) into egg biomass was calculated using the equation of Omkar and Pervez (2004) (Equation 3):

$$ECI = \frac{\text{Number of eggs laid}}{\text{Number of prey consumed}} \times 100 \quad (\text{Equation 3})$$

To evaluate the effects of prey density, prey developmental stage, and the presence of cassava exudate on predation, oviposition, and the efficiency of conversion of ingested food (ECI), data were analyzed using analysis of variance (ANOVA), and Tukey's Honestly Significant Difference (HSD) test was used for multiple comparisons at the 5% significance level ($\alpha = 0.05$).

Results

Functional response

Analysis of variance revealed that prey developmental stage ($F_{3,1213} = 515.5$, $P < 0.0001$), prey density ($F_{3,5665} = 2406.9$, $P < 0.0001$), and the presence of additional cassava exudate ($F_{1,820.9} = 348.7$, $P < 0.0001$) had significant effects on the predation activity of *T. aripo* (Table 1, Table 2). Across all developmental stages, predation rates increased with prey density. However, *T. aripo* consistently showed higher predation on earlier stages (eggs and larvae) compared to later ones (protonymphs and deutonymphs). Mean prey consumption ranged from 4.07 to 24.41 on eggs, 5.00 to 26.29 on larvae, 4.95 to 15.12 on protonymphs, and 4.57 to 14.15 on deutonymphs.

In general, the presence of cassava exudate significantly reduced predation across all prey stages and densities, although some exceptions were observed. The largest reduction (59%) was recorded on egg predation at a density of 15, while the smallest reduction (1.06%) occurred on deutonymphs at a density of 60 (Table 1).

The negative and significant linear coefficient from Equation 1 ($P < 0.05$) confirms that *T. aripo* exhibited a Type II functional response across all prey developmental stages (Table 3; Figure 1). The estimated attack rate (a) and handling time (T_h) are summarized in Table 4. The lowest attack rates were recorded on eggs and deutonymphs (0.0910 and 0.0894, respectively), while the highest values were observed on larvae and protonymphs (0.9301 and 0.7951, respectively). In contrast, handling time was about 2-fold longer for predators preying on protonymphs or deutonymphs than for those feeding on eggs or larvae (Table 1).

The presence of exudate markedly influenced *T. aripo*'s predatory performance. On eggs, the attack rate decreased by six-fold and the handling time increased four-fold. On deutonymphs, the attack rate was reduced by 1.4-fold, accompanied by an approximately one-fold increase in handling time. For protonymphs, exudate reduced the attack rate by 1.4-fold and increased the handling time by 1.1-fold. In contrast, when feeding on larvae, both parameters increased in the presence of exudate, with attack rate rising by 1.6-fold and handling time by 1.3-fold.

Numerical response and food conversion efficiency

Analysis revealed that prey developmental stage, prey density, and the presence of cassava exudate all had significant effects on the oviposition of *T. aripo*, as did most of their interactions, except for the interaction between exudate and prey density, which was not significant (Tables 1 & 2).

Oviposition increased progressively with prey developmental stage, with the lowest number of eggs laid on the egg stage (ranging from 0 eggs/female at density 5 to 0.69 eggs/female at density 60), and the highest on deutonymphs (from 0.62 eggs/female at density 5 to 1.71 eggs/female at density 30). With respect to prey density, oviposition exhibited a curvilinear pattern on early stages (eggs and larvae), increasing with density (Figure 2). On later stages (protonymphs and deutonymphs), oviposition peaked at intermediate density (30) and declined at the highest density

Table 1: Mean (\pm SE) daily consumption, oviposition, and efficiency of conversion onto egg biomass of the predatory mite *Typhlodromalus aripo* at different densities of *Mononychellus tanajoa* immature stages in the presence or absence of additional cassava foliar exudate

Density	Prey Stage	Mean consumption / day		Mean oviposition / day		Mean ECI (%)	
		Exudate absent	Exudate present	Exudate absent	Exudate present	Exudate absent	Exudate present
5	Egg	4.07 \pm 0.17 cA	3.67 \pm 0.16 cA	0 dA	0.08 \pm 0.02 bB	0 cB	2.34 \pm 0.60 bA
	Larva	5.00 \pm 0.00 aA	4.81 \pm 0.05 aA	0.05 \pm 0.01 cB	0.63 \pm 0.05 bA	1.05 \pm 0.39 cB	13.10 \pm 1.01 aA
	Protonymph	4.95 \pm 0.02 aA	4.80 \pm 0.03 aB	0.30 \pm 0.03 bB	0.48 \pm 0.05 bA	6.20 \pm 0.62 bB	10.10 \pm 0.93 aA
	Deutonymph	4.57 \pm 0.11 bA	4.12 \pm 0.09 bB	0.62 \pm 0.05 aA	0.62 \pm 0.08 aA	13.82 \pm 1.32 aA	14.86 \pm 2.30 aA
15	Egg	9.46 \pm 0.59 bA	6.87 \pm 0.41 cB	0.02 \pm 0.01 dA	0.16 \pm 0.04 cB	0.14 \pm 0.09 dB	2.03 \pm 0.57 dA
	Larva	15.00 \pm 0.00 aA	13.64 \pm 0.15 aB	0.39 \pm 0.04 cB	0.83 \pm 0.04 bA	2.59 \pm 0.30 cB	6.07 \pm 0.31 cA
	Protonymph	14.19 \pm 0.12 aA	12.30 \pm 0.16 bB	0.82 \pm 0.06 bB	1.09 \pm 0.04 aA	5.78 \pm 0.44 bB	8.91 \pm 0.36 bA
	Deutonymph	8.50 \pm 0.25 bA	7.46 \pm 0.19 cB	1.18 \pm 0.06 aA	1.11 \pm 0.08 aA	14.08 \pm 0.65 aA	14.78 \pm 1.04 aA
30	Egg	20.91 \pm 0.83 bA	8.80 \pm 0.37 dB	0.23 \pm 0.03 dA	0.39 \pm 0.06 cA	1.12 \pm 0.17 dB	4.25 \pm 0.74 dA
	Larva	24.66 \pm 0.51 aA	22.10 \pm 0.30 aB	1.13 \pm 0.07 cB	1.61 \pm 0.05 bA	4.61 \pm 0.26 cB	7.36 \pm 0.24 cA
	Protonymph	14.70 \pm 0.43 cA	12.50 \pm 0.36 bB	1.39 \pm 0.07 bB	1.63 \pm 0.04 bA	9.48 \pm 0.39 bB	13.27 \pm 0.67 bA
	Deutonymph	13.35 \pm 0.50 cA	10.75 \pm 0.24 cB	1.76 \pm 0.05 aA	1.86 \pm 0.06 aA	13.39 \pm 0.57 aB	17.52 \pm 0.60 aA
60	Egg	24.41 \pm 0.98 bA	21.14 \pm 0.35 aB	0.69 \pm 0.09 cB	1.18 \pm 0.08 bA	2.79 \pm 0.42 dB	5.62 \pm 0.37 cA
	Larva	26.29 \pm 0.42 aA	20.08 \pm 0.50 aB	1.71 \pm 0.46 bB	1.55 \pm 0.04 aA	4.60 \pm 0.25 cB	7.81 \pm 0.27 bA
	Protonymph	15.12 \pm 0.43 cA	13.85 \pm 0.41 bA	1.36 \pm 0.08 abB	1.71 \pm 0.05 aA	8.96 \pm 0.47 bB	12.55 \pm 0.50 aA
	Deutonymph	14.15 \pm 0.33 cA	14.30 \pm 0.34 bA	1.56 \pm 0.05 aA	1.61 \pm 0.06 aA	11.15 \pm 0.47 aA	11.36 \pm 0.47 aA

Means followed by different letters are statistically different (Turkey HSD test, $P < 0.05$); small letters and capital letters signify differences among prey stages within each density and between exudate treatments for each prey stage, respectively

Table 2: Analysis of variance of the effects of prey stage, prey density and additional exudate on the predation, oviposition and efficiency of conversion of *Typhlodromalus aripo* in functional response experiment under laboratory conditions

Source	df	Predation			Oviposition rate			Efficiency of conversion		
		MS	F value	P value	MS	F value	P value	MS	F value	P value
Exudate	1	479	203.03	<0.0001	7.35	46.14	<0.0001	0.1387	137.52	<0.0001
Prey stage	3	1178	499.41	<0.0001	23.16	145.32	<0.0001	34.78	344.93	<0.0001
Prey density	3	5615	2380.35	<0.0001	37.03	232.35	<0.0001	0.0122	12.11	<0.0001
Exudate x Prey density	3	117	49.69	<0.0001	0.03	0.184	0.9070	0.0066	6.52	0.0002
Exudate x Prey stage	3	70	29.59	<0.0001	0.68	4.26	0.0054	0.076	7.49	0.0052
Prey density x Prey stage	9	356	151.11	<0.0001	1.51	9.46	<0.0001	0.0099	9.84	<0.0001
Exudate x Prey stage x Prey density	9	61	25.72	<0.0001	0.47	2.92	0.0021	0.0061	6.07	0.0022
Residuals	554	2	-	-	0.16	-	-	0.0010	-	-

Table 3: Maximum-likelihood estimates from logistic regression of the proportion of *Mononychellus tanajoa* stages consumed by *Typhlodromalus aripo* as a function of initial prey density in the absence or presence of additional exudate

Prey stage	Parameter	Exudate absent				Exudate present			
		Estimate	SE	Z-value	P-value	Estimate	SE	Z-value	P-value
Egg	P_0	2.5420	0.5739	4.430	<0.0001	1.4222	0.2432	5.847	<0.0001
	P_1	-0.2588	0.0768	-3.370	0.0007	-0.1219	0.0159	-7.661	<0.0001
	P_2	0.0095	0.0028	3.510	0.0004	0.0014	0.0002	7.135	<0.0001
	P_3	-0.0001	0.00002	-3.748	0.0002	-	-	-	-
Larva	P_0	7.6932	2.6443	2.909	0.0036	3.8913	0.4504	8.639	<0.0001
	P_1	-0.2773	0.1045	-2.652	0.0080	-0.1145	0.0250	-4.577	<0.0001
	P_2	-	-	-	-	0.0006	0.0002	2.132	0.0330
	P_3	-	-	-	-	-	-	-	-
Protonymph	P_0	7.1092	0.6913	10.288	<0.0001	4.2831	0.3702	11.567	<0.0001
	P_1	-0.3395	0.0364	-9.314	<0.0001	-0.2163	0.0211	-10.231	<0.0001
	P_2	0.0033	0.00004	8.030	<0.0001	0.0021	0.0002	8.099	<0.0001
	P_3	-	-	-	-	-	-	-	-
Deutonymph	P_0	4.2350	0.7564	5.599	<0.0001	2.8650	0.5688	5.036	<0.0001
	P_1	-0.4378	0.0951	-4.600	<0.0001	-0.3035	0.0755	-4.016	<0.0001
	P_2	0.0134	0.0033	3.999	<0.0001	0.0086	0.0027	3.114	0.0018
	P_3	-0.0001	0.00003	-3.842	0.0002	-0.00007	0.00002	-2.80	0.0051

Table 4: Estimates from logistic regression of the proportion of *Mononychellus tanajoa* stages consumed by *Typhlodromalus aripo* as a function of initial prey density in the absence or presence of cassava foliar exudate

Prey stage	Parameter	Exudate absent			Exudate present		
		Estimate±SE	T/T _h	R ²	Estimate ± SE	T/Th	R ²
Egg	a (h ⁻¹)	0.0910 ± 0.0142 (0.0626–0.1194)			0.0150 ± 0.0017 (0.0154–0.0225)		
	T_h (h)	0.7129 ± 0.0584 (0.5961–0.8296)	33.6	0.85	2.7881 ± 0.1188 (2.5512–3.0250)	8.6	0.96
Larva	a (h ⁻¹)	0.7951 ± 0.2417 (0.3128–1.2773)			1.2762 ± 0.4920 (0.2966–2.2556)		
	T_h (h)	0.8852 ± 0.0174 (0.8505–0.9199)	27.1	0.86	1.1179 ± 0.0212 (1.0758–1.1601)	24.4	0.84
Protonymph	a (h ⁻¹)	0.9301 ± 0.3361 (0.2596–1.6005)			0.6353 ± 0.2323 (0.1727–1.0979)		
	T_h (h)	1.5579 ± 0.0320 (1.4940–1.6218)	15.4	0.72	1.7627 ± 0.0400 (1.6832–1.8423)	13.6	0.77
Deutonymph	a (h ⁻¹)	0.0894 ± 0.0105 (0.0685–0.1103)			0.0521 ± 0.0040 (0.0440–0.0601)		
	T_h (h)	1.4463 ± 0.0530 (1.3405–1.5521)	16.5	0.79	1.3457 ± 0.0504 (1.2451–1.4463)	17.8	0.83

(60). Regression analysis showed significant and moderate correlations between prey density and oviposition ($P < 0.05$, $0.5 < R^2 < 0.75$), except for the deutonymph stage, where a weaker but significant relationship was found ($R^2 = 0.413$). The efficiency of conversion of ingested food (ECI) by *T. aripo* increased with the prey's developmental stage across all densities. Overall, ECI was lowest for females fed on eggs and highest for those fed on deutonymphs. For early stages (eggs, larvae, and protonymphs), ECI tended to increase with prey density, whereas for deutonymphs, it generally declined as density increased (Table 1). The presence of cassava exudate resulted in a statistically significant but modest increase in oviposition, and a substantial enhancement of ECI, particularly when *T. aripo* fed on eggs, larvae, or protonymphs. The largest increase in oviposition (71%) was observed on eggs at density 60, while the greatest ECI enhancement (63%) occurred on larvae. In contrast, no significant effect of exudate was detected on either oviposition or ECI when predators fed on deutonymphs (Table 1).




Figure 1: Functional response of *Typhlodromalus aripo* to eggs, larvae, protonymphs, and deutonymphs of its prey *Mononychellus tanajoa* in the absence or presence of cassava foliar exudate




Figure 2: Numerical response of *Typhlodromalus aripo* to eggs, larvae, protonymphs, and deutonymphs of its prey *Mononychellus tanajoa*, in the absence or presence of cassava foliar exudate

Discussion

This study addresses a critical knowledge gap regarding the functional and numerical responses of the predatory mite *T. aripo*, a key biological control agent against the cassava green mite in sub-Saharan Africa. Beyond quantifying predation and reproductive performance, our findings underscore the influence of prey developmental stage and plant-associated foods such as cassava exudate, in shaping both the predator's functional and numeric responses. These insights contribute to a deeper understanding of predator-prey interactions and offer valuable implications for predictive evaluation of any predatory mite for use as a biological control agent in sub-Saharan Africa (Yaninek and Hanna, 2003). Overall, our findings underscore the complex interplay between prey stage, prey density, and plant-derived foods in shaping both the functional and numerical responses of *T. aripo*.

Our results demonstrate that the predatory mite *T. aripo* exhibits a Type II functional response when feeding on all immature stages of its prey, consistent with patterns reported for many phytoseiid mites (Piyani et al., 2021; Fathipur et al., 2017; Fathipur et al., 2020; Carrillo and Pena, 2011; Park et al., 2021). This functional response type, characterized by a decelerating prey consumption rate at higher prey densities, suggests that handling time imposes a limit on predation, as confirmed by our parameter estimates. The longer handling times recorded for protonymphs and deutonymphs compared with eggs and larvae likely reflect increased prey size, mobility and defensive behaviors in later developmental stages, which can slow down prey capture and processing (Sabelis, 1981). Moreover, Sabelis (1990) demonstrated that, rather than being constrained solely by time-budget rules, functional response parameters are largely determined by predator satiation, which depends on prey consumption and gut clearance.

Predatory mites are categorized into four lifestyle types, with most generalist predators classified as type III. These predators feed and reproduce on a wide variety of prey, but they can also thrive on alternative food sources such as pollen, plant exudates, and honeydew – either as survival food in the absence of prey or as a complementary food source that boosts reproduction when prey is present (McMurtry et al., 2013). Previous studies have shown that type III generalists may prefer *T. urticae* larvae over eggs, or exhibit no clear preference between the two (Blackwood et al., 2001). For instance, *Amblyseius eharai*, a type III generalist, was reported to prefer *T. urticae* larvae over other stages (Park et al., 2021). Similarly, *Neoseiulus californicus* consumed more larvae, whereas *N. cucumeris* consumed more eggs (Zhang et al., 2017).

Similarly, in our study, *T. aripo*, also classified a type III generalist predator (McMurtry et al., 2013), displayed higher predation rates on early developmental stages (eggs and larvae) than on later stages (protonymphs and deutonymphs), with larvae being the most consumed stage. Strikingly, all predator females (100%) consumed all available larvae at prey densities of 5 and 15, and 82% at higher densities, demonstrating a strong preference for the larval stage of its prey. Eggs and larvae are immobile and have softer integuments, facilitating capture and consumption, whereas protonymphs and deutonymphs are more mobile, better defended, and require longer handling times. The approximately two-fold longer handling times recorded on protonymphs and deutonymphs compared to eggs and larvae support this interpretation, and are consistent with earlier observations for other phytoseiids preying on spider mite nymphs or adults (Sabelis, 1986).

Numerical response patterns further revealed that *T. aripo*'s reproductive output is strongly influenced by prey stage. The highest oviposition rates on deutonymphs, despite their lower consumption rates compared with early stages, suggest that this prey stage provides the greatest nutritional benefit per individual consumed. This is supported by the observed trends in efficiency of conversion of ingested food (ECI), already peaking on deutonymphs at the lowest density. Similar stage-specific effects on predator fecundity have been reported for other phytoseiids, where later prey stages supply more biomass and higher-quality nutrients, promoting egg production (Fathipour and Maleknia, 2016; Fathipour et al., 2022).

The presence of cassava exudate had stage-specific effects on the predatory performance of *T. aripo*. On eggs, for instance, it reduced the attack rate six-fold and increased handling time four-fold. Similar patterns have been reported for *Amblyseius swirskii* feeding on spider mite eggs, where the addition of pollen reduced the attack rate 13-fold and increased handling time 1.4-fold (Fathipour et al., 2017). Moreover, while exudate generally reduced predation rates, it modestly increased oviposition and substantially enhanced ECI for early prey stages, consistent with earlier observations for other phytoseiids (Nomikou et al., 2010; Fathipour et al., 2020). This result might indicate that in the presence of exudate, *T. aripo* invests more efficiently in reproduction per prey consumed, by allocating energy toward egg production rather than increased foraging activity. For deutonymphs, however, exudate had no significant effect on either oviposition or ECI, suggesting that the large size and nutritional content of this

later developmental stage buffer the predator's reproductive performance against foraging interference from exudate.

Conclusion

The predatory mite *T. aripo* exhibits a type II functional response against all immature stages of its prey *M. tanajoa*, demonstrating strong potential as an effective biological control agent against this pest, given its high attack rates, preference for vulnerable prey stages, and capacity to convert prey consumption into reproductive output. While cassava exudate can reduce predation rate in certain contexts, it also enhances oviposition and food conversion efficiency on early prey stages, suggesting it plays an important role in sustaining predator populations in period of prey scarcity. This dual effect highlights the importance of integrating plant traits, such as exudate production, into biological control strategies to optimize predator performance and ensure long-term suppression of pest populations in cassava systems.

Acknowledgments

We thank R. Houndafoche, P. Sovimi, for their assistance in maintaining the predator colonies and conducting experiments. This research was supported by the International Institute of Tropical Agriculture (IITA).

References

- Bakker, F., Klein, M. (1990). The significance of cassava exudate for predacious mites. Symp. Biol. Hung. 39: 437-439.
- Blackwood, J.S., Schausberger, P., Croft. B.A. (2001). Prey-stage preference in generalist and specialist phytoseiid mites (Acari: Phytoseiidae) when offered *Tetranychus urticae* (Acari: Tetranychidae) eggs and larvae. Environmental Entomology 30: 1103–1111.
- Carrillo, D., Pena, J.A. (2011). Prey-stage preferences and functional and numerical responses of *Amblyseius largoensis* (Acari: Phytoseiidae) to *Raoiella indica* (Acari: Tenuipalpidae). Experimental and Applied Acarology 55(3-4): 361-372, DOI 10.1007/s10493-011-9488-7
- Cohen, J. (1988). Statistical power analysis for the behavioral sciences (2nd Ed.). New York: Routledge.
- Dalir, S., Hajiqanbar, H., Fathipour, Y., Khanamani, M. (2020). Age-Dependent Functional and Numerical Responses of *Neoseiulus cucumeris* (Acari: Phytoseiidae) on Two-Spotted Spider Mite (Acari: Tetranychidae). Journal of Economic Entomology, 114(1), 50–61. <https://doi.org/10.1093/jee/toaa266>
- Ding-Xu, L., Juan, T., Zuo-Rui, S. (2007). Functional response of the predator *Scolothrips takahashii* to hawthorn spider mite, *Tetranychus viennensis*: effect of age and temperature. BioControl 52: 41–61.
- Fan, Y., Petitt. F. L. (1994). Functional response of *Neoseiulus barkeri* Hughes on two-spotted spider mite (Acari: Tetranychidae). Experimental and Applied Acarology 18: 613–621.
- Farazmand, A., Fathipour, Y., Kamali, K. (2012). Functional response and mutual interference of *Neoseiulus californicus* and *Typhlodromus bagdasarjani* (Acari: Phytoseiidae) on *Tetranychus urticae* (Acari: Tetranychidae). International Journal of Acarology 38: 369–376.
- Fathipour, Y., Maleknia, B., Soufbaf, M., Reddy, G.V.P. (2020). Functional and numerical responses, mutual interference and resource switching of *Amblyseius swirskii* on two-spotted spider mite. Biological Control 146: 104266.
- Fathipour, Y., Karimi, M., Farazmand, A., Talebi, A.A. (2017). Age-specific functional response and predation rate of *Amblyseius swirskii* (Phytoseiidae) on two-spotted spider mite. Systematic and Applied Acarology 22(2), 159–169. <https://doi.org/10.11158/saa.22.2.1>
- Fathipour, Y., Maleknia, B. (2016). Mite Predators. In: Omkar (Ed.), Ecofriendly Pest Management for Food Security. Elsevier, San Diego, USA, pp. 329-366
- Forero, G., Rodríguez, M., Cantor, F., Rodríguez, D., Cure, J.R. (2008). Critérios para el manejo de *Tetranychus urticae* Koch (Acari: Tetranychidae) com elácaro de predador *Amblyseius* (*Neoseiulus*) sp. (Acari: Phytoseiidae) encultivos de rosas. Agronomía Colombiana 26:78-86.
- Friese, D.D., Mégevand, B., Yaninek, J.S. (1987). Culture maintenance and mass production of exotic phytoseiids. Insect Science and its Application 8: 875-878
- Gerson, U., Smiley, R.L., Ochoa, R. (2003) Mites (Acari) for pest control. Blackwell Science Ltd., UK, 539 pp.
- Gnanvossou, D., Hanna, R., Toko, M. (2005). Comparative life history traits of three neotropical phytoseiid mites maintained on plant-based diets. Biological control 35(1): 32-39

- Hanna, R., Onzo, A. (2009). Within-plant migration of the predatory mite *Typhlodromalus aripo* from the apex to the leaves of cassava: response to day–night cycle, prey location and prey density. *Journal of Insect behavior* 22: 186-195.
- Gnanvossou, D., Yaninek, S., Hanna, R., Dicke, M. (2003). Effects of prey mite species on life history of the phytoseiid predators *Typhlodromalus manihoti* and *Typhlodromalus aripo*. *Experimental and Applied Acarology* 30: 265-278
- Holling, C. S. (1961). Principles of insect predation. *Annual Review of Entomology* 6, 163-182
- Holling, C. S. 1959. Some characteristics of simple types of predation and parasitism. *The Canadian Entomologist* 91: 385–398.
- Jafari, S., Fathipour, Y., Faraji, F. (2012). The influence of temperature on the functional response and prey consumption of *Neoseiulus barkeri* (Phytoseiidae) on two-spotted spider mite. *Journal of Entomological Society of Iran* 31: 39–52
- Juliano, S. A. 2001. Nonlinear curve fitting: predation and functional response curves, pp. 178–196. In S. M. Scheiner and J. Gurevitch (eds.), *Design and analysis of ecological experiments*. Oxford University Press, New York
- Klein, M. (1990). Is cassava exudate a factor in the persistence of *Typhlodromalus limonicus* (Garman & McGregor) in the Weld? M.Sc. Thesis, University of Amsterdam, Amsterdam, The Netherlands, 34 pp.
- Madadi, H., Enkegaard, A., Brodsgaard, H.F., Kharrazi-Pakdel, A., Mohaghegh, J. Ashouri, A. (2007). Host plant effects on the functional response of *Neoseiulus cucumeris* to onion thrips larvae. *Journal of Applied Entomology* 131: 728–733.
- Marafeli, P.P., Reis, P.R., Silveira, E.C., Toledo, M.A., Souza-Pimentel, G.C. (2011). *Neoseiulus californicus* (McGregor, 1954) preying in different Life stages of *Tetranychus urticae* Koch, 1836 (Acari: Phytoseiidae, Tetranychidae). *Acarologia* 51: 499-506.
- McMurtry, J.A., Scriven, G.T. (1966). Effects of pollen and prey density on the number of prey consumed by *Amblyseius hibisci* (Acarina: Phytoseiidae). *Annals of Entomological Society of America* 59: 147-149.
- McMurtry J.A., de Moraes, G.J., Famah Sourassou, N. (2013). Revision of the lifestyles of phytoseiid mites (Acari: Phytoseiidae) and implications for biological control strategies. *Systematic & Applied Acarology* 18(4): 297–320. <http://dx.doi.org/10.11158/saa.18.4.1>
- Megevand, B., Klay, A., Gnanvossou, D., Paraiso, G. (1993). Maintenance and mass rearing of phytoseiid predators of the cassava green mite. *Experimental and Applied Acarology* 17: 115-128.
- Nikbin, R., Sahrsgard, A., Hosseini. M. (2014). Age-specific functional response of *Trichogramma brassicae* (Hymenoptera: Trichogrammatidae) parasitizing different egg densities of *Ephestia kuehniella* (Lepidoptera: Pyralidae). *Journal of Agriculture Science and Technology* 16: 1205–1216.
- Nomikou, M., Sabelis, M.W., Arne Janssen, A. (2010). Pollen subsidies promote whitefly control through the numerical response of predatory mites. *BioControl* 55:253–260
- Omkar, O., Pervez, A. (2004). Functional and numerical responses of *Propylea dissecta* (Col., Coccinellidae). *Journal of Applied Entomology* 128: 140–146.
- Onzo A, Sabelis M.W., Hanna R. (2003). Interactions in an acarine predator guild: impact on *Typhlodromalus aripo* abundance and biological control of cassava green mite in Benin, West Africa. *Experimental and Applied Acarology* 31(3-4) 225-231
- Onzo A., Sabelis, M.W., Hanna, R. (2014) Single versus multiple enemies and the impact on biological control of spider mites in cassava fields in West-Africa. *Experimental and Applied Acarology* 62: 293-311.
- Onzo, A., Hanna, R., Sabelis, M.W. (2012) The predatory mite *Typhlodromalus aripo* prefers green-mite induced plant odours from pubescent cassava varieties. *Experimental and Applied Acarology* 58:359–370
- Onzo, A., Hanna, R., Toko, M. (2013). Mix-planting pubescent and glabrous cassava affects abundance of *Typhlodromalus aripo* and its prey mite *Mononychellustana joa*. *Journal of Applied Entomology* 138: 297-306.
- Park, Y.-G., Lee, J.-H., Lim, U.T. (2011). Functional response of *Amblyseius eharai* (Acari: Phytoseiidae) on *Tetranychus urticae* (Acari: Tetranychidae). *PLoS ONE* 16(12): e0260861.
- Patel, K., Zhang. Z.Q. (2017). Functional and numerical responses of *Amblydromalus limonicus* and *Neoseiulus cucumeris* to eggs and first instar nymph of tomato/potato psyllid (*Bactericera cockerelli*). *Systematic and Applied Acarology* 22: 1476–1488.
- Reis, P.R., Teodoro, A.V., Pedro Neto, M. (2000). Predatory cctivity of phytoseiid mites on developmental stages of coffee ring spot mite (Acari: Phytoseiidae:Tenuipalpidae). *Neotropical Entomology*29: 547-553.
- Rogers. D. (1972). Random search and insect population models. *Journal of Animal Ecology* 41(2): 369–383.

- Sabelis, M.W. (1986). The functional response of predatory mites to the density of two-spotted spider mites. In: Dynamics of Physiologically Structured Populations Metz JAJ, Diekmann O (eds) Lecture Notes in Biomathematics Springer, Berlin Heidelberg New York pp 298-321
- Sabelis, M.W. (1990). How to analyze prey preference when prey densities varies? A new method to discriminate between effects of gut fullness and prey type composition. *Oecologia* 82: 289-298.
- Samaras, K., Pappas, M.L., Pekas, A. (2022). Benefits of a balanced diet? Mixing prey with pollen is advantageous for the phytoseiid predator *Amblydromalus limonicus*. *Biological control* 155, 104531
- Toko, M., O'Neil, R.J., Yaninek, J.S. (1994). Effects of cassava exudate and prey densities on the survival and the reproduction of *Typhlodromalus limonicus* (Acari: Phytoseidae), a predator of cassava green mite *Mononychellus tanajoa* (Acari: Tetranychidae). *Experimental and Applied Acarology* 18: 221-231
- Yaninek S., Hanna, R. (2003). Cassava green mite in Africa - a unique example of successful classical
- Zheng, Y., De Clercq, P., Song, Z.-W., Li, D., Zhang, B.-X. (2017). Functional response of two *Neoseiulus* species preying on *Tetranychus urticae* Koch. *Systematic & Applied Acarology* 22(7): 1059–1068