



RESEARCH ARTICLE

INCESSANT CRYING: THINK ABOUT HAIR TOURNIQUET SYNDROME

S. Saghir¹, I. Laatfa², F. Charaf², A. Ayad¹, M. Sellouti¹ and R. Abilkassem¹

1. Pediatrics Department, Mohammed V Military Hospital, Rabat, Morocco.

2. Pediatrics Department, Children's Hospital CHIS Rabat, Morocco.

Manuscript Info

Manuscript History

Received: 4 August 2025

Final Accepted: 7 September 2025

Published: October 2025

Key words:-

Hair-thread tourniquet syndrome; pain;
Incessant crying.

Abstract

Tourniquet syndrome, also known as hair-thread tourniquet syndrome, is a rare but potentially serious condition. It is defined as the strangulation of a body part (toe, finger, penis, clitoris, nipple) by a thread-like object, most often a hair or textile thread. This phenomenon causes progressive circulatory obstruction, first venous and lymphatic, then arterial, which can progress to irreversible ischemia and tissue necrosis if diagnosis and treatment are delayed. First described by Quinn in 1971, this syndrome occurs primarily in infants and young children, a period when postpartum maternal hair loss is common and promotes the presence of loose hair in the child's environment. The most commonly reported locations are the toes (particularly the 2nd and 3rd), followed by the fingers and, more rarely, the genitals. In our paper, we describe two cases of this syndrome with an isolated and non-specific clinical presentation that could be the cause of a diagnostic delay.

"© 2025 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

Hair tourniquet syndrome is a rare but potentially serious pediatric emergency. It occurs when a hair or thread wraps around a finger, toe, penis, or other part of the body, causing progressive strangulation. Diagnosis is often delayed because the injury is subtle and masked by edema, which can compromise the viability of the affected segment.

Although rare, tourniquet syndrome is probably widely underdiagnosed because its initial clinical signs can be subtle and confused with other conditions such as local infection, trauma, or inflammatory reaction. In infants, it often manifests as incessant crying, unexplained irritability, or refusal to feed, making a thorough clinical examination essential. Its importance lies in the fact that it is a medical and surgical emergency: rapid recognition and complete removal of the foreign body restore blood flow and prevent irreversible sequelae, including amputation. Therefore, raising awareness among parents and healthcare professionals is essential to improve early diagnosis and prevent complications.

Clinical Observation:

Case 1:

Our clinical case concerns a 2-month-old infant brought in for incessant crying, agitation, and refusal to feed. The initial examination was inconclusive, but upon closer inspection, a hair was found wrapped around a toe, causing

distal edema and painful erythema. Treatment was based on removing the hair, and the outcome was marked by the gradual disappearance of pain and clinical improvement.



Figure 1

Case 2:

A 3-month-old infant with no particular medical history was admitted for incessant crying that could not be comforted. Clinical examination revealed a circular inflammatory groove on the third toe of the right foot, secondary to strangulation by a wrapped hair, suggestive of digital tourniquet syndrome.



Figure 2



Figure 3

In both cases, the evolution was favorable after the removal of the hair fragment without complications during follow-up.

Discussion:-

Tourniquet syndrome is widely documented in the literature through case series and isolated reports, but it remains a little-known condition. According to Quinn (1971), who first described it, the most common locations are the toes, particularly the second and third toes. Several subsequent studies, notably those by Barton and Sloan (1993) and Golshevsky et al. (2005), confirm this predominance in the feet, followed by the fingers and, more rarely, the genitals. In our observation, the location was on the toe, which is consistent with these data.

Clinically, the authors describe initially nonspecific symptoms, dominated by unexplained crying or unusual irritability in infants. This often leads to a delay in diagnosis, with some cases only being identified after several hours or even several days. In our case, the delay in consultation was four hours, which is consistent with that reported in most series.

With regard to management, the literature emphasizes the importance of immediate release. Most of the cases described benefited from simple mechanical extraction, sometimes assisted by a magnifying glass, while complex situations required surgical incision to release the buried hair. Golshevsky et al. also mentioned the use of thioglycolate-based depilatory creams, but with reservations due to the risk to the skin. In our observation, treatment consisted of immediate release by mechanical extraction, which is in line with standard recommendations and led to a favorable outcome.

The literature emphasizes the predominance of toe involvement, but atypical locations exist and should be known to clinicians. Management is based on immediate and complete removal of the foreign body, sometimes requiring surgical incision. The prognosis is generally favorable if the diagnosis is made early, whereas delay exposes the patient to the risk of irreversible ischemia, necrosis, and amputation. Our observation illustrates the importance of increased vigilance in the face of any unusual clinical picture in children. Finally, prevention, through raising awareness among parents and training healthcare professionals, remains the best weapon for reducing the incidence and complications of this syndrome.

Finally, the prognosis reported in the literature depends on the time to diagnosis. Cases treated quickly have a favorable outcome, while delays are responsible for necrosis and amputations, as reported by Pomeranz (2000). In our case, the outcome was favorable, illustrating the importance of early recognition of this syndrome.

Conclusion:-

Tourniquet syndrome is a rare but serious condition characterized by the strangulation of a body part by a hair or thread. Its significance lies in the severity of potential complications, contrasting with the simplicity of treatment when performed early. The clinical presentation is often misleading and can delay diagnosis, especially in infants

where symptoms are nonspecific. Careful inspection of any painful and unexplained swelling therefore remains the key step in quickly identifying this condition.

This cases should be kept in mind as an etiology in infants with incessant crying with normal examination, These babies should be examined undressed to prevent any irreversible complication including appendage loss.

Figures:

Figure 1: Swelling and redness of the toe surrounded by hair fragments

Figure 2: Strangulation of the 4th Toe with hair fragment

Figure 3: Hair fragment after removal

Conflict of Interest: The authors declare that they have no competing interests.

Authorship:

The corresponding author contributed to the diagnosis and management of the patient.

The other authors contributed to the writing and editing of the manuscript..

All authors read and approved the final manuscript.

Acknowledgments:-

Not applicable.

References:-

1. Arkoubi & Salati (2024): comprehensive and updated review of tourniquet syndrome.
2. Belani et al. (2019): case of emergency surgical exploration in an infant.
3. Dunphy et al. (2018): report of a rare case of surgical emergency in an infant.
4. Turkish J Plast Surg (2020): two clinical cases illustrating the diversity of locations.
5. Djokic et al. (2023): recent review highlighting frequent diagnostic delays.
6. Case reports (2023, ScienceDirect): example of surgical management under a magnifying glass.
7. Hebal & Hunter (2016): retrospective study proposing a therapeutic algorithm.
8. Kuiper & de Korte (2015): pediatric case with toe involvement.