



Journal Homepage: - www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI: 10.21474/IJAR01/21913

DOI URL: <http://dx.doi.org/10.21474/IJAR01/21913>



RESEARCH ARTICLE

INTEGRATED CARE APPROACHES FOR DIABETES MANAGEMENT IN PREGNANCY: MULTIDISCIPLINARY STRATEGIES AND OUTCOMES

Faiza Parveen Baloch and Ntsoaki Betty Mopane

1. Primary Health Care Corporation, Al Mina Street, Doha, Qatar.

Manuscript Info

Manuscript History

Received: 07 August 2025

Final Accepted: 09 September 2025

Published: October 2025

Key words:-

Diabetes in pregnancy; gestational diabetes; integrated care; multidisciplinary care; maternal health; telemedicine.

Abstract

Background: Diabetes in pregnancy, including pre-existing Type 1 and Type 2 diabetes as well as gestational diabetes mellitus (GDM), is a major global health challenge. It is associated with increased risks of maternal complications, adverse neonatal outcomes, and long-term metabolic disorders in both mother and child. Fragmented healthcare delivery often limits effective management, underscoring the need for integrated and multidisciplinary approaches.

Objective: This review aimed to synthesize current evidence on integrated care approaches for diabetes management during pregnancy, emphasizing key components, implementation models, outcomes, challenges, and future directions.

Methods: A narrative review of contemporary literature was conducted, focusing on integrated care models and multidisciplinary strategies applied to diabetes in pregnancy. Sources included peer-reviewed articles, clinical guidelines, and systematic reviews published in recent years. Key themes analyzed were components of integrated care, care delivery models, clinical outcomes, cost-effectiveness, patient satisfaction, and barriers to implementation.

Results: Integrated care models incorporating medical management, nutritional therapy, blood glucose monitoring, fetal surveillance, and mental health support demonstrated significant improvements in maternal and neonatal outcomes. Evidence shows reductions in HbA1c levels, hospitalizations, neonatal intensive care admissions, and pregnancy related complications. Multidisciplinary clinics and telehealth solutions improved care coordination, patient engagement, and accessibility. Integrated approaches were also associated with higher patient satisfaction and long-term cost-effectiveness. However, persistent challenges include limited access in underserved populations, health literacy deficits, and fragmented provider communication.

"© 2025 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Conclusion: Integrated care approaches for diabetes in pregnancy provide measurable benefits in clinical outcomes, cost-effectiveness, and patient satisfaction. Overcoming barriers related to access, health literacy, and inter-provider

Corresponding Author:- Faiza Parveen Baloch

Address:- Primary Health Care Corporation, Al Mina Street, Doha, Qatar.

coordination is essential for scaling these models. Future strategies should focus on personalized multidisciplinary care, technology-driven solutions, preventive interventions, and supportive policy frameworks to build a sustainable, patient-centered healthcare ecosystem for pregnant individuals with diabetes.

Introduction:-

Diabetes during pregnancy, encompassing pre-existing Type 1, Type 2, and gestational diabetes mellitus (GDM), represents a substantial global health concern, impacting 5–20% of pregnancies worldwide (Scavini & Secchi, 2019). The physiological adaptations of gestation inherently exacerbate glucose intolerance, thereby imposing dynamic challenges to glycemic control. GDM, characterized by its typical onset in the second or third trimester, fundamentally stems from an inadequate pancreatic beta-cell compensatory response to the amplified insulin resistance of pregnancy (Lende & Rijhsinghani, 2020). Identified risk factors for GDM include maternal obesity, advanced maternal age, specific ethnic backgrounds, and a family history of Type 2 diabetes. Unmanaged GDM escalates maternal risks such as preeclampsia and increased rates of C-sections, alongside a 7-10-fold heightened likelihood of postpartum Type 2 diabetes (Adam et al., 2023; McCance, 2015). Neonatal complications include macrosomia, hypoglycemia, and respiratory distress. Critically, both mother and offspring face enduring risks of Type 2 diabetes and metabolic syndrome, underscoring an intergenerational transmission of metabolic dysfunction (Adam et al., 2023; McCance, 2015).

Pregestational diabetes carries more severe outcomes, including higher risks of miscarriage, major congenital anomalies, and perinatal mortality, directly correlated with suboptimal first-trimester glycemic control (Alexopoulos et al., 2019; Mackin et al., 2018; Malaza et al., 2022). Missed preconception care opportunities are common (Scavini & Secchi, 2019; Alexopoulos et al., 2019). Persistent disparities in screening and management lead to suboptimal outcomes (Lende & Rijhsinghani, 2020). Fragmented healthcare models are insufficient for this complex interplay of factors. Integrated care, offering coordinated, person-centered approaches across primary, specialized, social, and community support systems (Goodwin et al., 2012), is urgently needed to provide clinically effective, accessible, efficient, equitable, and patient-centered care (Bashir et al., 2024; Glasgow, 2003). Systematic approaches, including quality improvement, are essential for better diabetes care (O'Connor et al., 2011). This review explores integrated care principles, components, models, and outcomes, examining challenges and proposing future directions for this vulnerable population.

Types of Diabetes in pregnancy:

Pathophysiology, risks, and management:-

Diabetes during pregnancy, including gestational diabetes mellitus (GDM) and pre-existing Type 1 and Type 2 diabetes, each requires specific management approaches.

Gestational Diabetes Mellitus (GDM):-

GDM is a form of glucose intolerance first identified during pregnancy (Alexopoulos et al., 2019), affecting about 16.9% of live births (McCance, 2015). It typically resolves after childbirth but significantly raises a woman's lifetime risk of developing Type 2 diabetes (McCance, 2011).

Pathophysiology and diagnosis: GDM usually develops in the second or third trimester (around 24-28 weeks). due to pregnancy hormones inducing insulin resistance and inadequate beta-cell compensation (Lende & Rijhsinghani, 2020). Screening often involves a 75-g oral glucose tolerance test (OGTT) or a two-step approach (American Diabetes Association Professional Practice Committee, 2022).

Maternal and Neonatal risks: Maternal risks include preeclampsia and increased cesarean delivery, with a 7-10-fold higher likelihood of developing Type 2 diabetes postpartum. Neonatal complications include macrosomia, hypoglycemia, and respiratory distress syndrome. Long-term, these children have an increased risk of childhood obesity and Type 2 diabetes (Adam et al., 2023; McCance, 2015).

Management: Initial management for GDM involves medical nutrition therapy (MNT) and regular physical activity. If these measures don't achieve glycemic targets (e.g., fasting blood sugar below 95 mg/dL, 1-hour post-meal below 140 mg/dL, 2-hour post-meal below 120 mg/dL), insulin is the preferred medication. Metformin might also be considered in certain situations (American Diabetes Association Professional Practice Committee, 2022).

Pregestational Diabetes: Type 1 and Type 2:-

Type 1 Diabetes in pregnancy (T1DM): Women with T1DM face complex glycemic control challenges due to fluctuating insulin sensitivity and increased insulin requirements (McCance, 2015), heightened risk for severe hypoglycemia and DKA (McCance, 2011). Risks: High rates of major congenital malformations, spontaneous abortion, stillbirth, preterm delivery, and macrosomia (Mackin et al., 2018; McCance, 2015). Maternal risks include accelerated retinopathy/nephropathy, severe hypoglycemia, DKA, and increased preeclampsia risk (McCance, 2011). Management: Intensive insulin therapy (MDI or insulin pump), frequent SMBG or CGM. Rigorous preconception counseling (HbA1c < 6.5%) is paramount for reducing congenital anomaly risk (American Diabetes Association Professional Practice Committee, 2022).

Type 2 Diabetes in pregnancy (T2DM): Increasing T2DM prevalence (Alexopoulos et al., 2019; McCance, 2015) leads to outcomes often as poor as or worse than T1DM, with increased risks for preeclampsia, cesarean delivery, and macrosomia (Malaza et al., 2022). Risks: Similar to GDM/T1DM, including congenital anomalies (lower than T1DM), macrosomia, preterm birth, and neonatal hypoglycemia. Maternal risks include accelerated progression of diabetic complications and heightened gestational hypertension/preeclampsia risk (McCance, 2015). Management: Often necessitates a shift from oral agents to insulin. Preconception planning is vital: glycemic optimization, cessation of teratogenic medications, and screening for existing complications (American Diabetes Association Professional Practice Committee, 2022).

Table 1: Comparison of Gestational Diabetes Mellitus (GDM) and Pregestational Diabetes (Type 1 & Type 2) in Pregnancy

Feature	Gestational Diabetes Mellitus (GDM)	Type 1 Diabetes in Pregnancy	Type 2 Diabetes in Pregnancy
Definition	Glucose intolerance with onset or first recognition during pregnancy [4, 28, 43].	Autoimmune beta-cell destruction, absolute insulin deficiency, diagnosed pre-pregnancy [4, 43-44].	Progressive loss of insulin secretion on background of insulin resistance, diagnosed pre-pregnancy [4, 43-44].
Onset	Typically 2nd or 3rd trimester [28].	Usually childhood/adolescence, but any age.	Usually adulthood, often associated with obesity/sedentary lifestyle [34].
Pathophysiology	Pregnancy-induced insulin resistance, inadequate beta-cell compensation [28].	Autoimmune beta-cell destruction.	Insulin resistance and progressive beta-cell dysfunction.
Maternal Risks (Short-term)	Preeclampsia, cesarean delivery, future Type 2 diabetes [1, 4, 29, 31, 34].	Severe hypoglycemia, DKA, accelerated retinopathy/nephropathy, preeclampsia [4].	Preeclampsia, cesarean delivery, accelerated complications [31, 34].
Neonatal Risks (Short-term)	Macrosomia, neonatal hypoglycemia, respiratory distress syndrome [1, 28, 31, 34].	Congenital anomalies (high risk), macrosomia, preterm birth, perinatal mortality [4, 29, 31].	Congenital anomalies (moderate risk), macrosomia, preterm birth, neonatal hypoglycemia [31, 34].
Long-term Implications (Mother)	High risk (7-10x) of developing Type 2 diabetes [1, 34].	Worsening microvascular complications, glycemic control challenges.	Worsening microvascular/macrovacular complications.
Long-term Implications (Offspring)	Increased risk of childhood obesity, Type 2 diabetes [1,	Increased risk of obesity, metabolic syndrome [44].	Increased risk of obesity, metabolic syndrome [44].

	34].		
Primary Management	MNT, exercise; insulin if targets not met [6, 28, 43].	Intensive insulin therapy (MDI/pump), frequent monitoring [4, 6].	Insulin often required; lifestyle changes; metformin sometimes continued [4, 6, 43].
Preconception Care Importance	Not applicable.	Crucial for optimizing glycemic control and reducing congenital anomaly risk [4, 6, 44].	Crucial for optimizing glycemic control, medication review, and complication screening [4, 6, 43].

All diabetes types in pregnancy require a multidisciplinary approach (obstetricians, endocrinologists, diabetes educators, dietitians) (Alexopoulos et al., 2019; McCance, 2015). Preconception planning for pregestational diabetes is crucial for glycemic optimization ($HbA1c < 6.5\%$) and complication management (American Diabetes Association Professional Practice Committee, 2022; McCance, 2015). Suboptimal uptake of preconception care remains a significant gap (Scavini & Secchi, 2019).

Components of integrated care in Diabetes management during pregnancy:-

Integrated care for diabetes in pregnancy optimizes maternal and neonatal outcomes through continuity, coordination, and patient-centeredness, with several key components.

Medical management: Achieving glycemic control:-

Medical management is foundational for preventing complications, focusing on precise blood glucose control, often with intensive insulin therapy due to its efficacy and fetal safety (American Diabetes Association Professional Practice Committee, 2022). Insulin doses are dynamically adjusted for changing physiological needs. Management also includes monitoring and managing comorbidities like gestational hypertension, preeclampsia, retinopathy, and nephropathy. Regular lab assessments ($HbA1c$, renal/thyroid function) are crucial (American Diabetes Association Professional Practice Committee, 2022).

Nutritional therapy: The foundation of glycemic control:-

Nutritional therapy (MNT) is a first-line component, involving individualized meal planning guided by a registered dietitian nutritionist (RD/RDN) (American Diabetes Association Professional Practice Committee, 2022). Goals include adequate caloric/nutrient intake, healthy fetal growth, strict glycemic targets, and appropriate gestational weight gain. Recommendations emphasize complex carbohydrates, lean proteins, and healthy fats, limiting simple sugars. Carbohydrate intake is strategically distributed to minimize postprandial glucose excursions and prevent ketosis (American Diabetes Association Professional Practice Committee, 2022). RDs provide culturally sensitive education for adherence.

Blood glucose monitoring: The compass for management:-

Regular and accurate blood glucose monitoring is indispensable for optimal glycemic control. Both fasting and postprandial self-monitoring (SMBG) are routinely recommended (American Diabetes Association Professional Practice Committee, 2022). Continuous glucose monitoring (CGM) systems are increasingly valuable, especially for Type 1 diabetes, providing real-time data and identifying glycemic excursions, reducing macrosomia and neonatal hypoglycemia (American Diabetes Association Professional Practice Committee, 2022).

Fetal monitoring: Safeguarding fetal well-being:-

Fetal monitoring is critical for identifying and managing complications. Intensive surveillance is warranted due to increased risks of stillbirth, IUGR, and macrosomia. This includes regular ultrasounds (from 28 weeks) for fetal growth and amniotic fluid volume. Formal antepartum fetal monitoring (NSTs, BPPs) often begins around 32 weeks, guiding optimal timing and mode of delivery (American Diabetes Association Professional Practice Committee, 2022).

Mental health support: Addressing the psychosocial burden:-

Dedicated mental health support is crucial for managing the significant psychosocial burden of diabetes in pregnancy. The demanding nature of daily diabetes management exacerbates anxiety, stress, and diabetes distress (Guo et al., 2021). Integrated care models effectively incorporating psychological screening and targeted support, especially those with dedicated mental health professionals or online-offline strategies, reduce anxiety/depression, improve self-management, and enhance quality of life (Guo et al., 2021). Psychosocial assessment is essential throughout pregnancy and postpartum (American Diabetes Association Professional Practice Committee, 2022).

This holistic support, via counseling, peer groups, or coping strategies, empowers patients, significantly improves adherence to complex treatment plans, and fosters emotional well-being, leading to better glycemic outcomes and a more positive pregnancy experience (De Hert et al., 2011, De Hert et al., 2011).

Integrated care for diabetes in pregnancy is a multifaceted approach combining medical/nutritional management, blood glucose/fetal monitoring, and mental health support. This collaborative strategy is crucial for mitigating risks and optimizing outcomes for mother and child.

Models of integrated care: Multidisciplinary clinics, digital telehealth solutions, and national case studies:-

Integrated care models enhance outcomes for mothers and infants by streamlining services and improving coordination.

Multidisciplinary Clinics: The Hub of Collaborative Care:-

Multidisciplinary clinics serve as a hub for integrated care, bringing together obstetricians, endocrinologists, diabetes educators, dietitians, and mental health professionals to create individualized care plans for pregnant individuals with diabetes. This integrated approach has also been successful in pediatric diabetes clinics, where regular assessments of emotional well-being resulted in improved quality of life and reduced emotional stress. Such structured, holistic collaboration clearly enhances patient outcomes, satisfaction, and adherence to treatment (Brodar et al., 2022, Theofilou et al, 2023, Lannon et al., 2024). The principles are highly relevant to pregnant individuals (Versloot et al., 2023). These clinics also enhance inter-provider communication and consistent patient messaging through regular team meetings (Rushforth et al., 2016).

Digital and telehealth-based care: Extending reach and continuity:-

Digital health and telehealth solutions revolutionize integrated care, particularly for chronic disease management in underserved populations. Telehealth provides remote clinical healthcare, bridging distances and improving access. Miller (2019) found telemonitoring in home healthcare reduced emergency visits and readmissions by enabling early intervention. Nasir et al. (2018) identified telehealth as a facilitator of timely follow-up, patient education, and efficient care coordination. For diabetes in pregnancy, digital platforms enable remote glucose monitoring, virtual consultations, and continuous support, benefiting women facing logistical challenges, enhancing self-management and adherence (Miller, 2019; Nasir et al., 2018).

Case studies and national programs: Demonstrating scalability and impact:-

Integrated care models have shown high scalability and effectiveness across various healthcare settings. For instance, the Canadian Mental Health–Integrated Diabetes Clinic expanded successfully at the national level, significantly improving patient engagement and enhancing the quality of life among young individuals with diabetes. This illustrates how such comprehensive, integrated approaches can be effectively implemented on a larger scale to achieve meaningful health improvements (Bentz et al., 2023, de Wit et al., 2022). In the U.S., the VA's integrated care model (Serper et al., 2023), although studied for cirrhosis, highlights principles like interdisciplinary collaboration and telehealth, directly applicable to diabetes in pregnancy. Telehealth, particularly during the COVID-19 pandemic, demonstrated its capacity to make integrated care more accessible and resilient. A systematic review also identified integrated maternal care strategies in low- and middle-income countries (Van der Werf et al., 2022).

Successful implementation relies on robust technological infrastructure, continuous provider training, consistent policy support, and sustainable funding (Serper et al., 2023). Integrated care is considered the optimal model, necessitating robust specialist services (Greenwood et al., 2005). The strong evidence from studies (Berg et al., 2025; Haque et al., 2024; Miller, 2019; Nasir et al., 2018; Serper et al., 2023) confirms that these patient-centered approaches lead to superior clinical outcomes, higher patient satisfaction, cost-effectiveness, and sustainable healthcare for pregnant individuals with diabetes.

Outcomes of integrated care: Demonstrating value across multiple dimensions:-

Integrated care models for diabetes in pregnancy aim to improve maternal/neonatal health, demonstrate cost-effectiveness, and enhance patient satisfaction through coordinated care.

Maternal and neonatal outcomes: Improved health and reduced complications:-

Integrated care approaches show substantial benefits for mothers and newborns. A systematic review found integrated care associated with a 0.5 percentage point HbA1c reduction in diabetes patients (Dorling et al., 2015),

clinically significant for reducing complications. Integrated care programs also showed a 19% reduction in hospital admission rates across various chronic conditions (Dorling et al., 2015). For neonatal outcomes, studies in integrated health systems consistently show decreased NICU admission rates and patient-days, without increased readmissions or mortality (Braun et al., 2020). This suggests efficient resource utilization while improving indicators like macrosomia and neonatal hypoglycemia. In LMICs, integrated approaches improved care follow-up, coordination, and interprofessional collaboration, crucial for reducing maternal/neonatal morbidity/mortality (van der Werf et al., 2022).

Cost-Effectiveness: Economic benefits for sustainable healthcare:-

Integrated care models can be highly cost-effective, particularly over longer follow-up periods (Rocks et al., 2020). Initial investments can lead to substantial long-term savings through reduced complications, fewer hospitalizations, and decreased emergency department visits. For maternal/newborn health in LMICs, strategies improving service utilization have been cost-effective, involving community-based interventions and quality improvement (Mangham-Jefferies et al., 2014). An initiative in Ukraine led to cost savings per birth (Mangham-Jefferies et al., 2014). While methodological variations exist (Mangham-Jefferies et al., 2014; Martin et al., 2023), evidence points to integrated care as a financially prudent strategy for managing diabetes in pregnancy, optimizing resource utilization.

Patient satisfaction: Fostering engagement and empowerment:-

Patient satisfaction is a vital outcome reflecting quality and patient-centered healthcare. Integrated care models prioritize patient involvement, clear communication, shared decision-making, and individualized support, all recognized drivers of enhanced patient satisfaction and improved experience. This collaborative environment fosters trust and adherence, aligning with patient-centered care principles in diabetes that focus on individuality, engagement, and empowerment (Chen et al., 2024). A comprehensive program for Type 2 diabetes patients in Mexico demonstrated significant reductions in anxiety, depression, and distress, leading to improved quality of life and high patient satisfaction (García-Ulloa et al., 2024). The inherent principles of integrated care—multidisciplinary collaboration, personalized plans, access to information, and psychosocial support—contribute synergistically to a positive patient experience (Dorling et al., 2015), enhancing adherence and long-term health outcomes. An integrative review of birth centers also found high patient satisfaction with comprehensive, personalized care (Alliman & Phillippi, 2016). Overall, integrated care for diabetes in pregnancy yields multifaceted positive outcomes, including improved maternal/neonatal health, cost-effectiveness, and substantially enhanced patient satisfaction, strongly supporting its continued implementation.




Figure I. Outcomes before and after implementation of integrated care: Comparison of clinical and patient-centered outcomes before and after integrated care implementation. Metrics include HbA1c reduction, hospital

admissions, neonatal intensive care unit (NICU) admissions, and patient satisfaction. Integrated care is associated with reduced hospital/NICU admissions, improved glycemic control, and higher patient satisfaction. References: Dorling et al., 2015; Braun et al., 2020; García-Ulloa et al., 2024

Challenges and barriers in integrated care for Diabetes management in pregnancy: navigating complexities:-

Despite its promise, integrated care for diabetes in pregnancy faces persistent challenges: access to care, health literacy, and coordination between providers. These impact patient engagement, clinical outcomes, and system efficiency.

Access to care: Bridging the divide:-

Access to consistent, quality healthcare is a major barrier, especially for pregnant individuals in underserved communities. Logistical and financial obstacles (lack of insurance, transportation, unaffordability of services/supplies) are prominent (Chin et al., 2001). Delays in prenatal care initiation are detrimental for diabetes, where early intervention is crucial. Addressing access requires expanded community-based services, digital health platforms, and policy changes to improve insurance and reduce costs (Chin et al., 2001).

Health literacy: Empowering informed self-management:-

Health literacy is crucial for effective diabetes self-management, enabling informed health decisions. Inadequate health literacy correlates with suboptimal glycemic control, increased complications, and poor self-care understanding (Schillinger et al., 2002; Al Sayah et al., 2012). This challenge is particularly acute during pregnancy, where complex self-care routines are essential. Despite general education, many struggle due to information overload or language/cultural barriers, affecting vulnerable populations disproportionately (Ahola & Groop, 2013). Integrated care must provide culturally tailored, interactive, and consistent education, moving beyond passive materials to ensure comprehension and practical application. Empowering individuals through improved health literacy is a critical solution to self-management barriers (Mahmoodi & Khanjani, 2020), leading to better outcomes.

Coordination between providers: The interdisciplinary imperative:-

Effective coordination among diverse healthcare professionals is a cornerstone of integrated care for diabetes in pregnancy, yet it's often hindered by unclear roles and fragmented communication. Primary care clinicians, crucial first contacts, experience significant burden from time constraints, patient loads, and ambiguous professional boundaries (Rushforth et al., 2016). They find diabetes uniquely challenging due to its complexity and perceived lack of systemic support (Larme & Pugh, 1998). Poor communication leads to inconsistent patient messaging, eroding trust and undermining adherence (Nam et al., 2011). Inefficient resource use and delayed interventions result, exacerbated by non-interoperable electronic health records that limit real-time data access. To overcome this, integrated care teams require clearly defined roles, robust real-time communication platforms, regular interdisciplinary meetings, and dedicated care coordinators (Rushforth et al., 2016; Nam et al., 2011). Communication is a critical influencing factor for integrated care outcomes (Baxter et al., 2018). Without strong coordination, integrated care remains an unfulfilled aspiration.

Table 2: Common Challenges in Implementing Integrated Care and Proposed Solutions

Challenge	Impact on Care	Proposed Solutions
Access to Care	Missed appointments, delayed diagnosis/treatment, suboptimal outcomes for vulnerable populations, increased emergency visits.	Expand community-based services (mobile clinics, satellite offices), strategic telehealth/digital platforms, improve insurance coverage, address transportation barriers (e.g., ride-share programs, public transit support), culturally competent outreach [1, 15].
Health Literacy	Poor self-management, medication non-adherence, increased complications, patient disengagement, limited understanding of risks.	Culturally tailored education programs, use of plain language and visual aids, "teach-back" method for comprehension, interactive digital tools, dedicated diabetes educators for personalized coaching, involving family/support networks [1-3, 22, 30].
Coordination Between Providers	Fragmented care, inconsistent patient messaging, duplicated efforts, delayed critical interventions, provider burnout, incomplete patient records.	Clear role delineation and protocols, fully interoperable Electronic Health Records (EHRs), secure and real-time communication platforms (e.g., shared messaging), regular interdisciplinary team meetings/case conferences, dedicated care coordinators/navigators, joint training

		initiatives [1, 10, 22,27,37, 42].
Funding & Reimbursement	Limited resources for integrated models, lack of incentives, sustainability issues, undervaluation of non-physician services.	Implement value-based care models (linking payment to outcomes), bundled payments for episodes of care, adequate reimbursement for non-physician services (e.g., dietitians, diabetes educators, social workers), grant funding for pilot programs and research on cost-effectiveness, policy changes to support team-based care [7,39].
Data Integration & Sharing	Incomplete patient information, difficulty tracking longitudinal outcomes, hindered research and quality improvement initiatives, compromised patient safety.	Standardized data collection protocols, robust and interoperable health information exchange systems, secure data sharing agreements with clear privacy guidelines, investment in robust IT infrastructure and cybersecurity, common data models across institutions [10].
Provider Training & Education	Lack of skills in interdisciplinary collaboration, limited understanding of specific diabetes-in-pregnancy challenges, burnout.	Mandatory interprofessional education, specialized training in diabetes in pregnancy for all team members, continuous professional development on integrated care principles, fostering a culture of mutual respect and shared learning [7, 10, 39].

Addressing access, health literacy, and care coordination is essential for maximizing integrated care's effectiveness. Success hinges on equitable service, tailored education, and seamless provider communication, requiring policy reform, funding, and a cultural shift towards patient-centered care.

Future directions and recommendations: Forging a resilient and patient-centered healthcare ecosystem:-

Addressing the escalating prevalence and complexity of diabetes in pregnancy requires a proactive, forward-thinking approach. Integrated care models are crucial for improving maternal and neonatal outcomes, reducing healthcare burdens, and enhancing patient experience. Firstly, future care must center on truly personalized, comprehensive multidisciplinary models. Beyond obstetricians and endocrinologists, this expanded team must routinely include mental health professionals and social workers. As the American Diabetes Association (2009) highlighted, effective integrated care addresses not only glycemic control and medical comorbidities but also profound psychosocial stressors. A holistic approach integrating mental health screening, counseling, and social services for determinants of health (e.g., food insecurity) is crucial for well-being and adherence.

Pharmacogenomics can further refine personalized care, though nascent in pregnancy (Schaefer-Graf et al., 2018). Secondly, technology offers transformative enhancements. With diabetes prevalence projected to increase significantly (Rowley et al., 2017), scalable digital tools are essential. Advanced remote glucose monitoring, intuitive mobile apps, and telehealth platforms can manage conditions efficiently, especially in underserved areas. These technologies must prioritize user-friendliness, cultural sensitivity, data security, and seamless EHR integration. Leveraging AI and machine learning could revolutionize risk prediction and generate personalized treatment algorithms, shifting towards proactive management (Schaefer-Graf et al., 2018; Coman et al., 2024). Thirdly, robust preventive strategies and public health interventions must be aggressively prioritized. Aggressive population-level prevention, including universal preconception and early pregnancy diabetes screening, lifestyle modification programs, and community-based education, can substantially reduce disease incidence (Rowley et al., 2017). This includes promoting healthy behaviors pre-conception and addressing social determinants of health. Prevention of GDM development through lifestyle has had varying success, emphasizing earlier intervention (Schaefer-Graf et al., 2018; Malaza et al., 2022).

Lastly, consistent policy support and pragmatic reimbursement frameworks are essential for sustaining and expanding integrated care. Current fee-for-service models inadvertently incentivize fragmented care. Funding mechanisms must incentivize team-based care, adequately reimburse non-physician professionals, and cover digital health services. Policy reforms are needed for seamless data sharing and interoperability. Advocating for value-based care, where providers are reimbursed for outcomes, can align incentives. A "single level health care" approach is advocated for equitable access (Bagchee, 2005). These four pillars—integrated teams, technology, prevention, and policy support—will forge a resilient, responsive, and equitable healthcare ecosystem, optimizing outcomes for mothers and children.

Conclusion:-

Diabetes during pregnancy, encompassing pre-existing Type 1 and Type 2 diabetes and gestational diabetes mellitus (GDM), is a significant and escalating global public health challenge. Its rising prevalence directly links to adverse maternal/neonatal outcomes and profound long-term health implications for both mother and offspring. The complex management of glycemic control and potential complications underscores the urgent need for comprehensive, coordinated, and patient-centered care models that transcend traditional fragmented approaches.

This review explored the critical role of integrated care approaches in optimizing diabetes management throughout pregnancy. We delineated the distinct characteristics and risks of GDM, Type 1, and Type 2 diabetes in pregnancy, emphasizing meticulous glycemic control and early intervention. The core, synergistic components of effective integrated care—medical management, nutritional therapy, blood glucose monitoring, fetal surveillance, and mental health support—contribute to demonstrably improved clinical outcomes.

Furthermore, we critically analyzed various integrated care models, from multidisciplinary clinics fostering direct collaboration to innovative digital and telehealth solutions enhancing accessibility. Case studies from national programs provide compelling empirical evidence of scalability and sustained effectiveness in diverse settings, demonstrating their potential to improve clinical outcomes, enhance patient satisfaction, and achieve greater cost-effectiveness. Despite these benefits, widespread and equitable implementation faces significant barriers, including access to care (socioeconomic/geographical disparities), varying health literacy levels, and issues in inter-provider coordination. Addressing these multifaceted challenges is fundamental, requiring systemic reforms, targeted educational interventions, and robust technological infrastructure.

Looking towards the future, recommendations emphasize personalized, holistic care models integrating psychosocial support. Further leveraging cutting-edge technological innovations and prioritizing aggressive, widespread preventive strategies are crucial. Establishing robust, supportive policy and sustainable reimbursement frameworks is also vital. By fostering seamless collaboration, deploying technological advancements, championing preventive health, and implementing supportive policies, the healthcare ecosystem can be fundamentally transformed. This will enable it to effectively address the growing burden of diabetes in pregnancy, delivering high-quality, patient-centered, compassionate, and efficient care. Ultimately, a future-focused, integrated approach holds the profound potential to ensure healthier outcomes, mitigate long-term risks, and foster a brighter future for mothers and their children globally, breaking the vicious intergenerational cycle of metabolic disease and improving public health on a grand scale.

Conflict of Interest Statement:-

The authors declare that there is no conflict of interest regarding the publication of this review article.

Funding Statement:-

This review article did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Acknowledgment:-

The authors would like to acknowledge the valuable contributions of researchers and clinicians whose work has been cited in this review. We also thank PHCC for providing access to resources and literature essential to the completion of this manuscript.

References:-

1. Adam, Sumaiya, Harold David McIntyre, Kit Ying Tsoi, Anil Kapur, Ronald C. Ma, Stephanie Dias, Pius Okong et al. "Pregnancy as an opportunity to prevent type 2 diabetes mellitus: FIGO Best Practice Advice." *International Journal of Gynecology & Obstetrics* 160 (2023): 56-67.
2. Ahola, Aila J., and P-H. Groop. "Barriers to self-management of diabetes." *Diabetic medicine* 30, no. 4 (2013): 413-420.
3. Al Sayah, Fatima, Sumit R. Majumdar, Beverly Williams, Sandy Robertson, and Jeffrey A. Johnson. "Health literacy and health outcomes in diabetes: a systematic review." *Journal of general internal medicine* 28 (2013): 444-452.
4. Alexopoulos, Anastasia-Stefania, Rachel Blair, and Anne L. Peters. "Management of preexisting diabetes in pregnancy: a review." *Jama* 321, no. 18 (2019): 1811-1819.

5. Alliman, Jill, and Julia C. Phillippi. "Maternal outcomes in birth centers: an integrative review of the literature." *Journal of midwifery & women's health* 61, no. 1 (2016): 21-51.
6. American Diabetes Association Professional Practice Committee, and American Diabetes Association Professional Practice Committee. "15. Management of diabetes in pregnancy: Standards of Medical Care in Diabetes—2022." *Diabetes Care* 45, no. Supplement_1 (2022): S232-S243.
7. American Diabetes Association. "Standards of medical care in diabetes—2009." *Diabetes care* 32, no. Suppl 1 (2009): S13.
8. Bagchee, Shyamal. "Private health sector in India: single level health care is the only solution." *BMJ: British Medical Journal* 331, no. 7528 (2005): 1339.
9. Bashir, Mohammed, Yassin Fagier, Badreldeen Ahmed, and Justin C. Konje. "An overview of diabetes mellitus in pregnant women with obesity." *Best Practice & Research Clinical Obstetrics & Gynaecology* 93 (2024): 102469.
10. Baxter, S. K., Maxine Johnson, Duncan Chambers, Anthea Sutton, Elizabeth Goyder, and Andrew Booth. "Understanding new models of integrated care in developed countries: a systematic review." *Health Services and Delivery Research* 6, no. 29 (2018).
11. Bentz, Jamie Anne, Rebecca Hancock-Howard, Zal Press, Francesca Brundisini, and Sarah Berglas. "CADTH Health Technology Review Living With Type 2 Diabetes." *Can J Health Technol* 3, no. 3 (2023).
12. Braun, David, Eric Braun, Vicki Chiu, Anthony E. Burgos, Mandhir Gupta, Marianna Volodarskiy, and Darios Getahun. "Trends in neonatal intensive care unit utilization in a large integrated health care system." *JAMA network open* 3, no. 6 (2020): e205239.
13. Brodar, Kaitlyn E., Rafael O. Leite, Daniella Marchetti, Manuela Jaramillo, Eileen Davis, Janine Sanchez, Alan M. Delamater, Patrice G. Saab, and Annette M. La Greca. "Psychological screening and consultation in a pediatric diabetes clinic: Medical providers' perspectives." *Clinical Practice in Pediatric Psychology* 10, no. 2 (2022): 164-179.
14. Chen, Tsung-Tai, Wei-Chih Su, and Mei-I. Liu. "Patient-centered care in diabetes care-concepts, relationships and practice." *World Journal of Diabetes* 15, no. 7 (2024): 1417.
15. Chin, Marshall H., Sandy Cook, Lei Jin, Melinda L. Drum, James F. Harrison, Julie Koppert, Fay Thiel et al. "Barriers to providing diabetes care in community health centers." *Diabetes care* 24, no. 2 (2001): 268-274.
16. Coman, Laura-Ioana, Marilena Ianculescu, Elena-Anca Paraschiv, Adriana Alexandru, and Ioana-Anca Bădăraș. "Smart solutions for diet-related disease management: Connected care, remote health monitoring systems, and integrated insights for advanced evaluation." *Applied Sciences* 14, no. 6 (2024): 2351.
17. De Hert, Marc, Christoph U. Correll, Julio Bobes, Marcelo Cetkovich-Bakmas, D. A. N. Cohen, Itsuo Asai, Johan Detraux et al. "Physical illness in patients with severe mental disorders. I. Prevalence, impact of medications and disparities in health care." *World psychiatry* 10, no. 1 (2011): 52.
18. De Hert, Marc, D. A. N. Cohen, Julio Bobes, Marcelo Cetkovich-Bakmas, Stefan Leucht, David M. Ndeti, John W. Newcomer et al. "Physical illness in patients with severe mental disorders. II. Barriers to care, monitoring and treatment guidelines, plus recommendations at the system and individual level." *World psychiatry* 10, no. 2 (2011): 138.
19. de Wit, Maartje, Katarzyna A. Gajewska, Eveline R. Goethals, Vincent McDarby, Xiaolei Zhao, Given Hapunda, Alan M. Delamater, and Linda A. DiMeglio. "ISPAD clinical practice consensus guidelines 2022: psychological care of children, adolescents and young adults with diabetes." *Pediatric Diabetes* 23, no. 8 (2022): 1373.
20. Dorling, Grail, Tim Fountaine, SORCHA McKenna, and Brindan Suresh. "The evidence for integrated care." *McKinsey and Company* (2015).
21. García-Ulloa, Ana Cristina, José Luis Cárdenas-Fragoso, Diana Hernández-Juárez, Nancy Haydée Serrano-Pérez, Paula Blancarte-Jaber, Carlos Aguilar-Salinas, Nancy R. Mejía-Domínguez, Alejandro Zentella-Dehesa, and Sergio Hernández-Jiménez. "A case study on the implementation of quality and comprehensive indicators set in routine diabetes care." *Archives of Public Health* 82, no. 1 (2024): 121.
22. Glasgow, Russell E. "Translating research to practice: lessons learned, areas for improvement, and future directions." *Diabetes care* 26, no. 8 (2003): 2451-2457.
23. Goodwin, Nick, Judith Smith, Alisha Davies, Claire Perry, Rebecca Rosen, Anna Dixon, Jennifer Dixon, and Chris Ham. "Integrated care for patients and populations: improving outcomes by working together." *London: King's Fund* (2012).
24. Greenwood, Richard, Ken Shaw, and Peter Winocour. "Diabetes and the quality and outcomes framework: integrated care is best model for diabetes." *BMJ: British Medical Journal* 331, no. 7528 (2005): 1340.

25. Guo, Ying, Ling Zhou, Bei Sun, Chenhui Wang, and Jinling Zhang. "Application of online-offline integrated medical care management in patients with gestational diabetes." *Ginekologia Polska* 92, no. 10 (2021): 720-725.
26. Haque, Mohammad M., W. Kathy Tannous, William H. Herman, Jincy Immanuel, William M. Hague, Helena Teede, Joanne Enticott et al. "Cost-effectiveness of diagnosis and treatment of early gestational diabetes mellitus: economic evaluation of the TOBOGM study, an international multicenter randomized controlled trial." *EClinicalMedicine* 71 (2024).
27. Lannon, Carole M., Christine L. Schuler, LaCrecia Thomas, Emily Gehring, Keith J. Mann, and Laurel K. Leslie. "Integrating Emotional Health Assessments into Pediatric Care: Initial Learnings from an MOC Part 4 Activity." *Pediatric Quality & Safety* 9, no. 5 (2024): e768.
28. Larme, Anne C., and Jacqueline A. Pugh. "Attitudes of primary care providers toward diabetes: barriers to guideline implementation." *Diabetes care* 21, no. 9 (1998): 1391-1396.
29. Lende, Michelle, and Asha Rijhsinghani. "Gestational diabetes: overview with emphasis on medical management." *International journal of environmental research and public health* 17, no. 24 (2020): 10846.
30. Mackin, Sharon T., Scott M. Nelson, Joannes J. Kerssens, Rachael Wood, Sarah Wild, Helen M. Colhoun, Graham P. Leese, Sam Philip, and Robert S. Lindsay. "Diabetes and pregnancy: national trends over a 15 year period." *Diabetologia* 61, no. 5 (2018): 1081-1088.
31. Mahmoodi, Mohammad Reza, and Narges Khanjani. "Barriers and Limitations to Obstacle Diabetes Self-Management with a Focus on Nutritional Literacy: Solutions and Opportunities. Critical Review and Research Synthesis." *Critical Comments in Biomedicine* (2020).
32. Malaza, Nompumelelo, Matladi Masete, Sumaiya Adam, Stephanie Dias, Thembeke Nyawo, and Carmen Pheiffer. "A systematic review to compare adverse pregnancy outcomes in women with pregestational diabetes and gestational diabetes." *International journal of environmental research and public health* 19, no. 17 (2022): 10846.
33. Mangham-Jefferies, Lindsay, Catherine Pitt, Simon Cousens, Anne Mills, and Joanna Schellenberg. "Cost-effectiveness of strategies to improve the utilization and provision of maternal and newborn health care in low-income and lower-middle-income countries: a systematic review." *BMC pregnancy and childbirth* 14 (2014): 1-23.
34. Martin, Elizabeth, Bassel Ayoub, and Yvette D. Miller. "A systematic review of the cost-effectiveness of maternity models of care." *BMC pregnancy and childbirth* 23, no. 1 (2023): 859.
35. McCance, David R. "Diabetes in pregnancy." *Best practice & research Clinical obstetrics & gynaecology* 29, no. 5 (2015): 685-699.
36. McCance, David R. "Pregnancy and diabetes." *Best practice & research Clinical endocrinology & metabolism* 25, no. 6 (2011): 945-958.
37. Miller, Melissa. "Telehealth Integration Influencing Success and Sustainability." PhD diss., Walden University, 2019.
38. Nam, Soohyun, Catherine Chesla, Nancy A. Stotts, Lisa Kroon, and Susan L. Janson. "Barriers to diabetes management: patient and provider factors." *Diabetes research and clinical practice* 93, no. 1 (2011): 1-9.
39. Nasir, Jamal Abdul, Shahid Hussain, and Chuangyin Dang. "An integrated planning approach towards home health care, telehealth and patients group based care." *Journal of network and computer applications* 117 (2018): 30-41.
40. O'Connor, Patrick J., Noni L. Bodkin, Judith Fradkin, Russell E. Glasgow, Sheldon Greenfield, Edward Gregg, Eve A. Kerr et al. "Diabetes performance measures: current status and future directions." *Diabetes care* 34, no. 7 (2011): 1651.
41. Rocks, Stephen, Daniela Berntson, Alejandro Gil-Salmerón, Mudathira Kadu, Nieves Ehrenberg, Viktoria Stein, and Apostolos Tsiachristas. "Cost and effects of integrated care: a systematic literature review and meta-analysis." *The European Journal of Health Economics* 21 (2020): 1211-1221.
42. Rowley, William R., Clement Bezold, Yasemin Arian, Erin Byrne, and Shannon Krohe. "Diabetes 2030: insights from yesterday, today, and future trends." *Population health management* 20, no. 1 (2017): 6-12.
43. Rushforth, Bruno, Carolyn McCrorie, Liz Glidewell, Eleanor Midgley, and Robbie Foy. "Barriers to effective management of type 2 diabetes in primary care: qualitative systematic review." *British Journal of General Practice* 66, no. 643 (2016): e114-e127.
44. Scavini, Marina, and Antonio Secchi. "Diabetes in pregnancy." *Acta Diabetologica* 56 (2019): 719-721.
45. Schaefer-Graf, Ute, Angela Napoli, Christopher J. Nolan, and Diabetic Pregnancy Study Group. "Diabetes in pregnancy: a new decade of challenges ahead." *Diabetologia* 61 (2018): 1012-1021.

46. Schillinger, D., K. Grumbach, and J. D. Piette. "Association of functional health literacy with glycemic control among diabetes patients." *JAMA* 288 (2002).
47. Serper, Marina, Aneza Agha, Patrik A. Garren, Tamar H. Taddei, David E. Kaplan, Peter W. Groeneveld, Rachel M. Werner, and Judy A. Shea. "Multidisciplinary teams, efficient communication, procedure services, and telehealth improve cirrhosis care: A qualitative study." *Hepatology Communications* 7, no. 6 (2023): e0157.
48. Theofilou, Paraskevi, and Dimitris D. Vlastos. "The psychological burden of families with diabetic children: A literature review focusing on quality of life and stress." *Children* 10, no. 6 (2023): 937.
49. van den Berg, Maud, Julia Spaan, Jacoba Van Der Kooy, Monique Klerkx, Charlotte Krol, Arie Franx, Kees TB Ahaus, and Hilco J. van Elten. "Value-based evaluation of gestational diabetes mellitus care pathway redesign by using cost and outcome data." *BMC Pregnancy and Childbirth* 25, no. 1 (2025): 1-11.
50. van der Werf, Laura, Silvia Evers, Laura Prieto-Pinto, Daniel Samacá-Samacá, and Aggie Paulus. "Integrated maternal care strategies in low-and middle-income countries: a systematic review." *International Journal of Integrated Care* 22, no. 2 (2022): 26.
51. Versloot, Judith, Hana Saab, Simona C. Minotti, Amna Ali, Julia Ma, Robert J. Reid, Sheryl Parks, and Ian Zenlea. "An integrated care model to support adolescents with diabetes-related quality-of-life concerns: an intervention study." *Canadian Journal of Diabetes* 47, no. 1 (2023): 3-10.